

WJG 20th Anniversary Special Issues (3): Inflammatory bowel disease**Anti-TNF alpha in the treatment of ulcerative colitis: A valid approach for organ-sparing or an expensive option to delay surgery?**

Gianluca Rizzo, Daniela Pugliese, Alessandro Armuzzi, Claudio Coco

Gianluca Rizzo, Claudio Coco, Department of Surgical Sciences, Complesso Integrato Columbus, Catholic University of the Sacred Heart, 00168 Rome, Italy

Daniela Pugliese, Alessandro Armuzzi, Internal Medicine and Gastroenterology Unit, Complesso Integrato Columbus, Catholic University of the Sacred Heart, 00168 Rome, Italy

Author contributions: Rizzo G, Pugliese D, Armuzzi A and Coco C equally participated in the conception, design, analysis, interpretation of data and drafting of this article; all the authors revised the article critically for important intellectual content and gave final approval of the version to be published.

Correspondence to: Gianluca Rizzo, MD, Department of Surgical Sciences, Complesso Integrato Columbus, Catholic University of the Sacred Heart, Via F. Moscati 31-33, 00168 Rome, Italy. gianluca.rizzo1979@libero.it

Telephone: +39-6-3503618-622 Fax: +39-6-3503344

Received: November 1, 2013 Revised: February 6, 2014

Accepted: March 12, 2014

Published online: May 7, 2014

Abstract

Ulcerative colitis (UC) is an inflammatory bowel disease affecting large bowel with variable clinical course. The history of disease has been modified by the introduction of biologic therapy, in particular Infliximab (IFX), that has demonstrated efficacy in inducing fast symptoms remission, promoting mucosal healing and maintaining long-term remission. However, surgery is still needed for UC patients: in case of failure of medical therapy and if acute complications or a malignancy occurred. Surgical treatment is associated with a short-term post-operative mortality and morbidity respectively of 0%-4% and 30%. In this study we systematically analyzed: the role of IFX in reducing the colectomy rate, the risk of post-operative morbidity in pre-operatively IFX-treated patients and the cost-effectiveness of IFX therapy. Four of 5 analyzed randomized controlled trials demonstrated

that therapy with IFX significantly reduces the colectomy rate. Moreover, pre-operative treatment with IFX doesn't seem to increase post-operative infectious complications. By an economic point of view, the cost-effectiveness of IFX-therapy was demonstrated for UC patients suffering from moderate to severe UC in a study based on a cost estimation of the National Health Service of England and Wales. However, the argument is debated.

© 2014 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: Ulcerative colitis; Infliximab; Colectomy; Post-operative complications; Cost-effectiveness; Inflammatory bowel disease

Core tip: The introduction of biologic therapy with Infliximab (IFX) has significantly modified the clinical course of ulcerative colitis. In this study we systematically analyzed the role of IFX in reducing the colectomy rate, the risk of post-operative morbidity in pre-operatively IFX-treated patients and the cost-effectiveness of IFX therapy.

Rizzo G, Pugliese D, Armuzzi A, Coco C. Anti-TNF alpha in the treatment of ulcerative colitis: A valid approach for organ-sparing or an expensive option to delay surgery? *World J Gastroenterol* 2014; 20(17): 4839-4845 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v20/i17/4839.htm> DOI: <http://dx.doi.org/10.3748/wjg.v20.i17.4839>

INTRODUCTION

Ulcerative colitis (UC) is a chronic inflammatory disease, of an unknown etiology, affecting the large bowel. It is characterized by a contiguous mucosal inflammation

starting in the rectum and proximally progressing in continuity in the colon for a different distance. According to the Montreal Classification, which describes the maximal macroscopic extent of the disease at colonoscopy, the distribution of UC is commonly classified as: proctitis, left-sided and extensive colitis^[1-4]. Disease activity is grouped into remission, mild, moderate, or severe but the clinical course of UC is variable and can range from a long-standing remitting to a refractory or fulminant disease^[5-11]. Solberg *et al.*^[5], in a population-based cohort study of 843 patients with inflammatory bowel disease, enrolled in South-Eastern Norway and systematically followed-up at 1, 5 and 10 years after diagnosis, identified 4 different clinical patterns: (1) initial high activity to remission or mild severity (55%); (2) chronic intermittent symptoms (37%); (3) continuous symptoms (6%); and (4) initial low activity to increased severity (1%)^[12]. The main symptoms of UC are bloody diarrhea and abdominal pain, associated with urgency and tenesmus. UC is conventionally treated with a step-up approach, based on the severity and extent of the disease, including various agents such as 5-aminosalicylates, corticosteroids and immunosuppressants (including thiopurines and cyclosporine). The primary aims of medical therapy in UC should be inducing and maintaining long-term remission, achieving mucosal healing, minimizing steroid-dependence, avoiding serious complications (hospitalization and surgery) and improving patients' quality of life^[13-17]. However, standard therapy is not always able to achieve these goals and patients become steroid-dependent or experience frequent or severe relapse, with consequent increased risk of hospitalization and surgery. The history of disease has been partially modified by the introduction of biologic therapies. Infliximab (IFX) has demonstrated efficacy in inducing fast symptoms-remission, promoting mucosal healing and maintaining long-term remission^[17-20]. It is currently approved for patients with moderate to severe UC who have incomplete response, are intolerant or have any medical contraindications to corticosteroids and/or immunomodulators^[21-24]. It is also recommended as rescue therapy in severe steroid-refractory disease^[25-27] and in steroid-dependent patients^[28,29].

Surgery is still needed for UC patients, in case of failure of medical therapy, occurrence of acute complications (such as fulminant colitis, toxic megacolon and bowel perforation) or development of malignancy. Since its first description (1978), restorative proctocolectomy with ileal pouch anal anastomosis (IPAA) represents the gold standard of surgical treatment of UC: all the colon and rectum are removed and a J pouch is created with terminal ileum and anatomized to the anal canal. This restorative operation, avoiding a permanent stoma, maintains intestinal continuity and preserves the patient's body image^[30-33]. Moreover, the introduction of minimally invasive approach significantly contributed in improving the acceptance and tolerability of this procedure^[34-37], reducing the rate of post-operative adhesions and post-operative hospital stay and improving cosmetic results^[38-41].

However, even if in skilled hands, proctocolectomy with IPAA is not without risk and is associated with an estimated short-term mortality ranged between 0% and 4% and a morbidity rate of about 30%, with an incidence of pelvic sepsis ranging from 5% to 24% and a re-surgery risk of about 16%^[42,43].

If surgery represents a definitive solution for cessation of symptoms, withdrawing medical therapies and reducing the cancer-risk, it is not free of long-term post-operative morbidity (as pouchitis, fecal incontinence, female fecundity or fertility) with a relevant impact on patients' quality of life^[44-46]. Population-based studies have reported a 10-years cumulative risk of colectomy ranging between 9%-30%, with some differences among countries. Approximately, 4% to 9% of UC patients will require colectomy within the first year of diagnosis and, subsequently, the risk of colectomy increases of 1% per year^[47-50].

It is still under debate whether, in the long-term, the biological therapies could be a valid approach for organ-sparing, rather than an expensive option to delay surgery. Aim of this review was to evaluate the real impact of biological therapy on the rate of colectomy in UC patients.

A review of the literature searching for the terms "anti-tumor necrosis factor- α (TNF- α)", "infliximab" matched with the terms "ulcerative colitis" and "surgery" was performed, using PubMed, MEDLINE, EMBASE and Cochrane databases. All relevant articles (both experimental and observational studies) in English between January 2000 and July 2013 were reviewed.

Therapy with infliximab and rate of colectomy

We identified five randomized controlled trials^[51-54], one meta-analysis^[55] and six observational studies^[56-61] following literature search. Characteristics of the studies are summarized in Table 1.

RANDOMIZED CONTROLLED TRIALS AND META-ANALYSIS

Two randomized, double-blind, placebo-controlled trials, ACT 1 and ACT 2, demonstrated the efficacy of IFX for induction (week 8) and maintenance (week 30 and week 54 for ACT1) of clinical response and remission in patients with moderate-to-severe UC, despite the use of conventional therapy^[51]. Further analysis, from ACT 1-2 open-label extension phase, focused on colectomy and hospitalization rates during follow-up to 54 wk. Compared with placebo, the cumulative incidence of colectomy through 54 wk for IFX was significantly lower (10% *vs* 17%, $P = 0.02$), with an absolute risk reduction of 7% (95%CI: 0.01-0.12, HR = 0.59). Moreover, in IFX-treated patients were recorded fewer (compared to placebo group) UC-related hospitalizations and surgical procedures per 100 patient-years of treatment (40 *vs* 20, $P = 0.003$; 34 *vs* 21, $P = 0.03$ respectively)^[52].

Previous controlled smaller studies have addressed the risk of colectomy in patients with severe UC treated

Table 1 Rate of colectomy after therapy with Infliximab

Ref.	RCT	Pts (n)	Type of disease	FU	Rate of colectomy in IFX Pts	Rate of colectomy in control Pts	P value
Rutgeerts <i>et al</i> ^[51]	Y	364 (IFX)	Moderate to severe UC	54 wk (ACT1)	9.5%	14.7%	< 0.05
Sandborn <i>et al</i> ^[52]		364 (control)		30 wk (ACT2)			
Sands <i>et al</i> ^[53]	Y	8 (IFX)	Severe active steroid-refractory	2 wk	50%	100%	NS
		3 (control)					
Järnerot <i>et al</i> ^[25]	Y	24 (IFX)	Severe to moderate UC not responding to conventional therapy	3 mo	29.2%	66.7%	0.017
		21 (control)					
Gustavsson <i>et al</i> ^[54]	Y	24 (IFX)	Severe to moderate UC not responding to conventional therapy	36 mo	50%	76%	0.012
		21 (control)					
Aratari <i>et al</i> ^[56]	N	15 (IFX)	Severe steroid-refractory UC	26 mo	18%	-	-
Teisner <i>et al</i> ^[57]	N	52 (IFX)	Acute, severe and chronic refractory UC	22 mo	27%	-	-
Ferrante <i>et al</i> ^[58]	N	121 (IFX)	Acute severe refractory UC	33 mo	17%	-	-
Oussalah <i>et al</i> ^[59]	N	191 (IFX)	UC	18 mo	18.8%	-	-
		Multicenter					
Desmond <i>et al</i> ^[60]	N	21 (IFX)	UC	14 mo	9.5%	-	-
Garcia-Planella <i>et al</i> ^[61]	N	22 (IFX)	UC	84 mo	27.3%	-	-

RCT: Randomized controlled trial; Pts: Patients; FU: Follow-up; IFX: Infliximab; UC: Ulcerative colitis; Y: Yes; N: Not.

with IFX as rescue therapy. In 2001 Sands reported data on 11 patients with severe steroid-refractory disease, of whom 8 treated with IFX and 3 with placebo. After 2 wk, all patients treated with placebo underwent to surgery, while only 50% of patients receiving IFX needed surgery; however, the sample size was too small to detect a statistically significant benefit^[53]. Later, 45 patients with moderate to severe UC were randomized to IFX or placebo (24 *vs* 21 respectively) both after four day from the start of corticosteroid treatment. In the placebo group more patients (14/21, 66.7%) than IFX group (7/24, 29.2%) had a colectomy ($P = 0.017$; OR = 4.9; 95%CI: 1.4-17) within 3 mo after randomization^[25]. After a follow-up of 3 years, 50% of patients in the IFX group and 76% in the placebo group had a colectomy ($P = 0.012$)^[54].

Recently, Costa *et al*^[55] presented data from a meta-analysis on the benefit of IFX in reducing hospitalization and/or major surgeries in patients with inflammatory bowel disease. They analyzed 11 studies: 5 randomized controlled trials (RCTs) and 6 observational studies. In the RCTs, IFX treatment was associated with a significant 43% odds reduction of overall major surgery risk (OR = 0.57; 95%CI: 0.37-0.88) with a number-to-treat to avoid colectomy of 11 (95%CI: 6-51) for 1.2 years. However, a not significant increase was found in pooled results from observational studies (OR = 1.43; 95%CI: 0.65-3.13). The authors concluded that this discrepancy could be explained by the heterogeneity of observational studies, including patients at high risk of colectomy due to more severe disease and refractoriness to previous treatment.

OBSERVATIONAL STUDY

The first data on the long-term risk of colectomy were reported in a study of 314 UC patients from Italy. Among them, 52 (16.5%) patients had severe UC and were treated with intravenous corticosteroids for a median of 7 days. Of 15 patients who did not respond,

11 received IFX with short-term clinical benefit and 4 underwent urgent colectomy. In the long-term follow-up, another 6 patients underwent elective colectomy for a disease relapse, with a total colectomy rate, following the acute flare-up, of 19%. The long-term colectomy risk was not different between patients treated with IFX and steroid-responsive patients (18% *vs* 11%, respectively), as IFX was able to avoid urgent colectomy, but not to reduce the risk of elective surgery^[56]. The risk of long-term colectomy in severe UC was also evaluated in a smaller Danish study of 52 UC patients. Nineteen (37%) patients had severe UC and 7 of them (37%) underwent colectomy after a median follow-up of 22 mo (range 4-57 mo). Among the remaining patients with a chronic refractory UC, the colectomy rate was 21%. The authors concluded that IFX can avoid colectomy in two-thirds of the patients with acute, severe UC, but the beneficial effect on colectomy rate in chronic, refractory UC seems less convincing^[57]. Long-term data on colectomy in UC patients treated with IFX come from referral centers studies. In the Leuven's cohort of 121 refractory UC patients (patients with acute severe attack, refractory to intravenous steroids were excluded), 21 patients (17%) came to colectomy and 68% of initial responders achieved a sustained clinical response during a median follow-up of 33 mo (IQR 17-49.8). Lack of short-term clinical benefit, high values of baseline C-reactive protein (CRP) and previous intravenous steroid or cyclosporine treatment were identified as independent predictors of colectomy^[58]. These results are similar to those reported in a French multicenter study, in which, among 191 patients who received at least one IFX infusion, 36 patients (18.8%) underwent colectomy during a median follow-up of 18 mo (IQR 8-32). Independent predictors of colectomy were: no clinical response after IFX induction, high baseline CRP value, previous treatment with cyclosporine and IFX indication for acute severe UC^[59]. Furthermore, other experience supported history of hospital admission as an indepen-

Table 2 Rate of overall post-operative morbidity and post-operative infectious complications in patients pre-operatively treated with Infliximab

Ref.	RCT	Pts (IFX vs control) (n)	PO morbidity (IFX vs control)	P value	PO infectious complications (IFX vs control)	P value
Schluender <i>et al</i> ^[64]	N	17 vs 134	36% vs 28%	> 0.05	18% vs 8%	> 0.05
Selvasekar <i>et al</i> ^[62]	N	47 vs 254	62% vs 49%	0.10	28% vs 10%	< 0.01
Mor <i>et al</i> ^[63]	N	46 vs 46	34.8% vs 15.2%	0.004	21.7% vs 2.2%	0.011
Ferrante <i>et al</i> ^[65]	N	22 vs 119	11.1% vs 28.6%	> 0.05	9% vs 24%	0.161
Gainsbury <i>et al</i> ^[66]	N	29 vs 52	44.8% vs 44.2%	0.96	17.2% vs 26.9%	0.32
Rizzo <i>et al</i> ^[67]	N	16 vs 22	37.5% vs 22.7%	> 0.05	18.7% vs 18.2%	> 0.05

RCT: Randomized controlled trial; Pts: Patients; PO: Post-operative; IFX: Infliximab; Y: Yes.

dent predictor of the need of colectomy^[60,61].

Peri-operative infliximab and post-operative outcome

An increasing number of patients undergo to surgery after experienced biologic therapy. There is an emerging concern on the safety profile of IFX in the peri-operative setting in potentially pre-surgical patients. Many groups have reported their experiences for UC patients and there is not an agreement on the impact of these drugs on post-operative complications^[62-67]. The main characteristics of the studies are summarized in Table 2. Recently, Yang *et al*^[68] performed a high quality meta-analysis based on five studies, including 706 patients, who were treated with IFX before restorative procto-colectomy with IPAA. The authors did not find a strong association between pre-operative treatment with IFX and short-term infectious complications (OR = 2.24), but it was associated with a significantly increased risk of short-term overall post-operative complications (OR = 1.80). However, these results need to be interpreted with caution. The subgroup analysis was underpowered to assess the nature of these complications because of the small sample size and heterogeneity of the included studies (end-points, patients' characteristics and indication, type and timing of surgery).

Infliximab and surgery: Cost-effectiveness

An important issue of IFX therapy is its cost-effectiveness. IFX is often perceived to be an expensive treatment option for patients with IBD. Tsai *et al*^[69] made a cost-effectiveness analysis in UC patients based on a cost estimation of the National Health Service of England and Wales for the year 2006-07. At analysis of responders patients only, therapy with IFX was associated at an additional 0.753 quality-adjusted life year (QALYs) at an additional cost of £20662 compared to standard care without IFX; the estimated incremental cost per QALY gained for IFX against standard care was £27424. At analysis of remission patients, therapy with IFX derived an additional 0.387 QALYs at an additional cost of £7615 compared with standard care without IFX. The estimated incremental cost per QALY gained for IFX against standard care was £19696. The authors conclude that therapy with IFX appears to be a cost-effective treatment option for

adult patients suffering from moderate to severe UC. In a recent study, Park *et al*^[70] created a Markov model simulating 2 cohorts of 21-year-old patients with severe UC, following them until 100 years of age or death, comparing early colectomy with IPAA strategy to the standard medical therapy strategy (including IFX). In this study standard medical therapy accrued a discounted lifetime cost of \$236370 per patient; in contrast, early colectomy with IPAA accrued a discounted lifetime cost of \$147763 per patient. QALY-gained for standard medical therapy was 20.78, while QALY-gained for early colectomy with IPAA was 20.72; the resulting incremental cost-effectiveness ratio was approximately \$1.5 million per QALY-gained. So, the authors concluded that early colectomy with IPAA after diagnosis of severe UC reduce health care expenditures and provides comparable quality of life compared exhaustive standard medical therapy. Only an extremely low quality of life after IPAA could maintain the standard medical therapy strategy as the optimal management strategy in severe UC.

CONCLUSION

IFX has demonstrated efficacy in inducing and maintaining clinical and endoscopic remission in the long run. IFX can also avoid urgent colectomy in patients with severe acute UC refractory to intravenous steroids. The real impact of biological therapy on the natural history of UC is still controversial, whereas it is not clear if it allows avoidance of colectomy or rather than a delay. The median colectomy risk for UC patients treated with IFX is about 10%-20% in both RCTs and observational studies, with higher rate for patients with severe acute attack. Data from RCTs support the efficacy of IFX in reducing the risk of surgeries in the long-term, but none was designed to assess IFX effects on surgeries. Real life data from referral centers do not confirm this issue, but each study includes patients with different baseline characteristics and risks of colectomy. From these evidences, it seems that patients with more severe disease, high inflammation burden, refractoriness to intravenous steroids and/or cyclosporine and history of hospitalizations, have higher risk of colectomy. So, it is necessary for physicians taking in account the risks and the benefits of medical

versus surgical therapy, concerning about cost and side effects of medications, cost and morbidity of surgery and patients' quality of life. Prospective, specifically designed studies are necessary to assess the long-term risk of colectomy in UC patients treated with IFX.

REFERENCES

- Dignass A, Eliakim R, Magro F, Maaser C, Chowers Y, Geboes K, Mantzaris G, Reinisch W, Colombel JF, Vermeire S, Travis S, Lindsay JO, Van Assche G. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 1: definitions and diagnosis. *J Crohns Colitis* 2012; **6**: 965-990 [PMID: 23040452 DOI: 10.1016/j.crohns.2012.09.003]
- Stange EF, Travis SP, Vermeire S, Beglinger C, Kupcinkas L, Geboes K, Barakauskiene A, Villanacci V, Von Herbay A, Warren BF, Gasche C, Tilg H, Schreiber SW, Schölmerich J, Reinisch W. European evidence based consensus on the diagnosis and management of Crohn's disease: definitions and diagnosis. *Gut* 2006; **55** Suppl 1: i1-15 [PMID: 16481628]
- Ordás I, Eckmann L, Talamini M, Baumgart DC, Sandborn WJ. Ulcerative colitis. *Lancet* 2012; **380**: 1606-1619 [PMID: 22914296 DOI: 10.1016/S0140-6736(12)60150-0]
- Beaugerie L, Sokol H. Clinical, serological and genetic predictors of inflammatory bowel disease course. *World J Gastroenterol* 2012; **18**: 3806-3813 [PMID: 22876031 DOI: 10.3748/wjg.v18.i29.3806]
- Solberg IC, Lygren I, Jahnsen J, Aadland E, Høie O, Cvancarova M, Bernklev T, Henriksen M, Sauar J, Vatn MH, Moum B. Clinical course during the first 10 years of ulcerative colitis: results from a population-based inception cohort (IBSEN Study). *Scand J Gastroenterol* 2009; **44**: 431-440 [PMID: 19101844 DOI: 10.1080/00365520802600961]
- Magro F, Rodrigues A, Vieira AI, Portela F, Cremers I, Cotter J, Correia L, Duarte MA, Tavares ML, Lago P, Ministro P, Peixe P, Lopes S, Garcia EB. Review of the disease course among adult ulcerative colitis population-based longitudinal cohorts. *Inflamm Bowel Dis* 2012; **18**: 573-583 [PMID: 21793126 DOI: 10.1002/ibd.21815]
- Solberg IC, Lygren I, Cvancarova M, Jahnsen J, Stray N, Sauar J, Schreiber S, Moum B, Vatn MH. Predictive value of serologic markers in a population-based Norwegian cohort with inflammatory bowel disease. *Inflamm Bowel Dis* 2009; **15**: 406-414 [PMID: 19009607 DOI: 10.1002/ibd.20781]
- Satsangi J, Silverberg MS, Vermeire S, Colombel JF. The Montreal classification of inflammatory bowel disease: controversies, consensus, and implications. *Gut* 2006; **55**: 749-753 [PMID: 16698746]
- Vermeire S, Van Assche G, Rutgeerts P. Classification of inflammatory bowel disease: the old and the new. *Curr Opin Gastroenterol* 2012; **28**: 321-326 [PMID: 22647554 DOI: 10.1097/MOG.0b013e328354be1e]
- Latella G, Papi C. Crucial steps in the natural history of inflammatory bowel disease. *World J Gastroenterol* 2012; **18**: 3790-3799 [PMID: 22876029 DOI: 10.3748/wjg.v18.i29.3790]
- Blonski W, Buchner AM, Lichtenstein GR. Inflammatory bowel disease therapy: current state-of-the-art. *Curr Opin Gastroenterol* 2011; **27**: 346-357 [PMID: 21654383]
- Silverberg MS, Satsangi J, Ahmad T, Arnott ID, Bernstein CN, Brant SR, Caprilli R, Colombel JF, Gasche C, Geboes K, Jewell DP, Karban A, Loftus EV, Peña AS, Riddell RH, Sachar DB, Schreiber S, Steinhardt AH, Targan SR, Vermeire S, Warren BF. Toward an integrated clinical, molecular and serological classification of inflammatory bowel disease: report of a Working Party of the 2005 Montreal World Congress of Gastroenterology. *Can J Gastroenterol* 2005; **19** Suppl A: 5A-36A [PMID: 16151544]
- Panaccione R, Rutgeerts P, Sandborn WJ, Feagan B, Schreiber S, Ghosh S. Review article: treatment algorithms to maximize remission and minimize corticosteroid dependence in patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 2008; **28**: 674-688 [PMID: 18532990 DOI: 10.1111/j.1365-2036.2008.03753.x]
- Ford AC, Sandborn WJ, Khan KJ, Hanauer SB, Talley NJ, Moayyedi P. Efficacy of biological therapies in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol* 2011; **106**: 644-659, quiz 660 [PMID: 21407183 DOI: 10.1038/ajg.2011.73]
- Burger D, Travis S. Conventional medical management of inflammatory bowel disease. *Gastroenterology* 2011; **140**: 1827-1837.e2 [PMID: 21530749 DOI: 10.1053/j.gastro.2011.02.045]
- Meier J, Sturm A. Current treatment of ulcerative colitis. *World J Gastroenterol* 2011; **17**: 3204-3212 [PMID: 21912469 DOI: 10.3748/wjg.v17.i27.3204]
- Lawson MM, Thomas AG, Akobeng AK. Tumour necrosis factor alpha blocking agents for induction of remission in ulcerative colitis. *Cochrane Database Syst Rev* 2006; **(3)**: CD005112 [PMID: 16856078]
- Wilhelm SM, McKenney KA, Rivait KN, Kale-Pradhan PB. A review of infliximab use in ulcerative colitis. *Clin Ther* 2008; **30**: 223-230 [PMID: 18343261 DOI: 10.1016/j.clinthera.2008.02.014]
- Magro F, Portela F. Management of inflammatory bowel disease with infliximab and other anti-tumor necrosis factor alpha therapies. *BioDrugs* 2010; **24** Suppl 1: 3-14 [PMID: 21175228 DOI: 10.2165/11586290-000000000-00000]
- Van Assche G, Dignass A, Bokemeyer B, Danese S, Gionchetti P, Moser G, Beaugerie L, Gomollón F, Häuser W, Herlinger K, Oldenburg B, Panes J, Portela F, Rogler G, Stein J, Tilg H, Travis S, Lindsay JO. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 3: special situations. *J Crohns Colitis* 2013; **7**: 1-33 [PMID: 23040453 DOI: 10.1016/j.crohns.2012.09.005]
- Dignass A, Lindsay JO, Sturm A, Windsor A, Colombel JF, Allez M, D'Haens G, D'Hoore A, Mantzaris G, Novacek G, Oresland T, Reinisch W, Sans M, Stange E, Vermeire S, Travis S, Van Assche G. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 2: current management. *J Crohns Colitis* 2012; **6**: 991-1030 [PMID: 23040451 DOI: 10.1016/j.crohns.2012.09.002]
- Velayos F, Mahadevan U. Management of steroid-dependent ulcerative colitis: immunomodulatory agents, biologics, or surgery? *Clin Gastroenterol Hepatol* 2007; **5**: 668-671 [PMID: 17544992]
- Gisbert JP, Gomollón F, Hinojosa J, López San Román A. Adherence of gastroenterologists to European Crohn's and Colitis Organisation consensus on ulcerative colitis: a real-life survey in Spain. *J Crohns Colitis* 2010; **4**: 567-574 [PMID: 21122561 DOI: 10.1016/j.crohns.2010.06.001]
- Kountouras J, Zavos C, Chatzopoulos D. Anti-tumor necrosis factor therapy for ulcerative colitis. *Gastroenterology* 2005; **129**: 1138-1139 [PMID: 16143158]
- Järnerot G, Hertervig E, Friis-Liby I, Blomquist L, Karlén P, Grännö C, Vilien M, Ström M, Danielsson A, Verbaan H, Hellström PM, Magnuson A, Curman B. Infliximab as rescue therapy in severe to moderately severe ulcerative colitis: a randomized, placebo-controlled study. *Gastroenterology* 2005; **128**: 1805-1811 [PMID: 15940615]
- Yamamoto-Furusho JK, Uzcanga LF. Infliximab as a rescue therapy for hospitalized patients with severe ulcerative colitis refractory to systemic corticosteroids. *Dig Surg* 2008; **25**: 383-386 [PMID: 19005257 DOI: 10.1159/000170882]
- Kohn A, Daperno M, Armuzzi A, Cappello M, Biancone L, Orlando A, Viscido A, Annese V, Riegler G, Meucci G, Marrollo M, Sostegni R, Gasbarrini A, Peralta S, Prantera C. Infliximab in severe ulcerative colitis: short-term results of different infusion regimens and long-term follow-up. *Ali-*

- ment *Pharmacol Ther* 2007; **26**: 747-756 [PMID: 17697208]
- 28 **Armuzzi A**, Pugliese D, Danese S, Rizzo G, Felice C, Marzo M, Andrisani G, Fiorino G, Sociale O, Papa A, De Vitis I, Rapaccini GL, Guidi L. Infliximab in steroid-dependent ulcerative colitis: effectiveness and predictors of clinical and endoscopic remission. *Inflamm Bowel Dis* 2013; **19**: 1065-1072 [PMID: 23448790 DOI: 10.1097/MIB.0b013e3182802909]
- 29 **Hultén L**. Proctocolectomy and ileostomy to pouch surgery for ulcerative colitis. *World J Surg* 1998; **22**: 335-341 [PMID: 9523513]
- 30 **Melville DM**, Ritchie JK, Nicholls RJ, Hawley PR. Surgery for ulcerative colitis in the era of the pouch: the St Mark's Hospital experience. *Gut* 1994; **35**: 1076-1080 [PMID: 7926909]
- 31 **Bennis M**, Tiret E. Surgical management of ulcerative colitis. *Langenbecks Arch Surg* 2012; **397**: 11-17 [PMID: 21922296 DOI: 10.1007/s00423-011-0848-x]
- 32 **Metcalf AM**. Elective and emergent operative management of ulcerative colitis. *Surg Clin North Am* 2007; **87**: 633-641 [PMID: 17560416]
- 33 **Boller AM**, Larson DW. Laparoscopic restorative proctocolectomy for ulcerative colitis. *J Gastrointest Surg* 2007; **11**: 3-7 [PMID: 17390179]
- 34 **Peters WR**. Laparoscopic total proctocolectomy with creation of ileostomy for ulcerative colitis: report of two cases. *J Laparoendosc Surg* 1992; **2**: 175-178 [PMID: 1535812]
- 35 **Wexner SD**, Cera SM. Laparoscopic surgery for ulcerative colitis. *Surg Clin North Am* 2005; **85**: 35-47, viii [PMID: 15619527]
- 36 **Mathis KL**, Boostrom SY, Pemberton JH. New developments in colorectal surgery. *Curr Opin Gastroenterol* 2013; **29**: 72-78 [PMID: 23207599 DOI: 10.1097/MOG.0b013e32835a34ea]
- 37 **Lichtenstein GR**, Cohen R, Yamashita B, Diamond RH. Quality of life after proctocolectomy with ileoanal anastomosis for patients with ulcerative colitis. *J Clin Gastroenterol* 2006; **40**: 669-677 [PMID: 16940876]
- 38 **Dunker MS**, Bemelman WA, Slors JF, van Duijvendijk P, Gouma DJ. Functional outcome, quality of life, body image, and cosmesis in patients after laparoscopic-assisted and conventional restorative proctocolectomy: a comparative study. *Dis Colon Rectum* 2001; **44**: 1800-1807 [PMID: 11742165]
- 39 **Heikens JT**, de Vries J, van Laarhoven CJ. Quality of life, health-related quality of life and health status in patients having restorative proctocolectomy with ileal pouch-anal anastomosis for ulcerative colitis: a systematic review. *Colorectal Dis* 2012; **14**: 536-544 [PMID: 21176062 DOI: 10.1111/j.1463-1318.2010.02538.x]
- 40 **Bugra D**. Laparoscopic restorative proctocolectomy technical notes and postoperative results. *Acta Chir Iugosl* 2012; **59**: 39-45 [PMID: 23373357]
- 41 **McLaughlin SD**, Clark SK, Tekkis PP, Ciclitira PJ, Nicholls RJ. Review article: restorative proctocolectomy, indications, management of complications and follow-up--a guide for gastroenterologists. *Aliment Pharmacol Ther* 2008; **27**: 895-909 [PMID: 18266993 DOI: 10.1111/j.1365-2036.2008.03643.x]
- 42 **McGuire BB**, Brannigan AE, O'Connell PR. Ileal pouch-anal anastomosis. *Br J Surg* 2007; **94**: 812-823 [PMID: 17571291]
- 43 **Beliard A**, Prudhomme M. Ileal reservoir with ileo-anal anastomosis: long-term complications. *J Visc Surg* 2010; **147**: e137-e144 [PMID: 20832385 DOI: 10.1016/j.jvisurg.2010.07.001]
- 44 **Huetting WE**, Gooszen HG, van Laarhoven CJ. Sexual function and continence after ileo pouch anal anastomosis: a comparison between a meta-analysis and a questionnaire survey. *Int J Colorectal Dis* 2004; **19**: 215-218 [PMID: 14564464]
- 45 **Heikens JT**, Gooszen HG, Teepen JL, Huetting WE, Oostvogel HJ, van Vroonhoven TJ, van Krieken JH, van Laarhoven CJ. The ileo neo rectal anastomosis: long-term results of surgical innovation in patients after ulcerative colitis and familial adenomatous polyposis. *Int J Colorectal Dis* 2013; **28**: 111-118 [PMID: 22885881 DOI: 10.1007/s00384-012-1545-0]
- 46 **Tariverdian M**, Leowardi C, Hinz U, Welsch T, Schmidt J, Kienle P. Quality of life after restorative proctocolectomy for ulcerative colitis: preoperative status and long-term results. *Inflamm Bowel Dis* 2007; **13**: 1228-1235 [PMID: 17567871]
- 47 **Hoie O**, Wolters FL, Riis L, Bernklev T, Aamodt G, Clofent J, Tsianos E, Beltrami M, Odes S, Munkholm P, Vatn M, Stockbrügger RW, Moum B. Low colectomy rates in ulcerative colitis in an unselected European cohort followed for 10 years. *Gastroenterology* 2007; **132**: 507-515 [PMID: 17258717]
- 48 **Langholz E**, Munkholm P, Davidsen M, Binder V. Course of ulcerative colitis: analysis of changes in disease activity over years. *Gastroenterology* 1994; **107**: 3-11 [PMID: 8020674]
- 49 **Langholz E**, Munkholm P, Davidsen M, Nielsen OH, Binder V. Changes in extent of ulcerative colitis: a study on the course and prognostic factors. *Scand J Gastroenterol* 1996; **31**: 260-266 [PMID: 8833356]
- 50 **Van Assche G**, Dignass A, Reinisch W, van der Woude CJ, Sturm A, De Vos M, Guslandi M, Oldenburg B, Dotan I, Marteau P, Ardizzone A, Baumgart DC, D'Haens G, Gionchetti P, Portela F, Vucelic B, Söderholm J, Escher J, Koletzko S, Kolho KL, Lukas M, Mottet C, Tilg H, Vermeire S, Carbonnel F, Cole A, Novacek G, Reinshagen M, Tsianos E, Herrlinger K, Oldenburg B, Bouhnik Y, Kiesslich R, Stange E, Travis S, Lindsay J. The second European evidence-based Consensus on the diagnosis and management of Crohn's disease: Special situations. *J Crohns Colitis* 2010; **4**: 63-101 [PMID: 21122490]
- 51 **Rutgeerts P**, Sandborn WJ, Feagan BG, Reinisch W, Olson A, Johanss J, Travers S, Rachmilewitz D, Hanauer SB, Lichtenstein GR, de Villiers WJ, Present D, Sands BE, Colombel JF. Infliximab for induction and maintenance therapy for ulcerative colitis. *N Engl J Med* 2005; **353**: 2462-2476 [PMID: 16339095]
- 52 **Sandborn WJ**, Rutgeerts P, Feagan BG, Reinisch W, Olson A, Johanss J, Lu J, Horgan K, Rachmilewitz D, Hanauer SB, Lichtenstein GR, de Villiers WJ, Present D, Sands BE, Colombel JF. Colectomy rate comparison after treatment of ulcerative colitis with placebo or infliximab. *Gastroenterology* 2009; **137**: 1250-1260; quiz 1520 [PMID: 19596014 DOI: 10.1053/j.gastro.2009.06.061]
- 53 **Sands BE**, Tremaine WJ, Sandborn WJ, Rutgeerts PJ, Hanauer SB, Mayer L, Targan SR, Podolsky DK. Infliximab in the treatment of severe, steroid-refractory ulcerative colitis: a pilot study. *Inflamm Bowel Dis* 2001; **7**: 83-88 [PMID: 11383595]
- 54 **Gustavsson A**, Järnerot G, Hertervig E, Friis-Liby I, Blomquist L, Karlén P, Grännö C, Villien M, Ström M, Verbaan H, Hellström PM, Magnuson A, Halfvarson J, Tysk C. Clinical trial: colectomy after rescue therapy in ulcerative colitis - 3-year follow-up of the Swedish-Danish controlled infliximab study. *Aliment Pharmacol Ther* 2010; **32**: 984-989 [PMID: 20937043 DOI: 10.1111/j.1365-2036.2010.04435.x]
- 55 **Costa J**, Magro F, Caldeira D, Alarcão J, Sousa R, Vaz-Carneiro A. Infliximab reduces hospitalizations and surgery interventions in patients with inflammatory bowel disease: a systematic review and meta-analysis. *Inflamm Bowel Dis* 2013; **19**: 2098-2110 [PMID: 23860567 DOI: 10.1097/MIB.0b013e31829936c2]
- 56 **Aratari A**, Papi C, Clemente V, Moretti A, Luchetti R, Koch M, Capurso L, Caprilli R. Colectomy rate in acute severe ulcerative colitis in the infliximab era. *Dig Liver Dis* 2008; **40**: 821-826 [PMID: 18472316 DOI: 10.1016/j.dld.2008.03.014]
- 57 **Teisner AS**, Ainsworth MA, Brynskov J. Long-term effects and colectomy rates in ulcerative colitis patients treated with infliximab: a Danish single center experience. *Scand J Gastroenterol* 2010; **45**: 1457-1463 [PMID: 20701434 DOI: 10.3109/00365521.2010.510572]
- 58 **Ferrante M**, Vermeire S, Fidder H, Schnitzler F, Noman M, Van Assche G, De Hertogh G, Hoffman I, D'Hoore A, Van Steen K, Geboes K, Penninckx F, Rutgeerts P. Long-term outcome after infliximab for refractory ulcerative colitis. *J Crohns Colitis* 2008; **2**: 219-225 [PMID: 21172214 DOI: 10.1016/j.crohns.2008.03.004]
- 59 **Oussalah A**, Evesque L, Laharie D, Roblin X, Boschetti G,

- Nancey S, Filippi J, Flourié B, Hebuterne X, Bigard MA, Peyrin-Biroulet L. A multicenter experience with infliximab for ulcerative colitis: outcomes and predictors of response, optimization, colectomy, and hospitalization. *Am J Gastroenterol* 2010; **105**: 2617-2625 [PMID: 20736936 DOI: 10.1038/ajg.2010.345]
- 60 **Desmond AN**, Shanahan F. Managing chronic disease in Ireland: hospital admission rates and clinical outcomes in a large ulcerative colitis population. *Ir J Med Sci* 2012; **181**: 65-71 [PMID: 21947686 DOI: 10.1007/s11845-011-0760-y]
- 61 **Garcia-Planella E**, Mañosa M, Van Domselaar M, Gordillo J, Zabana Y, Cabré E, López San Román A, Domènech E. Long-term outcome of ulcerative colitis in patients who achieve clinical remission with a first course of corticosteroids. *Dig Liver Dis* 2012; **44**: 206-210 [PMID: 22079262 DOI: 10.1016/j.dld.2011.10.004]
- 62 **Selvasekar CR**, Cima RR, Larson DW, Dozois EJ, Harrington JR, Harmsen WS, Loftus EV, Sandborn WJ, Wolff BG, Pemberton JH. Effect of infliximab on short-term complications in patients undergoing operation for chronic ulcerative colitis. *J Am Coll Surg* 2007; **204**: 956-962; discussion 962-963 [PMID: 17481518]
- 63 **Mor IJ**, Vogel JD, da Luz Moreira A, Shen B, Hammel J, Remzi FH. Infliximab in ulcerative colitis is associated with an increased risk of postoperative complications after restorative proctocolectomy. *Dis Colon Rectum* 2008; **51**: 1202-1207; discussion 1207-1210 [PMID: 18536964 DOI: 10.1007/s10350-008-9364-7]
- 64 **Schluender SJ**, Ippoliti A, Dubinsky M, Vasiliauskas EA, Papadakis KA, Mei L, Targan SR, Fleshner PR. Does infliximab influence surgical morbidity of ileal pouch-anal anastomosis in patients with ulcerative colitis? *Dis Colon Rectum* 2007; **50**: 1747-1753 [PMID: 17704969 DOI: 10.1007/s10350-007-9008-3]
- 65 **Ferrante M**, D'Hoore A, Vermeire S, Declerck S, Noman M, Van Assche G, Hoffman I, Rutgeerts P, Penninckx F. Corticosteroids but not infliximab increase short-term postoperative infectious complications in patients with ulcerative colitis. *Inflamm Bowel Dis* 2009; **15**: 1062-1070 [PMID: 19161179 DOI: 10.1002/ibd.20863]
- 66 **Gainsbury ML**, Chu DI, Howard LA, Coukos JA, Farraye FA, Stocchi AF, Becker JM. Preoperative infliximab is not associated with an increased risk of short-term postoperative complications after restorative proctocolectomy and ileal pouch-anal anastomosis. *J Gastrointest Surg* 2011; **15**: 397-403 [PMID: 21246415 DOI: 10.1007/s11605-010-1385-6]
- 67 **Rizzo G**, Armuzzi A, Pugliese D, Verbo A, Papa A, Mattana C, Rapaccini GL, Guidi L, Coco C. Anti-TNF-alpha therapies do not increase early postoperative complications in patients with inflammatory bowel disease. An Italian single-center experience. *Int J Colorectal Dis* 2011; **26**: 1435-1444 [PMID: 21594668 DOI: 10.1007/s00384-011-1236-2]
- 68 **Yang Z**, Wu Q, Wang F, Wu K, Fan D. Meta-analysis: effect of preoperative infliximab use on early postoperative complications in patients with ulcerative colitis undergoing abdominal surgery. *Aliment Pharmacol Ther* 2012; **36**: 922-928 [PMID: 23002804 DOI: 10.1111/apt.12060]
- 69 **Tsai HH**, Puneekar YS, Morris J, Fortun P. A model of the long-term cost effectiveness of scheduled maintenance treatment with infliximab for moderate-to-severe ulcerative colitis. *Aliment Pharmacol Ther* 2008; **28**: 1230-1239 [PMID: 18729845 DOI: 10.1111/j.1365-2036.2008.03839.x]
- 70 **Park KT**, Tsai R, Perez F, Cipriano LE, Bass D, Garber AM. Cost-effectiveness of early colectomy with ileal pouch-anal anastomosis versus standard medical therapy in severe ulcerative colitis. *Ann Surg* 2012; **256**: 117-124 [PMID: 22270693 DOI: 10.1097/SLA.0b013e3182445321]

P- Reviewers: Martin FPJ, Wasko-Czopnik D

S- Editor: Zhai HH **L- Editor:** A **E- Editor:** Wang CH

