

Protein Kinase B (PknB) of *Mycobacterium tuberculosis* Is Essential for Growth of the Pathogen *in Vitro* as well as for Survival within the Host*

Received for publication, March 10, 2014; Published, JBC Papers in Press, April 4, 2014; DOI 10.1074/jbc.M114.563536

Yogesh Chawla^{†1}, Sandeep Upadhyay^{†1}, Shazia Khan^{‡2}, Sathya Narayanan Nagarajan[‡], Francesca Forti[§], and Vinay Kumar Nandicoori^{‡3}

From the [†]National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi 110067, India and the [§]Dipartimento di Scienze Biomolecolari e Biotecnologie, Università degli Studi di Milano, Via Celoria 26, 20133 Milano, Italy

Background: *Mycobacterium tuberculosis* PknB plays a critical role in modulating cell division and cell wall synthesis.

Results: We present a comprehensive evaluation of the importance of various domains of PknB in modulating cell survival.

Conclusion: The intracellular kinase domain and extracytoplasmic PASTA domains of PknB are essential for cell survival.

Significance: PknB is essential for both *in vitro* growth and survival of the pathogen *in vivo*.

The *Mycobacterium tuberculosis* protein kinase B (PknB) comprises an intracellular kinase domain, connected through a transmembrane domain to an extracellular region that contains four PASTA domains. The present study describes the comprehensive analysis of different domains of PknB in the context of viability in avirulent and virulent mycobacteria. We find stringent regulation of PknB expression necessary for cell survival, with depletion or overexpression of PknB leading to cell death. Although PknB-mediated kinase activity is essential for cell survival, active kinase lacking the transmembrane or extracellular domain fails to complement conditional mutants not expressing PknB. By creating chimeric kinases, we find that the intracellular kinase domain has unique functions in the virulent strain, which cannot be substituted by other kinases. Interestingly, we find that although the presence of the C-terminal PASTA domain is dispensable in the avirulent *M. smegmatis*, all four PASTA domains are essential in *M. tuberculosis*. The differential behavior of PknB vis-à-vis the number of essential PASTA domains and the specificity of kinase domain functions suggest that PknB-mediated growth and signaling events differ in virulent compared with avirulent mycobacteria. Mouse infection studies performed to determine the role of PknB in mediating pathogen survival in the host demonstrate that PknB is not only critical for growth of the pathogen *in vitro* but is also essential for the survival of the pathogen in the host.

Protein phosphorylation/dephosphorylation is a critical regulatory mechanism by which extracellular cues are transduced into cellular responses. Traditionally, two-component systems comprising a histidine kinase sensor and the associated response regulator were thought to be responsible for phosphor-

ylation-mediated signaling in prokaryotes (1, 2). Two-component pathways are rare in eukaryotes, and phosphotransfer-based signaling predominantly involves phosphorylation/dephosphorylation on serine, threonine, or tyrosine residues, often in a cascade. The presence of a eukaryotic-like serine/threonine protein kinase (STPK)⁴ in a prokaryote was first reported in 1991 (*pkn1* in *Myxococcus xanthus*) (3). These “eukaryotic-like” STPKs play important roles in bacterial cellular processes, including cell division, cell wall synthesis, cell metabolism, and dormancy exit (4–7). Analysis of the *Mycobacterium tuberculosis* genome sequence suggested the presence of 11 putative eukaryotic-like STPKs and three protein phosphatases (8). Except for PknG and PknK, all of these kinases were predicted to have a transmembrane domain (8, 9). All of the kinases possessed the protein kinase “signature” motifs, including 11 conserved subdomains as per Hanks’ criteria, and amino acid sequence alignment of these STPK family members revealed that 15 catalytically important residues were conserved across all of them (9, 10). The *M. tuberculosis* STPKs affect key mycobacterial processes: signal transduction mediated by PknA and PknB plays an important role in determining cell shape, morphology, and possibly cell division (7); PknG and PknH influence *M. tuberculosis* virulence, adaptation, and growth within the host (11, 12); and PknF affects cell division, growth rate, morphology, and glucose transport (13). Recently, Ortega *et al.* (14) proposed a role for PknB as a replication switch in response to hypoxia. They demonstrated that PknB activity is necessary for reactivation of cells from the hypoxic state.

Protein kinases A and B, encoded by *pknA* and *pknB*, respectively, are part of the operon carrying cistrons coding for protein phosphatase *pstP*, *rodA* (involved in cell shape control), and *pbpA* (involved in peptidoglycan synthesis). This locus also

* This work was supported by grants from the National Institute of Immunology (New Delhi, India) and the Department of Biotechnology, India (to V. K. N.).

¹ Senior Research Fellow of the Council of Scientific and Industrial Research.

² Present address: Wellman Center for Photomedicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114.

³ To whom correspondence should be addressed. Tel.: 91-11-26703789; Fax: 91-11-26742125; E-mail: Vinaykn@nii.ac.in.

⁴ The abbreviations used are: STPK, serine/threonine protein kinase; JM, juxtamembrane; TM, transmembrane; PASTA domain, penicillin-binding protein and serine threonine kinase-associated domain; FHA domain, forkhead-associated domain; MyBP, myelin-binding protein; MBP, maltose-binding protein; aa, amino acids; ATc, anhydrotetracycline; IVN, isovaleronitrile; BAC, bacterial artificial chromosome.

includes two FHA (forkhead-associated) domain-containing genes and is found near the origin of replication throughout the *Mycobacterium* genus (15). As in most other kinases, PknA and PknB consist of a kinase domain, a juxtamembrane region, a short transmembrane domain, and an extracellular region (9, 16). Both kinases have been found to be essential based on transposon-insertion experiments (17), and the *pknB* gene can be disrupted by allelic replacement in *M. tuberculosis* and *M. smegmatis* only in the presence of a second functional copy of the gene (18). In 2009, Forti *et al.* (19) generated the *M. tuberculosis* H37Rv-p_{pptr}-*pknB* conditional mutant in which the genomic copy of *pknB* was converted to a pristinamycin-inducible copy and demonstrated its essentiality, reaffirming the previous findings.

PknB is among the highly characterized serine/threonine protein kinases of mycobacteria. The kinase domain of PknB (aa 1–279) by itself has been shown to be sufficient for its activity (20); however, the extent of its activity in comparison with full-length protein has not been determined. Crystal structure analysis of the PknB kinase domain found it to be a two-lobed structure (N- and C-terminal lobes) showing conservation of protein fold and catalytic machinery to eukaryotic STPK homologs (21, 22). The N-terminal lobe contains the ATP binding site, whereas the C-terminal lobe is involved in rendering an active state and in stabilizing interactions with the substrate (23). Based on the structures, PknB is proposed to form both back-to-back and front-to-front dimers (24). The dimerization of PknB was shown to be essential for autophosphorylation and activation of kinase through an allosteric mechanism (23). The extracellular domain of PknB is predicted to have four conserved PASTA (penicillin-binding protein and serine/threonine kinase-associated) domains (22, 25). This domain has been suggested to play a role in the recognition of D-alanyl-D-alanine dipeptides used to build up the peptidoglycan layers (25). Using nuclear magnetic resonance and small angle x-ray scattering, the four PASTA domains have been shown to be organized in a linear fashion, in contrast to penicillin-binding protein PBP2x (from *Streptococcus pneumoniae*), where the two PASTA domains fold over in a compact arrangement (26, 27). The extracellular domain of *M. tuberculosis* PknB has also been shown to bind mucopeptides, and the specific amino acids in the stem peptide that impact the efficacy of the interaction have been identified (28).

Although the PknB structure and the mode of activation *in vitro* are now understood, structure-function relationships of the various domains have not been investigated in the context of mycobacterial growth and survival. The present study was undertaken to comprehensively evaluate the importance of the different domains in modulating PknB function *in vivo*. Using *pknB* conditional mutants in *Mycobacterium smegmatis* and *M. tuberculosis* strains, we observe that variable expression of PknB regulates cell growth and morphology, with both overexpression and depletion leading to cell death. Interestingly, we find that the minimum number of PASTA domains required in the extracytoplasmic region varies between avirulent and virulent mycobacteria. Infection studies with a mouse model reveal that depletion of PknB results in clearance of pathogen from the

host tissues, indicating definitively that PknB is essential for survival of the pathogen within the host.

EXPERIMENTAL PROCEDURES

Bacterial Strains, Reagents, and Radioisotopes—Bacterial strains used in the study are listed in Table 1. Cloning and expression vectors were purchased or obtained from their respective sources: pENTR-D-TOPO vector (Invitrogen), pMAL-c2x expression vector (New England Biolabs), pNit-1 vector (29), pST-HiT vector (30), and p2Nil suicide delivery vector (31). [γ -³²P]ATP (6000 Ci/mmol) was purchased from PerkinElmer Life Sciences. Pristinamycin 1A was purchased from Molcan Corp. All medium components were purchased from BD Biosciences. Restriction/modification enzymes were purchased from New England Biolabs and MBI Fermentas. DNA oligonucleotides and analytical grade chemicals and reagents were purchased from Sigma-Aldrich or GE Healthcare.

Creation of *pknB* Conditional Gene Replacement Mutant in *M. smegmatis* (*mc*² Δ *pknB*)—The integration-proficient HiT-PknB construct expressing *M. tuberculosis pknB* under a tetracycline-inducible promoter was electroporated into *M. smegmatis* *mc*²155 to generate the merodiploid strain *mc*²*HiT-pknB*. Upstream and downstream flanks of *pknB* were amplified using specific primers, and the flanks were cloned into HpaI-HindIII sites in the p2Nil vector. This was followed by the cloning of a 6-kb PacI cassette from pGOAL17 into the corresponding site on p2Nil to generate the p2Nil- Δ *pknB* construct. The two-step recombination procedure described by Parish and Stoker was used to disrupt the *pknB* gene at its native locus in the genome (31). The mutant *mc*² Δ *pknB* (clone numbers 23 and 26) surviving only in the presence of the inducer anhydrotetracycline (ATc) was used for further studies (Fig. 1, A–C). The creation of the *M. tuberculosis* H37Rv pristinamycin-inducible *pknB* mutant p_{pptr}-*pknB* (Rv-p_{pptr}-B) used in this study was reported earlier (19).

Generation of Plasmid Constructs—Full-length *M. tuberculosis pknA*, *pknB*, and *pknH* genes were amplified from BAC clones (8) using gene-specific primers and Phusion DNA polymerase (New England Biolabs), and the amplicons were cloned into pENTR-D-TOPO, pMAL-c2X, or pNit vector. HiT-PknB was created by subcloning the NdeI-HindIII fragment from pENTR-PknB into the corresponding sites in pST-HiT. The PknB-KD (aa 1–279), PknB-JM (aa 1–331), PknB-TM (aa 1–352), PknB-PASTA1 (aa 1–422), PknB-PASTA2 (aa 1–490), and PknB-PASTA3 (aa 1–557) fragments were amplified from pENTR-PknB, using PknB-specific forward and domain-specific reverse primers, and the amplicons were cloned into pMAL-c2X and pNit vectors. All point mutations were made by overlapping PCR using appropriate mutagenic primers, and the mutations were confirmed by DNA sequencing. To make the PknA-B and PknH-B chimeric proteins, the intracellular domains of PknA or PknH were fused to the transmembrane and extracellular domains of PknB using overlapping PCR.

Protein Expression, Purification, and *In Vitro* Kinase Assay—pMAL-c2X constructs expressing PknB or its mutants were transformed into *Escherichia coli* BL21 (DE3) Codon Plus cells

M. tuberculosis PknB Is Essential for Pathogen Survival

TABLE 1

Strains used in the study

Strains	Description	Source
DH5 α	<i>E. coli</i> strain used for cloning experiments	Invitrogen
BL21 (DE3) codon plus	<i>E. coli</i> strain used for protein expression	Stratagene
mc ² 155	Wild type <i>M. smegmatis</i> strain	ATCC, 700084
mc ² HiT-PknB	mc ² 155 strain transformed with integrative vector containing ATc inducible PknB at L5 site; Hyg ^r	This study
mc ² ::pNit	mc ² 155 strain transformed with pNit vector	This study
mc ² ::PknB	mc ² 155 strain transformed with pNit PknB	This study
mc ² Δ pknB	<i>M. smegmatis</i> pknB conditional gene replacement mutant	This study
mc ² Δ pknB::pNit	mc ² Δ pknB transformed with pNit-1 vector	This study
mc ² Δ pknB::PknB	mc ² Δ pknB transformed with pNit-PknB construct	This study
mc ² Δ pknB::PknB-K40M	mc ² Δ pknB transformed with pNit-PknB-K40M construct	This study
mc ² Δ pknB::PknB-KD	mc ² Δ pknB transformed with pNit-PknB-KD (aa 1–279) construct	This study
mc ² Δ pknB::PknB-JM	mc ² Δ pknB transformed with pNit-PknB-JM (aa 1–331) construct	This study
mc ² Δ pknB::PknB-TM	mc ² Δ pknB transformed with pNit-PknB-TM (aa 1–352) construct	This study
mc ² Δ pknB::PknB-(A-TM)	mc ² Δ pknB transformed with pNit-PknB-(A-TM) construct	This study
mc ² Δ pknB::PknB-(H-TM)	mc ² Δ pknB transformed with pNit-PknB-(H-TM) construct	This study
mc ² Δ pknB::PknB- Δ TM	mc ² Δ pknB transformed with pNit-PknB- Δ TM construct	This study
mc ² Δ pknB::PknB-PASTA1	mc ² Δ pknB transformed with pNit-PknB-PASTA1 (aa 1–422) construct	This study
mc ² Δ pknB::PknB-PASTA2	mc ² Δ pknB transformed with pNit-PknB-PASTA2 (aa 1–490) construct	This study
mc ² Δ pknB::PknB-PASTA3	mc ² Δ pknB strain transformed with pNit-PknB-PASTA3 (aa 1–557) construct	This study
mc ² Δ pknB::PknA-B	mc ² Δ pknB transformed with pNit PknA-B (aa 1–338 of PknA fused with aa 332–626 of PknB) construct	This study
mc ² Δ pknB::PknH-B	mc ² Δ pknB transformed with pNit PknH-B (aa 1–402 of PknH fused with aa 332–626 of PknB) construct	This study
H37Rv	Wild type <i>M. tuberculosis</i> strain	ATCC
H37Rv::pNit	H37Rv strain transformed with pNit-1 vector	This study
H37Rv::PknB	H37Rv strain transformed with pNit-PknB construct	This study
Rv-pptr-B	<i>M. tuberculosis</i> H37Rv pknB conditional expression strain under regulation of pptr promoter	Ref. 19
Rv-pptr-B::pNit	Rv-pptr-B transformed with pNit-1 vector	This study
Rv-pptr-B::PknB	Rv-pptr-B transformed with pNit PknB construct	This study
Rv-pptr-B::PknB-K40M	Rv-pptr-B transformed with pNit-PknB-K40M construct	This study
Rv-pptr-B::PknB-KD	Rv-pptr-B transformed with pNit-PknB-KD construct	This study
Rv-pptr-B::PknB-JM	Rv-pptr-B transformed with pNit-PknB-JM construct	This study
Rv-pptr-B::PknB-TM	Rv-pptr-B transformed with pNit-PknB-TM construct	This study
Rv-pptr-B::PknB-(A-TM)	Rv-pptr-B transformed with pNit PknB-(A-TM) construct	This study
Rv-pptr-B::PknB-(H-TM)	Rv-pptr-B transformed with pNit PknB-(H-TM) construct	This study
Rv-pptr-B::PknB- Δ TM	Rv-pptr-B transformed with pNit PknB- Δ TM construct	This study
Rv-pptr-B::PknB-PASTA1	Rv-pptr-B transformed with pNit PknB-PASTA1 construct	This study
Rv-pptr-B::PknB-PASTA2	Rv-pptr-B transformed with pNit PknB-PASTA2 construct	This study
Rv-pptr-B::PknB-PASTA3	Rv-pptr-B transformed with pNit PknB-PASTA3 construct	This study
Rv-pptr-B::PknA-B	Rv-pptr-B transformed with pNit PknA-B construct	This study
Rv-pptr-B::PknH-B	Rv-pptr-B transformed with pNit PknH-B construct	This study

(Stratagene), and the MBP-tagged proteins were purified as described (32). His-GarA was expressed and purified as described earlier (33). An *in vitro* kinase assay was performed as described previously using 2.5 pmol of PknB kinase or its mutants and using 100 pmol of GarA or MyBP as the substrate (33).

Growth Pattern Analysis—The strategy used for creation of the pknB conditional gene replacement mutant in *M. smegmatis* (mc² Δ pknB) is described above. Electrocompetent cells of mc²155 or H37Rv were transformed with pNit or pNit-PknB to generate mc²::pNit, mc²::PknB, Rv::pNit, and Rv::PknB. To analyze the growth patterns of *M. smegmatis* mc²155, mc²::pNit, mc²::PknB, and mc² Δ pknB, each strain was grown in LB medium containing 0.05% Tween 80, 0.2% glycerol, and appropriate antibiotic to A₆₀₀ ~0.8, either in the absence of inducer isovaleronitrile (IVN) in the case of mc²::pNit and mc²::PknB or in the presence of inducer ATc in the case of mc² Δ pknB mutant. *M. tuberculosis* transformants Rv::pNit and Rv::PknB as well as the mutant strain Rv-pptr-B were grown to A₆₀₀ ~0.8 in 7H9-ADC containing 0.05% Tween 80, 0.2% glycerol, and appropriate antibiotic in the presence of inducer pristinamycin 1A in the case of Rv-pptr-B or in the absence of inducer IVN in the case of H37Rv, Rv::pNit and Rv::PknB. The cultures were washed twice in PBS containing 0.05% Tween 80 and were used to seed fresh cultures at an initial A₆₀₀ of 0.02 (in the case of *M. smegmatis*) or 0.1 (in the case of *M. tuberculosis*) in the pres-

ence or absence of inducer. To analyze growth in *M. smegmatis* strains, A₆₀₀ was measured every 3–30 h after initiation. To analyze growth in *M. tuberculosis* strains, A₆₀₀ was measured every 24 h to 8 days after initiation. Experiments were performed in triplicates, and the average A₆₀₀ values were plotted as a function of time. S.E. values were calculated using Sigma-Plot for each time point. For Western blot analysis, cultures of *M. smegmatis* strains were initiated at an A₆₀₀ of 0.2 and were grown for 9 h in the absence or presence of ATc. Cultures of *M. tuberculosis* strains were likewise initiated and grown in the absence or presence of inducer for 6 days. In order to determine whether the absence of PknB has bacteriostatic or bacteriocidal effects, cultures were withdrawn on days 0, 2, and 5, and different dilutions were spotted on plates containing the inducer pristinamycin 1A. In growth analysis of complementation mutants, mc² Δ pknB or Rv-pptr-B strains were electroporated with pNit or pNit-PknB or pNit constructs containing various PknB mutants, including chimeric proteins, and growth patterns were analyzed as described above.

Scanning Electron Microscopy—In order to perform scanning electron microscopy, cells were incubated for 3 h in fixative (2.5% glutaraldehyde, 4% paraformaldehyde in 0.1 M sodium cacodylate buffer, pH 7.3). Fixed cells were then treated with increasing concentrations of ethanol (ranging from 25 to 100%) stepwise, followed by hexamethyldisilazane treatment. This

was followed by coating with gold particles and visualizing under the microscope at $\times 10,000$ magnification (Carl Zeiss, Evo LS scanning electron microscope).

Infection of Mice—*M. tuberculosis* H37Rv or Rv-pptr-B or Rv-pptr-B::PknB were grown in the presence of pristinamycin until midlogarithmic phase (A_{600} of 0.6). The bacilli were harvested by centrifugation at $3000 \times g$, washed twice with PBS, and resuspended in saline. To evenly disperse the bacteria, the suspension was passed through a 27.5-gauge needle 10 times. C57BL/6 mice (6–8 weeks old of either sex) were housed in individually ventilated cages at the Tuberculosis Aerosol Challenge Facility, International Centre for Genetic Engineering and Biotechnology (New Delhi, India). Mice ($n = 6$) were infected with 2×10^8 colony-forming units of either H37Rv, Rv-pptr-B, or Rv-pptr-B::PknB by the aerosol route using the Madison Aerosol Chamber (University of Wisconsin, Madison, WI) precalibrated to deliver ~ 80 – 150 bacilli/mice. Two mice each were sacrificed at 24 h postexposure, and the lung homogenates were plated on nutrient 7H11 agar in triplicates for assessing the bacterial load. Bacterial loads for each set were again assessed 4 weeks and 8 weeks after aerosol exposure. This was achieved by homogenizing aseptically removed organs in 2 ml of sterile saline containing 0.05% Tween 80 (Sigma-Aldrich). Serial dilutions of homogenates thus obtained were plated on 7H11 plates supplemented with 10% OADC (BD Biosciences), and colonies were counted after 21 days.

Histopathology—Harvested organs were fixed in 10% neutral buffered formalin. The tissue was embedded in paraffin wax and cut into 5- μ m sections using a microtome. Sections were then stained with hematoxylin and eosin (H&E) stain and subjected to histopathological analyses. The tissue samples were coded, and a pathologist having no prior knowledge of the experimental groups evaluated lung sections for granulomatous architecture, and the numbers of granulomas were counted. Each granuloma was graded by the following criteria: (a) granulomas with necrosis were given a score of 5; (b) granulomas without necrosis were given a score of 2.5; (c) granulomas with fibrous connective tissue were given a score of 1. Total granuloma score was calculated by multiplying the number of granulomas of each type by the score and then adding them up to obtain a total granuloma score for each sample.

RESULTS

Stringent Regulation of PknB Expression in the Cell Is Necessary for Cell Growth—In order to investigate the role of PknB in modulating physiological events in mycobacteria, we created a PknB conditional knock-out strain. We adopted the approach of integrating an inducible copy of the gene in the genome prior to deleting the *pknB* gene in *M. smegmatis* at its native locus. A comparison of the PCR amplicons obtained in potential deletion mutants, wild type strain, merodiploid strain, and negative clone revealed that the native *pknB* gene had been successfully disrupted (Fig. 1, A–C). In a previous study, *M. tuberculosis* H37Rv-pptr-B (Rv-pptr-B) mutant strain was created with the help of a suicide delivery plasmid, in which the *pknB* gene in the genome was replaced with a pristinamycin-inducible copy (19) (Fig. 1F). To characterize the functions of PknB, we used

the *M. smegmatis* $\Delta pknB$ ($mc^2\Delta pknB$) strain created in this study and the Rv-pptr-B mutant from the previous study.

PknB is the last gene of the operon, which, in addition to others, carries the *pknA* gene, the only other essential serine/threonine protein kinase in *M. tuberculosis*. We therefore ascertained whether the expression of *pknA*, which lies immediately upstream of *pknB*, is affected in the mutants. We found that although the down-regulation of PknB expression in the absence of ATc was evident even after 6 h, it takes up to 9 h for substantial depletion (Figs. 1D and 2C). The expression of PknA was unaffected by the depletion of PknB (Fig. 2C). Conversely, PknB was considerably overexpressed upon induction with IVN in strain $mc^2::PknB$ (Fig. 1D). Analysis carried out with *M. tuberculosis* strains revealed a considerable diminution in the levels of PknB when Rv-pptr-B was grown in the absence of pristinamycin and substantial overexpression of PknB in Rv::PknB strain upon induction with IVN (Figs. 1G and 2A). Here too PknA expression was unaffected by depletion or overexpression of PknB (Fig. 1G). These results suggest that disrupting the *pknB* at its native location did not have any impact on the expression of other genes in the operon.

We found that the growth of mycobacterial strains, $mc^2::PknB$ and Rv::PknB, was not significantly different from that of wild type cells in the absence of inducer IVN (Fig. 1, E and H). Importantly, we found that the overexpression of PknB dramatically compromised growth in the case of both $mc^2::PknB$ and Rv::PknB strains (Fig. 1, E and H). Similarly, $mc^2\Delta pknB$ and Rv-pptr-B growth patterns did not significantly diverge from those of the corresponding wild type strains in the presence of inducers. Depletion of PknB had strong effects on mycobacterial growth in both *M. smegmatis* and *M. tuberculosis* (Fig. 1, E and H). Although it was apparent that mycobacteria showed compromised growth upon PknB depletion, it was not clear if the absence of PknB was leading to a bacteriostatic or bactericidal phenotype. In order to differentiate between these possibilities, H37Rv and Rv-pptr-B cultures grown in the presence or absence of inducer for 0, 2, or 5 days were spotted at different dilutions on plates containing pristinamycin (Fig. 2B). Western blot analysis of lysates of Rv-pptr-B prepared from cultures grown with or without pristinamycin after 2, 4, or 6 days showed that efficient depletion of PknB could only be observed after 4–6 days of growth (Fig. 2A). In agreement with this observation, inducer depletion for 2 days had no obvious effect on cell growth (Fig. 2B). However, when cultures were spotted on plates after 5 days of growth in the absence of inducer, we clearly observed drastic reduction (~ 3 log fold) in the titer (Fig. 2B). These results strongly suggest that depletion of PknB eventually leads to cell death.


The overexpression of PknB in *M. smegmatis* and *Mycobacterium bovis* BCG has been reported to alter the appearance of the cells, making them wider and bulging (7). Because efficient reduction of PknB expression could only be observed 9 h after the removal of inducer (Fig. 2C), scanning electron microscopy was performed after 9 and 12 h of overexpression or removal of inducer. PknB overexpression for 9 h results in cells of uneven width, with certain parts of the cells bulging out (Fig. 2D). Overexpression of PknB for 12 h resulted in the lysis of a majority of the cells, causing an accumulation of cell debris (Fig. 2D).

M. tuberculosis PknB Is Essential for Pathogen Survival

Depletion of PknB over a period of 9 h had an extensive impact on the morphology of the cells. Cells appeared shrunken, and some of the cells seemed to be on the verge of lysis (Fig. 2D). When cells were visualized after depletion over a period of 12 h, extensive lysis was observed in a majority of the cells. Taken together, we conclude that PknB is absolutely essential in both avirulent and virulent strains of mycobacteria, and its depletion

or overexpression alters cell morphology, eventually leading to cell death.

Kinase Activity of PknB Is Essential for Cell Survival—PknB is a 626-aa protein with four distinct domains: an N-terminal kinase domain spanning 1–279 aa, a juxtamembrane domain located between 279–331 aa, a 20-amino acid transmembrane domain (residues 332–353 aa), and an extracellular segment



273 amino acids in length, containing four PASTA domains (Fig. 3A). The invariant lysine residue in the kinase domain that interacts with the glutamate residue of α C helix and stabilizes ATP by making contacts with α - and β -phosphates is present at position 40. We expressed MBP-tagged PknB and PknB-K40M (putative kinase-dead mutant) in *E. coli* and purified the recombinant proteins for *in vitro* kinase assays using GarA (an FHA domain containing protein that has previously been identified as a substrate of PknG and PknB (34, 35)) as substrate. Results obtained (Fig. 3, B and C) unequivocally demonstrated the necessity of the invariant lysine for PknB kinase activity, in agreement with previous data (36, 37). To examine the importance of PknB kinase activity in mediating cell survival, Rv-pptr-B and $mc^2\Delta pknB$ strains were transformed with pNit vector, pNit-PknB, or pNit-PknB-K40M, and growth patterns were analyzed. Western blot analysis of whole cell extracts prepared from $mc^2\Delta pknB$ transformants in the absence of IVN and in the presence and absence of ATc demonstrated robust expression of PknB and PknB-K40M (Fig. 3D). Western blot analysis of Rv-pptr-B transformants also demonstrated robust expression of proteins (Fig. 3F), suggesting that leaky expression from the IVN-inducible promoter is adequate. Analysis of growth (Fig. 3, E and G) showed that conditional mutants of mycobacteria transformed with vector failed to survive in the absence of inducers. Rv-pptr-B and $mc^2\Delta pknB$ transformed with pNit-PknB grew very well, demonstrating that wild type PknB expressed episomally was capable of functional complementation. On the other hand, mycobacterial strains transformed with plasmids expressing the kinase-dead PknB-K40M did not show growth recovery, thus underlining the importance of PknB kinase activity in mediating cell survival.

The Extracellular Domain of PknB Is Necessary for Survival—The kinase domain of PknB is capable of autophosphorylation as well as phosphorylation of substrates *in vitro* (38, 39). However, to date, the relative activities of different deletion constructs of PknB in comparison with the full-length PknB have not been determined. Accordingly, PknB-KD, PknB-JM, and PknB-TM deletion mutants (Fig. 4A) were purified, and their ability to phosphorylate substrates *in vitro* was determined at a limiting concentration of enzyme (Fig. 4, B and C). The activity of PknB deletion mutants ranged from 25 to 50% when compared with full-length PknB, with PknB-JM being the least active.

Next we sought to examine the possibility of various PknB deletion mutants rescuing growth of mycobacteria PknB conditional mutants. Western blot analysis of cell lysates from $mc^2\Delta pknB$ transformants showed IVN-inducible expression of PknB fragments (Fig. 4D). Growth analysis of $mc^2\Delta pknB$ transformants revealed that although the wild type PknB was able to rescue growth in cells depleted of endogenous PknB, all of the PknB deletion mutant proteins failed to do so (Fig. 4E). To determine the ability of PknB mutants to complement the Rv-pptr-B, transformants were streaked on solid media in the presence or absence of pristinamycin. Although all of the transformed strains grew robustly in the presence of inducer, full-length PknB alone was able to rescue growth in absence of inducer (Fig. 4F). Thus, whereas PknB kinase activity was essential for functional complementation (Fig. 3), the catalytic domain alone (PknB-KD or PknB-JM) or anchoring of the active kinase to the cell membrane (PknB-TM) was insufficient for rescuing the growth defects. These results suggest that in addition to PknB kinase activity, the extracellular segment is essential for cell survival.

The Localization of PknB to the Cell Membrane Is Essential for Cell Survival—The intracellular and extracellular domains of PknB are connected through a transmembrane (TM) domain of ~ 20 amino acids. However, the precise role of the TM in targeting PknB to the cell membrane has not been elucidated. We reasoned that, if the only function of TM domain is to interact with the lipid bilayer, then replacing it with TM domains of other transmembrane kinases should not have any impact on PknB function. To test this hypothesis, we created plasmid constructs in which the PknB transmembrane domain was either deleted or swapped with that of PknA (control), PknA, or PknH (Fig. 5A). Western blot analysis of $mc^2\Delta pknB$ cells transformed with the various constructs demonstrated proficient expression of the PknB-TM deletion mutant as well as the PknB proteins carrying TM domains of other kinases (Fig. 5B). Growth analysis of transformants showed that swapping the TM domain of PknB with that of PknA or PknH did not affect their ability to complement $\Delta pknB$ (Fig. 5C), although PknB-(A-TM) grew marginally slower. However, deletion of the TM domain was not tolerated. This was reflected in the inability of PknB- Δ TM to complement $mc^2\Delta pknB$ (Fig. 5C), suggesting that localization of PknB to the membrane is essential for cell viability. Results obtained with Rv-pptr-B trans-


FIGURE 1. PknB depletion and PknB over expression affects growth of mycobacteria. A, schematic representation of $mc^2\Delta pknB$ conditional mutant. The primers used for PCR-based confirmation are depicted as F1 and F2 (forward primers) and R1 and R2 (reverse primers). B, agarose gels showing PCR amplification using the respective genomic DNA of different *M. smegmatis* strains, with various primer sets as indicated. Expected sizes of PCR amplicons are as follows: for the F1-R1 primer pair: wild type, 0.95 kb; mutants, no amplicon; for the F1-R2 primer pair: wild type, 1.98 kb; mutants, 0.49 kb; for the F2-R2 primer pair: wild type, 2.9 kb; mutants, 1.42 kb. Leftmost lane, gene ruler 1-kb DNA ladder; lane 1, mc^2155 (wild type); lane 2, mc^2 HiT-PknB (merodiploid PknB strain); lane 3, $mc^2\Delta pknB$ -23 (conditional depletion strain); lane 4, $mc^2\Delta pknB$ -26 (conditional depletion strain); lane 5, negative conditional depletion strain. C, growth of *M. smegmatis* mc^2155 , $mc^2::pNit$ -PknB, negative mutant clone, and $mc^2\Delta pknB$ clones 23 and 26 on 7H10 agar plates containing either no antibiotic, hygromycin only, or hygromycin with ATc inducer. D, all of the cultures were seeded at an initial A_{600} of 0.2 in order to obtain sufficient cell pellets for whole cell lysate (WCL) preparations. WCLs from *M. smegmatis* mc^2155 (mc^2155), mc^2155 transformed with pNit ($mc^2::pNit$), or pNit-PknB ($mc^2::PknB$) grown for 6 h in the absence or presence of 5 μ M IVN inducer and PknB conditional mutant strain *M. smegmatis* $\Delta pknB$ ($mc^2\Delta pknB$) grown for 6 h in the absence or presence of ATc were subjected to Western blotting with rabbit polyclonal α -PknB and α -GroEL1 antibodies. GroEL1 was used as a loading control to demonstrate equal loading of lysates. E, *in vitro* growth pattern analysis. All of the cultures were seeded at an initial A_{600} of 0.02. Growth of the different strains was monitored as described under "Experimental Procedures." F, schematic representation of *M. tuberculosis* pptr-pknB (Rv-pptr-B) mutant. G, *M. tuberculosis* H37Rv (H37Rv), H37Rv transformed with pNit (Rv::pNit), or pNit-PknB (Rv::PknB) grown to $A_{600} \sim 0.8$ were used to seed fresh cultures. Cultures were seeded at an initial A_{600} of 0.2 in order to obtain sufficient cell pellets for WCL preparations. Cultures were grown in the absence or presence of inducer (5 μ M IVN) for 4 days. In the case of Rv-pptr-B, freshly seeded cultures (A_{600} of 0.2) were grown in the presence or absence of pristinamycin for 4 days. WCLs were prepared and subjected to Western blotting with α -PknB, α -PknA, and α -GroEL1 (loading control) antibodies. H, *in vitro* growth pattern analysis of *M. tuberculosis* H37Rv cells that are either overexpressing PknB or depleted of PknB. All of the cultures were seeded at an initial A_{600} of 0.1. Error bars, S.E.

M. tuberculosis PknB Is Essential for Pathogen Survival

formed with the various engineered PknB constructs reflected those obtained with *M. smegmatis*, thus indicating that the function of the TM domain is conserved between the avirulent and virulent strains (Fig. 5D). These data suggest that although the presence of the TM domain is essential, perhaps for target-

ing the protein to the cell membrane, specific sequences do not dictate the outcome.

All Four Extracellular PASTA Domains of PknB Are Essential in M. tuberculosis—PASTA domains have been shown to interact with penicillin and are believed to be involved in sensing the



environment (25). Further PASTA domains along with the TM domain were shown to be necessary and sufficient for appropriate localization of PknB (28). The data presented in Fig. 4 demonstrate that the extracellular PASTA domains are essential for PknB function. PknB has four extracellular PASTA domains, which have been denoted here as PASTA1 to -4 (Fig. 6A). In order to determine the minimal number of PASTA domains required, we created three constructs, wherein PknB has one (aa 1–421), two (aa 1–490), or three (aa 1–557) PASTA domains (Fig. 6A). Whereas PknB-PASTA2 and PknB-PASTA3 expression was robust, we could not detect PknB-PASTA1 expression, suggesting that the protein may be unstable (Fig. 6, B and D). Growth analysis of $mc^2\Delta pknB$ transformants unambiguously demonstrated that only PknB-PASTA3 could complement $\Delta pknB$ phenotype. PknB-PASTA1 and PknB-PASTA2 failed to rescue $mc^2\Delta pknB$ growth defects in the absence of ATc (Fig. 6C). In order to determine whether the same holds true in *M. tuberculosis*, Rv-pptR-B cells were transformed with the various constructs, and the growth patterns of transformants in liquid culture were analyzed. To our surprise, in Rv-pptR-B, none of the PASTA deletion mutants could complement the conditional mutant phenotype (Fig. 6E). The data described above clearly demonstrate that whereas the presence of the PASTA4 domain is dispensable in *M. smegmatis*, in *M. tuberculosis*, all four PASTA domains are necessary for PknB function and cell viability.

The Intracellular Domain of *M. tuberculosis* PknB Has Unique Functions—Extracellular PASTA domains have been shown to interact with muopeptides *in vitro* (5, 28). Our data establish that PASTA domains are critical to PknB function in mycobacteria. Taken together, it is possible that the extracellular PASTA domains may work akin to a sensor, which, upon interaction with muopeptides, transmits signals to the intracellular kinase domain, leading to the phosphorylation of specific substrates. We sought to address the question of specificity in such a communication between the PASTA and the kinase domains. For this, we created chimeric kinases wherein the intracellular kinase and juxtamembrane domains of other kinases were fused to the transmembrane and extracellular PASTA domains of PknB (Fig. 7A). InhA, KasA, KasB, FabH, and EmbR are phosphorylated by PknA, PknB, and PknH (32, 40–43). Because the three kinases have overlapping substrates, we selected PknA and PknH for the domain swap experiments. Both kinases were selected because one (PknA) belongs to the same clade as PknB, whereas the other (PknH) belongs to a different clade (15).

First, we sought to determine relative activities of chimeric proteins with respect to PknB. To accomplish this, MBP-tagged PknB, PknA-B, and PknH-B proteins were purified, and their ability to phosphorylate universal substrate myelin basic pro-

tein (MyBP) was determined *in vitro* at a limiting concentration of enzymes (Fig. 7, B and C). Although all three proteins were catalytically active, the quantitation of activity per μM enzyme showed that the activity of PknH-B was $\sim 34\%$ as compared with PknB. In contrast, the PknA-B chimera was found to be almost 60% active in comparison with PknB (Fig. 7C). Western blot analysis of extracts prepared from $mc^2\Delta pknB$ transformants showed that wild type PknB as well as the PknA-B and PknH-B chimeric proteins showed robust expression, PknH-B migrating more slowly as expected due to its larger size (Fig. 7D). Interestingly, $mc^2\Delta pknB$ transformants expressing PknH-B displayed growth patterns similar to $mc^2\Delta pknB$ transformants expressing full-length wild type PknB (Fig. 7E). This suggests that PknB PASTA/TM domains are able to communicate with the kinase/JM domains of PknH, and the chimeric kinase is able to phosphorylate PknB substrates, enabling cell survival in *M. smegmatis* (Fig. 7E). This result was quite interesting because the PknH-B chimera, which had substantially lower activity, showed successful complementation. We speculate that higher expression levels of PknH-B chimera probably compensated for its inefficient activity. On the other hand, the PknA-B chimeric protein, which was an efficient enzyme *in vitro*, failed to rescue $mc^2\Delta pknB$ growth defects, indicating either lack of communication between the extracellular and intracellular domains of the chimeric protein or unique substrate recognition/specificity of the PknA intracellular/JM domains (Fig. 7E). This result was somewhat surprising because PknA and PknB both belong to the same clade, and both play a role in modulation of cell shape and cell division.

A similar analysis was carried out in the Rv-pptR-B mutant, wherein transformants were streaked on plates with or without pristinamycin. However, unlike in the $mc^2\Delta pknB$ mutant, both PknH-B and PknA-B chimeric proteins failed to complement Rv-pptR-B (Fig. 7G). Further, we observed that even in the presence of pristinamycin (which induces expression of full-length PknB from the chromosomal copy), growth of Rv-pptR-B::PknA-B was compromised (Fig. 7G). Thus, even basal level of expression of PknA-B appears to be toxic to these cells. This was also supported by the observation that we obtained very few transformants, and they were smaller in size and grew much slower. These results reflect unique functions of intracellular domains of kinases in the virulent strain compared with the avirulent strain.


PknB Is Essential for Survival of Pathogen in the Host—In order to elucidate the importance of PknB in the growth and survival of *M. tuberculosis* inside the host, we have employed the murine model of experimental tuberculosis. Data in Fig. 2A demonstrated that in the absence of pristinamycin, PknB protein in Rv-pptR-B strain would be depleted in 4 days. Data in Fig. 3E suggested that PknB protein levels are restored even in the

FIGURE 2. Depletion of PknB results in cell lysis. A, in order to monitor the extent of PknB depletion over a period of 6 days, 100 ml of fresh Rv-pptR-B cultures were seeded at an initial A_{600} of 0.1 (in order to obtain sufficient quantities) and grown in the absence or presence of Pristinamycin 1A for 2, 4, and 6 days. WCLs were subjected to Western blotting with α -PknB and α -GroEL1 antibodies. B, H37Rv or Rv-pptR-B were grown to an A_{600} of ~ 0.8 in presence of inducer were washed twice in PBS containing 0.05% Tween 80 and were used to seed fresh cultures at an initial A_{600} of 0.1. Cultures were grown in the presence or absence of pristinamycin for 0, 2 or 5 days. Cultures were serially diluted and spotted on 7H10 agar plates containing pristinamycin. Drastic reduction in the titer of Rv-pptR-B grown in the absence of inducer could be observed between days 2 and 5. C, Western blot analysis to check PknB expression in *M. smegmatis* (mc^2155), merodiploid ($mc^2\text{Hit-PknB}$), and conditional depletion ($mc^2\Delta pknB$) strains. Whole cell lysates were prepared and probed with α -PknB, α -PknA, and α -GroEL1 antibodies. D, morphology of mc^2155 , $mc^2\Delta pknB$, and $mc^2::\text{PknB}$, observed by scanning electron microscopy. Cultures were grown in the absence of ATc in the case of $mc^2\Delta pknB$ and in the presence of 5 μM IVN in the case of $mc^2::\text{PknB}$ for 9 or 12 h. Scale bar, 1 μm .

M. tuberculosis PknB Is Essential for Pathogen Survival

absence of IVN when the Rv-pptr-B strain was transformed with pNit-PknB (Rv-pptr-B::PknB). To investigate the role of PknB *in vivo*, C57BL/6 mice were infected with H37Rv, Rv-pptr-B, or Rv-pptr-B::PknB strains independently through the aerosol route. The animals were euthanized after day 1 or after

4 and 8 weeks of infection, and the impact of disruption of PknB on virulence was evaluated by bacterial enumeration in lungs and spleens (Fig. 8A). Aerosol infection in mice led to implantation of ~100 bacilli in lungs at day 1 postinfection (Fig. 8B). Gross evaluation of lungs and spleen morphology 8 weeks



postinfection revealed discrete tubercles distributed throughout the tissue in the lungs of mice infected with H37Rv and Rv-pptr-B::PknB. On the other hand, markedly reduced inflammation was observed in lungs of animals infected with Rv-pptr-B strain (Fig. 8A). Moreover, the spleens of H37Rv- and Rv-pptr-B::PknB-infected mice exhibited substantial enlargement (splenomegaly) in comparison with that of Rv-pptr-B-infected mice (Fig. 8A).

At 4 weeks postinfection, the bacillary load in the lungs of mice infected with H37Rv and Rv-pptr-B::PknB was 5.12 and 4.63 log₁₀ cfu, respectively (Fig. 8B). Likewise, the bacillary load in the lungs of mice infected with H37Rv and Rv-pptr-B::PknB 8 weeks postinfection was 4.54 and 4.25 log₁₀ cfu, respectively (Fig. 8B). However, no bacilli were recovered from the lungs of mice infected with Rv-pptr-B strain at both time points, even after incubation of the plates at 37 °C for a prolonged period. Correspondingly, at 4 weeks postinfection, the splenic bacillary load of H37Rv- and Rv-pptr-B::PknB-infected mice corresponded to 5.6 × 10³ and 4.75 × 10³ cfu, respectively (data not shown). In accordance with lung data, no bacilli were recovered from the spleen of mice infected with Rv-pptr-B strain (data not shown).

We assessed the impact of disruption of PknB on the pathogenic ability of *M. tuberculosis* to establish infection and cause disease progression to pathological the state. The gross pathological changes observed were in agreement with observed pulmonary and splenic bacillary loads. At 8 weeks postinfection, the lungs of the animals infected with either H37Rv or Rv-pptr-B::PknB displayed substantial infection with numerous large tubercles and necrosis, whereas the lungs of animals infected with Rv-pptr-B exhibited significantly reduced gross pathological changes (Fig. 9A, arrows). The gross pathological scores observed for the lungs isolated from the Rv-pptr-B-infected mice were considerably lower than those seen with H37Rv- and Rv-pptr-B::PknB-infected animals (Fig. 9B). In contrast, mice infected with Rv-pptr-B displayed negligible lesions compared with those infected with the parental and complemented strains (Fig. 9, A and B). In agreement with the gross pathological scores, animals infected with the H37Rv and Rv-pptr-B::PknB exhibited enhanced histopathological damage as compared with animals infected with Rv-pptr-B. However, in the case of infection with Rv-pptr-B, lung parenchyma exhibited normal microarchitecture, with infiltration of a few leukocytes (Fig. 9C). The absence of significant lesions in the tissues of the animals infected with Rv-pptr-B demonstrate that PknB is crucial for the pathogenesis of *M. tuberculosis* in the host

tissues. Thus, the data obtained using the murine model of experimental tuberculosis clearly indicate that PknB is extremely crucial and plays an indispensable role in the pathogenesis of *M. tuberculosis*.


DISCUSSION

In this report, we describe the comprehensive evaluation of the importance of various domains of PknB in modulating its function in avirulent and virulent mycobacteria. To accomplish this objective, first, we sought to evaluate the effect of PknB overexpression and depletion on cell growth, survival, and morphology. Results presented in Fig. 2 demonstrate that both overexpression and depletion of PknB resulted in cells with varied morphology with eventual cell death. This is in line with the previous finding, wherein overexpression of PknA and PknB in *M. bovis* BCG was shown to result in a deviation from normal cell morphology (7). Although the morphology observed was not exactly the same, we observed that overexpression led to the appearance of bulging cells and cells with uneven width (Fig. 2). Results obtained upon PknB depletion are also consistent with earlier observations, wherein allelic disruption could only be achieved in the presence of a functional copy of the gene or when *pknB* was converted into an inducible gene (18, 19). Cell morphology observed upon depletion was similar to that observed consequent to overexpression of PknB (Fig. 2). Interestingly, the effect of overexpression and depletion of PknB was identical in both avirulent and virulent strains, suggesting a central role for PknB in all mycobacterial species. The lethality of PknB overexpression or depletion is probably due to the critical roles played by the kinase in modulating a number of cellular processes via its ability to phosphorylate multiple cellular substrates. Phosphorylation events do not have similar effects on the modulation of substrate enzymatic activities. For example, the activities of KasA, MabA, FabH, and InhA are negatively regulated by phosphorylation (32, 40–42, 44), whereas the activity of KasB is enhanced upon phosphorylation by the kinase (41). These results suggest that stringent regulation of PknB expression and substrate phosphorylation by PknB is necessary for maintaining cellular homeostasis and for appropriate responses to the multitude of intracellular and extracellular cues in the bacteria.


We found that although none of the deletion mutants are as efficient as full-length PknB, these proteins are proficient in phosphorylating GarA, with comparative activity ranging from 25 to 50% (Fig. 4). However, none of the PknB deletion mutants could complement the mc²Δ*pknB* or Rv-pptr-B mutants (Fig.

FIGURE 3. Kinase activity of PknB is essential for its functionality. A, schematic representation of PknB depicting the various domains and critical residues. B, *in vitro* kinase assays were carried out using 80 nM MBP or PknB or PknB-K40M, 3.33 μM GarA, 10 μM ATP, and 10 μCi [³²P]ATP. The band corresponding to the phosphorylated GarA (*p-GarA*) is indicated. C, bands corresponding to phospho-GarA were excised from the gel and quantified by liquid scintillation counting in three independent experiments. The activity was calculated as cpm in phospho-GarA/min/μM enzyme. Activity of PknB was normalized to 100% in each experiment, and the percentage of activity in other samples was calculated with respect to (*w.r.t.*) PknB activity. The results were plotted with percentage of activity on the y axis and the samples on the x axis in the form of histograms. D, mc²Δ*pknB* strain was transformed with pNit vector, pNit-PknB, or pNit-PknB-K40M constructs. Cultures grown to A₆₀₀ ~0.8 in the presence of inducer (ATc) were washed twice, and fresh cultures were seeded at an initial A₆₀₀ of 0.2. Cultures were grown in the presence or absence of ATc as indicated above for 6 h and WCLs were resolved and subjected to Western blotting with α-PknB and α-GroEL1 antibodies. E, growth pattern analysis of mc²Δ*pknB* transformants grown in the absence of ATc. F, Rv-pptr-B transformants grown to A₆₀₀ ~0.8 in the presence of inducer pristinamycin 1A were washed twice, and fresh cultures were seeded at an initial A₆₀₀ of 0.2. Cultures were grown for 4 days in the presence or absence of pristinamycin as indicated above, and WCLs were resolved from transformant strains and subjected to Western blotting with α-PknB and α-GroEL1 antibodies. G, growth pattern analysis of Rv-pptr-B transformants in the absence of pristinamycin. All of the cultures were seeded at an initial A₆₀₀ of 0.1. Error bars, S.E.


A.



B.



C.



D.




FIGURE 5. Anchoring of PknB to the cell membrane is essential for cell survival. *A*, schematic representation of PknB-TM deletion and swap mutations. *B*, *mc*²*ΔpknB* was transformed with pNit vector, pNit-PknB, pNit-PknB-(A-TM), pNit-PknB-(H-TM), or pNit-PknB- Δ TM constructs. Cultures were seeded at an initial A_{600} of 0.2 and grown in the presence or absence of ATc for 6 h. WCLs prepared were subjected to Western blotting with α -PknB and α -GroEL1 antibodies. *C*, *in vitro* growth analysis of *mc*²*ΔpknB* transformants performed in the absence of inducer ATc. Cultures were seeded at an initial A_{600} of 0.02. *D*, growth analysis of Rv-pptR-B transformants. Rv-pptR-B transformants were streaked on 7H10 agar plates in the presence or absence of 2 μ g/ml pristinamycin 1A. Error bars, S.E.

pptR-B mutants might be due to their failure to localize to the septum and poles. The fact that PknB Δ TM also failed to complement the conditional mutants supports this notion (Fig. 5). Although the TM domain *per se* is necessary for PknB function, the sequence of the TM domain does not seem to be vital

because both PknB-TM swap mutants successfully complemented mutant strains (Fig. 5).

Most of the substrates identified in *M. tuberculosis* are actually phosphorylated by multiple STPKs (45, 46), and thus a key feature among all of these kinase-substrate interactions is func-


FIGURE 4. Extracellular domain of PknB is essential for *in vitro* growth. *A*, schematic representation of PknB and deletion constructs. *B*, *in vitro* kinase assays performed using 80 nM MBP or PknB or PknB deletion mutants, 3.33 μ M GarA, 10 μ M ATP, and 10 μ Ci of [γ -³²P]ATP. Samples were resolved on 12% SDS-PAGE and autoradiographed. *C*, bands corresponding to phosphorylated GarA (*p*-GarA) were excised from the gel and quantified by liquid scintillation counting in three independent experiments. Activity of PknB was normalized to 100% in each experiment, and the percentage of activity in other samples was calculated with respect to (*w.r.t.*) PknB activity. The results were plotted with percentage of activity on the y axis and the samples on the x axis in the form of histograms. *D*, cultures were seeded at an initial A_{600} of 0.2. Cultures were grown in the presence or absence of ATc as indicated above for 6 h. For *mc*²*ΔpknB*::pNit, *mc*²*ΔpknB*::PknB-KD, *mc*²*ΔpknB*::PknB-JM, and *mc*²*ΔpknB*::PknB-TM strains, 0.2 μ M IVN inducer was added during depletion. WCLs prepared were subjected to Western blotting with α -PknB and α -GroEL1 antibodies. The arrows indicate bands corresponding to PknB deletion mutants. *E*, growth pattern analysis of *mc*²*ΔpknB* transformants grown in the absence of ATc. Cultures were seeded at an initial A_{600} of 0.02. *mc*²*ΔpknB*::PknB-KD, *mc*²*ΔpknB*::PknB-JM, and *mc*²*ΔpknB*::PknB-TM were grown in the absence of ATc and in the presence of 0.2 μ M inducer IVN. *F*, the Rv-pptR-B transformant strains were streaked on 7H10-agar plates containing 2 μ g/ml pristinamycin or 0.2 μ M IVN. Error bars, S.E.

M. tuberculosis PknB Is Essential for Pathogen Survival


culosis belong to the same genus and PknB *per se* is important in both organisms for growth and survival, the PknB kinase domain seems to have critical and non-redundant functions in the slow growing and virulent *M. tuberculosis*.

Considering the different environments the pathogen encounters, it should have the capability to adapt to a variety of different available nutrients (47, 48). The PASTA domains contained in protein kinases in various organisms are implicated to


A.




B.




C.




D.




E.



F.



G.



1. Rv-pptr-B::pNit
2. Rv-pptr-B::PknB
3. Rv-pptr-B::PknA-B
4. Rv-pptr-B::PknH-B

M. tuberculosis PknB Is Essential for Pathogen Survival




FIGURE 8. PknB is essential for *in vivo* survival. *A*, gross anatomy of lung and spleen from mice after 8 weeks of infection. *B*, C57Bl/6 mice were infected with H37Rv, Rv-pptr-B, or Rv-pptr-B::PknB as described under “Experimental Procedures.” Bacterial loads in the lungs of mice at different time points post infection were determined and represented as mean \pm S.D. (error bars) for each group ($n = 2$ for day 1 and $n = 6$ for weeks 4 and 8). The results were plotted with cfu log₁₀/lung on the y axis and the samples on the x axis in the form of a dot plot.

mediate important roles in cell division, growth, and development (49). Interactions of the extracellular PASTA domains of PrkC (a PknB ortholog in *Bacillus subtilis*) with mucopeptides have been shown to be essential for germination of spores (5). Findings by Mir *et al.* (28) suggested that exogenous mucopeptides might not be capable of penetrating through the lipid-rich mycobacterial outer membrane. In the absence of exogenous mucopeptides, the extracellular domain of PknB is likely to interact with either the peptidoglycan precursor mucopeptides or peptidoglycan hydrolysis mucopeptide products or with more complex mucopeptide ligands generated *in vivo* (28). PknB has been shown to phosphorylate PbpA (penicillin-binding protein), MurD (a ligase involved in the process of peptidoglycan synthesis), GlmU (*N*-acetylglucosamine-1-phosphate uridylyltransferase) (a key enzyme that synthesizes UDP-*N*-acetylglucosamine), and FhaA, an FHA domain-containing protein (46, 50–53). Recently, PknB-mediated phosphorylation

of the pseudokinase Mt-MviN was shown to regulate the interaction of Mt-MviN with FhaA (54). The phospho-MviN-FhaA complex was interpreted to regulate late stage peptidoglycan biosynthesis, thus establishing a regulatory feedback loop that can control peptidoglycan biosynthesis (54). In addition, Mur synthetases have been shown to interact with PknA and PknB (50). Based on our findings and previous data, we think that in addition to dictating PknB localization, the PASTA domain-ligand interaction also regulates the activity levels of PknB (23, 26).

An intriguing finding of this study is that, although PknB-PASTA3 lacking the C-terminal PASTA4 domain could rescue *mc*² Δ *pknB* mutant growth defects, it failed to complement the Rv-pptr-B mutant. Our data also showed that, in contrast to *M. smegmatis*, the PknB kinase domain has non-redundant functions in *M. tuberculosis* (Fig. 7). It is possible that the binding of mucopeptides is critical for PknB kinase domain function, and the signaling amplitude may be determined by the number of binding sites occupied at any point of time. Assuming that each PASTA domain contributes one binding site, the amplitude of signal generated by the binding of ligand to three PASTA domains may be sufficient to trigger a response in *M. smegmatis*. In *M. tuberculosis*, a response may be elicited only when all four PASTA domains are engaged. The notably distinct requirements of the number of PASTA domains and the non-redundant functions of the kinase domain in PknB-mediated signaling in *M. tuberculosis* reflect differential modulation of signaling events in virulent *versus* avirulent mycobacteria.

Our study has demonstrated that *M. tuberculosis* H37Rv or Rv-pptr-B::PknB exhibited normal growth and survival in the organs of infected mice as expected, although the growth of Rv-pptr-B::PknB was marginally lower than H37Rv at 4 and 8 weeks postinfection. On the contrary, no bacterial infection was apparent in organs harvested from the animals infected with Rv-pptr-B at both time points. These observations demonstrate for the first time that PknB is indispensable for the mycobacterium to initiate growth and to persist in the face of host innate and acquired immunity, a fact that is further corroborated by the observations that mice infected with Rv-pptr-B exhibited a significantly reduced gross pathological damage as compared with the animals infected with H37Rv or Rv-pptr-B::PknB. Thus, this study highlights the importance of PknB in the survival of *M. tuberculosis* both *in vitro* and *in vivo* and strongly suggests that PknB may be a potential target for therapeutic intervention against tuberculosis.

FIGURE 7. Intracellular domain of *M. tuberculosis* has a specific and distinct indispensable role. *A*, schematic representation of PknB chimeric proteins. *B*, MBP-tagged PknB, PknA-B, and PknH-B proteins were purified, and *in vitro* kinase assays were carried out using 25 nM MBP, PknB, PknA-B, or PknH-B; 6.25 μ M MyBP; and 10 μ Ci of [γ -³²P]ATP. *Auto-p*, autophosphorylation of respective kinases. *p-MyBP*, phosphorylated MyBP. Samples were resolved on 12% SDS-PAGE and autoradiographed. *C*, bands corresponding to phospho-MyBP were excised from the gel and quantified by liquid scintillation counting in three independent experiments. The activity was calculated as cpm of phospho-MyBP/min/ μ M enzyme. The activity of PknB was normalized to 100% in each experiment, and the percentage of activity in other samples was calculated with respect to (*w.r.t.*) PknB activity. The results were plotted with percentage of activity on the y axis and the samples on the x axis. *D*, *mc*² Δ *pknB* strain was transformed with pNit, pNit-PknB, pNit-PknA-B, or pNit-PknH-B constructs was seeded at A_{600} of 0.2 and grown in the presence or absence of ATc. WCLs were resolved and probed with α -PknB and α -GroEL1 antibodies. *E*, growth analysis of *mc*² Δ *pknB* transformants seeded at A_{600} of 0.02 in the absence of inducer ATc. *F*, Rv-pptr-B transformed with pNit, pNit-PknB, pNit-PknA-B, or pNit-PknH-B was seeded at an A_{600} of 0.2. WCLs prepared in the presence or absence of pristinamycin (as indicated) were resolved and probed with α -PknB and α -GroEL1 antibodies. *G*, Rv-pptr-B transformants were streaked on 7H10 agar plates with or without pristinamycin. Error bars, S.E.

M. tuberculosis PknB Is Essential for Pathogen Survival




FIGURE 9. PknB plays indispensable role in the pathogenesis. *A*, histopathological examination of six sections from each candidate mouse was carried out. The images shown are representative of typical lung sections in each experimental group. *B*, granuloma score of H&E-stained lung sections for each experimental group was performed as described. *C*, images depict H&E-stained lung sections of mice infected with H37Rv or Rv-pptr-B or Rv-pptr-B::PknB after 8 weeks. Images at $\times 100$ are shown in the left panels. Images of boxed regions at $\times 400$ are depicted in the corresponding right panels. G, granuloma; L, lymphocytes; FC, foamy histiocytes; AS, alveolar spaces; E, epithelioid cells.

Acknowledgments—We acknowledge Dr. Rajmani for help with animal infection experiments. We thank the Tuberculosis Aerosol Challenge Facility, International Centre for Genetic Engineering and Biotechnology (ICGEB) (New Delhi, India) for providing the facility to perform animal infection experiments. We acknowledge Dr. Amit Singh and Dr. Dhiraj Kumar (ICGEB) for scientific discussions. We thank Dr. Christopher M. Sassetti for providing pNit-1, Dr. Tanya Parish for providing p2Nil and pGOAL17 vectors, and Dr. Cole for providing the *M. tuberculosis* BAC library. We thank Rekha Rani (National Institute of Immunology) for support in obtaining scanning electron microscopy images and Aditya Yadav for generating pMAL constructs.

REFERENCES

1. Stock, J. B., Ninfa, A. J., and Stock, A. M. (1989) Protein phosphorylation and regulation of adaptive responses in bacteria. *Microbiol. Rev.* **53**, 450–490
2. Robinson, V. L., Buckler, D. R., and Stock, A. M. (2000) A tale of two components: a novel kinase and a regulatory switch. *Nat. Struct. Biol.* **7**, 626–633
3. Muñoz-Dorado, J., Inouye, S., and Inouye, M. (1991) A gene encoding a protein serine/threonine kinase is required for normal development of *M. xanthus*, a Gram-negative bacterium. *Cell* **67**, 995–1006
4. Echenique, J., Kadioglu, A., Romao, S., Andrew, P. W., and Trombe, M. C. (2004) Protein serine/threonine kinase StkP positively controls virulence and competence in *Streptococcus pneumoniae*. *Infect. Immun.* **72**,

M. tuberculosis PknB Is Essential for Pathogen Survival

- 2434–2437
- Shah, I. M., Laaberki, M. H., Popham, D. L., and Dworkin, J. (2008) A eukaryotic-like Ser/Thr kinase signals bacteria to exit dormancy in response to peptidoglycan fragments. *Cell* **135**, 486–496
 - Leiba, J., Hartmann, T., Cluzel, M. E., Cohen-Gonsaud, M., Delolme, F., Bischoff, M., and Molle, V. (2012) A novel mode of regulation of the *Staphylococcus aureus* catabolite control protein A (CcpA) mediated by Stk1 protein phosphorylation. *J. Biol. Chem.* **287**, 43607–43619
 - Kang, C. M., Abbott, D. W., Park, S. T., Dascher, C. C., Cantley, L. C., and Husson, R. N. (2005) The *Mycobacterium tuberculosis* serine/threonine kinases PknA and PknB: substrate identification and regulation of cell shape. *Genes Dev.* **19**, 1692–1704
 - Cole, S. T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S. V., Eiglmeier, K., Gas, S., Barry, C. E., 3rd, Tekai, F., Badcock, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R., Devlin, K., Feltwell, T., Gentles, S., Hamlin, N., Holroyd, S., Hornsby, T., Jagels, K., Krogh, A., McLean, A., Moule, S., Murphy, L., Oliver, K., Osborne, J., Quail, M. A., Rajandream, M. A., Rogers, J., Rutter, S., Seeger, K., Skelton, J., Squares, R., Squares, S., Sulston, J. E., Taylor, K., Whitehead, S., and Barrell, B. G. (1998) Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* **393**, 537–544
 - Av-Gay, Y., and Everett, M. (2000) The eukaryotic-like Ser/Thr protein kinases of *Mycobacterium tuberculosis*. *Trends Microbiol.* **8**, 238–244
 - Hanks, S. K., and Hunter, T. (1995) Protein kinases 6: the eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. *FASEB J.* **9**, 576–596
 - Cowley, S., Ko, M., Pick, N., Chow, R., Downing, K. J., Gordhan, B. G., Betts, J. C., Mizrahi, V., Smith, D. A., Stokes, R. W., and Av-Gay, Y. (2004) The *Mycobacterium tuberculosis* protein serine/threonine kinase PknG is linked to cellular glutamate/glutamine levels and is important for growth *in vivo*. *Mol. Microbiol.* **52**, 1691–1702
 - Walburger, A., Koul, A., Ferrari, G., Nguyen, L., Prescianotto-Baschong, C., Huygen, K., Klebl, B., Thompson, C., Bacher, G., and Pieters, J. (2004) Protein kinase G from pathogenic mycobacteria promotes survival within macrophages. *Science* **304**, 1800–1804
 - Deol, P., Vohra, R., Saini, A. K., Singh, A., Chandra, H., Chopra, P., Das, T. K., Tyagi, A. K., and Singh, Y. (2005) Role of *Mycobacterium tuberculosis* Ser/Thr kinase PknF: implications in glucose transport and cell division. *J. Bacteriol.* **187**, 3415–3420
 - Ortega, C., Liao, R., Anderson, L. N., Rustad, T., Ollodart, A. R., Wright, A. T., Sherman, D. R., and Grundner, C. (2014) *Mycobacterium tuberculosis* Ser/Thr protein kinase B mediates an oxygen-dependent replication switch. *PLoS Biol.* **12**, e1001746
 - Narayan, A., Sachdeva, P., Sharma, K., Saini, A. K., Tyagi, A. K., and Singh, Y. (2007) Serine threonine protein kinases of mycobacterial genus: phylogeny to function. *Physiol. Genomics* **29**, 66–75
 - Chakraborti, P. K., Matange, N., Nandicoori, V. K., Singh, Y., Tyagi, J. S., and Visweswariah, S. S. (2011) Signalling mechanisms in Mycobacteria. *Tuberculosis* **91**, 432–440
 - Sasseti, C. M., Boyd, D. H., and Rubin, E. J. (2003) Genes required for mycobacterial growth defined by high density mutagenesis. *Mol. Microbiol.* **48**, 77–84
 - Fernandez, P., Saint-Joanis, B., Barilone, N., Jackson, M., Gicquel, B., Cole, S. T., and Alzari, P. M. (2006) The Ser/Thr protein kinase PknB is essential for sustaining mycobacterial growth. *J. Bacteriol.* **188**, 7778–7784
 - Forti, F., Crosta, A., and Ghisotti, D. (2009) Pristinamycin-inducible gene regulation in mycobacteria. *J. Biotechnol.* **140**, 270–277
 - Durán, R., Villarino, A., Bellinzoni, M., Wehenkel, A., Fernandez, P., Boitel, B., Cole, S. T., Alzari, P. M., and Cerveñansky, C. (2005) Conserved autophosphorylation pattern in activation loops and juxtamembrane regions of *Mycobacterium tuberculosis* Ser/Thr protein kinases. *Biochem. Biophys. Res. Commun.* **333**, 858–867
 - Ortiz-Lombardía, M., Pompeo, F., Boitel, B., and Alzari, P. M. (2003) Crystal structure of the catalytic domain of the PknB serine/threonine kinase from *Mycobacterium tuberculosis*. *J. Biol. Chem.* **278**, 13094–13100
 - Young, T. A., Delagoutte, B., Endrizzi, J. A., Falick, A. M., and Alber, T. (2003) Structure of *Mycobacterium tuberculosis* PknB supports a universal activation mechanism for Ser/Thr protein kinases. *Nat. Struct. Biol.* **10**, 168–174
 - Lombana, T. N., Echols, N., Good, M. C., Thomsen, N. D., Ng, H. L., Greenstein, A. E., Falick, A. M., King, D. S., and Alber, T. (2010) Allosteric activation mechanism of the *Mycobacterium tuberculosis* receptor Ser/Thr protein kinase, PknB. *Structure* **18**, 1667–1677
 - Mieczkowski, C., Iavarone, A. T., and Alber, T. (2008) Auto-activation mechanism of the *Mycobacterium tuberculosis* PknB receptor Ser/Thr kinase. *EMBO J.* **27**, 3186–3197
 - Yeats, C., Finn, R. D., and Bateman, A. (2002) The PASTA domain: a β -lactam-binding domain. *Trends Biochem. Sci.* **27**, 438
 - Barthe, P., Mukamolova, G. V., Roumestand, C., and Cohen-Gonsaud, M. (2010) The structure of PknB extracellular PASTA domain from mycobacterium tuberculosis suggests a ligand-dependent kinase activation. *Structure* **18**, 606–615
 - Dessen, A., Mouz, N., Gordon, E., Hopkins, J., and Dideberg, O. (2001) Crystal structure of PBP2x from a highly penicillin-resistant *Streptococcus pneumoniae* clinical isolate: a mosaic framework containing 83 mutations. *J. Biol. Chem.* **276**, 45106–45112
 - Mir, M., Asong, J., Li, X., Cardot, J., Boons, G. J., and Husson, R. N. (2011) The extracytoplasmic domain of the *Mycobacterium tuberculosis* Ser/Thr kinase PknB binds specific muopeptides and is required for PknB localization. *PLoS Pathog.* **7**, e1002182
 - Pandey, A. K., Raman, S., Proff, R., Joshi, S., Kang, C. M., Rubin, E. J., Husson, R. N., and Sasseti, C. M. (2009) Nitrile-inducible gene expression in mycobacteria. *Tuberculosis* **89**, 12–16
 - Parikh, A., Kumar, D., Chawla, Y., Kurthkoti, K., Khan, S., Varshney, U., and Nandicoori, V. K. (2013) Development of a new generation of vectors for gene expression, gene replacement, and protein-protein interaction studies in mycobacteria. *Appl. Environ. Microbiol.* **79**, 1718–1729
 - Parish, T., and Stoker, N. G. (2000) Use of a flexible cassette method to generate a double unmarked *Mycobacterium tuberculosis* tlyA plcABC mutant by gene replacement. *Microbiology* **146**, 1969–1975
 - Khan, S., Nagarajan, S. N., Parikh, A., Samantaray, S., Singh, A., Kumar, D., Roy, R. P., Bhatt, A., and Nandicoori, V. K. (2010) Phosphorylation of enoyl-acyl carrier protein reductase InhA impacts mycobacterial growth and survival. *J. Biol. Chem.* **285**, 37860–37871
 - Tiwari, D., Singh, R. K., Goswami, K., Verma, S. K., Prakash, B., and Nandicoori, V. K. (2009) Key residues in *Mycobacterium tuberculosis* protein kinase G play a role in regulating kinase activity and survival in the host. *J. Biol. Chem.* **284**, 27467–27479
 - Villarino, A., Duran, R., Wehenkel, A., Fernandez, P., England, P., Brodin, P., Cole, S. T., Zimny-Arndt, U., Jungblut, P. R., Cerveñansky, C., and Alzari, P. M. (2005) Proteomic identification of *M. tuberculosis* protein kinase substrates: PknB recruits GarA, a FHA domain-containing protein, through activation loop-mediated interactions. *J. Mol. Biol.* **350**, 953–963
 - O'Hare, H. M., Durán, R., Cerveñansky, C., Bellinzoni, M., Wehenkel, A. M., Pritsch, O., Obal, G., Baumgartner, J., Vialaret, J., Johnsson, K., and Alzari, P. M. (2008) Regulation of glutamate metabolism by protein kinases in mycobacteria. *Mol. Microbiol.* **70**, 1408–1423
 - Carrera, A. C., Alexandrov, K., and Roberts, T. M. (1993) The conserved lysine of the catalytic domain of protein kinases is actively involved in the phosphotransfer reaction and not required for anchoring ATP. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 442–446
 - Spitaler, M., Villunger, A., Grunicke, H., and Uberall, F. (2000) Unique structural and functional properties of the ATP-binding domain of atypical protein kinase C- ι . *J. Biol. Chem.* **275**, 33289–33296
 - Av-Gay, Y., Jamil, S., and Drews, S. J. (1999) Expression and characterization of the *Mycobacterium tuberculosis* serine/threonine protein kinase PknB. *Infect. Immun.* **67**, 5676–5682
 - Boitel, B., Ortiz-Lombardía, M., Durán, R., Pompeo, F., Cole, S. T., Cerveñansky, C., and Alzari, P. M. (2003) PknB kinase activity is regulated by phosphorylation in two Thr residues and dephosphorylation by PstP, the cognate phospho-Ser/Thr phosphatase, in *Mycobacterium tuberculosis*. *Mol. Microbiol.* **49**, 1493–1508
 - Molle, V., Gulten, G., Vilchère, C., Veyron-Churlet, R., Zanella-Cléon, I., Sacchettini, J. C., Jacobs, W. R., Jr., and Kremer, L. (2010) Phosphorylation of InhA inhibits mycolic acid biosynthesis and growth of *Mycobacterium tuberculosis*. *Mol. Microbiol.* **78**, 1591–1605

41. Molle, V., Brown, A. K., Besra, G. S., Cozzone, A. J., and Kremer, L. (2006) The condensing activities of the *Mycobacterium tuberculosis* type II fatty acid synthase are differentially regulated by phosphorylation. *J. Biol. Chem.* **281**, 30094–30103
42. Veyron-Churlet, R., Molle, V., Taylor, R. C., Brown, A. K., Besra, G. S., Zanella-Cléon, I., Fütterer, K., and Kremer, L. (2009) The *Mycobacterium tuberculosis* β -ketoacyl-acyl carrier protein synthase III activity is inhibited by phosphorylation on a single threonine residue. *J. Biol. Chem.* **284**, 6414–6424
43. Sharma, K., Gupta, M., Pathak, M., Gupta, N., Koul, A., Sarangi, S., Baweja, R., and Singh, Y. (2006) Transcriptional control of the mycobacterial embCAB operon by PknH through a regulatory protein, EmbR, *in vivo*. *J. Bacteriol.* **188**, 2936–2944
44. Veyron-Churlet, R., Zanella-Cléon, I., Cohen-Gonsaud, M., Molle, V., and Kremer, L. (2010) Phosphorylation of the *Mycobacterium tuberculosis* β -ketoacyl-acyl carrier protein reductase MabA regulates mycolic acid biosynthesis. *J. Biol. Chem.* **285**, 12714–12725
45. Priscic, S., Dankwa, S., Schwartz, D., Chou, M. F., Locasale, J. W., Kang, C. M., Bemis, G., Church, G. M., Steen, H., and Husson, R. N. (2010) Extensive phosphorylation with overlapping specificity by *Mycobacterium tuberculosis* serine/threonine protein kinases. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 7521–7526
46. Grundner, C., Gay, L. M., and Alber, T. (2005) *Mycobacterium tuberculosis* serine/threonine kinases PknB, PknD, PknE, and PknF phosphorylate multiple FHA domains. *Protein Sci.* **14**, 1918–1921
47. Jayachandran, R., BoseDasgupta, S., and Pieters, J. (2013) Surviving the macrophage: tools and tricks employed by *Mycobacterium tuberculosis*. *Curr. Top. Microbiol. Immunol.* **374**, 189–209
48. Hett, E. C., and Rubin, E. J. (2008) Bacterial growth and cell division: a mycobacterial perspective. *Microbiol. Mol. Biol. Rev.* **72**, 126–156
49. Ruggiero, A., De Simone, P., Smaldone, G., Squeglia, F., and Berisio, R. (2012) Bacterial cell division regulation by Ser/Thr kinases: a structural perspective. *Curr. Protein Pept. Sci.* **13**, 756–766
50. Munshi, T., Gupta, A., Evangelopoulos, D., Guzman, J. D., Gibbons, S., Keep, N. H., and Bhakta, S. (2013) Characterisation of ATP-dependent Mur ligases involved in the biogenesis of cell wall peptidoglycan in *Mycobacterium tuberculosis*. *PLoS One* **8**, e60143
51. Parikh, A., Verma, S. K., Khan, S., Prakash, B., and Nandicoori, V. K. (2009) PknB-mediated phosphorylation of a novel substrate, *N*-acetylglucosamine-1-phosphate uridylyltransferase, modulates its acetyltransferase activity. *J. Mol. Biol.* **386**, 451–464
52. Dasgupta, A., Datta, P., Kundu, M., and Basu, J. (2006) The serine/threonine kinase PknB of *Mycobacterium tuberculosis* phosphorylates PBPA, a penicillin-binding protein required for cell division. *Microbiology* **152**, 493–504
53. Roumestand, C., Leiba, J., Galophe, N., Margeat, E., Padilla, A., Bessin, Y., Barthe, P., Molle, V., and Cohen-Gonsaud, M. (2011) Structural insight into the *Mycobacterium tuberculosis* Rv0020c protein and its interaction with the PknB kinase. *Structure* **19**, 1525–1534
54. Gee, C. L., Papavinasundaram, K. G., Blair, S. R., Baer, C. E., Falick, A. M., King, D. S., Griffin, J. E., Venghatakrishnan, H., Zukauskas, A., Wei, J. R., Dhiman, R. K., Crick, D. C., Rubin, E. J., Sasseti, C. M., and Alber, T. (2012) A phosphorylated pseudokinase complex controls cell wall synthesis in mycobacteria. *Sci. Signal.* **5**, ra7