

Comparative Analysis of Pathogenic Organisms in Cockroaches from Different Community Settings in Edo State, Nigeria

Clement Isaac^{1,*}, Philip Ogbeide Orue², Mercy Itohan Iyamu², Joy Imuetiyan Ehiaghe³, Osesojie Isaac²

¹Department of Zoology, ²Department of Microbiology, Ambrose Alli University, Ekpoma, Nigeria; ³Lahor Laboratory, Benin, Nigeria

Abstract: Cockroaches are abundant in Nigeria and are seen to harbour an array of pathogens. Environmental and sanitary conditions associated with demographic/socio-economic settings of an area could contribute to the prevalence of disease pathogens in cockroaches. A total of 246 cockroaches (*Periplaneta americana*) in urban (Benin, n=91), semi-urban (Ekpoma, n=75) and rural (Emuhi, n=70) settings in Edo State, Nigeria were collected within and around households. The external body surfaces and alimentary canal of these cockroaches were screened for bacterial, fungal, and parasitological infections. *Bacillus* sp. and *Escherichia coli* were the most common bacteria in cockroaches. However, *Enterococcus faecalis* could not be isolated in cockroaches trapped from Ekpoma and Emuhi. *Aspergillus niger* was the most prevalent fungus in Benin and Ekpoma, while *Mucor* sp. was predominant in Emuhi. Parasitological investigations revealed the preponderance of *Ascaris lumbricoides* in Benin and Emuhi, while *Trichuris trichura* was the most predominant in Ekpoma. The prevalence and burden of infection in cockroaches is likely to be a reflection of the sanitary conditions of these areas. Also, cockroaches in these areas making incursions in homes may increase the risk of human infections with these disease agents.

Key words: cockroach, pathogen, sanitary condition, Nigeria

INTRODUCTION

Cockroaches such as *Periplaneta americana* and *Blattella germanica* have filthy habits with an ability to spoil food, transfer pathogens, and cause allergic reactions and psychological distress [1]. Typical of an urban settlement in most developing countries is overcrowding, pollution, and lack of basic services such as clean water and sanitation. These conditions could enhance the infestation rate of cockroaches to a number of pathogens [2].

Over 100 species of bacteria have been isolated from domestic cockroaches [3] of which some could be potential mechanical transmitters of pathogenic bacteria [4]. A study in Osogbo, urban Nigeria, and Ekpoma, a semi-urban area revealed an array of bacteria harboured by cockroaches [5,6]. Similarly, fungal infections have been associated with both the external

body parts and feces of cockroaches [5,6]. Protozoan parasites and gastro-intestinal helminths have also been reported in cockroaches collected from both suburban and peri-urban areas [5,7] being widely implicated as reservoir hosts of medically important parasites [8,9].

The distribution of bacteria, fungi, and parasites in cockroaches could differ from one location to the other based on both environmental and sanitary conditions which could be used to assess the likely infectious diseases that pose great risk as we consider the capacity of cockroaches in spreading diseases in a complementary manner. Hence, this paper reports a comparative analysis of the prevalence of pathogenic organisms present in cockroaches sampled from urban, semi-urban, and rural areas of Edo State, Nigeria.

MATERIALS AND METHODS

Study Area

This study was carried out in urban (Benin), Semi-urban (Ekpoma), and rural (Emuhi) areas in Edo State, Nigeria between February and July 2013. The capital of the state is Benin. Benin (61°9'N, 5°36'E) is a city approximately 25 miles north

•Received 6 September 2013, revised 21 October 2013, accepted 31 October 2013.

*Corresponding author (cle21200@yahoo.co.uk)

© 2014, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

of the Benin River, and it is the center of Nigeria's rubber industry, with a traditional industry of palm nuts processing for oil production. Ekpoma (64°5'N, 60°8'E), a semi-urban settlement, is the headquarters of the Esan West Local Government Area and a host to a tertiary institution (Ambrose Alli University). It is mostly populated by the Esan people and students from various parts of Nigeria. Sanitary conditions in this area are almost underdeveloped. Fecal disposal and water supply are very elementary in the sense that the majority of the population relies on pit toilets for fecal disposal and hand-dug wells for water storage. Emuhi (60°40'N 60°10'E), a rural community [10] is also under the administration of Esan West Local Government Area. The community is popularly reputed in the region for the production of pineapple fruit. Fecal and waste disposal are done in crude manners due to the lack of modern toilet and waste disposal facilities.

Sample collection and analysis

A total of 246 adult cockroaches (*P. americana*) were trapped from different homes in Benin (n=91), Ekpoma (n=85), and Emuhi (n=70) for bacterial, fungal, and parasitological studies. Cockroaches were collected in different areas of individual homes using sweep net and insecticide spray from the parlours, bedrooms, verandas, kitchens, toilets, and bathrooms, and then stored in sterile universal containers and transported to the laboratories.

Pathogens on the external body of cockroaches were isolated by filling the containers with 5 ml of sterile normal saline and then shaken vigorously. Afterwards, a loopful of each suspension was cultured for bacteriological and fungal investigations. Furthermore, the alimentary canal of each cockroach was emptied in a different container of 5 ml of saline for an-

other round of bacteriological and fungal identification. Plates with blood and McConkey agar were used for bacteria culture, while Sabouraud and potato dextrose agar were used for fungi culture. These plates were incubated at 37°C for 24 hr or more. The organisms were identified using standard bacteriological and fungal techniques as described by Mackie and McCartney [11]. A 10-fold dilution was carried out on each suspension to determine the total viable counts per millilitre on each cockroach using the pour-plate method. These counts were made on plates showing discrete and evenly distributed colonies.

For parasitological studies, samples were transferred to a conical test tube and spun at 3,000 rpm for 5 min. The deposits were observed on clean glass slides and viewed under a microscope. The eggs of the parasites were identified and the number counted per ml. Results were entered in excel spreadsheet, and InStat statistical package was used to analyze the data.

RESULTS

Table 1 shows the different pathogens isolated from both the external body surface and alimentary canals of cockroaches. A total of 246 cockroaches were examined, and it was demonstrated that more microbes and parasites were collected in the alimentary canal than the body surface of cockroaches.

A comparison of the distribution of bacteria in cockroaches collected from urban, semi-urban, and rural communities are shown in Table 2. In Benin, 10 species of bacteria were isolated, whereas fewer numbers were identified in Ekpoma (7 species) and Emuhi (8 species). Clearly, *Bacillus* sp. was the most prevalent bacterial infection with Emuhi community showing the highest infection rate in cockroaches (OR, 8.23; $P < 0.05$).

Table 1. Pathogens isolated from the body surface and internal organs of cockroaches (n=246)

Bacteria		Fungus		Parasite (eggs)	
Body surface	Alimentary canal	Body surface	Alimentary canal	Body surface	Alimentary canal
<i>Bacillus</i> sp.	<i>Bacillus</i> sp.	<i>Aspergillus niger</i>	<i>Aspergillus niger</i>	<i>Ascaris lumbricoides</i>	<i>A. lumbricoides</i>
<i>Escherichia coli</i>	<i>E. coli</i>	-	<i>Candida</i> sp.	<i>Trichuris trichiura</i>	<i>T. trichiura</i>
<i>Proteus mirabilis</i>	<i>P. mirabilis</i>	<i>Mucor</i> sp.	<i>Mucor</i> sp.	-	<i>Enterobius vermicularis</i>
-	<i>Pseudomonas aeruginosa</i>	-	<i>Rhizopus</i>	-	<i>Schistosoma haematobium</i>
-	<i>Proteus vulgaris</i>	<i>Saccharomyces cerevisiae</i>	<i>S. cerevisiae</i>	<i>Coccidia</i>	-
<i>Staphylococcus aureus</i>	<i>Staphylococcus aureus</i>	-	<i>Fusarium</i> sp.	-	<i>Balantidium coli</i>
-	<i>Staphylococcus epidermis</i>	-	<i>Penicillium</i> sp.	<i>Entamoeba histolytica</i>	<i>E. histolytica</i>
-	<i>Enterococcus faecalis</i>				
<i>Citrobacter freundii</i>	-				
	<i>Salmonella</i> sp.				

Table 2. Prevalence and bacterial load in cockroaches from urban, semi-urban, and rural communities

Bacteria	Urban (Benin)		Semi-urban (Ekpoma)		Rural (Emuhi)	
	N (%)	Load (organisms/ml)	N (%)	Load (organisms/ml)	N (%)	Load (organisms/ml)
<i>Bacillus</i> sp.	47 (51.6)	8.0 x 10 ⁹	38 (41.2)	9.0 x 10 ⁹	56 (80.0)	3.1 x 10 ⁹
<i>Escherichia coli</i>	7 (7.7)	10 x 10 ⁹	13 (15.3)	3.3 x 10 ⁷	12 (17.1)	5.2 x 10 ⁹
<i>Proteus mirabilis</i>	10 (1.1)	4.0 x 10 ⁹	4 (4.7)	5.2 x 10 ⁹	2 (2.9)	3.8 x 10 ⁹
<i>Pseudomonas aeruginosa</i>	7 (7.7)	6.0 x 10 ⁹	4 (4.7)	3.4 x 10 ⁹	7 (10.0)	3.5 x 10 ⁹
<i>Proteus vulgaris</i>	1 (1.1)	11 x 10 ⁹	1 (1.2)	2.4 x 10 ⁶	-	-
<i>Staphylococcus aureus</i>	7 (7.7)	4.0 x 10 ⁹	4 (4.7)	5.3 x 10 ⁹	10 (14.3)	3.9 x 10 ⁸
<i>Staphylococcus epidermidis</i>	2 (2.2)	3.4 x 10 ⁹	-	-	1 (1.4)	3.0 x 10 ⁹
<i>Enterococcus faecalis</i>	2 (2.2)	5.0 x 10 ⁸	-	-	-	-
<i>Citrobacter freundii</i>	2 (2.2)	6.0 x 10 ⁹	2 (2.4)	3.4 x 10 ⁹	2 (2.9)	4.4 x 10 ⁵
<i>Salmonella</i> sp.	2 (2.2)	3.0 x 10 ⁹	-	-	1 (1.4)	4.9 x 10 ⁵

Table 3. Prevalence and fungal load in cockroaches from urban, semi-urban, and rural communities

Fungi	Urban (Benin)		Semi-urban (Ekpoma)		Rural (Emuhi)	
	N (%)	Load (fungi/ml)	N (%)	Load (fungi/ml)	N (%)	Load (fungi/ml)
<i>Aspergillus niger</i>	30 (33.0)	5.2 x 10 ⁹	21 (24.7)	4.3 x 10 ⁹	4 (5.7)	7.8 x 10 ⁹
<i>Candida</i> sp.	5 (5.5)	3.0 x 10 ⁷	1 (1.2)	7.0 x 10 ⁹	1 (1.4)	9.7 x 10 ⁸
<i>Mucor</i> sp.	17 (18.7)	5.7 x 10 ⁸	14 (16.5)	9.1 x 10 ⁹	12 (17.1)	4.3 x 10 ⁹
<i>Rhizopus</i> sp.	5 (5.5)	3.9 x 10 ⁹	-	-	-	-
<i>Saccharomyces cerevisiae</i>	5 (5.5)	8.1 x 10 ⁶	1 (1.2)	2.0 x 10 ⁶	7 (10.0)	6.6 x 10 ⁸
<i>Fusarium</i> sp.	3 (3.3)	4.0 x 10 ⁹	3 (3.5)	5.2 x 10 ⁹	2 (2.9)	4.5 x 10 ⁹
<i>Penicillium</i> sp.	2 (2.2)	3.1 x 10 ⁸	5 (5.9)	8.2 x 10 ⁸	5 (7.1)	5.7 x 10 ⁹

Table 4. Prevalence and parasitic load in cockroaches from urban, semi-urban, and rural communities

Parasites	Urban (Benin)		Semi-urban (Ekpoma)		Rural (Emuhi)	
	N (%)	Load (eggs/ml)	N (%)	Load (eggs/ml)	N (%)	Load (eggs/ml)
<i>Ascaris lumbricoides</i>	12 (13.2)	17	3 (3.5)	20	2 (2.9)	5
<i>Trichuris trichiura</i>	4 (4.4)	39	4 (4.7)	18	-	-
<i>Enterobius vermicularis</i>	3 (3.3)	30	1 (1.2)	4	1 (1.4)	6
<i>Schistosoma haematobium</i>	3 (3.3)	4	1 (1.2)	6	-	-
Coccidia	3 (3.3)	2	-	-	-	-
<i>Balantidium coli</i>	1 (1.1)	15	1 (1.2)	44	-	-
<i>Entamoeba histolytica</i>	2 (2.2)	4	1 (1.2)	3	-	-

Similarly, our data shows that *Escherichia coli* was the second most dominant bacterial isolate, with Emuhi recording the highest infection rate. *Enterococcus faecalis* could not be isolated in cockroaches from Ekpoma and Emuhi.

Table 3 summarizes a comparative data of fungal infections in cockroaches. Seven fungi were isolated in Benin, while 6 each for Ekpoma and Emuhi. Comparatively, *Aspergillus niger* had the highest prevalence in Benin and Ekpoma than other fungi except for the preponderance of *Mucor* sp. in cockroaches collected from Emuhi. *Rhizopus* sp. were seen in cockroaches from Benin only.

Parasites were isolated from cockroaches in urban, semi-urban, and rural settlements as presented in Table 4. Our record

shows that *Ascaris lumbricoides* was predominant in the 3 areas and particularly highest in cockroaches from Benin (OR= 3.12; $P < 0.05$). In addition, *Trichuris trichiura* was the most frequently occurring parasites in Ekpoma. We did not isolate *A. lumbricoides* and *Enterobius vermicularis* eggs in Emuhi, whereas coccidian parasites were recovered from Benin only.

DISCUSSION

This study isolated 10 species of bacteria from Benin; seven and 8 species from Ekpoma and Emuhi respectively. There was the predominance of *Bacillus* sp. and *E. coli* in the cockroaches collected from the 3 settlements. Also, the prevalence of *Staph-*

Staphylococcus aureus was higher in Emuhi than in the other areas. Additionally, *E. faecalis* was not isolated in cockroaches trapped from Ekpoma and Emuhi but was seen in Benin. The distribution of bacterial organisms in Ekpoma was not so different from the report of Tاتفeng et al. [5] except that, in this work, *Bacillus* sp. was the most frequently seen microorganism in cockroaches which was not the case in the previous investigation from Ekpoma. *Bacillus* sp. and *E. coli* being harboured by cockroaches can cause diarrhea [12]. *S. aureus* causes nosocomial infections, and cockroaches have been implicated as carriers [13,14]. Generally, spread of *S. aureus* is through human-to-human contact, although recently some veterinarians discovered that the infection can spread through pets [15]. The presence of *E. faecalis* in Benin aligns with the assertion that this organism survives highly unfriendly environmental conditions which characterizes some urban cities in Nigeria [16].

A total of 7 fungal species were isolated from cockroaches in Benin, while 6 species were present in cockroaches collected in Ekpoma and Emuhi. The most prevalent species in the 3 areas were *A. niger* and *Mucor* sp. *A. niger* causes a disease called black mold on certain fruits and vegetables, and it is ubiquitous in soil. Being regarded as an opportunistic pathogen [17], *A. niger* has been associated with otomycosis infection among Nigerians [18]. *Mucor* is a microbial genus of approximately 3,000 species of moulds commonly found in soil, digestive systems, plant surfaces, and rotten vegetable matter. However, some *Mucor* sp. have been reported as infectious agents, although their inability to grow at temperatures above 32°C raises doubt as to their validity as human pathogens, and their pathogenic role may be limited to cutaneous infections [19,20]. *Rhizopus* organism was not seen in Ekpoma and Emuhi but present in Benin. This could be attributed to the number of cockroaches analyzed against previous investigation of relatively larger sample size in Ekpoma where this fungus was isolated in cockroaches [5]. *Rhizopus* spp. are found on a wide variety of organic substrates while some are opportunistic agents of human zygomyces.

More parasites were recovered in Benin in comparison with the other communities. Also, we observed that *A. lumbricoides* was the most occurring parasite across the sampled areas; and this is similar to the observation of Etim et al. [7]. This also corroborates previous reports regarding the epidemiology of intestinal helminths in Edo State where *A. lumbricoides* was recorded to be the most prevalent intestinal parasite across age groups [21,22]. *Schistosoma haematobium* ova were identified in

a few cockroaches. We strongly suspect that some dwellers may be infected with this parasite which might have been picked up by these cockroaches probably in the toilet or bathroom. A study in Sokoto, Nigeria has also isolated *S. haematobium* eggs from cockroaches [23]. Furthermore, coccidian parasites were isolated in Benin only, an indication of cockroaches having contact with fecal droppings of birds, dogs, and/or cats. Poor fecal and garbage disposal systems observed in Benin City may have contributed in no small way to the parasitic contamination of these cockroaches.

The urban area of Benin had the highest burden of disease agents in cockroaches. This is quite instructive in the sense that it simply shows the sanitary situation of this area. Interestingly, this is contrary to our expectation considering the circumstances that rural dwellers are subjected to lack/inadequate basic infrastructural facilities which include waste disposal facilities. Consequently, it is assumed that these rural dwellers exhibit relatively poorer sanitary habits that predispose the environment to a higher proliferation of disease agents. Additionally, rural areas are being adjudged to be least developed socio-economically, with its attendant challenges of non-provision of waste disposal facilities, resulting in the habit of indiscriminately disposing of waste materials leading inevitably to generation of pockets of litter-spots that obviously constitute breeding sites for most disease agents; and when cockroaches visit these sites, they pick up all kinds of pathogens. However, the results we have for Benin is strongly adduced to some urban challenges other authors have considered as risk factors to disease outbreaks [2,24]. Moreover, we observed that in Benin City, there is inadequate waste disposal facility and agency, and therefore wastes are allowed to remain for a long time at collection points before they are disposed of. We observed therefore that between the period when the waste bins are filled to capacity and the time of disposal, cockroaches feast within these exposed bins and pick up all manners of pathogens that might contaminate food materials, kitchen utensils and other household items through its activities of fecal droppings [25] or mere body contacts with these items, thereby posing great health risks to dwellers.

In conclusion, cockroaches in Edo State could be mechanical transmitters of medically important pathogens, and inhabitants of urban centres are at the greatest risk of infection. Our results also indicate that the burden of disease pathogens in cockroaches could be related to both environmental conditions and sanitary habits of dwellers. Since cockroaches col-

lected from a representation of urban, semi-urban, and rural communities had issues with disease pathogen infestation, it is strongly recommended that control measures should be greatly improved especially by driving adequate and effective institution of waste management systems in Edo State.

CONFLICT OF INTEREST

We declare that we have no conflict of interest related to this work.

REFERENCES

- Brenner RJ. Economics and medical importance of German cockroaches, In Rust MK, Owens JM, Reiersen DA eds, Understanding and Controlling the German Cockroach. New York, NY, USA. Oxford University Press. 1995, p 77-92.
- Feizhaddad MH, Kassiri H, Sepand M, Ghasemi, F. Bacteriological survey of American cockroaches in hospitals. Middle-East J Sc Res 2012; 12: 985-989.
- Cruden DL, Markovetz, AJ. Microbiological study of domestic cockroaches in microbial ecology of the cockroach gut. Annual review of human dwelling localities. Afr J Microbiol Res Microbiol 1987; 41: 617-643.
- Pai HH. Multidrug resistant bacteria isolated from cockroaches in long-term care facilities and nursing homes. Acta Trop 2013; 125: 18-22.
- Tatfeng YM, Usuanlele MU, Orukpe, A, Digban, AK, Okodua M, Oviasogie F, Turay AA. Mechanical transmission of pathogenic organisms: the role of cockroaches. J Vect Borne Dis 2005; 42: 129-134.
- Adeleke MA, Akatah HA, Hassan AWO, Adebimp WO. Microbial load and multiple drug resistance of pathogenic bacteria isolated from faeces and body surfaces in an urban area of southwestern. J Microbiol Biotech Food Sc 2012; 1: 1448-1461.
- Etim SEI, Okon OE, Akpan PA, Ukpung GI, Oku EE. Prevalence of cockroaches (*Periplaneta americana*) in households in Calabar: public health implications. J Pub Health Epidemiol 2013; 5: 149-152.
- Thyseen PJ, Moretti T, Dic Ueta MT, Ribeiro OB. The role of insects (Blattodea, Diptera and Hymoptera) as possible mechanical vectors of helminths in the domiciliary and pre-domiciliary environment. Cad Saude Publica 2004; 20: 1096-1102.
- Chan OT, Lee TK, Hardman JM, Navin JJ. The cockroach as a host for *Trichinella and Enterobius vermicularis*: implications for public health. Hawaii Med J 2004; 63: 74-77.
- Ehisuoria SE. The Role of Non-agro Based Industries in Rural development in Esan land, Edo State, Nigeria. Unpublished Ph.D. Thesis. Department of Geography and Regional Planning, Ambrose Alli University, Ekpoma, Nigeria. 2012.
- Mackie TJ, McCartney JE. Laboratory strategies in diagnosis. In Colle JG, Duguid JP, Fraser AG, Marmion BP, eds, Practical Medical Microbiology, XIIIth ed. London, UK. Churchill Livingstone. 1989; p 590-624.
- Bouamama L, Sorlozano A, Laglaoui A, Lebbadi M, Aarab A, Gutierrez J. Antibiotic resistance patterns of bacterial strains isolated from *Periplaneta americana* and *Musca domestica* in Tangier, Morocco. J Infect Dev Ctries 2010; 4: 194-201.
- Akindele AA, Adewuyi IK, Adefioye, OA, Kassiri H, Kazemi SA, Adedokun SA, Olaolu AO. Antibiogram and beta-lactamase production of Dictyoptera (Blattidae) as carriers of bacterial *Staphylococcus aureus* isolates from different human pathogens in Ile-ife, Nigeria. American-Eurasian J Sci Res 2012; 5: 230-233.
- Tachbele, E, Erku W, Gebre-Michael T, Ashenafi M. Cockroach associated food-borne bacterial pathogens from some hospitals and restaurants in Addis Ababa, Ethiopia: distribution and antibiograms. J Rural Trop Public Health 2006; 5: 34-41.
- Cook H, Furuya E, Larson E, Vasquez G, Lowy F. Heterosexual transmission of community-associated methicillin-resistant *Staphylococcus aureus*. Clin Infect Dis 2007; 44: 410-413.
- Tendolkar PM, Baghdayan AS, Shankar N. Pathogenic enterococci: new developments in the 21st century. Cell Mol Life Sci 2003; 60: 2622-2636.
- Walsh TJ, Pizzo PA. Nosocomial fungal infections: a classification for hospital acquired fungal infections and mycoses arising from endogenous flora or reactivation. Ann Rev Microbiol 1988; 42: 517-545.
- Gugnani AC, Okafor BC, Nzelibe F, Njoku OA. Etiological agents of otomycosis in Nigeria. Mycoses 1989; 32: 224-229.
- Kwon-Chung KJ, Bennett JE. Candidiasis. Medical Mycology. Philadelphia, USA. Lea & Febiger. 1992, p 280-336.
- Hoog GS de, Queiroz Telles F, Haase G, Fernandez-Zeppenfeldt G, Attili-Angelis D, Gerrits van den Ende AHG, Matos T, Peltroche-Llacsahuanga H, Pizzirani-Kleiner AA, Rainer J, Richard-Yegres N, Vicente VA, Yegres F. Black fungi: clinical and pathogenic approaches. Med Mycol 2000; 38: 240-250.
- Omorodion AO, Nmorsi OPG, Isaac C, Umukoro DO, Akhile AO. Distribution of intestinal parasites among school-age children in Delta and Edo States of Nigeria. Parasitologists Union J 2012; 5: 1-6.
- Omorodion OA, Isaac C, Nmorsi OPG, Ogoya EM, Agholor KN. Prevalence of intestinal parasitic infection among tertiary institution students and pregnant women in south-south, Nigeria. J Microbiol Biotech Res 2012; 2: 815-819.
- Bala AY, Sule H. Vectorial potential of cockroaches in transmitting parasites of medical importance in Arkilla, Sokoto, Nigeria. Nigerian J Basic Appl Sci 2012; 20: 111-115.
- Cloarec A, Rivault C, Fontaine F, Gugander LA. Cockroaches as carriers of bacteria in multi-family dwellings. Epidemiol Infect 1992; 109: 483-490.
- Koura EA, Kamel EG. A study of protozoa associated with some harmful insects in the local environment. J Egypt Soc Parasitol 1990; 20: 105-115.

