



OPEN ACCESS

Regulation of 5-oxo-EETE synthesis by nitric oxide in human polymorphonuclear leucocytes upon their interaction with zymosan and *Salmonella typhimurium*

Galina M. VIRYASOVA*, Svetlana I. GALKINA*, Tatjana V. GAPONOVA†, Julia M. ROMANOVA‡ and Galina F. SUD'INA*¹

*Lomonosov Moscow State University, Belozersky Institute of Physico-Chemical Biology, Moscow 119234, Russia

†FGBU Hematology Research Centre, Russia Federation Ministry of Public Health, Moscow 125167, Russia

‡Gamaleya Research Institute of Epidemiology and Microbiology, Moscow 123098, Russia

Synopsis

In the present study we have presented data on the regulation of LT (leukotriene) and 5-oxo-EETE (5-oxo-6,8,11,14-eicosatetraenoic acid) syntheses in human neutrophils upon interaction with OZ (opsonized zymosan) or *Salmonella typhimurium*. Priming of neutrophils with PMA (phorbol 12-myristate 13-acetate) and LPS (lipopolysaccharide) elicits 5-oxo-EETE formation in neutrophils exposed to OZ, and the addition of AA (arachidonic acid) significantly increases 5-oxo-EETE synthesis. We found that NO (nitric oxide)-releasing compounds induce 5-oxo-EETE synthesis in neutrophils treated with OZ or *S. typhimurium*. Exposure of neutrophils to zymosan or bacteria in the presence of the NO donor DEA NONOate (1,1-diethyl-2-hydroxy-2-nitroso-hydrazine sodium) considerably increased the conversion of endogenously formed 5-HETE (5S-hydroxy-6,8,11,14-eicosatetraenoic acid) to 5-oxo-EETE. To our knowledge, this study is the first to demonstrate that NO is a potent regulator of 5-oxo-EETE synthesis in human polymorphonuclear leucocytes exposed to *Salmonella typhimurium* and zymosan.

Key words: leukotriene, lipoxygenase, neutrophil, nitric oxide, 5-oxo-EETE, *Salmonella*.

Cite this article as: Viryasova, G.M., Galkina, S.I., Gaponova, T.V., Romanova, J.M. and Sud'ina, G.F. (2014) Regulation of 5-oxo-EETE synthesis by nitric oxide in human polymorphonuclear leucocytes upon their interaction with zymosan and *Salmonella typhimurium*. Biosci. Rep. 34(3), art:e00108.doi:10.1042/BSR20130136


INTRODUCTION

The 5-LOX (5-lipoxygenase) pathway converts AA (arachidonic acid) to 5-HETE (5S-hydroxy-6,8,11,14-eicosatetraenoic acid) and LTs (leukotrienes). The 5-LOX pathway in neutrophils metabolizes AA to LTB₄ (leukotriene B₄), which mediates host defence and inflammatory responses [1]. 5-LOX is activated during the phagocytosis of microorganisms and foreign particles by neutrophils [2], which is accompanied by a considerable increase in NADPH-oxidase activity that generates superoxide anion O₂⁻ [3]. Simultaneously, neutrophils produce an arsenal of regulatory

molecules such as chemokines and cytokines [4]. One branch of the 5-LOX pathway produces 5-oxo-EETE (5-oxo-6,8,11,14-eicosatetraenoic acid) [5], which is the most potent eosinophil chemoattractant among lipid mediators [6,7] and exerts the similar effects on neutrophils [8,9] and monocytes [10]. 5-oxo-EETE is a potent stimulator of calcium mobilization [8] and surface expression of integrins in neutrophils [11]. 5-oxo-EETE inhibits apoptosis in prostate cancer cell lines [12,13] and in neuroblastoma cell lines [14]. 5-oxo-EETE is formed by the oxidation of the 5-LOX product 5-HETE by 5-HEDH (5-hydroxyeicosanoid dehydrogenase) (Scheme 1). 5-HEDH is highly selective for this substrate and requires NADP⁺ as a cofactor [15]. Despite the

Abbreviations: 5-LOX, 5-lipoxygenase; 5-HEDH, 5-hydroxyeicosanoid dehydrogenase; 5-HETE, 5S-hydroxy-6,8,11,14-eicosatetraenoic acid; 5-oxo-EETE, 5-oxo-6,8,11,14-eicosatetraenoic acid; AA, arachidonic acid; ANF, atrial natriuretic peptide; DEA NONOate, 1,1-diethyl-2-hydroxy-2-nitroso-hydrazine sodium; Dnp-Cl, 1-chloro-2,4-dinitrobenzene; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte/macrophage colony-stimulating factor; HBSS, Hanks balanced salt solution; HBSS/Hepes, Hanks balanced salts medium containing 10 mM Hepes; G6PD, glucose-6-phosphate dehydrogenase; LPS, lipopolysaccharide; LT, leukotriene; LTB₄, leukotriene B₄; mGC, membrane-bound form of GC; NO, nitric oxide; PGB₂, prostaglandin B₂; PMA, phorbol 12-myristate 13-acetate; PMNL, polymorphonuclear leucocyte; OZ, opsonized zymosan; ROS, reactive oxygen species; sGC, soluble guanylyl cyclase; YC-1, 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole.

¹ To whom correspondence should be addressed (email sudina@genebee.msu.ru).



Schematic presentation of leukotriene and 5-oxo-EETE synthesis.

high level of 5-HEDH activity in the microsomes of neutrophils, intact cells convert 5-HETE to only trace amounts of 5-oxo-EETE, suggesting that the production of this substance is highly regulated [5].

Activation of the respiratory burst by PMA (phorbol 12-myristate 13-acetate) in phagocytic cells increases the rate of formation of 5-oxo-EETE [16]. Non-enzymatic conversion of intracellular NADPH to NADP^+ by the addition of phenazine methosulfate activates 5-oxo-EETE synthesis from exogenously added 5-HETE, suggesting that the stimulatory effect of NADPH oxidase activation is due to the generation of NADP^+ , the cofactor required by 5-HEDH [17]. Most of the data on 5-oxo-EETE synthesis have been obtained using PMNLs (polymorphonuclear leucocytes) stimulated with the calcium ionophore A23187 or from exogenously added 5-HETE. Serum-treated zymosan stimulates the conversion of exogenously added 5-HETE to 5-oxo-EETE in neutrophils [17]; however, to our knowledge, no data are available on the production of 5-oxo-EETE by neutrophils when they are engaged in phagocytizing a target.

Therefore we focused our research on the regulation of 5-oxo-EETE synthesis in PMNLs exposed to OZ (opsonized zymosan) or *Salmonella typhimurium*. We found that priming PMNLs with PMA or LPS (lipopolysaccharides), which activate ROS (reactive oxygen species) formation, induced 5-oxo-EETE synthesis upon interaction with *S. typhimurium* and zymosan. When 5-HETE synthesis was increased by adding AA, we observed a concomitant increase in 5-oxo-EETE formation. Only stimulation of PMNLs with zymosan or bacteria in the presence of an NO donor significantly increased the conversion of endogenously formed 5-HETE to 5-oxo-EETE.

MATERIAL AND METHODS

The virulent strain of *S. typhimurium* (C53) was a kind gift from Professor F. Norel (Pasteur Institute, France) [18]. Bacteria [stock, 1×10^9 CFU (colony-forming units)/ml] were grown in Luria–Bertani broth, washed twice by using a physiological salt solution, and collected by centrifugation at 2000 g. The bacteria were opsonized using fresh serum from the same donor whose

blood was used to prepare neutrophils. Serum was isolated by centrifuging the clotted blood. For opsonization, bacteria were incubated for 15 min in Dulbecco's salt solution containing 5% (v/v) serum. Bacteria were washed by repeated centrifugation in Dulbecco's solution. HBSS (Hanks balanced salt solution) with calcium and magnesium but without phenol red and sodium hydrogen carbonate, Dulbecco's PBS with magnesium but without calcium, Dnp-Cl (1-chloro-2,4-dinitrobenzene), diamide and LPSs from *Salmonella enterica* serovar *Typhimurium* (smooth form derived from the strain ATCC 7823, Ra mutant TV119 and Re mutant SL1181) were from Sigma-Aldrich. LPS from deep core mutant (Re), complete core (Ra) and smooth (S) phenotypes of *S. typhimurium* was used, as they differ in the ability to activate superoxide production in neutrophils [19]. YC-1 [3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole] and fibronectin were from Calbiochem. The NO (nitric oxide) donor DEA (diethylamine) NONOate (1,1-diethyl-2-hydroxy-2-nitroso-hydrazine sodium), 5-HETE, 5-HPETE and AA were from Cayman Chemical. Ficoll-Paque was purchased from Pharmacia. Zymosan A particles from *Saccharomyces cerevisiae* (Sigma) were boiled in PBS for 5 min and were opsonized by incubation with 30% (v/v) autologous human serum for 30 min at 37°C; they were then washed and resuspended in HBSS/Hepes (Hanks balanced salts medium containing 10 mM Hepes).

Ethics statement

We prepared neutrophils from the blood of healthy volunteers. Blood was collected via venous puncture as approved by the Ministry of Public Health Service of the Russian Federation. Experimental and subject consent procedures were approved by the Institutional Ethics Committee of the A. N. Belozersky Institute of Physico-Chemical Biology.

Isolation of PMNLs

PMNLs were isolated from freshly drawn citrated donor blood using the standard techniques [20]. Leucocyte-rich plasma was prepared by sedimenting erythrocytes through 3% (w/v) dextran T-500 at room temperature (22–24°C). Granulocytes were purified by centrifugation through Ficoll-Paque (density, 1.077 g/ml) followed by hypotonic lysis of the remaining erythrocytes. PMNLs were washed twice with PBS, resuspended at 10^7 cells/ml (purity, 96–97%; viability 98–99%) in Dulbecco's PBS containing 1 mg/ml glucose (without CaCl_2), and stored at room temperature.

Incubations for studies of AA metabolism

PMNLs (2×10^7 cells) were incubated in 6 ml HBSS/Hepes medium at 37°C with or without test compounds for 30 min and were treated with zymosan or bacteria for 15–60 min (see figure legends). The treatment was terminated by the addition of an equal volume of methanol at -20°C . PGB₂ (prostaglandin B₂) served as an internal standard. The samples were stored at -20°C . The denatured cell suspension was centrifuged, and the supernatants obtained were considered water/methanol extracts.

Analysis of lipoxygenase reaction products

The water/methanol extracts were purified by solid-phase extraction by using C₁₈ Sep-Paks (500 mg) that were first equilibrated with methanol and then with water. The metabolites of 5-LOX were extracted using 1.5 ml methanol, and the samples were evaporated and redissolved in 50 μ l methanol/water (2:1). The purified samples were injected into a 5 μ m, 250 \times 4.6 mm Nucleosil[®] C18 column (Macherey-Nagel GmbH) and subjected to RP (reverse-phase) HPLC. The products were eluted at flow rate of 0.7 ml/min with a 30-min linear gradient from 20 to 70% solvent B, followed by a 3-min isocratic elution, and then elution for 3 min by using a linear gradient of 70–100% solvent B. The eluents consisted of methanol/acetonitrile/water/acetic acid/triethylamine in the ratios (A) 25:25:50:0.05:0.075 and (B) 50:50:0:0.05:0.04. Products of the 5-LOX pathway included 5S,12R-dihydroxy-6,14-*cis*-8,10-*trans*-eicosatetraenoic acid (LTB₄), iso-LTB₄ (5S,12SR-all-*trans*-diHETE), ω -OH-LTB₄, 5S-hydroxy-6-*trans*-8,11,14-*cis*-eicosatetraenoic acid (5-HETE) and 5-oxo-EETE (5-oxo-6-*trans*-8,11,14-*cis*-eicosatetraenoic acid). They were identified by their co-elution with the standards. The compounds were quantified by comparing their peak areas with that of PGB₂ and were corrected for differences in extinction coefficients as follows: PGB₂, 27000; LTB₄, iso-LTB₄ and ω -OH-LTB₄, 40 000 at 280 nm; 5-oxo-EETE, 22 000 at 279 nm and 5-HETE, 27 000 at 236 nm.

Scanning electron microscopy (SEM)

Neutrophils were deposited on fibronectin-coated cover-slips for 20 min (control) or in the presence of PMA, DEA NONOate and OZ particles, as indicated in the figure legend. The cells were fixed for 30 min in 2.5% (w/v) glutaraldehyde, fixed again for 15 min with 1% (w/v) osmium tetroxide in 0.1 M cacodylate (pH 7.3), dehydrated using a series of acetone concentrations, subjected to critical point drying with liquid CO₂ as the transitional fluid in a Balzers apparatus, sputter-coated with gold-palladium and observed at 15 kV by using a Camscan S-2 (Tescan, Cranberry Twp) SEM.

Statistical analysis

Results have been reported as mean \pm S.D. Analysis of the statistical significance was performed by Student's *t* test using GraphPadPrism6 and SigmaPlot9.0 software. Differences with *P* values of <0.05 were considered statistically significant.

RESULTS

Priming of PMNLs with PMA and LPS elicits 5-oxo-EETE formation upon treatment with zymosan

Human neutrophils synthesize 5-LOX metabolites (LTB₄ and its isomers, ω -OH-LTB₄ and 5-HETE) during the phagocytosis of OZ or *S. typhimurium* (OS) [19,21]. We incubated PMNLs in HBSS/Hepes. In the absence of other additives, PMNLs exposed to zymosan synthesized LTB₄, its isomers, ω -OH-LTB₄ and mod-

est amounts of 5-HETE (Figure 1A). Preincubation (priming) of cells with PMA augmented the synthesis of 5-LOX products as well as the formation of 5-oxo-EETE during the interaction of PMNL with OZ. The maximum effect was observed in the presence of 2 nM PMA (Figure 1A). The accumulation of 5-oxo-EETE by PMNLs in response to zymosan lasted for 15 min (Figure 1B).

To exclude the possibility that the yields of 5-LOX products were limited by the availability of AA, we added 16 μ M (optimum concentration, results not shown) AA together with OZ. There was a significant increase in 5-HETE and 5-oxo-EETE synthesis by PMNLs with maximal 5-oxo-EETE production at 1–2 nM PMA (Figure 1C). Neither PMA nor AA induced LT and 5-oxo-EETE syntheses in PMNLs in the absence of zymosan (results not shown). The accumulation of 5-oxo-EETE ceased 15 min after exposure of PMNLs to OZ and AA (Figure 1D). AA increases the oxidation state of the cell [22], and the increase in 5-oxo-EETE may be related to this effect. Furthermore, NO is a potent regulator of the oxidative state of cells. A large amount of NO is generated during inflammation (review in [23]).

NO activates 5-oxo-EETE synthesis by human neutrophils upon their interaction with zymosan or *S. typhimurium*

To determine whether NO directly affected 5-oxo-EETE synthesis in human neutrophils, we tested whether the NO-donor DEA NONOate influenced 5-oxo-EETE formation. Addition of the NO-donor DEA NONOate simultaneously with OZ (NO_OZ) induced 5-oxo-EETE synthesis in PMNLs that were not primed (Figure 2A, control). Priming of neutrophils with PMA and LPS (LPS at concentrations when the effect is maximal) efficiently increased the 5-oxo-EETE release, which was particularly apparent when the neutrophils interacted with opsonized *S. typhimurium* (Figure 2B). Among various LPS forms Ra-LPS possessing the maximal ability to activate superoxide production demonstrated the most prominent effect on 5-oxo-EETE synthesis (Figure 2A). Adding DEA NONOate to the bacteria caused a significant increase in the synthesis of 5-LOX products and 5-oxo-EETE during the interactions of neutrophils with the bacteria. We added various amounts of DEA NONOate with OZ or OS and found that NO_OZ was maximally effective in the presence of 250 μ M NONOate (Figure 2C). The effects of 250 and 500 μ M NONOate were comparable in the action of NO_OS (Figure 2D). Accumulation of 5-oxo-EETE ceased 15 or 20 min after the PMNLs were treated with NO_OZ (Figure 2E) or NO_OS (Figure 2F), respectively.

We next added DEA NONOate to PMNLs for different fixed times before their exposure to OZ or OS as well as after adding a target of phagocytosis (Figure 3). Formation of 5-oxo-EETE was maximized when the NO donor was added 5 min after the addition of OZ (Figure 3A) or 1 min after the addition of OS (Figure 3B). To achieve optimal synthesis of 5-oxo-EETE, it was important to add the NO donor after the initial interaction of PMNLs with OZ or OS. 5-HETE or 5-HPETE (5S-hydroperoxy-eicosatetraenoic acid) added directly with DEA NONOate did not give 5-oxo-EETE (results not shown). Adding to neutro-




Figure 1 Effects of PMA concentration on LTB₄, 5-HETE and 5-oxo-EETE production by PMNLs exposed to OZ (A) or AA + OZ (C) and time course of the 5-LOX reaction in the presence of 2 nM PMA (B,D). PMNLs (2×10^7) were incubated for 30 min at 37 °C without additives or with the indicated concentration of PMA and then 2 mg/ml OZ was added to the cells for 15 min (A,C) or for the indicated time (B,D). AA (16 μ M) was added simultaneously with OZ (C,D). The products of the reaction catalysed by 5-LOX were extracted from the medium and separated using HPLC. The synthesis of all products of the 5-LOX pathway was analysed, and the data for LTB₄, 5-HETE, 5-oxo-EETE and ω -OH-LTB₄ are presented. Results (A,C) are means \pm S.D. of four independent experiments, * $P < 0.05$, ** $P < 0.01$ versus the corresponding control without PMA. Results of time studies (B,D) are means \pm S.D. of two independent experiments performed in duplicate.

phils of diamide or Dnp-Cl to increase the intracellular concentration of 5-HPETE [24,25] inhibited rather than increased the synthesis of 5-oxo-EETE (results not shown). We assume that a mechanism of 5-oxo-EETE formation by a free radical reaction of 5-HpETE with NO is hardly possible, so we propose enzymatic, 5-HEDH-catalysed conversion of 5-HETE to 5-oxo-EETE in the presence of NO donor.

The synthesis of 5-oxo-EETE requires the activation of 5-LOX and 5-HEDH (review in [26]). NO appears to activate and inhibit the formation of 5-oxo-EETE in neutrophils. NO-releasing compounds are potent activators of NO-sensitive sGC (soluble guanylyl cyclase) [27,28]. sGC and protein kinase G mediate NO suppression of 5-LOX metabolism in rat macrophages [29] and human neutrophils [19]. YC-1 is an NO-independent sGC

activator [30]. The mGC (membrane-bound form of GC) is activated by a series of ligands, including ANF (atrial natriuretic peptide), but not by NO. We observed that adding YC-1 30 min before adding zymosan inhibited LTB₄ and 5-oxo-EETE syntheses in PMNLs (Figure 4). ANF, an activator of mGC, increased the formation of LTB₄ and 5-oxo-EETE in the presence of AA/OZ. Likewise, NO inhibited the activity of 5-LOX by 30–40% when added 5 min (Figure 3A) before OZ or 30 min (results not shown) before OZ or OS. Therefore we assume that the positive effect of adding NO last is related to its inability to inhibit 5-LOX when PMNLs commence phagocytosis.

The second enzyme involved in the synthesis of 5-oxo-EETE, i.e., 5-HEDH, requires NADP⁺ as a cofactor. NO induces a rapid decrease in intracellular NADPH levels [31]. We propose




Figure 2 Effects of NO on the formation of 5-LOX reaction products and 5-oxo-EETE in human PMNLs (A,B) PMNLs (2×10^7) were incubated for 30 min at 37 °C without additives or with 1 μ M of different LPS forms or PMA (1–5 nM, as indicated), and then 2 mg/ml of zymosan was added for 15 min (A) or 5×10^8 *S. typhimurium* was added for 20 min (B). OZ or OS was added simultaneously with DEA NONOate (final concentration, 500 μ M). The results represent the means \pm S.D. of four independent experiments, * $P < 0.05$ and ** $P < 0.01$, compared with the corresponding controls without PMA or LPS. (C,D) PMNLs (2×10^7) were incubated for 30 min at 37 °C with 2 nM PMA, and then either 2 mg/ml OZ was added for 15 min (C) or 5×10^8 OS was added for 20 min (D). OZ or OS was added simultaneously with DEA NONOate at the indicated final concentration of NONOate (0.1–1 mM). The results represent the means \pm S.D. of four independent experiments, * $P < 0.05$ and ** $P < 0.01$, compared with the corresponding data without NO donor. (E,F) Time course of the 5-LOX reaction in PMNLs. PMNLs were incubated with 2 nM PMA for 30 min and then exposed to NO_OZ (E) or NO_OS (F) in a final concentration of 500 μ M NONOate. The products generated by 5-LOX were extracted from the medium and separated using HPLC. The results are means \pm S.D. of three independent experiments performed in duplicate.




Figure 3 Effect of the NO donor DEA NONOate on LTB₄, 5-HETE and 5-oxo-ETE syntheses in PMNLs

(A) PMNLs (2×10^7) were incubated for 30 min at 37 °C with 2 nM PMA, and then for 15 min with 2 mg/ml OZ; 250 μ M DEA NONOate was added before (5 and 1 min before), simultaneously (NO + OZ) or after OZ (1, 5 and 10 min after) as indicated. The results represent the means \pm S.D. of five independent experiments, * $P < 0.05$, compared with data when the NO donor was added simultaneously with OZ. (B) PMNLs (2×10^7) were incubated for 30 min at 37 °C with 2 nM PMA and then for 15 min with 5×10^8 OS. DEA NONOate (250 μ M) was added before (5 and 1 min before), simultaneously (NO + OS), and after OS (1, 5, and 10 min after) as indicated. The results represent the means \pm S.D. of four independent experiments, * $P < 0.05$, compared with data acquired when the NO donor added simultaneously with OS.

that NO activates 5-oxo-ETE formation in neutrophils through interactions of 5-LOX and 5-HEDH pathways that are regulated by numerous factors. Treatment of PMNLs with NO_OZ generated the maximal yield of 5-oxo-ETE from endogenously formed 5-HETE, compared with the yield in the absence of NO (Figures 1–3). PMNLs treated with NO efficiently converted available endogenously formed 5-HETE to 5-oxo-ETE.

DEA NONOate induced changes in 5-LOX and 5-HEDH activity that coincided with alterations in the interaction of human neutrophils with opsonised zymosan particles in the presence of the NO donor (Figure 5). Human neutrophils can phagocytose bacteria or OZ and bind them to their outer surface. We had previously demonstrated that extracellular binding occurs either

by direct attachment of bacteria or yeast to the neutrophil surface or by binding of these cells at a distance through cytonemes (dynamic thread-like membrane tubulovesicular extensions of neutrophils [32]). Our data indicate that NO donors induced cytoneme formation in human neutrophils and shifted the interaction between neutrophils and bacteria or zymosan particles towards extracellular binding instead of phagocytosis [19,33,34]. SEM observations showed that treatment with DEA NONOate stimulated extracellular attachment of OZ particles to the cell surface (Figure 5) in the presence or absence of PMA.

DISCUSSION

Neutrophils play a major role in the immune response to bacterial and fungal infections and eliminate pathogens through phagocytosis [3,35,36]. During phagocytosis, neutrophils produce ROS and release proteolytic enzymes that inflict local tissue damage. PMNLs undergo apoptosis and are removed from the site of inflammation. It was shown that dying neutrophils exhibit a dramatically increased ability to synthesize 5-oxo-ETE [37]. Here, we have shown that NO, a potent activator of apoptosis, uniquely induces 5-oxo-ETE synthesis in human neutrophils when they interact with their phagocytic target.

5-oxo-ETE synthesis increases in neutrophils and other phagocytic cells through activation of NADPH oxidase [16]. Oxidative stress increases 5-oxo-ETE formation in response to exogenously added 5-HETE [38]. Here, we observed a concomitant increase in LTB₄ synthesis and release of 5-oxo-ETE when neutrophils interacted with OZ when exposed to PMA or LPS. The effect was highest when zymosan particles were added together with the NO-releasing compound DEA NONOate (Figures 2 and 3).

The synthesis of 5-oxo-ETE requires the activation of 5-LOX but also requires a stimulus that increases the activity of the highly specific enzyme 5S-hydroxyeicosanoid dehydrogenase that requires NADP⁺. NADP⁺ is normally present at low intracellular concentrations, which limits the basal synthesis of 5-oxo-ETE. G6PD (glucose-6-phosphate dehydrogenase) is the key enzyme that maintains intracellular levels of NADPH [39]. AA is an inhibitor of G6PD activity [39]. Moreover, AA activates NADPH oxidase in PMNLs, leading to superoxide formation and oxidative stress [22]. Our present findings show that AA_OZ induced a significant net release of 5-oxo-ETE in the absence of PMA (Figure 1C).

The NADPH level in cells is rapidly reduced by the addition of an NO donor [40]. NO is lipophilic and migrates through cells, which allows it to interact with numerous molecules [41]. One may propose that 5-oxo-ETE can be made from 5-HPETE by a free radical reaction (as reported for macrophages [42]). However, observed patterns of 5-oxo-ETE synthesis indicate contradiction with this hypothesis. We did treatments for stimulation of increased formation of 5-HPETE in cells. According to the data of Hatzelmann and Ullrich [24,25], Dnp-Cl and diamide, the agents lowering the GSH-peroxidase activity in cells,




Figure 4 Influence of guanylate cyclase activators on 5-LOX product formation in OZ- and AA_OZ-stimulated PMNLs

The cells (2×10^7) were incubated for 30 min at 37 °C without additives or with 2 nM PMA, 3 μ M YC-1 or 30 nM ANF as indicated, followed by a 15-min incubation with 2 mg/ml OZ (A) or AA_OZ (16 μ M AA) (B). The 5-LOX products were extracted from the medium and separated using HPLC. The results represent the means \pm S.D. of three independent experiments, * $P < 0.05$ versus corresponding control.




Figure 5 Effect of NO on the interaction of human neutrophils with OZ

SEM images of human neutrophils that attached to fibronectin-coated substrata under the control conditions (A) or in the presence of 2 nM PMA (D) and were then exposed to opsonised zymosan (B,E) or DEA NONOate (final concentration, 500 μ M NONOate) together with OZ (C,F). Images are representative of the results of three independent experiments.

increased 5-HPETE level in neutrophils. We used Dnp-Cl and diamide in our experiments. We have found that Dnp-Cl and diamide rather inhibited than increased 5-oxo-EETE formation in PMNLs exposed to NO (results not shown).

When neutrophils are challenged with OZ or OS in the presence of an NO donor, they undergo a dramatic shift in 5-LOX metabolism that results in the formation of 5-oxo-EETE through a mechanism that remains to be defined in detail. Excessive and unregulated NO synthesis is implicated in many pathophysiological conditions, including cancer [43–45]. High neutrophil counts are associated with poor clinical outcomes for different types of cancer [46–49]. Systemic inflammation promotes the progression of malignancies through mechanisms mediated by neutrophils [50]. For example, treating neutrophils *in vitro* with 500 nM PMA significantly increases the adhesion of tumour cells to neutrophils [51].

NO is an important regulator of the interaction between macrophages and bacteria. For example, *Salmonella enterica* serovar *Enteritidis* effectively suppresses the production of NO by HD11 chicken macrophages [52], but an inhibitor of protein kinase A, H-89, dramatically reverses the suppressive effect of *Salmonella enteritidis* on the production of NO [53]. The bacteriostatic activity of H-89 augments the killing of *Salmonella enteritidis* by neutrophils [53]. We speculate that this effect may be mediated by 5-LOX metabolites and 5-oxo-EETE because the inhibition of protein kinase A by H-89 potentially activates 5-LOX [54].

In summary, we have shown here that exogenous and endogenous NO are intimately involved in the interaction between leucocytes and bacteria. NO plays a critical role in the synthesis of 5-oxo-EETE in neutrophils. In its turn, 5-oxo-EETE can enhance the survival of neutrophils by stimulating the release of GM-CSF (granulocyte/macrophage colony-stimulating factor) from monocytes [55]. During inflammation, neutrophil apoptosis can be markedly delayed by GM-CSF and G-CSF (granulocyte colony-stimulating factor) [56,57]. From other side, in PMNLs pretreated with GM-CSF or G-CSF, the 5-oxo-EETE becomes a potent activator of degranulation and superoxide anion production [58]. Tight regulation of the neutrophil lifespan is a critical process in resolving inflammation and the 5-oxo-EETE synthesis in neutrophil may be relevant to neutrophil survival.

AUTHOR CONTRIBUTION

Galina Viryasova, Svetlana Galkina and Galina Sud'ina designed and conducted the experiments. Galina Viryasova, Svetlana Galkina, Tatjana V. Gaponova, Julia M. Romanova and Galina Sud'ina analysed the data. Galina Viryasova, Svetlana Galkina, Tatjana V. Gaponova, Julia M. Romanova and Galina Sud'ina wrote the paper.

ACKNOWLEDGEMENT

We thank Elizaveta B. Pirozhkova for the technical assistance.

FUNDING

This work was supported by the Russian Foundation for Basic Research [grant numbers 13-04-00496 and 13-04-00393].

REFERENCES

- 1 Peters-Golden, M. and Henderson, W. R. (2007) Leukotrienes. *N. Engl. J. Med.* **357**, 1841–1854 [CrossRef](#)
- 2 Williams, J. D., Lee, T. H., Lewis, R. A. and Austen, F. (1985) Intracellular retention of the 5-lipoxygenase pathway product, leukotriene B₄, by human neutrophils activated with unopsonized zymosan. *J. Immunol.* **134**, 2624–2630 [PubMed](#)
- 3 Segal, A. W. (2005) How neutrophils kill microbes. *Annu. Rev. Immunol.* **23**, 197–223 [CrossRef](#) [PubMed](#)
- 4 Mantovani, A., Cassatella, M. A., Costantini, C. and Jaillon, S. (2011) Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat. Rev. Immunol.* **11**, 519–531 [CrossRef](#) [PubMed](#)
- 5 Powell, W. S. and Rokach, J. (2005) Biochemistry, biology and chemistry of the 5-lipoxygenase product 5-oxo-EETE. *Prog. Lipid Res.* **44**, 154–183 [CrossRef](#) [PubMed](#)
- 6 Powell, W. S., Chung, D. and Gravel, S. (1995) 5-Oxo-6,8,11,14-eicosatetraenoic acid is a potent stimulator of human eosinophil migration. *J. Immunol.* **154**, 4123–4132 [PubMed](#)
- 7 Schwenk, U. and Schroeder, J. M. (1995) 5-Oxo-eicosanoids are potent eosinophil chemotactic factors – functional characterization and structural requirements. *J. Biol. Chem.* **270**, 15029–15036 [CrossRef](#) [PubMed](#)
- 8 Powell, W. S., Gravel, S., MacLeod, R. J., Mills, E. and Hashefi, M. (1993) Stimulation of human neutrophils by 5-oxo-6,8,11,14-eicosatetraenoic acid by a mechanism independent of the leukotriene B₄ receptor. *J. Biol. Chem.* **268**, 9280–9286 [PubMed](#)
- 9 O'Flaherty, J. T., Cordes, J., Redman, J. and Thomas, M. J. (1993) 5-Oxo-eicosatetraenoate, a potent human neutrophil stimulus. *Biochem. Biophys. Res. Commun.* **192**, 129–134 [CrossRef](#) [PubMed](#)
- 10 Sozzani, S., Zhou, D., Locati, M., Bernasconi, S., Luini, W., Mantovani, A. and O'Flaherty, J. T. (1996) Stimulating properties of 5-oxo-eicosanoids for human monocytes: synergism with monocyte chemotactic protein-1 and -3. *J. Immunol.* **157**, 4664–4671 [PubMed](#)
- 11 Powell, W. S., Gravel, S., Halwani, F., Hii, C. S., Huang, Z. H., Tan, A. M. and Ferrante, A. (1997) Effects of 5-oxo-6,8,11,14-eicosatetraenoic acid on expression of CD11b, actin polymerization, and adherence in human neutrophils. *J. Immunol.* **159**, 2952–2959 [PubMed](#)
- 12 Ghosh, J. (2004) Rapid induction of apoptosis in prostate cancer cells by selenium: reversal by metabolites of arachidonate 5-lipoxygenase. *Biochem. Biophys. Res. Commun.* **315**, 624–635 [CrossRef](#) [PubMed](#)
- 13 Sarveswaran, S. and Ghosh, J. (2013) OXER1, a G protein-coupled oxoeicosatetraenoid receptor, mediates the survival-promoting effects of arachidonate 5-lipoxygenase in prostate cancer cells. *Cancer Lett.* **336**, 185–195 [CrossRef](#) [PubMed](#)
- 14 Bell, E., Ponthan, F., Whitworth, C., Westermann, F., Thomas, H. and Redfern, C. P. (2013) Cell survival signalling through PPAR δ and arachidonic acid metabolites in neuroblastoma. *PLoS ONE* **8**, e68859 [CrossRef](#) [PubMed](#)
- 15 Powell, W. S., Gravelle, F. and Gravel, S. (1992) Metabolism of 5(S)-hydroxy-6,8,11,14-eicosatetraenoic acid and other 5(S)-hydroxyeicosanoids by a specific dehydrogenase in human polymorphonuclear leukocytes. *J. Biol. Chem.* **267**, 19233–19241 [PubMed](#)
- 16 Powell, W. S., Zhang, Y. and Gravel, S. (1994) Effects of phorbol myristate acetate on the synthesis of 5-oxo-6,8,11,14-eicosatetraenoic acid by human polymorphonuclear leukocytes. *Biochemistry* **33**, 3927–3933 [CrossRef](#) [PubMed](#)

- 17 Powell, W. S., Gravelle, F. and Gravel, S. (1994) Phorbol myristate acetate stimulates the formation of 5-oxo-6,8,11,14-eicosatetraenoic acid by human neutrophils by activating NADPH oxidase. *J. Biol. Chem.* **269**, 25373–25380 [PubMed](#)
- 18 Kowarz, L., Coynault, C., Robbe-Saule, V. and Norel, F. (1994) The *Salmonella typhimurium* katF (rpoS) gene: cloning, nucleotide sequence, and regulation of spvR and spvABCD virulence plasmid genes. *J. Bacteriol.* **176**, 6852–6860 [PubMed](#)
- 19 Zagryazhskaya, A. N., Lindner, S. C., Grishina, Z. V., Galkina, S. I., Steinhilber, D. and Sud'ina, G. F. (2010) Nitric oxide mediates distinct effects of various LPS chemotypes on phagocytosis and leukotriene synthesis in human neutrophils. *Int. J. Biochem. Cell Biol.* **42**, 921–931 [CrossRef PubMed](#)
- 20 Sud'ina, G. F., Brock, T. G., Pushkareva, M. A., Galkina, S. I., Turutin, D. V., Peters-Golden, M. and Ullrich, V. (2001) Sulphatides trigger polymorphonuclear granulocyte spreading on collagen-coated surfaces and inhibit subsequent activation of 5-lipoxygenase. *Biochem. J.* **359**, 621–629 [CrossRef PubMed](#)
- 21 Golenkina, E. A., Galkina, S. I., Romanova, J. M., Lazarenko, M. I. and Sud'ina, G. F. (2011) Involvement of red blood cells in the regulation of leukotriene synthesis in polymorphonuclear leukocytes upon interaction with *Salmonella typhimurium*. *APMIS* **119**, 635–642 [CrossRef PubMed](#)
- 22 Hii, C. S. and Ferrante, A. (2007) Regulation of the NADPH oxidase activity and anti-microbial function of neutrophils by arachidonic acid. *Arch. Immunol. Ther. Exp. (Warsz)* **55**, 99–110 [CrossRef PubMed](#)
- 23 Pacher, P., Beckman, J. S. and Liaudet, L. (2007) Nitric oxide and peroxynitrite in health and disease. *Physiol. Rev.* **87**, 315–424 [CrossRef PubMed](#)
- 24 Hatzelmann, A. and Ullrich, V. (1987) Regulation of 5-lipoxygenase activity by the glutathione status in human polymorphonuclear leukocytes. *Eur. J. Biochem.* **169**, 175–184 [CrossRef PubMed](#)
- 25 Hatzelmann, A., Schatz, M. and Ullrich, V. (1989) Involvement of glutathione peroxidase activity in the stimulation of 5-lipoxygenase activity by glutathione-depleting agents in human polymorphonuclear leukocytes. *Eur. J. Biochem.* **180**, 527–533 [CrossRef PubMed](#)
- 26 Powell, W. S. and Rokach, J. (2013) The eosinophil chemoattractant 5-oxo-EETE and the OXE receptor. *Prog. Lipid Res.* **52**, 651–665 [CrossRef PubMed](#)
- 27 Gerzer, R., Böhme, E., Hofmann, F. and Schultz, G. (1981) Soluble guanylate cyclase purified from bovine lung contains heme and copper. *FEBS Lett.* **132**, 71–74 [CrossRef PubMed](#)
- 28 Friebe, A. and Koesling, D. (2003) Regulation of nitric oxide-sensitive guanylyl cyclase. *Circ Res.* **93**, 96–105 [CrossRef PubMed](#)
- 29 Coffey, M. J., Phare, S. M., Luo, M. and Peters-Golden, M. (2008) Guanylyl cyclase and protein kinase G mediate nitric oxide suppression of 5-lipoxygenase metabolism in rat alveolar macrophages. *Biochim. Biophys. Acta* **1781**, 299–305 [CrossRef PubMed](#)
- 30 Hwang, T. L., Hung, H. W., Kao, S. H., Teng, C. M., Wu, C. C. and Cheng, S. J. (2003) Soluble guanylyl cyclase activator YC-1 inhibits human neutrophil functions through a cGMP-independent but cAMP-dependent pathway. *Mol. Pharmacol.* **64**, 1419–1427 [CrossRef PubMed](#)
- 31 García-Nogales, P., Almeida, A. and Bolaños, J. P. (2003) Peroxynitrite protects neurons against nitric oxide-mediated apoptosis. A key role for glucose-6-phosphate dehydrogenase activity in neuroprotection. *J. Biol. Chem.* **278**, 864–874 [CrossRef PubMed](#)
- 32 Galkina, S. I., Sud'ina, G. F. and Ullrich, V. (2001) Inhibition of neutrophil spreading during adhesion to fibronectin reveals formation of long tubulovesicular cell extensions (cytonemes). *Exp. Cell Res.* **266**, 222–228 [CrossRef PubMed](#)
- 33 Galkina, S. I., Molotkovsky, J. G., Ullrich, V. and Sud'ina, G. F. (2005) Scanning electron microscopy study of neutrophil membrane tubulovesicular extensions (cytonemes) and their role in anchoring, aggregation and phagocytosis. The effect of nitric oxide. *Exp. Cell Res.* **304**, 620–629 [CrossRef PubMed](#)
- 34 Galkina, S. I., Romanova, J. M., Stadnichuk, V. I., Molotkovsky, J. G., Sud'ina, G. F. and Klein, T. (2009) Nitric oxide-induced membrane tubulovesicular extensions (cytonemes) of human neutrophils catch and hold *Salmonella enterica* serovar *Typhimurium* at a distance from the cell surface. *FEMS Immunol. Med. Microbiol.* **56**, 162–171 [CrossRef PubMed](#)
- 35 Cohn, Z. A. and Hirsch, J. G. (1960) The influence of phagocytosis on the intracellular distribution of granule-associated components of polymorphonuclear leucocytes. *J. Exp. Med.* **112**, 1015–1022 [CrossRef PubMed](#)
- 36 Iyer, G. Y. N., Islam, D. M. F. and Quastel, J. H. (1961) Biochemical aspects of phagocytosis. *Nature* **192**, 535–541 [CrossRef](#)
- 37 Graham, F. D., Erlemann, K. R., Gravel, S., Rokach, J. and Powell, W. S. (2009) Oxidative stress-induced changes in pyridine nucleotides and chemoattractant 5-lipoxygenase products in aging neutrophils. *Free Radic. Biol. Med.* **47**, 62–71 [CrossRef PubMed](#)
- 38 Erlemann, K. R., Cossette, C., Gravel, S., Lesimple, A., Lee, G. J., Saha, G., Rokach, J. and Powell, W. S. (2007) Airway epithelial cells synthesize the lipid mediator 5-oxo-EETE in response to oxidative stress. *Free Radic. Biol. Med.* **42**, 654–664 [CrossRef PubMed](#)
- 39 Stanton, R. C. (2012) Glucose-6-phosphate dehydrogenase, NADPH, and cell survival. *IUBMB Life* **64**, 362–369 [CrossRef PubMed](#)
- 40 Bian, K. and Murad, F. (2003) Nitric oxide (NO)–biogenesis, regulation, and relevance to human diseases. *Front. Biosci.* **8**, d264–d278 [CrossRef PubMed](#)
- 41 Wink, D. A., Hines, H. B., Cheng, R. Y., Switzer, C. H., Flores-Santana, W., Vitek, M. P., Ridnour, L. A. and Colton, C. A. (2011) Nitric oxide and redox mechanisms in the immune response. *J. Leukoc. Biol.* **89**, 873–891 [CrossRef PubMed](#)
- 42 Zarini, S. and Murphy, R. C. (2003) Biosynthesis of 5-oxo-6,8,11,14-eicosatetraenoic acid from 5-hydroperoxyeicosatetraenoic acid in the murine macrophage. *J. Biol. Chem.* **278**, 11190–11196 [CrossRef](#)
- 43 Grimm, E. A., Sikora, A. G. and Ekmekcioglu, S. (2013) Molecular pathways: inflammation-associated nitric-oxide production as a cancer-supporting redox mechanism and a potential therapeutic target. *Clin. Cancer Res.* **19**, 5557–5563 [CrossRef PubMed](#)
- 44 Bian, K., Ghassemi, F., Sotolongo, A., Siu, A., Shauger, L., Kots, A. and Murad, F. (2012) NOS-2 signaling and cancer therapy. *IUBMB Life* **64**, 676–683 [CrossRef PubMed](#)
- 45 Janakiram, N. B. and Rao, C. V. (2012) iNOS-selective inhibitors for cancer prevention: promise and progress. *Future Med. Chem.* **4**, 2193–2204 [CrossRef PubMed](#)
- 46 Shimada, H., Takiguchi, N., Kainuma, O., Soda, H., Ikeda, A., Cho, A., Miyazaki, A., Gunji, H., Yamamoto, H. and Nagata, M. (2010) High preoperative neutrophil-lymphocyte ratio predicts poor survival in patients with gastric cancer. *Gastric Cancer* **13**, 170–176 [CrossRef PubMed](#)
- 47 An, X., Ding, P. R., Wang, F. H., Jiang, W. Q. and Li, Y. H. (2011) Elevated neutrophil to lymphocyte ratio predicts poor prognosis in nasopharyngeal carcinoma. *Tumour Biol.* **32**, 317–324 [CrossRef PubMed](#)
- 48 Jensen, H. K., Donskov, F., Marcussen, N., Nordmark, M., Lundbeck, F. and von der Maase, H. (2009) Presence of intratumoral neutrophils is an independent prognostic factor in localized renal cell carcinoma. *J. Clin. Oncol.* **27**, 4709–4717 [CrossRef PubMed](#)



- 49 Li, Y. W., Qiu, S. J., Fan, J., Zhou, J., Gao, Q., Xiao, Y. S. and Xu, Y. F. (2011) Intratumoral neutrophils: a poor prognostic factor for hepatocellular carcinoma following resection. *J. Hepatol.* **54**, 497–505 [CrossRef PubMed](#)
- 50 McDonald, B., Spicer, J., Giannais, B., Fallavollita, L., Brodt, P. and Ferri, L. E. (2009) Systemic inflammation increases cancer cell adhesion to hepatic sinusoids by neutrophil mediated mechanisms. *Int. J. Cancer* **125**, 1298–1305 [CrossRef PubMed](#)
- 51 Cools-Lartigue, J., Spicer, J., McDonald, B., Gowing, S., Chow, S., Giannais, B., Bourdeau, F., Kubes, P. and Ferri, L. (2013) Neutrophil extracellular traps sequester circulating tumor cells and promote metastasis. *J. Clin. Invest.* **123**, 3446–3458 [CrossRef PubMed](#)
- 52 He, H., Genovese, K. J., Swaggerty, C. L., Nisbet, D. J. and Kogut, M. H. (2012) A comparative study on invasion, survival, modulation of oxidative burst, and nitric oxide responses of macrophages (HD11), and systemic infection in chickens by prevalent poultry *Salmonella* serovars. *Foodborne Pathog. Dis.* **9**, 1104–1110 [CrossRef PubMed](#)
- 53 He, H., Genovese, K. J., Swaggerty, C. L., Nisbet, D. J. and Kogut, M. H. (2013) Nitric oxide as a biomarker of intracellular *Salmonella* viability and identification of the bacteriostatic activity of protein kinase A inhibitor H-89. *PLoS ONE* **8**, e58873 [CrossRef PubMed](#)
- 54 Luo, M., Jones, S. M., Flamand, N., Aronoff, D. M., Peters-Golden, M. and Brock, T. G. (2005) Phosphorylation by protein kinase A inhibits nuclear import of 5-lipoxygenase. *J. Biol. Chem.* **280**, 40609–40616 [CrossRef PubMed](#)
- 55 Stamatou, P. B., Chan, C. C., Monneret, G., Ethier, D., Rokach, J. and Powell, W. S. (2004) 5-Oxo-6,8,11,14-eicosatetraenoic acid stimulates the release of the eosinophil survival factor granulocyte-macrophage colony stimulating factor from monocytes. *J. Biol. Chem.* **279**, 28159–28164 [CrossRef PubMed](#)
- 56 Kobayashi, S. D., Voyich, J. M., Whitney, A. R. and DeLeo, F. R. (2005) Spontaneous neutrophil apoptosis and regulation of cell survival by granulocyte macrophage-colony stimulating factor. *J. Leukoc. Biol.* **78**, 1408–1418 [CrossRef PubMed](#)
- 57 van Raam, B. J., Drewniak, A., Groenewold, V., van den Berg, T. K. and Kuijpers, T. W. (2008) Granulocyte colony-stimulating factor delays neutrophil apoptosis by inhibition of calpains upstream of caspase-3. *Blood* **112**, 2046–2054 [CrossRef PubMed](#)
- 58 O'Flaherty, J. T., Kuroki, M., Nixon, A. B., Wijkander, J., Yee, E., Lee, S. L., Smitherman, P. K., Wykle, R. L. and Daniel, L. W. (1996) 5-Oxo-eicosanoids and hematopoietic cytokines cooperate in stimulating neutrophil function and the mitogen-activated protein kinase pathway. *J. Biol. Chem.* **271**, 17821–17828 [CrossRef PubMed](#)

Received 23 December 2013/2 April 2014; accepted 9 April 2014

Published as Immediate Publication 9 April 2014, doi 10.1042/BSR20130136
