

Pituitary adenylate cyclase–activating polypeptide is reduced in Alzheimer disease

Pengcheng Han, MD,
PhD
Winnie Liang, PhD
Leslie C. Baxter, PhD
Junxiang Yin, MD, PhD
Zhiwei Tang, MD, PhD
Thomas G. Beach, MD
Richard J. Caselli, MD
Eric M. Reiman, MD
Jiong Shi, MD, PhD

Correspondence to
Dr. Shi:
Jiong.Shi@DignityHealth.org

ABSTRACT

Objectives: There is growing evidence that pituitary adenylate cyclase–activating polypeptide (PACAP) is associated with Alzheimer disease (AD) pathology in animal models, but human studies are needed.

Methods: We studied the brains of patients with pathologically confirmed late-onset AD and age-matched cognitively normal (CN) subjects to investigate the expression of PACAP messenger RNA (34 AD and 14 CN) and protein (12 AD and 11 CN) in a case-control study.

Results: We report that PACAP levels are reduced in multiple brain regions, including the entorhinal cortex, the middle temporal gyrus, the superior frontal gyrus, and the primary visual cortex. This reduction is correlated with higher amyloid burden (CERAD plaque density) in the entorhinal cortex and superior frontal gyrus but not in the primary visual cortex, a region spared in most cases of AD. PACAP expression is lower in advanced Braak stages (V and VI) than in moderate stages (III and IV). Increased PACAP levels are associated with decreased scores on the Dementia Rating Scale, a global cognitive measure. Finally, CSF levels paralleled brain levels in AD but not in Parkinson dementia or frontotemporal dementia brains.

Conclusions: The close relationship between PACAP reduction and the severity of AD pathology suggests that downregulation of PACAP may contribute to AD pathogenesis. *Neurology*® 2014;82:1724–1728

GLOSSARY

AD = Alzheimer disease; **CERAD** = Consortium to Establish a Registry for Alzheimer's Disease; **CN** = cognitively normal; **ENT** = entorhinal cortex; **MTG** = middle temporal gyrus; **PACAP** = pituitary adenylate cyclase–activating polypeptide; **PDD** = Parkinson disease with dementia; **PVC** = primary visual cortex; **SFG** = superior frontal gyrus.

Pituitary adenylate cyclase–activating polypeptide (PACAP) is a potent neurotrophic and neuroprotective peptide that binds to and activates G protein–coupled receptors.¹ In a gene expression survey study using 3 different mouse models of Alzheimer disease (AD), PACAP was one of the 3 downregulated genes.² Recent studies demonstrate that PACAP can stimulate nonamyloidogenic processing, inhibit amyloid deposition, facilitate β -amyloid clearance, and improve cognitive performance in amyloid precursor protein transgenic mice.³ PACAP can also promote synaptic transmission, long-term potentiation, and memory performance under physiologic conditions.⁴ However, the relevance of PACAP expression has not been studied in the human AD brain. We hypothesized that PACAP expression is reduced in human AD, and its reduction is associated with AD pathology and cognitive performance.

METHODS We obtained postmortem human brains from the Banner Sun Health Research Institute Brain and Body Donation Program. Frozen brain tissue was obtained from patients with a clinical and pathologic diagnosis of late-onset AD and from age-matched cognitively normal (CN) subjects. Brain donors all underwent extensive longitudinal clinical and neuropsychological assessment antemortem, as previously described.⁵ We selected AD cases as being “intermediate” or “high” probability for AD according to National Institute on Aging–Reagan criteria (National Institute on Aging–Alzheimer's Association criteria).⁶ CN subjects did not meet the criteria for AD or dementia. The AD cases and controls did not differ significantly in their age at death, sex, educational level,

From the Barrow Neurological Institute (P.H., L.C.B., J.Y., Z.T., J.S.), St. Joseph Hospital and Medical Center, Dignity Health Organization, Phoenix; Translational Genomics Research Institute (W.L.), Phoenix, AZ; Department of Neurosurgery (Z.T.), The First Hospital of Kunming Medical University, Kunming, China; Civin Laboratory for Neuropathology (T.G.B.), Banner Sun Health Research Institute, Sun City; Department of Neurology (R.J.C.), Mayo Clinic Arizona, Scottsdale; and Banner Alzheimer's Institute (E.M.R.), Phoenix, AZ.

Go to Neurology.org for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Table 1 Neurotranscriptome of ADCYAP1 gene expression

Gene	Brain region	Fold change	p Value
ADCYAP1	ENT	-3.702532581	0.127300685
ADCYAP1	MTG	-4.039581566	0.005626227
ADCYAP1	SFG	-8.911001119	0.008381943
ADCYAP1	PVC	-5.904469845	0.002280590

Abbreviations: ENT = entorhinal cortex; MTG = middle temporal lobe; PACAP = pituitary adenylate cyclase-activating polypeptide; PVC = primary visual cortex; SFG = superior frontal cortex.

ADCYAP1 is the PACAP gene.

or postmortem interval. A neuropathologist (T.G.B.) determined cortical β -amyloid neuritic plaque density (CERAD evaluation) and Braak tangle stage. We obtained postmortem cisternal CSF samples from the same cohort.

We quantified protein sample solution (RIPA buffer containing $1\times$ proteinase inhibitor cocktails, P8340; Sigma-Aldrich, St. Louis, MO) and CSF (AD = 12 and CN = 11) for PACAP expression using a standard ELISA kit (MBS160511; MyBioSource Inc., San Diego, CA) according to the manufacturer's protocol. Briefly, we loaded samples and incubated them with biotin-labeled PACAP antibody at 37°C for 60 minutes. We washed the plate with the washing buffer 5 times, followed by incubation with the chromogen solution at 37°C for 10 minutes. We stopped the reaction by adding a stop solution. We read the final results at OB 450 nm. We compared the ELISA results of PACAP levels in the brain with the Western blot results. The PACAP level measured was in the middle range of the standard curve and correlated closely with the Western blot results (CN: Pearson $r = 0.9093$, $p < 0.01$; AD: Pearson $r = 0.8243$, $p < 0.01$; data not shown).

We used another set of postmortem brain samples (AD = 34, CN = 14) for transcriptome study as described previously.⁷ Briefly, we stained brain sections with a combination of thioflavin-S (Sigma-Aldrich, Dallas, TX) and 1% neutral red (Fisher Scientific, Chicago, IL). We identified and laser-captured pyramidal neurons onto Arcturus CapSure Macro LCM Caps and extracted according to the manufacturer's protocol. We isolated total RNA from the neuronal cell lysate with the Arcturus PicoPure RNA Isolation Kit with DNase I treatment using Qiagen's RNase-free DNase Set (Valencia, CA). Isolated total RNA from each sample of approximately 500 neurons was double-round amplified, cleaned, and biotin-labeled with Affymetrix's GeneChip per the manufacturer's protocol. Amplified and labeled complementary RNA was quantified on a spectrophotometer

and run on a 1% Tris-acetate-EDTA (TAE) gel to check for an evenly distributed range of transcript sizes.

We used t tests to compare the values of the 2 groups. For comparing values across multiple groups, we used a one-way analysis of variance with post hoc Tukey test. We applied Pearson correlation tests for correlation analyses. We reported all results as mean \pm standard error and set $p < 0.05$ as statistically significant.

Standard protocol approvals, registrations, and patient consents.

We received approval from an ethical standards committee on human experimentation (the Western Institutional Review Board) for any experiments using human subjects. Written informed consent was obtained from all patients (or guardians of patients) participating in the study (consent for research).

RESULTS PACAP is reduced in multiple areas of human AD brain.

Neurons were laser-captured and microdissected from multiple brain regions of AD and CN subjects. Neuronal expression of *ADCYAP1* (the PACAP gene) was significantly reduced in AD brains overall, and regionally in the middle temporal gyrus (MTG), superior frontal gyrus (SFG), and primary visual cortex (PVC) (table 1). To validate this, we selected a different cohort of 12 cases of AD postmortem brain samples and 11 CN cases for neurotranscriptome-based screening. Using ELISA to quantify PACAP protein expression, PACAP protein levels were reduced in AD in the same 3 regions as well as the entorhinal cortex (ENT) (table 2).

PACAP reduction is associated with pathologic hallmarks of AD.

Amyloid plaques and neurofibrillary tangles are the 2 pathologic hallmarks of AD. PACAP protein levels were reduced with higher CERAD amyloid plaque scores in the ENT and SFG but not in the MTG or PVC (figure 1, A–D). Regarding neurofibrillary tangles, the AD cases had Braak stages ranging from IV to VI, whereas the stages in CN samples were III and IV. PACAP levels were reduced in Braak stages V and VI (all AD cases) compared with stages III and IV (figure 1E).

PACAP levels in CSF parallel brain levels. PACAP concentration in AD CSF was reduced relative to CN CSF


Table 2 ELISA quantification of PACAP

	CN		AD		p Value
	PACAP (mean \pm SEM, $\times 10^{-2}$ ng/mg protein)	No.	PACAP (mean \pm SEM, $\times 10^{-2}$ ng/mg protein)	No.	
ENT	2.13 \pm 0.25	11	1.41 \pm 0.19	12	0.035
MTG	1.24 \pm 0.11	8	0.73 \pm 0.05	11	0.007
SFG	6.35 \pm 0.33	8	3.63 \pm 0.22	10	0.001
PVC	5.98 \pm 0.35	7	3.86 \pm 0.45	11	0.004

Abbreviations: AD = Alzheimer disease; CN = cognitively normal controls; ENT = entorhinal cortex; MTG = middle temporal lobe; PACAP = pituitary adenylate cyclase-activating polypeptide; PVC = primary visual cortex; SFG = superior frontal cortex.

PACAP levels were normalized with total protein (mg) of the brain tissue.

Figure 1 PACAP levels are inversely related to AD pathology



(A–D) PACAP levels in AD brains were analyzed for correlation with amyloid plaque quantity (indicated by CERAD plaque score). Lower PACAP levels were correlated with higher CERAD in the ENT (Pearson $r = -0.6764$, $p < 0.05$, panel A) and SFG (Pearson $r = -0.7088$, $p < 0.05$, panel C) but not in the MTG (B) or PVC (D). PACAP level was quantified by ELISA and normalized with total protein (mg) of the brain tissue. (E) PACAP level in CSF was quantified and correlated with tau pathology (indicated by Braak stage). All CN cases were in stage III or IV. Four AD cases were in stage III or IV, while the other 5 AD cases were in stage V or VI. PACAP was lower in advanced Braak stages V and VI than in moderate Braak stages III and IV ($p < 0.05$). PACAP level was quantified by ELISA and normalized with CSF volume (mL). AD = Alzheimer disease (blue dot); CERAD = Consortium to Establish a Registry for Alzheimer’s Disease; CN = cognitively normal controls (black dot); ENT = entorhinal cortex; MTG = middle temporal lobe; PACAP = pituitary adenylate cyclase-activating polypeptide; PVC = primary visual cortex; SFG = superior frontal cortex.


(figure 2A). In contrast, CSF PACAP levels from patients with Parkinson disease with dementia (PDD) and frontotemporal lobar degeneration were similar to those of CN subjects (figure 2A). CSF PACAP concentrations in patients with AD strongly correlated with Mattis Dementia Rating Scale–Revised scores, a measure of global cognitive functioning (figure 2B).

DISCUSSION In the 4 brain areas we examined, both PACAP messenger RNA transcription and protein expression were significantly reduced in AD brains compared with matched CN controls. PACAP

levels in the CSF were also significantly decreased in AD vs controls. Our findings suggest that the reduced neurotrophic effects from PACAP may be an important factor contributing to Alzheimer pathology.

The CERAD amyloid plaque burden was correlated with lower PACAP expression in the ENT and SFG but not the MTG or PVC. Although the lack of correlation with the MTG was surprising, the PVC is often relatively spared (save in the variant syndrome of posterior cortical atrophy). The involvement of neurofibrillary tangles in AD follows the progression from the ENT to the temporal lobe and then

Figure 2 Reduction in PACAP levels in CSF is specific to AD



PACAP levels were quantified by ELISA as ng PACAP per mL undiluted CSF. (A) PACAP was reduced in AD but not in PDD or FTLD. (B) PACAP is correlated with the z score of the DRS (Pearson $r = 0.8197$, $p = 0.0001$). CN and AD cases were separated by the dotted line based on their DRS-R score. AD = Alzheimer disease (blue dot); CN = cognitively normal controls (black dot); DRS-R = Dementia Rating Scale-Revised, a global cognitive assessment; FTLD = frontotemporal lobe dementia (green dot); PACAP = pituitary adenylate cyclase-activating polypeptide; PDD = Parkinson disease with dementia (purple dot). * $p < 0.05$; ** $p < 0.01$.

to the distal neocortex. AD cases had higher Braak stages but lower PACAP levels, suggesting that PACAP could reflect tau pathology.

PACAP is an abundant neurotrophin in the brain. PACAP levels in the CSF may be a good surrogate for diagnosis and monitoring the disease progression because CSF is more readily available than brain biopsy. PACAP levels in the CSF are reduced in AD but not in PDD or frontotemporal lobar degeneration, suggesting that it may also be specific for AD, consistent with the close relationship between PACAP and pathologic markers of AD.

PACAP is reduced in the cortical regions that typically characterize the topography of AD pathology. In addition to its role as an effective biomarker, PACAP may have potential as a therapeutic target as has been shown in animal models of AD. Translation of PACAP research to human subjects may provide us with not only new early biomarkers but also exciting therapeutic options for AD. Additional studies are needed to clarify the nature of the relationship between PACAP reductions and the neuropathologic features of AD and the extent to which it might provide a target for therapeutic interventions.

AUTHOR CONTRIBUTIONS

Dr. Han: drafting and revising the manuscript, experiment design, and data acquisition and analysis. Dr. Liang: experiment design of the transcriptome part and critical revisions. Dr. Baxter: neuropsychologic data analysis and critical revisions. Dr. Yin and Dr. Tang: data acquisition and revision. Dr. Beach: neuropathologic examination, and data interpretation and revision. Dr. Caselli and Dr. Reiman: data interpretation and critical revision. Dr. Shi: concept and design, data analysis and interpretation, and critical revision.

ACKNOWLEDGMENT

The authors are deeply grateful for the patients and families who have participated in our brain donation program. The authors thank Dr. Geidy Serrano for preparing the human pathologic tissues.

STUDY FUNDING

Supported by Arizona Alzheimer's Disease Consortium and Barrow Neurological Foundation.

DISCLOSURE

P. Han and W. Liang report no disclosures relevant to the manuscript. L. Baxter receives funding from Arizona Alzheimer's Disease Consortium and Barrow Neurological Foundation. J. Yin and Z. Tang report no disclosures relevant to the manuscript. T. Beach receives grant support from the National Institute of Neurological Disorders and Stroke (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), and the Michael J. Fox Foundation for Parkinson's Research. He performs research services (payments to the institute, not to himself) as part of contractual agreements with Avid Radiopharmaceuticals/Eli Lilly Corporation, Bayer Healthcare, GE Healthcare, and Navidea Healthcare. R. Caselli receives research support from the NIH/National Institute on Aging and the Arizona Alzheimer's Research Consortium. E. Reiman reports serving as a scientific advisor to SYGNIS, AstraZeneca, Bayer, Eisai, Elan, Eli Lilly, GlaxoSmithKline, Intellect, Link Medicine, Novartis, Siemens, and Takeda. He has had research contracts with AstraZeneca and Avid/Eli Lilly; a patent pending for a biomarker strategy to evaluate preclinical AD treatments (through Banner Health); and research grants from NIH (National Institute on Aging), Anonymous Foundation, Nomis Foundation, Banner Alzheimer's Foundation, and the State of Arizona. J. Shi receives funding from Arizona Alzheimer's Disease Consortium and Barrow Neurological Foundation. Go to Neurology.org for full disclosures.

Received November 25, 2013. Accepted in final form February 10, 2014.

REFERENCES

- Vaudry D, Falluel-Morel A, Bourgault S, et al. Pituitary adenylate cyclase-activating polypeptide and its receptors: 20 years after the discovery. *Pharmacol Rev* 2009;61:283–357.
- Wu ZL, Ciallella JR, Flood DG, O'Kane TM, Bozyczko-Coyne D, Savage MJ. Comparative analysis of cortical gene expression in mouse models of Alzheimer's disease. *Neurobiol Aging* 2006;27:377–386.
- Rat D, Schmitt U, Tippmann F, et al. Neuropeptide pituitary adenylate cyclase-activating polypeptide (PACAP) slows down Alzheimer's disease-like pathology in amyloid precursor protein-transgenic mice. *FASEB J* 2011;25:3208–3218.

4. Roberto M, Brunelli M. PACAP-38 enhances excitatory synaptic transmission in the rat hippocampal CA1 region. *Learn Mem* 2000;7:303–311.
5. Caselli RJ, Walker D, Sue L, Sabbagh M, Beach T. Amyloid load in nondemented brains correlates with APOE e4. *Neurosci Lett* 2010;473:168–171.
6. Consensus recommendations for the postmortem diagnosis of Alzheimer's disease. The National Institute on Aging, and Reagan Institute Working Group on Diagnostic Criteria for the Neuropathological Assessment of Alzheimer's Disease. *Neurobiol Aging* 1997; 18:S1–S2.
7. Liang WS, Dunckley T, Beach TG, et al. Altered neuronal gene expression in brain regions differentially affected by Alzheimer's disease: a reference data set. *Physiol Genomics* 2008;33:240–256.

WriteClick: Rapid Online Correspondence

The editors encourage comments about recent articles through WriteClick:

Go to www.neurology.org and click on the “WriteClick” tab at the top of the page. Responses will be posted within 72 hours of submission.

Before using WriteClick, remember the following:

- WriteClick is restricted to comments about studies published in *Neurology* within the last eight weeks
- Read previously posted comments; redundant comments will not be posted
- Your submission must be 200 words or less and have a maximum of five references; reference one must be the article on which you are commenting
- You can include a maximum of five authors (including yourself)

Earn 20 CME Credits Toward MOC with New NeuroPISM Modules

Choose from the latest lineup of quality modules to join the AAN's exclusive performance improvement programs designed to help you address both the Performance in Practice (PIP) and Continuing Medical Education (CME) components of Maintenance of Certification (MOC).

- **NEW! Distal Symmetric Polyneuropathy (DSP)** includes **eight quality measures**, addressing accurate and appropriate evaluation/monitoring of DSP and associated symptoms to guide treatment options, patient safety, and best practices to assist patients in managing their pain and improving quality of life
- **Acute Stroke** addresses **six quality measures**, including deep vein thrombosis prophylaxis (DVT) for ischemic stroke or intracranial hemorrhage, discharged on antiplatelet therapy, dysphagia screening, rehabilitation service considerations, and more
- **Dementia** includes **10 quality measures** addressing underuse of effective services and patient-centered care strategies, and patient safety issues

Learn about all of the other available modules and purchase yours today:

www.aan.com/view/neuropi