



Published in final edited form as:

*Mol Ecol.* 2014 May ; 23(9): 2242–2259. doi:10.1111/mec.12733.

## Gene expression divergence between malaria vector sibling species *Anopheles gambiae* and *An. coluzzii* from rural and urban Yaoundé Cameroon

Bryan J. Cassone<sup>\*1</sup>, Colince Kamdem<sup>§2</sup>, Changde Cheng<sup>\*</sup>, John C. Tan<sup>\*</sup>, Matthew W. Hahn<sup>†</sup>, Carlo Costantini<sup>§,‡</sup>, and Nora J. Besansky<sup>\*</sup>

<sup>\*</sup>Eck Institute for Global Health & Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556-0369

<sup>§</sup>Laboratoire de Recherche sur le Paludisme, Organisation de Coordination pour la lutte Contre les Endémies en Afrique Centrale (OCEAC), Yaoundé, Cameroun

<sup>†</sup>Department of Biology & School of Informatics and Computing, Indiana University, Bloomington, IN 47405

<sup>‡</sup>Institut de Recherche pour le Développement (IRD), MIVEGEC (UM1, UM2, CNRS 5290, IRD 224), 34394 Montpellier Cedex 5, France

### Abstract

Divergent selection based on aquatic larval ecology is a likely factor in the recent isolation of two broadly sympatric and morphologically identical African mosquito species, the malaria vectors *Anopheles gambiae* and *An. coluzzii*. Population-based genome scans have revealed numerous candidate regions of recent positive selection, but have provided few clues as to the genetic mechanisms underlying behavioral and physiological divergence between the two species, phenotypes which themselves remain obscure. To uncover possible genetic mechanisms, we compared global transcriptional profiles of natural and experimental populations using gene-based microarrays. Larvae were sampled as second and fourth instars from natural populations in and around the city of Yaoundé, capital of Cameroon, where the two species segregate along a gradient of urbanization. Functional enrichment analysis of differentially expressed genes revealed that *An. coluzzii*—the species that breeds in more stable, biotically complex and potentially polluted urban water bodies—over-expresses genes implicated in detoxification and immunity relative to *An. gambiae*, which breeds in more ephemeral and relatively depauperate pools and puddles in suburbs and rural areas. Moreover, our data suggest that such over-expression by *An.*

---

Corresponding author: Nora J. Besansky, Eck Institute for Global Health & Department of Biological Sciences, 317 Galvin Life Sciences Bldg, University of Notre Dame, Notre Dame IN 46556-0369; tel: 574-631-9321; fax: 574-631-3996; nbesansk@nd.edu.

<sup>1</sup>Present address: Department of Plant Pathology, Ohio Agricultural Research and Development Center, The Ohio State University, Wooster, OH 44691

<sup>2</sup>Present address: Department of Entomology, University of California, Riverside, CA 92521

#### Data accessibility

Microarray data have been deposited with Array Express under accession E-MEXP-3147.

#### Author contributions

N.J.B., C. Costantini, and M.W.H. designed the study and obtained funding; C.K. and C. Costantini conducted field work; B.J.C. and J.C.T. performed molecular work; B.J.C., C. Cheng, M.W.H. and N.J.B. analyzed data; B.J.C., M.W.H. and N.J.B. wrote the paper.

*coluzzii* is not a transient result of induction by xenobiotics in the larval habitat, but an inherent and presumably adaptive response to repeatedly encountered environmental stressors. Finally, we find no significant overlap between the differentially expressed loci and previously identified genomic regions of recent positive selection, suggesting that transcriptome divergence is regulated by trans-acting factors rather than cis-acting elements.

## Keywords

comparative transcriptomics; cryptic species; ecological and environmental genomics; M and S molecular forms; microarrays; transcriptional variation

---

## Introduction

Cryptic species—those difficult or impossible to distinguish morphologically at any life stage—are widespread both geographically and across the tree of life (Bickford *et al.* 2007; Pfenninger & Schwenk 2007). Due to their morphological similarity, cryptic species are likely to be very similar ecologically (Leibold & McPeck 2006). Thus, for recently diverged cryptic species to coexist stably, competition is presumably limited by physiological and behavioral adaptations to alternative local habitats or resources, which confer some degree of specialization and niche partitioning (Chesson 2000; Montero-Pau *et al.* 2011). Identifying the genetic basis of such adaptive phenotypic divergence can offer insight into molecular mechanisms underlying ecological isolation, a grand challenge of ecological genomics.

If physiological or behavioral traits distinguishing cryptic species are known in advance and can be scored in controlled laboratory crosses, quantitative trait locus mapping is the most direct way to relate phenotype with an underlying genotype. However, where this approach is unfeasible, comparative genomic approaches involving genome scans of nucleotide divergence along chromosomes (Li *et al.* 2008; Nosil *et al.* 2009; Storz 2005) or gene expression profiling (Romero *et al.* 2012; Whitehead & Crawford 2006b; Zheng *et al.* 2011) allow biological discovery and hypothesis generation in the absence of *a priori* information about the nature of ecological divergence between species. There have long been good theoretical grounds for the expectation that transcriptional rewiring underlies many phenotypic changes between species (Britten & Davidson 1969, 1971). More recently, growing empirical evidence supports the important role of transcriptional variation in the evolution of physiological and behavioral differences between species (Borneman *et al.* 2007; Matzkin 2012; Nowick *et al.* 2009; Whitehead & Crawford 2006a; Wittkopp *et al.* 2008). Thus, whole-transcriptome profiling of sympatric and closely related cryptic species can be a powerful approach to discover the functional basis of their ecological isolation (Whitehead 2012).

Two of the principal malaria vector mosquitoes in tropical Africa belong to the *Anopheles gambiae* sibling species complex. Formerly designated as *An. gambiae* M and S molecular forms (della Torre *et al.* 2001), and recently named *An. coluzzii* and *An. gambiae* (Coetzee *et al.* 2013), these sister species are very recently diverged and strictly sympatric in many parts of West and Central Africa (della Torre *et al.* 2005). Despite the absence of intrinsic genetic

barriers to gene exchange in hybrids (Diabate *et al.* 2007; Hahn *et al.* 2012), assortative mating (Dabire *et al.* 2013; Diabate *et al.* 2006; Diabate *et al.* 2009), and ecologically based reproductive isolation (Diabate *et al.* 2008; Diabate *et al.* 2005) limit the amount of localized interspecific gene flow and introgression, such that both species appear to be evolving collectively on independent trajectories across much of their shared range (Reidenbach *et al.* 2012).

As adults, *An. coluzzii* and *An. gambiae* can be found resting in the same houses and blood feeding on the same human hosts. However, in the savannas of West Africa, the aquatic immature stages are associated with alternative breeding sites (Gimonneau *et al.* 2012a). Presumed to be the ancestral species, *An. gambiae* breeds in temporary rain dependent pools and puddles which are largely free of predators. In such sites, *An. gambiae* outcompetes *An. coluzzii* (Diabate *et al.* 2008; Lehmann & Diabate 2008). *An. coluzzii* is found in more stable sites such as rice fields and reservoirs, and is behaviorally more adept at avoiding the predators that abound in such habitats (Diabate *et al.* 2008; Gimonneau *et al.* 2010; Gimonneau *et al.* 2012b). In the rainforests of Central Africa, larval habitat partitioning also occurs, but here it appears to be associated with gradients of urbanization rather than rice agriculture (Kamdem *et al.* 2012). In the Cameroon capital Yaoundé, *An. gambiae* larvae are found in temporary rain dependent pools in the rural suburbs, but *An. coluzzii* breeds in more stable collections of water within the city center, some of which serve as dumping grounds, polluted by the decay of organic material into ammonia and other nitrogenous breakdown products (Kamdem *et al.* 2012) (Fig. 1). As predicted, dose-mortality tests of larvae collected in and around Yaoundé confirmed that *An. coluzzii* is more resistant to ammonia than *An. gambiae* (Tene Fossog *et al.* 2013a). Similarly, dose-mortality tests of larvae sampled along the coast of the Gulf of Guinea also have shown that *An. coluzzii* resists salinity better than *An. gambiae* (B. Tene Fossog, D. Ayala, P. Acevedo, P. Kengne, P. Awono-Ambene, F. Njiokou, C. Antonio-Nkondjio, M. Pombi, N. Besansky, C. Costantini, unpublished data). Taken together with evidence from ecological niche modeling (Costantini *et al.* 2009; Simard *et al.* 2009), the habitat segregation of these species is consistent with an ecological speciation process involving niche expansion by *An. coluzzii* into larval habitats of marginal quality.

Genome scans of divergence between *An. gambiae* and *An. coluzzii* have revealed numerous candidate regions involved in adaptive natural selection (Lawniczak *et al.* 2010; Neafsey *et al.* 2010; Reidenbach *et al.* 2012; White *et al.* 2011), but few clues as to the functional basis of their divergence. We reasoned that differences in gene regulation between these sister taxa may help uncover the molecular basis for species-specific traits. Previously, we compared transcriptional profiles between laboratory-adapted colonies of *An. gambiae* and *An. coluzzii*, and focused mainly on adult females (Cassone *et al.* 2008). Here, we sample co-occurring natural populations of both taxa, and examine mixed sexes at the larval stage. Given that larvae of *An. coluzzii* appear better able to withstand a variety of environmental stressors—both biotic and abiotic—we predicted that larval samples of this species might express stress-responsive genes (*e.g.* immune genes, detoxification genes) at higher levels than their *An. gambiae* counterparts. To explore this question, we performed whole transcriptome profiling of *An. gambiae* and *An. coluzzii* larvae at two developmental stages

(2<sup>nd</sup> and 4<sup>th</sup> instar), sampled from rural and urban habitats in and around Yaoundé, Cameroon that are exclusive to *An. gambiae* (rural sites) or *An. coluzzii* (urban sites) (Kamdem *et al.* 2012). Regardless of ontogenic stage, *An. gambiae* and *An. coluzzii* RNA samples showed differential gene expression in accord with our hypothesis—namely, elevation of transcripts implicated in immunity and detoxification in *An. coluzzii* relative to *An. gambiae*. Our results suggest that the successful exploitation of more stable but marginal larval breeding habitats by *An. coluzzii* may have been facilitated by adaptive transcriptional responses involving the upregulation of stress-responsive genes.

## Materials and methods

### Mosquito sampling

Collections were made in May 2009 from eight localities within a 50 × 50 km area encompassing Yaoundé, Cameroon (Figure 2; Table 1). In this area, intensive longitudinal larval and adult surveys of species composition conducted during one week monthly over the course of one year (May 2008 to April 2009) have revealed spatial segregation of *An. coluzzii* and *An. gambiae* along a gradient of urbanization (Kamdem *et al.* 2012). At opposite ends of the gradient, larval habitats were consistently exclusive to *An. coluzzii* (urban endpoint) or *An. gambiae* (rural endpoint). Based on these prior surveys, localities sampled during the present study (May 2009) at opposite ends of the urbanization gradient were expected to contain only one or the other species. The expected species composition at each locality was confirmed by molecular identifications performed individually on DNA extracted from a subset of 15 larvae, using an rDNA-based PCR assay (Santolamazza *et al.* 2004); all identifications agreed with expectation (Kamdem *et al.* 2012). All larval collections (and larval sacrifices for RNA sampling) were carried out at the same diel time to control for gene expression differences associated with photoperiod (Rund *et al.* 2011). At each locality, larvae were collected by dipper from multiple larval habitats.

### RNA sampling and processing

Individual larvae were killed at the intended sampling times by submersion in tubes containing 1.5 ml of RNAlater RNA stabilization reagent (QIAGEN), to preserve their RNA. All such tubes were transferred from Yaoundé to the University of Notre Dame, where they were stored at –20C prior to RNA extraction. Samples for RNA extraction consisted of 20–40 pooled larvae from the same locality, ontogenic stage, and treatment. Samples from the four rural (or urban) sampling locations (Figure 2; Table 1) were considered as four biological replicates. Total RNA was extracted using the RNeasy Mini Kit (QIAGEN). RNA integrity was assessed by electrophoresis of 1 µl aliquots on 1.5% agarose gels. Total RNA was treated with DNase I (Invitrogen) to remove any contaminating DNA, and concentrated. RNA was quantified using RiboGreen Assays (Molecular Probes/Invitrogen) and the SpectraMAX M2 microplate reader (Molecular Devices). RNA was amplified and converted into double-stranded cDNA using the TransPlex Whole Transcriptome Amplification Kit (Sigma-Aldrich). Quality and quantity of cDNA were assessed using the 2100 Bioanalyzer (Agilent).

## Nimblegen microarray processing

Labeling and amplification of 1 µg cDNA using validated Cy3 dye randomers (TriLink) followed the standard sample labeling protocol (Roche Nimblegen). Using the Nimblegen Hybridization Kit (Roche Nimblegen), 6 µg of labeled cDNA was hybridized to 12 × 135 custom Nimblegen arrays designed from genebuild AgamP3.5 (Cassone *et al.* 2011). Each array interrogated 13,254 predicted AgamP3.5 genes, using at least five duplicated 60-mer probes per gene. Probes were chosen to interrogate gene regions in which there were no nucleotide differences between the species, based on their reference genome assemblies (Lawniczak *et al.* 2010). Arrays were washed using the Nimblegen Wash Buffer Kit and scanned in the UND Genomics Core Facility using the MS 200 Microarray Scanner (Roche NimbleGen). Array image data quality was assessed and raw fluorescence intensity values generated using NimbleScan v2.5 software (Roche NimbleGen). A total of four 12-plex chips, totaling 48 arrays (2 species × 2 ontogenic stages × 3 treatments × 4 replicates) were run (see Experimental design, below). Labeled cDNA samples from the different treatments and instar stages were evenly distributed across the 12-plex chips, and no batch effects were detected. Data from 3 arrays were omitted prior to analysis due to poor data quality (Table 2). The data have been deposited with Array Express under accession #E-MEXP-3147.

XYX files containing the raw expression intensity values were imported into Bioconductor (Gentleman *et al.* 2004). Using custom R scripts (Supplementary File S1), a filter was applied to omit oligos that were misannotated or affected by physical imperfections, as previously described (Cassone *et al.* 2011). The oligo sets were further filtered to contain only those interrogating expressed genes, defined as having an average oligo set intensity value greater than 1500 in any two replicates for at least one condition (species × ontogenic stage × treatment). The final set of filtered expressed oligos was background adjusted, normalized, and summarized using the RMA function. Normalized oligo sets were converted to their respective *An. gambiae* gene ID based on the AgamP3.5 gene build, with oligo sets interrogating the same gene collapsed into a single value based on mean oligo intensity.

## Experimental design and statistical analysis

Larval RNA samples—each with four biological replicates—were derived from two species (*An. gambiae* and *An. coluzzii*), two ontogenic stages (2<sup>nd</sup> and 4<sup>th</sup> instar), and three larval treatments (Table 2). Treatments refer to larval environmental conditions prior to sacrifice for RNA extraction. Larvae that were sacrificed immediately upon collection in the field were considered to be “untreated.” The other two treatments involve larvae that were transferred to the OCEAC insectary in Yaoundé, where they were cultured for 24 h under standard insectary conditions (27±2°C, 85±5% RH, and 12 h light/dark). The water in which they were cultured for 24 h was derived either from the larval habitats from which they were collected (a treatment referred to as “auto-translocation”) or was derived from larval habitats exclusive to the other species (a treatment referred to as “allo-translocation”).

Bayes-moderated *t*-tests in the limma package of R were used to examine gene expression differences between untreated 2<sup>nd</sup>- or 4<sup>th</sup>-instar *An. coluzzii* and *An. gambiae* larvae. Ontogenic stages were considered separately given profound transcriptomic changes during development (Koutsos *et al.* 2007). The significance threshold was defined at  $P < 0.05$

(FDR<0.25)(Benjamini & Hochberg 1995), as our aim was to explore functional enrichment rather than an assessment of individual genes. In addition, we set a minimum threshold log<sub>2</sub>-fold difference of 1.8 between species.

To assess whether short-term exposure to alternative water sources significantly impacted gene expression, we evaluated the 2<sup>nd</sup> and 4<sup>th</sup> instar larval samples cultured for 24 h in an “auto” or “allo” water source (excluding the untreated samples). For each instar stage, we conducted Bayes-moderated *t*-tests to assay treatment-specific differences. We ran two *t*-tests per larval stage that employed slightly different definitions of treatment. In one, treatment was defined as a 24 h exposure to water from breeding sites of the same or opposite species; the hypothesis was that exposure of either species to the opposite water source would perturb gene expression profiles relative to samples maintained in water from their normal breeding sites. In the other, treatment was defined as a 24 h exposure to water from *An. coluzzii* or *An. gambiae* larval habitats; here, the hypothesis was that exposure of either species to potentially polluted water in which *An. coluzzii* was breeding would perturb expression profiles relative those from samples exposed to water from relatively pristine *An. gambiae* habitats.

Functional enrichment of GO and other annotation terms in candidate gene lists was investigated for 2<sup>nd</sup>- and 4<sup>th</sup>-instar larvae using the DAVID functional annotation tool (Huang *et al.* 2009). The DAVID clustering module combines functionally related gene groups into annotation clusters. The clusters are assigned an enrichment score which represents the minus log-transformed geometric mean of the modified Fisher Exact (EASE) Scores within the cluster (Hosack *et al.* 2003; Huang *et al.* 2007). Significantly enriched annotation clusters were defined as those containing at least five genes and a minimum enrichment score of 1.3 ( $P<0.05$ ).

### Quantitative RT-PCR validation

Total RNA from untreated 2<sup>nd</sup> instar larvae of both species was also used for quantitative real-time reverse transcription PCR (qRT-PCR) to validate differential expression detected by microarray. For this purpose, single-stranded cDNA was synthesized from DNase-treated total RNA using TaqMan Reverse Transcription Reagents (Applied Biosystems, Foster City, CA) with random hexamers instead of oligo d(T)<sub>16</sub>. A subset of four genes were tested that by microarray were up-regulated in *An. coluzzii* with a log<sub>2</sub>-fold difference of at least 2.5—(AGAP0)09728, -00424, -04433 and -09380 (Supplementary Table S1). Primers to these targets were designed using Primer Express 3.0 (Applied Biosystems, Foster City, CA) (Supplementary Table S1). Ribosomal protein S7 (AGAP010592) was the internal control reference gene for relative quantification, amplified in a separate tube using forward and reverse primers as given in Dong *et al.* (2006). After optimization, qRT-PCR reactions (15 µl) were performed in triplicate using SYBR Green PCR Master Mix (Applied Biosystems, Foster City, CA), 300 nm of each primer, and an AB7500 Real Time PCR System. After an initial step at 50C for 2 min, cycling conditions were 95C for 15 s, 60C for 1 min, and 95C for 15 s, for 40 cycles. Relative expression was calculated using the comparative C<sub>T</sub> method, after evaluating the kinetic real-time PCR efficiency of target and reference amplifications to ensure they were approximately equal (ABI user bulletin, <http://>

[docs.appliedbiosystems.com/pebiiodocs/04303859.pdf](https://docs.appliedbiosystems.com/pebiiodocs/04303859.pdf)). The qRT-PCR results confirmed the microarray data (Supplementary Table S1). The qRT-PCR data are available as Supplementary Table S2.

### Reanalysis of Affymetrix array data

For reanalysis of the gene expression data of Cassone *et al.* (2008) based on the Affymetrix *Anopheles/Plasmodium* GeneChip, we restricted our consideration to the 4<sup>th</sup> instar larval samples, and performed two *An. coluzzii*-*An. gambiae* contrasts. In the first, we compared the *An. coluzzii* colony from Cameroon (*An. coluzzii* CAM, a derivative of the parental Yaoundé colony established from a district of Yaoundé) to its nearest available geographic counterpart, the *An. gambiae* Pimperena colony (established from Pimperena, Mali). In the second, two colonies from Mali were compared: the same *An. gambiae* Pimperena colony and *An. coluzzii* Mali-NIH. Using the BioMart tool from VectorBase (Megy *et al.* 2012), probes from the Affymetrix array were mapped onto the current AgamP3.7 annotation. Starting with the filtered, background adjusted, normalized and summarized 4<sup>th</sup> instar data, genes with a minimum threshold log<sub>2</sub>-fold difference of 1.8 between species were subject to Bayes-moderated *t*-tests to identify significant differential expression ( $P < 0.05$ ) for each contrast.

### Comparison to differentiated regions of the genome

To ask whether differentially expressed genes also showed signals of nucleotide differentiation, we collected data on such genes from two previous studies: Neafsey *et al.* (2010) and Reidenbach *et al.* (2012). Significantly differentiated genes from Neafsey *et al.* were taken from the supplementary information of that paper, and significantly differentiated genes from Reidenbach *et al.* were supplied by the author. In order to determine whether there was a significant overlap between the genes found to be differentially expressed and those found to be differentiated at the nucleotide level, we used a binomial expectation where the probability of success was defined as the proportion of genes found to be differentially expressed with  $P < 0.05$ . This expectation was used to determine the fraction of all genes with significant nucleotide differentiation that are also expected to show significant differential expression. *P*-values were calculated based on the observed overlap and the probability mass function of a binomial distribution with the given probability of success, calculated separately for each set of differentially expressed genes considered (*e.g.* 2<sup>nd</sup> instar, 4<sup>th</sup> instar, etc.).

## Results

Of the 48 microarrays that were hybridized, 45 passed quality control filters (Table 2). Each microarray interrogated 13,254 genes (based on the AgamP3.5 gene build). Of these, 692 genes were eliminated from further consideration due to missing data (blank intensity values for one or more samples), and an additional 2,579 genes were omitted because they were not detected as expressed (see Methods). Expression patterns from the remaining 9,983 genes were compared between *An. coluzzii* and *An. gambiae* larvae sampled from natural breeding sites in and around Yaoundé, Cameroon.

## Transcriptional divergence between *An. coluzzii* and *An. gambiae* larvae from natural populations

Initially, we examined differential gene expression between *An. coluzzii* and *An. gambiae* larvae that were sacrificed immediately upon sampling from their natural larval habitats. The results are summarized in a volcano plot (Figure 3). Of the 9,983 expressed genes, 8.9% ( $n = 890$ ) differed significantly between species at the 2<sup>nd</sup> instar stage and displayed log<sub>2</sub>-fold-changes between 1.8 and 4.7 (Supplementary File S2). A comparable number of genes (957; 9.6%) were significantly differentially expressed between 4<sup>th</sup> instar larvae of *An. coluzzii* and *An. gambiae* with log<sub>2</sub>-fold-change between 1.8 and 4.4 (Supplementary File S2). The pattern of differential gene expression reflected a bias toward upregulation in *An. coluzzii* at both ontogenic stages, particularly in the younger (2<sup>nd</sup> instar) age class (Figure 3). At the log<sub>2</sub>-fold-change threshold of 1.8, the trend was marginal; approximately 56% ( $n = 499$ ) and 52% ( $n = 501$ ) of genes were more highly expressed in *An. coluzzii* larvae at the 2<sup>nd</sup> and 4<sup>th</sup> instar stage, respectively. However, this trend was more pronounced at higher log<sub>2</sub>-fold-change thresholds. For example, above a fold-change threshold of 2.0, 67% ( $n = 383$ ) and 55% ( $n = 372$ ) of genes in 2<sup>nd</sup> and 4<sup>th</sup> instar larvae were overexpressed in *An. coluzzii*. Increasing the threshold to 3.0, 96% ( $n = 48$ ) and 75% ( $n = 45$ ) of genes were more highly expressed in *An. coluzzii*.

The DAVID data mining tool (Huang *et al.* 2009; Huang *et al.* 2007) was used to classify the genes differentially expressed between *An. coluzzii* and *An. gambiae* into functionally related groups (annotation clusters). For both ontogenic stages we partitioned candidate genes into two lists (up-regulated in *An. coluzzii* or *An. gambiae*) and mined each for annotation clusters with enrichment scores greater than 1.3 containing at least 5 genes (Figure 4).

In *An. coluzzii* larvae at the 2<sup>nd</sup> instar, four significantly enriched annotation clusters were identified among the 499 genes significantly upregulated at this stage. All four clusters contained genes that potentially function in stress responses. In the cofactor metabolism cluster were nine genes, some of which are predicted to encode iron-sulfur or metallo-sulfur proteins with potential roles in an oxidative stress response. Also in this cluster were genes encoding enzymes from the citric acid cycle (TCA), including the pace-making enzyme citrate synthase, as well as aconitase and succinate dehydrogenase (AGAP0012048, -07852, and -00618, respectively). The glutathione S-transferase (*GST*) cluster contained 18 genes, among them *GSTs* whose products are considered vital for the detoxification of xenobiotic compounds (Board & Menon 2013). In addition, there were genes encoding a molecular chaperone and proteins responsible for protein turnover and removal of damaged and abnormal proteins, via the ubiquitin-proteasome system (a ubiquitin ligase, and components of the proteasome complex). The large proteolysis cluster also contained some genes that may function in protein turnover and removal. These include ubiquitin carriers, ubiquitin ligase, and ubiquitin hydrolases. Additionally, nine of the 49 genes in this annotation cluster are immune-related, including clip domain serine proteases (AGAP003250, -03251, -04719, -11789, -11791), serine protease inhibitors (AGAP001375, -03139, -09670) and an autophagy gene (AGAP004023). A connection to immune response is also hinted by the fibrinogen annotation cluster, which had an enrichment score of 7.76—far higher than any



other clusters identified in the gene list from 2<sup>nd</sup> instar *An. coluzzii* larvae. Fibrinogen domain-containing proteins are required for coagulation in vertebrates, but in invertebrates they appear to play a role in defense against pathogens (Hanington & Zhang 2011). In this regard, it is intriguing that two eupolytin genes were overexpressed in 2<sup>nd</sup> instar *An. coluzzii* larvae (and three at the 4<sup>th</sup> instar stage). In the ground beetle *Eupolyphaga sinensis*, the *eupolytin-1* gene encodes a protease that has been shown to hydrolyze fibrinogen and activate plasminogen (Yang *et al.* 2011); in *An. coluzzii*, cleavage of fibrinogen by eupolytin may function in the activation of these defense molecules. One other annotation cluster had a significant enrichment score of 1.33 ( $P < 0.05$ ), although it did not pass our filter requiring membership of at least five genes, and thus is not shown in Figure 4. The three genes in this cluster, identified by DAVID through their relationship to arginine and glutamine metabolism, encode enzymes that catalyze the last three steps in the urea cycle, responsible for the disposal of ammonia in many animal species. These are argininosuccinate synthase, argininosuccinate lyase, and arginase (AGAP003015, -08141, and -08783, respectively). It is noteworthy that in older (4<sup>th</sup> instar) *An. coluzzii* larvae, the only two significantly enriched annotation clusters identified matched two of the clusters from the 2<sup>nd</sup> instar stage (“proteolysis” and “fibrinogen”) and had similar enrichment scores, suggesting that these functions may be upregulated throughout larval development.

In contrast to *An. coluzzii*, functional annotation clustering of candidate genes upregulated in *An. gambiae* identified no significantly enriched functions in 2<sup>nd</sup> instar larvae, among the 391 upregulated genes. The functions enriched in 4<sup>th</sup> instar larvae among 456 upregulated genes—“translation”, “cuticle”, and “ribosome”—may relate to growth and development. Overall, there was no overlap of significantly enriched functions between *An. gambiae* and *An. coluzzii*.

The fact that two of the annotation clusters identified among upregulated genes in *An. coluzzii* at both ontogenic stages contain immune-related genes with potential roles in immune (or environmental stress) response prompted us to manually inspect the candidate gene lists for all genes assigned as immune-related in the reference genome assembly (Bartholomay *et al.* 2010; Waterhouse *et al.* 2007; Waterhouse *et al.* 2010), whether or not they were assigned to clusters of enriched annotation terms by DAVID. Of the 419 immune-related genes in the PEST (AgamP3.7) assembly (and updates to AgamP3.7 by R. Waterhouse, personal communication), 340 were detected as expressed in our experiments. Comparing 2<sup>nd</sup> instar larvae of *An. coluzzii* and *An. gambiae*, 46 were significantly differentially expressed, of which 41 (89%) were upregulated in *An. coluzzii* (Table 3). Although the pattern was not as pronounced in 4<sup>th</sup> instar larvae, of the 42 immune-related genes that were significantly differentially expressed between species, most (30; 71%) were upregulated in *An. coluzzii* (Table 3). The overrepresentation of upregulated immune genes in *An. coluzzii* is significant at both ontogenic stages [2<sup>nd</sup> instar,  $\chi^2 = 35.08$  (1),  $P \ll 0.001$ ; 4<sup>th</sup> instar,  $\chi^2 = 10.16$  (1),  $P = 0.001$ ].

## Short exposure to alternative aqueous environments has no significant impact on the larval transcriptome

Different transcriptional patterns between untreated larvae (*i.e.* sacrificed immediately upon sampling from native habitats) of *An. coluzzii* and *An. gambiae* may have been the result of a plastic response to distinctive abiotic (or biotic) features of their native larval habitats. In particular, the stress responsive genes overexpressed in *An. coluzzii* (*e.g.* immune-related genes, *GSTs*) may have been induced by the presence of xenobiotic chemicals or higher concentrations of organic waste such as ammonia in water from its larval habitat, pollutants that should be largely absent from the *An. gambiae* breeding sites that we sampled. None of the four sampling localities where *An. coluzzii* was breeding were in close proximity to areas used for intense urban farming (in contrast to, *e.g.*, Nkolondom; Tene Fossog *et al.* 2013b), nor were the larval habitats created by the practice of agriculture (*e.g.* furrows), and thus the urban breeding sites were unlikely to be contaminated by insecticides commonly applied by farmers. However, urban larval breeding sites are frequently associated with domestic waste and organic pollutants (Antonio-Nkondjio *et al.* 2011). As an initial assessment of whether the observed transcriptional differences may have resulted from a plastic response by *An. coluzzii* to pollutant exposure, we performed a short-term semi-field translocation experiment.

Upon sampling from their native habitats, *An. coluzzii* and *An. gambiae* larvae at the 2<sup>nd</sup> and 4<sup>th</sup> instar stages were transported to the OCEAC insectary and reared for 24 h, either in water taken from their native habitats (auto-translocation), or in water taken from the habitat of the other species (allo-translocation), before being sacrificed and subject to global transcriptome analysis. We tested for significantly different patterns of gene expression by Bayes-moderated *t*-tests, repeating the test to cover two possible definitions of “treatment”: water source “auto” (native) versus “allo” (foreign), or water source from “*coluzzii*” versus “*gambiae*” larval habitat (see Methods). Strikingly, treatment—regardless of its definition and the ontogenic stage—showed no significant differential expression, with all genes having an FDR = 1. Our results suggest that, at least on the basis of the short (24 h) exposure measured in this experiment, transcriptional differences between species are not adequately explained by a plastic response to acute pollutant exposure in *An. coluzzii* larval habitats.

## Transcriptome divergence between *An. coluzzii* and *An. gambiae* larvae is similar between field and laboratory samples

Previously, we measured transcriptional profiles between laboratory colonies of *An. coluzzii* and *An. gambiae*, including 4<sup>th</sup> instar larvae (Cassone *et al.* 2008). Comparison between those results and the present study is difficult, not only due to the genetic effects of colonization and relaxed selection under artificial rearing conditions, but also due to diverse geographic origins of the mosquitoes and technical differences between microarray platforms. The 2008 study was based on the *Anopheles/Plasmodium* Affymetrix GeneChip, which aside from inherent differences (such as probe length) relative to Nimblegen chips, was designed from the first *An. gambiae* genome assembly and annotation, and thus fails to interrogate numerous genes predicted in more recent annotations. Despite these important caveats, we reasoned that if basic trends in transcriptomic profiles are indeed inherent to species differences, and not merely a reflection of recent local adaptation or transcriptional

plasticity, we should see these same general trends in contrasts between colonies as well as natural populations.

We began by comparing 4<sup>th</sup> instar transcriptomes of the *An. coluzzii* colony from Yaoundé, Cameroon (CAM) to its geographically nearest available *An. gambiae* counterpart, from Mali (Pimperena). Of the 529 genes significantly differentially expressed between 4<sup>th</sup> instar larvae of *An. coluzzii* CAM and *An. gambiae* Pimperena colonies above a log<sub>2</sub>-fold-change of 1.8 (Supplementary File S2), 291 (55%) were up-regulated in *An. coluzzii* CAM. As seen with field-collected larvae, increasing the fold-change threshold increased the proportion of genes that were upregulated in this species relative to *An. gambiae* Pimperena.

A functional enrichment analysis of the genes upregulated in *An. coluzzii* CAM larvae revealed six significant annotation clusters pointing to detoxification and proteolysis functions (Figure 5A; Supplementary File S2). Although genes clustered by these shared annotation terms do not necessarily all function in proteolysis or detoxification, the common enrichment of these themes in field and colony *An. coluzzii* larvae is striking. Of the six annotation clusters, four are associated with detoxification processes: an ion/metal-binding cluster containing numerous cytochrome P450s (*CYP450s*), superoxide dismutase and alkaline phosphatase (45 total genes, enrichment score 3.299), a transferase cluster containing five UDP-glucuronosyl transferases (*UGTs*) (11 genes, score 1.85), a carboxylesterase cluster (7 genes, score 1.85), and finally, a cluster featuring five *GSTs* and a chaperone involved in protein turnover (13 genes, score 1.54). The large proteolysis cluster included immune-related genes such as caspases, clip-domain serine proteases, and a serine protease inhibitor (25 genes, score 1.43). Indeed, the serine protease inhibitor gene *SRPN2* was the most highly over-expressed gene detected in *An. coluzzii* CAM larvae (log<sub>2</sub>-fold change 7.88). As observed for field-collected *An. coluzzii* larvae, the CAM proteolysis cluster also contained eupolytin genes that may function in activation of fibrinogen. Notable for its absence was any detectable enrichment of fibrinogen genes among those significantly upregulated in 4<sup>th</sup> instar larvae from *An. coluzzii* CAM, although four fibrinogen genes were among those upregulated, showing large fold-changes ranging from 2.6 to 4.1 relative to *An. gambiae* Pimperena. Inspection of the probe sets present on the Affymetrix array revealed that 12 of 43 fibrinogen genes annotated in the current *An. gambiae* reference assembly are not interrogated by any probe on this outdated array, suggesting that the absence of enrichment for fibrinogen is inconclusive. Of note, an annotation cluster of C-type lectins that—like fibrinogens—mediate several invertebrate immune responses just missed our significance threshold (8 genes, score 1.2). Moreover, additional genes with potential roles in innate immunity were also over-expressed in *An. coluzzii*, although not clustered by DAVID: two thioester-containing proteins (*TEP1*, *TEP9*), gambicin (*GAMI*), defensin (*DEF3*), a glucan binding protein (*GGBP2*), a scavenger receptor (*SCRBQ1*), and an inhibitor of apoptosis (*IAP7*).

In contrast to the strong enrichment of detoxification and proteolysis annotation terms among 291 genes upregulated in *An. coluzzii* CAM larvae, only one small annotation cluster involving peptidase inhibition was detected among 244 genes upregulated in *An. gambiae* Pimperena larvae (5 genes, enrichment score 1.8). Taken together, the transcriptional profiles of 4<sup>th</sup> instar larvae from CAM and Pimperena colonies are broadly similar to those

measured from field-collected larvae: *An. coluzzii* larvae seem to be characterized by inherently higher levels of expression from classes of genes that may play a role in helping this species overcome the stress of more marginal habitats. This result is striking not merely because this trend seems to have persisted through the colonization process, but also because it has persisted for so long. The *An. coluzzii* CAM colony was derived from a parental Yaoundé colony established in 1988 (Tchuinkam *et al.* 1993), and in the 25 years since then (>250 generations) has been reared under benign laboratory conditions.

Population structure of *An. gambiae* across Africa appears to be quite shallow (Lehmann *et al.* 2003), but *An. coluzzii* populations are notably structured between the forest ecozone of Cameroon and West African savanna populations (Lee *et al.* 2009; Pinto *et al.* 2013; Slotman *et al.* 2007). Despite this, 4<sup>th</sup> instar larvae from the *An. coluzzii* Mali-NIH colony also overexpress immune defense genes—including fibrinogens—relative to their *An. gambiae* Pimperena counterparts (Figure 5B, Supplementary File S2), in keeping with our findings from natural populations in Cameroon. In addition to several fibrinogens, immune-related genes over-expressed in *An. coluzzii* Mali-NIH larvae included six caspases, several clip-domain serine proteases, C-type lectins, and five thioester-containing proteins (including *TEPI*). Among the genes in the enriched “Proteolysis” annotation cluster were four CYP450s. By contrast, and consistent with our results from other contrasts between colonies and population samples from this pair of species, only two annotation clusters were enriched in the set of genes overexpressed in *An. gambiae* Pimperena, and these had no obvious relationship to stress response, neither immune defense nor detoxification.

## Discussion

The results of ecological niche modeling and distribution mapping of *An. gambiae* and *An. coluzzii* in both West and Central Africa are consistent with a process of ecological speciation, in which the more recently derived taxon, *An. coluzzii*, has undergone niche expansion into anthropogenic larval habitats of marginal quality, promoted by larval competition in the original, optimal habitat (Costantini *et al.* 2009; Simard *et al.* 2009). In the savanna, *An. coluzzii* habitats are irrigated rice fields, which would have been a notable landscape feature only after the domestication of African rice from a wild ancestor within the last 2,000–3,000 years (Li *et al.* 2011; Porter 1976). Unlike the temporary pools and puddles exploited by *An. gambiae*, rice fields are relatively stable, biotically complex and rich in a notable environmental stressor: predators. The ecological variables associated with larval habitat segregation in the rainforests of Cameroon differ, but relatively recent niche expansion by *An. coluzzii* into a more stable larval habitat of lower (marginal) quality appears to be a common denominator. In the equatorial forest region, *An. coluzzii* is associated with densely populated urban centers such as Douala and Yaoundé (Antonio-Nkondjio *et al.* 2011; Kamdem *et al.* 2012), whose explosive growth—characterized by inferior housing and poor urban planning, sanitation and drainage of surface water (Antonio-Nkondjio *et al.* 2011)—is a recent phenomenon in the decades since the end of World War II and independence (Franqueville 1984). In these cities, *An. coluzzii* breeds in both temporary and permanent water bodies that, while largely free from top predators (larvivorous fish), may contain variable amounts of other biotic and abiotic environmental stressors, such as competition from *Culex* mosquito larvae and exposure to pollutants from

decaying organic matter, household and industrial waste, or insecticide run-off from urban agriculture (Antonio-Nkondjio *et al.* 2011). Urban breeding sites vary greatly in water quality and other physicochemical characteristics, but on average they contain 2- to 100-fold higher concentrations of ammonia, nitrates, phosphate, sulfate, and other chemicals, have higher turbidity, conductivity and total hardness, and tend to be more alkaline than rural breeding sites (Antonio-Nkondjio *et al.* 2011; Tene Fossog *et al.* 2013a). Particularly in the dry season when suitable temporary and less polluted larval habitats become scarce, *An. coluzzii* can occupy large organically polluted breeding sites in association with competing larvae of *Culex* species, mainly *Cx. quinquefasciatus* (Antonio-Nkondjio *et al.* 2011; Kamdem *et al.* 2012).

The capacity of *An. coluzzii* to invade and exploit newly available anthropogenic habitats of poorer quality, both in the rural savanna and urbanized equatorial forest, prompted the hypothesis that this taxon has rapidly, and perhaps repeatedly, evolved a tolerance to stressors in the environment, mediated by altered gene expression. Our transcriptome profiling of larvae sampled from the field in Cameroon and from laboratory colonies is consistent with this prediction. Differential gene expression between species generally reflected overexpression in *An. coluzzii* relative to *An. gambiae*, and functional annotations enriched among upregulated genes included detoxification and immune defense. Between natural populations from Yaoundé, gene classes overrepresented in *An. coluzzii* included *GSTs*, serine proteases, serine protease inhibitors, and a wide spectrum of other immune-related genes, notably fibrinogens. Between laboratory colonies not subject to the same environmental stresses, immune-related and detoxification gene classes were similarly overexpressed. Previous studies have shown that short-term (as little as 24 h) exposure of susceptible mosquito larvae to sub-lethal doses of various xenobiotics (heavy metal, herbicide, or insecticide) induces overproduction of transcripts encoding detoxification enzymes that can confer larval tolerance and potentially cross-tolerance (David *et al.* 2010; Poupardin *et al.* 2008). However—although not definitive—our data suggest that the upregulation of detoxification and immune gene expression observed in *An. coluzzii* was unlikely to have been a transient (plastic) response to xenobiotics present in water from the larval breeding habitat. First, reciprocal translocation for 24 h in water from larval habitats of the opposite species had no impact on gene expression patterns of either species. Physicochemical data were not recorded from the sampled breeding sites during our experiments; however, such data have been collected previously from a variety of larval habitats in and around Yaoundé, and significant differences were few (Antonio-Nkondjio *et al.* 2011; Tene Fossog *et al.* 2013a; Tene Fossog *et al.* 2012). Importantly, none of the larval habitats sampled for this study were near cultivated areas, thus contamination with insecticides used in farming was very unlikely. Moreover, had the presence of other pollutants in *An. coluzzii* habitats been directly responsible for inducing gene upregulation in this species, our expectation would be that exposure of *An. gambiae* larvae to water from these habitats should have resulted in upregulation of stress-responsive genes in this species (assuming that 24 h was a sufficiently long exposure), yet this was not observed. Second, the expression profiles of wild and colonized *An. coluzzii* from Yaoundé were broadly similar, despite laboratory culture of the Yaoundé colony since 1988 (Tchuinkam *et al.* 1993). Thus, our working hypothesis is that the higher expression of detoxification and immune defense

genes in *An. coluzzii* relative to *An. gambiae* reflects a constitutive increase in basal expression levels, as observed recently for immune activation genes in *Aedes aegypti* populations refractory to Dengue virus (Sim *et al.* 2013). More definitive evidence for this working hypothesis could come from common garden experiments in the laboratory, using the F1 progeny of field-collected females of both species.

To the extent that expression of detoxification and immune defense genes is inherently higher in *An. coluzzii* relative to *An. gambiae*, it may be an adaptive response to environmental conditions that have been repeatedly encountered. If this is the case, it is important to stress that there is a dearth of information to support inferences about the ultimate sources of selective pressures acting on *An. coluzzii* historically, as so little is known of its biogeographic history, ecology, or behavior. One obvious recent (and potentially confounding) factor, certainly the one that has commanded the most attention, is selection by insecticides used to control agricultural pests. Through runoff, insecticides intended to target pests of crops also contaminate the furrows and irrigation ditches associated with agriculture (“cultivated sites”) where mosquitoes can breed. The emergence of insecticide resistance in non-target insects such as mosquitoes has long been recognized (Lines 1988). With respect to the malaria vectors *An. gambiae*, *An. coluzzii*, and *An. arabiensis* (another sibling species), this association has been observed not only with traditional agriculture (Diabate *et al.* 2002; Muller *et al.* 2008), but also with the recent and rapidly expanding practice of intensive urban agriculture in African cities (Antonio-Nkondjio *et al.* 2011; Jones *et al.* 2012; Yadouleton *et al.* 2009). Urban *An. coluzzii* populations are resistant to multiple insecticide classes, and an important mechanism of insecticide tolerance appears to be metabolic, mediated by enzyme-based detoxification (Djouaka *et al.* 2008; Nwane *et al.* 2013; Oduola *et al.* 2012).

Previous microarray analyses of DDT-resistant or permethrin-resistant adult *An. coluzzii*, collected as larvae from a variety of breeding sites in urban and agricultural areas of southern Benin, Nigeria, and Cameroon, revealed over-expression of *CYP450s*, *GSTs*, carboxylesterases, and *UGTs* (Djouaka *et al.* 2008; Tene Fossog *et al.* 2013b). The association of over-expressed detoxification genes with insecticide resistance implicates these genes in metabolic resistance, but as only a handful of individual genes have been shown to metabolize insecticides and many of them may have unrelated functions, caution is merited and firm conclusions await direct experimental validation. Table 4 presents a compilation of the detoxification genes found in the present study to be over-expressed in field-collected or colonized *An. coluzzii* larvae. It is noteworthy that only one *CYP450* (*CYP325C2*, over-expressed in 4<sup>th</sup> instar field-collected larvae) has been associated previously with insecticide resistance in *An. coluzzii* (Tene Fossog *et al.* 2013b), and of all the detoxification genes over-expressed in our field collections, only one (*GSTE2*) has been validated to metabolize DDT and permethrin (Daborn *et al.* 2012; Ranson *et al.* 2001). The transcriptional profile of the field-collected *An. coluzzii* samples does not appear to match that expected for a highly resistant strain (in contrast to the *An. coluzzii* CAM colony, which over-expresses *CYP6P3* and *CYP6M2*, among others). We did not test our larval samples for insecticide resistance, but two recent surveys conducted in Cameroon (in Yaoundé and the coastal city of Douala) reported insecticide resistance bioassays on mosquitoes collected

from cultivated sites or polluted/non-polluted larval habitats away from cultivated sites. A panel of insecticides of different classes were applied, either to larvae (Tene Fossog *et al.* 2012) or to emerged adults (Antonio-Nkondjio *et al.* 2011). Importantly, both studies identified substantial heterogeneity in levels of resistance to DDT, permethrin, deltamethrin, and bendiocarb, based on the type of larval habitat. In good agreement with a previous survey of permethrin resistance in Benin and Nigeria (Djouaka *et al.* 2008), the level of insecticide resistance to diagnostic concentrations was much higher in mosquitoes collected from urban agricultural sites versus other urban sites polluted (or not) with organic waste or petroleum products—even if the cultivated versus non-cultivated larval habitats were separated by less than 10 km. Taken together, these data suggest that agricultural insecticides play a very important selective role, affecting all *An. coluzzii* in Yaoundé to some degree, even those derived from uncultivated breeding sites. However, it is not clear that insecticides are the only selective factor, or more importantly, the initial driver responsible for overexpression of metabolic detoxification genes in *An. coluzzii*. Alternative environmental factors, biotic and/or abiotic, may have played supporting or even primary roles (Nkya *et al.* 2013).

A second selective factor that may play a role in transcriptional divergence between *An. coluzzii* and *An. gambiae*, especially in detoxification genes, could be the toxic effects of ammonia, a byproduct of the breakdown of organic waste present in polluted urban larval habitats. Indeed, in a previous study we measured higher levels of ammonia in such habitats, and higher tolerance of ammonia by *An. coluzzii* larvae breeding in those habitats in urban Yaoundé, relative to *An. gambiae* breeding in pristine larval habitats in the rural outskirts of Yaoundé (Tene Fossog *et al.* 2013a). Disposal of dietary ammonia is a major and universal problem faced by all adult female mosquitoes during digestion of the proteinaceous blood meal, and by larvae that ingest excess protein relative to their growth requirements (Scaraffia *et al.* 2005; von Dungern & Briegel 2001; Weihrauch *et al.* 2012). Mosquitoes utilize multiple metabolic routes for the disposal of nitrogenous waste, including the synthesis of urea catalyzed by arginase (Isoe & Scaraffia 2013; Scaraffia *et al.* 2005; Scaraffia *et al.* 2008; von Dungern & Briegel 2001), one of three enzymes from the urea (ornithine) cycle found to be significantly overexpressed in *An. coluzzii* larvae in this study. However, because the typical urea cycle is apparently absent in mosquitoes due to lack of the ornithine carbamoyltransferase gene (Zdobnov *et al.* 2002), this pathway has been considered non-functional for ammonia disposal (Scaraffia *et al.* 2005). Instead, environmental ammonia ingested by larvae may be transported and directly excreted from the Malpighian tubules, hindgut, and anal papillae (Donini & O'Donnell 2005; Weihrauch *et al.* 2012). Neither putative ammonium transporter proteins nor aquaporins were detected as overexpressed in *An. coluzzii*. However, V-type H<sup>+</sup>-ATPase (V-ATPase) is known to play a crucial role in transepithelial ammonia transport in many systems (Weihrauch *et al.* 2012). In both 2<sup>nd</sup> and 4<sup>th</sup> instar samples of *An. coluzzii*, the V-ATPase 21kDa subunit (AGAP009334) was significantly over-expressed by log<sub>2</sub>-fold change of ~3. Consistent with a role for this V-ATPase in excretion, MozAtlas (Baker *et al.* 2011) reports statistically significant enrichment of its expression in the adult Malpighian tubules. These observations merit more detailed follow-up study.

A third selective factor potentially contributing to transcriptional divergence between *An. coluzzii* and *An. gambiae* may be differential exposure to pathogenic fungi, bacteria, and parasites found in their distinctive larval breeding sites. In a previous study employing genome scans of divergence between natural populations of these species, we uncovered a region of exceptional divergence on chromosome 3L in samples from West Africa, and implicated the *TEPI* gene, whose complement C3-like product has antiparasitic and antibacterial activity (White *et al.* 2011). DNA sequence analysis of these population samples suggested that an allelic variant of *TEPI* unique to *An. coluzzii* had swept to fixation in Mali and Burkina Faso, and was spreading into neighboring Ghana. This allele, closely related but distinct from previously characterized *TEPI* resistance alleles, conferred resistance to *Plasmodium* parasites that invade and replicate within the adult female mosquito (White *et al.* 2011). However, given the relatively low rates and intensities of *Plasmodium* infection in nature for both *An. coluzzii* and *An. gambiae* adult female populations, we speculated that the most likely source of pathogen-mediated selection for resistance came from the *An. coluzzii* larval habitat—a longer-lasting and more biotically diverse aquatic milieu than that exploited by *An. gambiae*, thus presumably harboring distinctive pathogen populations. Although the particular West African resistance allele described in White *et al.* (2011) is apparently absent from Central Africa, the *TEPI* gene was over-expressed in our 2<sup>nd</sup> instar samples of *An. coluzzii* from Cameroon (log<sub>2</sub>-fold change 2.4), together with a plethora of other immune-related genes, most notably fibrinogens (Table 3, Supplementary File S2). The *TEPI* gene was also the second-ranked over-expressed gene in 4<sup>th</sup> instar larvae from the CAM colony (compared to *An. gambiae* Pimperena; log<sub>2</sub>-fold change 7.1) and the top-ranked over-expressed gene in the Mali-NIH colony (compared to *An. gambiae* Pimperena; log<sub>2</sub>-fold change 8.8), closely followed by four other *TEP* genes (log<sub>2</sub>-fold changes from 6.1-4.8 in Mali-NIH) and two C-type lectins (log<sub>2</sub>-fold changes from 6.1-4.0 in Mali-NIH). A non-coincidental relationship between immune and stress responses has recently been highlighted in the model organism *Drosophila* (Davies *et al.* 2012). Emerging evidence from the fruitfly suggests that apparently unrelated stresses (*e.g.* immune, oxidative, osmotic, xenobiotic) are both sensed and countered through overlapping cellular responses mediated by the Malpighian tubule, the insect equivalent of the vertebrate kidney and liver (Davies *et al.* 2012). If this immune and stress “cross talk” applies more broadly to mosquitoes as expected, it may provide a conceptual framework for understanding the functional enrichment of both immune-related and detoxification genes among those over-expressed in *An. coluzzii*.

A previous study of transcriptional divergence between *An. coluzzii* and *An. gambiae* was conducted on adults that emerged in the OCEAC insectary from larval collections made at two localities 10 km apart in Yaounde (Nkolbisson) and on its outskirts (Nkolondom) (Aguilar *et al.* 2010). These localities differ from those sampled in the present study in that both, particularly Nkolondom, are sites of urban farming (*e.g.* cultivated sites where larval habitats are subject to contamination by insecticides). Both *An. coluzzii* and *An. gambiae* were collected at each location, although they were present in highly skewed proportions. Based on competitive hybridizations to an Agilent glass slide microarray, the authors reported that transcriptional divergence between the same species across the two localities was ten times greater than transcriptional divergence between different species across the



same two localities. This conclusion apparently contradicts our findings in the present study, but the source of the discrepancy is difficult to discern, not least because these previous microarray data are not in a public database. However, the basis of their conclusion seems to rest on the observation that 61 transcripts were in common between different species across sites, while only 6 transcripts were in common between samples of the same species across sites (Aguilar *et al.* 2010), a finding that may not be robust. Possible factors contributing to the different conclusions may be: probes on the Agilent array that do not interrogate the same genes as our Nimblegen array; technical differences between platforms; differences in statistical methods, analytical approach, and data interpretation; differences in life stage analyzed; and differences in sampling localities (cultivated versus uncultivated). Two common findings between the studies are that in the urban Yaoundé locality, *An. coluzzii* shows a higher proportion of over-expressed genes, and those genes are enriched for immune functions.

Given our results on the suite of genes that are differentially expressed between *An. coluzzii* and *An. gambiae*—genes that are presumably evolving in response to environmental stressors—we asked whether these same genes were identified in previous genome scans for regions under recent positive selection. In our earlier study of gene expression in laboratory colonies (Cassone *et al.* 2008), we found a slight enrichment for differentially expressed genes in the pericentromeric region of the X chromosome, a large area that had been identified as under selection by some of the first genome scans in *An. gambiae* (Stump *et al.* 2005; Turner & Hahn 2007; Turner *et al.* 2005). However, using two more recent, high-resolution studies of nucleotide differentiation between these two species (Neafsey *et al.* 2010; Reidenbach *et al.* 2012), we found no significant enrichment for differentially expressed genes among those genes in windows of high population divergence ( $P > 0.10$  for all sets of differentially expressed genes, for comparisons to both datasets). We also repeated these analyses using the genes adjacent to the previously identified targets of natural selection, to account for the possibility that *cis*-acting changes were not located immediately within the genes they regulated. Again, we found no significant enrichment for differentially expressed genes. Even if differences in gene expression between the two species are driven by natural selection, the mutations responsible for regulating such differences may occur in the coding regions of genes that show no differential expression (i.e. *trans*-acting regulators). Therefore, we would only expect a significant overlap between these two types of data if most differences are driven by *cis*-regulation (cf. Fay & Wittkopp 2008). Our results suggest that this is not the case.

Adaptation in the mosquito species considered here may be representative of range expansions in other systems. For instance, *Drosophila mojavensis* is a cactophilic species of the North American desert, comprised of four recently diverged (<0.5 MYA) subspecies adapted to different cactus hosts with differing sets of allelochemicals (Matzkin 2014, and refs therein). Similar to our findings, transcriptional changes in detoxification genes appear to be associated with cactus host shifts—including those known to confer resistance to DDT (e.g. *GSTE2*) (Matzkin 2012; Matzkin & Markow 2013). However, as was also true for the *D. mojavensis* transcriptomic analysis, the fraction of detoxification (and in our case, immune) genes among all differentially expressed genes is small, and it is likely that the

adaptive divergence of *An. coluzzii* and *An. gambiae* involved selection on many additional genes. Although the designation of *An. coluzzii* and *An. gambiae* as separate species may be controversial due to ongoing hybridization and introgression, known differences between them at the larval stage (in preferred larval habitat, ammonia tolerance, development time, and predator avoidance behavior) and at the adult stage (in mate recognition and swarming behaviors) (Dabire *et al.* 2013; Diabate *et al.* 2009; Gimonneau *et al.* 2010; Gimonneau *et al.* 2012a; Gimonneau *et al.* 2012b; Kamdem *et al.* 2012; Lehmann & Diabate 2008; Pennetier *et al.* 2010; Roux *et al.* 2013; Tene Fossog *et al.* 2013a) suggest that their divergence involved multifarious selection on multiple genetic targets that presumably include chemosensory genes and regulators of development. Even for closely related species with few morphological differences, these examples suggest that there can—or must—be multiple adaptive differences between them in order to maintain niche differentiation and ecological specialization.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank R. Waterhouse for providing an updated set of immune-related genes, D. Lawson for assistance with VectorBase, I. Morlais for providing information about OCEAC colonies, and three anonymous reviewers. We are grateful to the OCEAC Malaria Research Laboratory staff for excellent technical and field support. Financial support was provided by NIH grant R01 AI63508 to NJB.

## References

- Aguilar R, Simard F, Kamdem C, et al. Genome-wide analysis of transcriptomic divergence between laboratory colony and field *Anopheles gambiae* mosquitoes of the M and S molecular forms. *Insect Molecular Biology*. 2010; 19:695–705. [PubMed: 20738426]
- Antonio-Nkondjio C, Fossog BT, Ndo C, et al. *Anopheles gambiae* distribution and insecticide resistance in the cities of Douala and Yaounde (Cameroon): influence of urban agriculture and pollution. *Malar J*. 2011; 10:154. [PubMed: 21651761]
- Baker DA, Nolan T, Fischer B, et al. A comprehensive gene expression atlas of sex- and tissue-specificity in the malaria vector, *Anopheles gambiae*. *BMC Genomics*. 2011; 12:296. [PubMed: 21649883]
- Bartholomay LC, Waterhouse RM, Mayhew GF, et al. Pathogenomics of *Culex quinquefasciatus* and meta-analysis of infection responses to diverse pathogens. *Science*. 2010; 330:88–90. [PubMed: 20929811]
- Benjamini Y, Hochberg Y. Controlling the false discovery rate – A practical and powerful approach to multiple testing. *J Roy Stat Soc B*. 1995; 57:289–300.
- Bickford D, Lohman DJ, Sodhi NS, et al. Cryptic species as a window on diversity and conservation. *Trends in ecology & evolution*. 2007; 22:148–155. [PubMed: 17129636]
- Board PG, Menon D. Glutathione transferases, regulators of cellular metabolism and physiology. *Biochimica et Biophysica Acta*. 2013; 1830:3267–3288. [PubMed: 23201197]
- Borneman AR, Gianoulis TA, Zhang ZD, et al. Divergence of Transcription Factor Binding Sites Across Related Yeast Species. *Science*. 2007; 317:815–819. [PubMed: 17690298]
- Britten RJ, Davidson EH. Gene regulation for higher cells: a theory. *Science*. 1969; 165:349–357. [PubMed: 5789433]

- Britten RJ, Davidson EH. Repetitive and non-repetitive DNA sequences and a speculation on the origins of evolutionary novelty. *The Quarterly review of biology*. 1971; 46:111–138. [PubMed: 5160087]
- Cassone BJ, Molloy MJ, Cheng C, et al. Divergent transcriptional response to thermal stress by *Anopheles gambiae* larvae carrying alternative arrangements of inversion 2La. *Molecular Ecology*. 2011; 20:2567–2580. [PubMed: 21535279]
- Cassone BJ, Mouline K, Hahn MW, et al. Differential gene expression in incipient species of *Anopheles gambiae*. *Molecular Ecology*. 2008; 17:2491–2504. [PubMed: 18430144]
- Chesson P. Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics*. 2000; 31:343–366.
- Coetzee M, Hunt RH, Wilkerson R, et al. *Anopheles coluzzii* and *Anopheles amharicus*, new members of the *Anopheles gambiae* complex. *Zootaxa*. 2013; 3619:246–274.
- Costantini C, Ayala D, Guelbeogo WM, et al. Living at the edge: biogeographic patterns of habitat segregation conform to speciation by niche expansion in *Anopheles gambiae*. *BMC Ecol*. 2009; 9:16. [PubMed: 19460144]
- Dabire, KR.; Sawadodgo, S.; Diabate, A., et al. Medical and Veterinary Entomology. 2013. Assortative mating in mixed swarms of the mosquito *Anopheles gambiae* s.s. M and S molecular forms, in Burkina Faso, West Africa.
- Daborn PJ, Lumb C, Harrop TW, et al. Using *Drosophila melanogaster* to validate metabolism-based insecticide resistance from insect pests. *Insect Biochemistry and Molecular Biology*. 2012; 42:918–924. [PubMed: 23023059]
- David JP, Coissac E, Melodelima C, et al. Transcriptome response to pollutants and insecticides in the dengue vector *Aedes aegypti* using next-generation sequencing technology. *BMC Genomics*. 2010; 11:216. [PubMed: 20356352]
- David JP, Ismail HM, Chandor-Proust A, Paine MJ. Role of cytochrome P450s in insecticide resistance: impact on the control of mosquito-borne diseases and use of insecticides on Earth. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2013; 368:20120429.
- Davies SA, Overend G, Sebastian S, et al. Immune and stress response ‘cross-talk’ in the *Drosophila* Malpighian tubule. *Journal of insect physiology*. 2012; 58:488–497. [PubMed: 22306292]
- della Torre A, Fanello C, Akogbeto M, et al. Molecular evidence of incipient speciation within *Anopheles gambiae* s.s. in West Africa. *Insect Molecular Biology*. 2001; 10:9–18. [PubMed: 11240632]
- della Torre A, Tu ZJ, Petrarca V. On the distribution and genetic differentiation of *Anopheles gambiae* s.s. molecular forms. *Insect Biochemistry and Molecular Biology*. 2005; 35:755–769. [PubMed: 15894192]
- Diabate A, Baldet T, Chandre F, et al. The role of agricultural use of insecticides in resistance to pyrethroids in *Anopheles gambiae* s.l. in Burkina Faso. *American Journal of Tropical Medicine and Hygiene*. 2002; 67:617–622. [PubMed: 12518852]
- Diabate A, Dabire KR, Heidenberger K, et al. Evidence for divergent selection between the molecular forms of *Anopheles gambiae*: role of predation. *BMC Evol Biol*. 2008; 8:5. [PubMed: 18190719]
- Diabate A, Dabire RK, Kengne P, et al. Mixed swarms of the molecular M and S forms of *Anopheles gambiae* (Diptera: Culicidae) in sympatric area from Burkina Faso. *Journal of Medical Entomology*. 2006; 43:480–483. [PubMed: 16739404]
- Diabate A, Dabire RK, Kim EH, et al. Larval development of the molecular forms of *Anopheles gambiae* (Diptera: Culicidae) in different habitats: a transplantation experiment. *Journal of Medical Entomology*. 2005; 42:548–553. [PubMed: 16119542]
- Diabate A, Dabire RK, Millogo N, Lehmann T. Evaluating the effect of postmating isolation between molecular forms of *Anopheles gambiae* (Diptera: Culicidae). *Journal of Medical Entomology*. 2007; 44:60–64. [PubMed: 17294921]
- Diabate A, Dao A, Yaro AS, et al. Spatial swarm segregation and reproductive isolation between the molecular forms of *Anopheles gambiae*. *Proc Biol Sci*. 2009; 276:4215–4222. [PubMed: 19734189]

- Djouaka RF, Bakare AA, Coulibaly ON, et al. Expression of the cytochrome P450s, CYP6P3 and CYP6M2 are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae* s.s. from Southern Benin and Nigeria. *BMC Genomics*. 2008; 9:538. [PubMed: 19014539]
- Dong Y, Aguilar R, Xi Z, et al. *Anopheles gambiae* immune responses to human and rodent *Plasmodium* parasite species. *PLoS Pathog*. 2006; 2:e52. [PubMed: 16789837]
- Donini A, O'Donnell MJ. Analysis of Na<sup>+</sup>, Cl<sup>-</sup>, K<sup>+</sup>, H<sup>+</sup> and NH<sub>4</sub><sup>+</sup> concentration gradients adjacent to the surface of anal papillae of the mosquito *Aedes aegypti*: application of self-referencing ion-selective microelectrodes. *The Journal of experimental biology*. 2005; 208:603–610. [PubMed: 15695753]
- Fay JC, Wittkopp PJ. Evaluating the role of natural selection in the evolution of gene regulation. *Heredity*. 2008; 100:191–199. [PubMed: 17519966]
- Franqueville, A. Collection Mémoires. Vol. 104. ORSTOM; Paris: 1984. Yaoundé, construire une capitale.
- Gentleman RC, VJC, DMB, et al. Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol*. 2004; 5:R80. [PubMed: 15461798]
- Gimonneau G, Bouyer J, Morand S, et al. A behavioral mechanism underlying ecological divergence in the malaria mosquito *Anopheles gambiae*. *Behav Ecol*. 2010; 21:1087–1092. [PubMed: 22476108]
- Gimonneau G, Pombi M, Choisy M, et al. Larval habitat segregation between the molecular forms of the mosquito, *Anopheles gambiae* in a rice field area of Burkina Faso, West Africa. *Medical and Veterinary Entomology*. 2012a; 26:9–17. [PubMed: 21501199]
- Gimonneau G, Pombi M, Dabire RK, et al. Behavioural responses of *Anopheles gambiae* sensu stricto M and S molecular form larvae to an aquatic predator in Burkina Faso. *Parasit Vectors*. 2012b; 5:65. [PubMed: 22463735]
- Hahn MW, White BJ, Muir CJ, Besansky NJ. No evidence for biased co-transmission of speciation islands in *Anopheles gambiae*. *Phil Trans R Soc B*. 2012; 367:374–384. [PubMed: 22201167]
- Hanington PC, Zhang S-M. The primary role of fibrinogen-related proteins in invertebrates is defense, not coagulation. *J Innate Immun*. 2011; 3:17–27. [PubMed: 21063081]
- Hosack DA, Dennis G Jr, Sherman BT, Lane HC, Lempicki RA. Identifying biological themes within lists of genes with EASE. *Genome Biol*. 2003; 4:R70. [PubMed: 14519205]
- Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*. 2009; 4:44–57. [PubMed: 19131956]
- Huang DW, Sherman BT, Tan Q, et al. The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol*. 2007; 8:R183. [PubMed: 17784955]
- Isoe J, Scaraffia PY. Urea synthesis and excretion in *Aedes aegypti* mosquitoes are regulated by a unique cross-talk mechanism. *PloS one*. 2013; 8:e65393. [PubMed: 23755226]
- Jones CM, Toe HK, Sanou A, et al. Additional selection for insecticide resistance in urban malaria vectors: DDT resistance in *Anopheles arabiensis* from Bobo-Dioulasso, Burkina Faso. *PloS one*. 2012; 7:e45995. [PubMed: 23049917]
- Kamdem C, Tene Fossog B, Simard F, et al. Anthropogenic habitat disturbance and ecological divergence between incipient species of the malaria mosquito *Anopheles gambiae*. *PloS one*. 2012; 7:e39453. [PubMed: 22745756]
- Koutsos AC, Blass C, Meister S, et al. Life cycle transcriptome of the malaria mosquito *Anopheles gambiae* and comparison with the fruitfly *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104:11304–11309. [PubMed: 17563388]
- Lawniczak MK, Emrich SJ, Holloway AK, et al. Widespread divergence between incipient *Anopheles gambiae* species revealed by whole genome sequences. *Science*. 2010; 330:512–514. [PubMed: 20966253]
- Lee Y, Cornel AJ, Meneses CR, et al. Ecological and genetic relationships of the Forest-M form among chromosomal and molecular forms of the malaria vector *Anopheles gambiae sensu stricto*. *Malar J*. 2009; 8:75. [PubMed: 19383163]

- Lehmann T, Diabate A. The molecular forms of *Anopheles gambiae*: a phenotypic perspective. *Infect Genet Evol.* 2008; 8:737–746. [PubMed: 18640289]
- Lehmann T, Licht M, Elissa N, et al. Population Structure of *Anopheles gambiae* in Africa. *Journal of Heredity.* 2003; 94:133–147. [PubMed: 12721225]
- Leibold MA, McPeck MA. Coexistence of the niche and neutral perspectives in community ecology. *Ecology.* 2006; 87:1399–1410. [PubMed: 16869414]
- Li YF, Costello JC, Holloway AK, Hahn MW. “Reverse ecology” and the power of population genomics. *Evolution.* 2008; 62:2984–2994. [PubMed: 18752601]
- Li ZM, Zheng XM, Ge S. Genetic diversity and domestication history of African rice (*Oryza glaberrima*) as inferred from multiple gene sequences. *Theor Appl Genet.* 2011; 123:21–31. [PubMed: 21400109]
- Lines JD. Do agricultural insecticides select for insecticide resistance in mosquitoes? A look at the evidence. *Parasitology today.* 1988; 4:S17–S20. [PubMed: 15463085]
- Matzkin LM. Population transcriptomics of cactus host shifts in *Drosophila mojavensis*. *Molecular Ecology.* 2012; 21:2428–2439. [PubMed: 22512269]
- Matzkin LM. Ecological Genomics of Host Shifts in *Drosophila mojavensis*. *Advances in Experimental Medicine and Biology.* 2014; 781:233–247. [PubMed: 24277303]
- Matzkin, LM.; Markow, TA. Transcriptional differentiation across the four cactus host races of *Drosophila mojavensis*. In: Michalak, P., editor. *Speciation: Natural Processes, Genetics and Biodiversity*. Nova Science Publishers Inc; New York: 2013. p. 119-135.
- Megy K, Emrich SJ, Lawson D, et al. VectorBase: improvements to a bioinformatics resource for invertebrate vector genomics. *Nucleic Acids Research.* 2012; 40:D729–734. [PubMed: 22135296]
- Montero-Pau J, Ramos-Rodriguez E, Serra M, Gomez A. Long-term coexistence of rotifer cryptic species. *PloS one.* 2011; 6:e21530. [PubMed: 21738691]
- Muller P, Chouaibou M, Pignatelli P, et al. Pyrethroid tolerance is associated with elevated expression of antioxidants and agricultural practice in *Anopheles arabiensis* sampled from an area of cotton fields in Northern Cameroon. *Molecular Ecology.* 2008; 17:1145–1155. [PubMed: 18179425]
- Neafsey DE, Lawniczak MK, Park DJ, et al. SNP genotyping defines complex gene-flow boundaries among African malaria vector mosquitoes. *Science.* 2010; 330:514–517. [PubMed: 20966254]
- Nkya TE, Akhouayri I, Kisinza W, David JP. Impact of environment on mosquito response to pyrethroid insecticides: facts, evidences and prospects. *Insect Biochemistry and Molecular Biology.* 2013; 43:407–416. [PubMed: 23123179]
- Nosil P, Funk DJ, Ortiz-Barrientos D. Divergent selection and heterogeneous genomic divergence. *Molecular Ecology.* 2009; 18:375–402. [PubMed: 19143936]
- Nowick K, Gernat T, Almaas E, Stubbs L. Differences in human and chimpanzee gene expression patterns define an evolving network of transcription factors in brain. *Proceedings of the National Academy of Sciences of the United States of America.* 2009; 106:22358–22363. [PubMed: 20007773]
- Nwane P, Etang J, Chouasmall yi UM, et al. Multiple insecticide resistance mechanisms in *Anopheles gambiae* s.l. populations from Cameroon, Central Africa. *Parasites & vectors.* 2013; 6:41. [PubMed: 23433176]
- Oduola AO, Idowu ET, Oyebola MK, et al. Evidence of carbamate resistance in urban populations of *Anopheles gambiae* s.s. mosquitoes resistant to DDT and deltamethrin insecticides in Lagos, South-Western Nigeria. *Parasites & vectors.* 2012; 5:116. [PubMed: 22686575]
- Pennetier C, Warren B, Dabire KR, Russell IJ, Gibson G. “Singing on the wing” as a mechanism for species recognition in the malarial mosquito *Anopheles gambiae*. *Current Biology.* 2010; 20:131–136. [PubMed: 20045329]
- Pfenninger M, Schwenk K. Cryptic animal species are homogeneously distributed among taxa and biogeographical regions. *BMC evolutionary biology.* 2007; 7:121. [PubMed: 17640383]
- Pinto J, Egyir-Yawson A, Vicente J, et al. Geographic population structure of the African malaria vector *Anopheles gambiae* suggests a role for the forest-savannah biome transition as a barrier to gene flow. *Evolutionary applications.* 2013; 6:910–924. [PubMed: 24062800]


- Porteres, R. African cereals: *Eleusine*, fonio, black fonio, teff, *Brachiaria*, *Paspalum*, *Pennisetum* and African rice. In: Harlan, JR.; de Wet, JMJ.; Stemler, ABL., editors. *Origins of African plant domestication*. Mouton Publishers; The Hague: 1976. p. 409-452.
- Poupardin R, Reynaud S, Strode C, et al. Cross-induction of detoxification genes by environmental xenobiotics and insecticides in the mosquito *Aedes aegypti*: impact on larval tolerance to chemical insecticides. *Insect Biochemistry and Molecular Biology*. 2008; 38:540–551. [PubMed: 18405832]
- Ranson H, Rossiter L, Ortelli F, et al. Identification of a novel class of insect glutathione S-transferases involved in resistance to DDT in the malaria vector *Anopheles gambiae*. *The Biochemical journal*. 2001; 359:295–304. [PubMed: 11583575]
- Reidenbach KR, Neafsey DE, Costantini C, et al. Patterns of genomic differentiation between ecologically differentiated M and S forms of *Anopheles gambiae* in West and Central Africa. *Genome Biology and Evolution*. 2012; 4:1202–1212. [PubMed: 23132896]
- Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the evolution of gene regulation. *Nature reviews Genetics*. 2012; 13:505–516.
- Roux O, Diabate A, Simard F. Divergence in threat-sensitivity among aquatic larvae of cryptic mosquito species. *The Journal of animal ecology*. 2013
- Rund SS, Hou TY, Ward SM, Collins FH, Duffield GE. Genome-wide profiling of diel and circadian gene expression in the malaria vector *Anopheles gambiae*. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108:E421–430. [PubMed: 21715657]
- Santolamazza F, Della Torre A, Caccone A. Short report: A new polymerase chain reaction-restriction fragment length polymorphism method to identify *Anopheles arabiensis* from *An. gambiae* and its two molecular forms from degraded DNA templates or museum samples. *American Journal of Tropical Medicine and Hygiene*. 2004; 70:604–606. [PubMed: 15210999]
- Scaraffia PY, Isoe J, Murillo A, Wells MA. Ammonia metabolism in *Aedes aegypti*. *Insect Biochemistry and Molecular Biology*. 2005; 35:491–503. [PubMed: 15804581]
- Scaraffia PY, Tan G, Isoe J, et al. Discovery of an alternate metabolic pathway for urea synthesis in adult *Aedes aegypti* mosquitoes. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105:518–523. [PubMed: 18182492]
- Sim S, Jupatanakul N, Ramirez JL, et al. Transcriptomic profiling of diverse *Aedes aegypti* strains reveals increased basal-level immune activation in dengue virus-refractory populations and identifies novel virus-vector molecular interactions. *PLoS neglected tropical diseases*. 2013; 7:e2295. [PubMed: 23861987]
- Simard F, Ayala D, Kamdem GC, et al. Ecological niche partitioning between the M and S molecular forms of *Anopheles gambiae* in Cameroon: the ecological side of speciation. *BMC Ecol*. 2009; 9:17. [PubMed: 19460146]
- Slotman MA, Tripet F, Cornel AJ, et al. Evidence for subdivision within the M molecular form of *Anopheles gambiae*. *Molecular Ecology*. 2007; 16:639–649. [PubMed: 17257119]
- Storz JF. Using genome scans of DNA polymorphism to infer adaptive population divergence. *Molecular Ecology*. 2005; 14:671–688. [PubMed: 15723660]
- Stump AD, Fitzpatrick MC, Lobo NF, et al. Centromere-proximal differentiation and speciation in *Anopheles gambiae*. *Proc Natl Acad Sci U S A*. 2005; 102:15930–15935. [PubMed: 16247019]
- Tchuinkam T, Mulder B, Dechering K, et al. Experimental infections of *Anopheles gambiae* with *Plasmodium falciparum* of naturally infected gametocyte carriers in Cameroon: factors influencing the infectivity to mosquitoes. *Tropical Medicine and Parasitology*. 1993; 44:271–276. [PubMed: 8134766]
- Tene Fossog B, Antonio-Nkondjio C, Pierre Kengne P, et al. Physiological correlates of ecological divergence along an urbanization gradient: differential tolerance to ammonia among molecular forms of the malaria mosquito *Anopheles gambiae*. *BMC Ecol*. 2013a; 13:1. [PubMed: 23294940]
- Tene Fossog B, Kopya E, Ndo C, et al. Water Quality and *Anopheles gambiae* Larval Tolerance to Pyrethroids in the Cities of Douala and Yaounde (Cameroon). *Journal of tropical medicine*. 2012; 2012:429817. [PubMed: 22719775]
- Tene Fossog B, Poupardin R, Costantini C, et al. Resistance to DDT in an urban setting: common mechanisms implicated in both M and S forms of *Anopheles gambiae* in the city of Yaounde Cameroon. *PloS one*. 2013b; 8:e61408. [PubMed: 23626680]

- Turner TL, Hahn MW. Locus- and population-specific selection and differentiation between incipient species of *Anopheles gambiae*. *Molecular Biology and Evolution*. 2007; 24:2132–2138. [PubMed: 17636041]
- Turner TL, Hahn MW, Nuzhdin SV. Genomic islands of speciation in *Anopheles gambiae*. *PLoS Biol*. 2005; 3:e285. [PubMed: 16076241]
- von Dungern P, Briegel H. Protein catabolism in mosquitoes: ureotely and uricotely in larval and imaginal *Aedes aegypti*. *Journal of insect physiology*. 2001; 47:131–141. [PubMed: 11064020]
- Waterhouse RM, Kriventseva EV, Meister S, et al. Evolutionary dynamics of immune-related genes and pathways in disease-vector mosquitoes. *Science*. 2007; 316:1738–1743. [PubMed: 17588928]
- Waterhouse RM, Povelones M, Christophides GK. Sequence-structure-function relations of the mosquito leucine-rich repeat immune proteins. *BMC Genomics*. 2010; 11:531. [PubMed: 20920294]
- Weihrauch D, Donini A, O'Donnell MJ. Ammonia transport by terrestrial and aquatic insects. *Journal of insect physiology*. 2012; 58:473–487. [PubMed: 22100291]
- White BJ, Lawniczak MK, Cheng C, et al. Adaptive divergence between incipient species of *Anopheles gambiae* increases resistance to Plasmodium. *Proc Natl Acad Sci U S A*. 2011; 108:244–249. [PubMed: 21173248]
- Whitehead A. Comparative genomics in ecological physiology: toward a more nuanced understanding of acclimation and adaptation. *The Journal of experimental biology*. 2012; 215:884–891. [PubMed: 22357582]
- Whitehead A, Crawford DL. Neutral and adaptive variation in gene expression. *Proceedings of the National Academy of Sciences of the United States of America*. 2006a; 103:5425–5430. [PubMed: 16567645]
- Whitehead A, Crawford DL. Variation within and among species in gene expression: raw material for evolution. *Molecular Ecology*. 2006b; 15:1197–1211. [PubMed: 16626448]
- Wittkopp PJ, Haerum BK, Clark AG. Regulatory changes underlying expression differences within and between *Drosophila* species. *Nature Genetics*. 2008; 40:346–350. [PubMed: 18278046]
- Yadouleton AW, Asidi A, Djouaka RF, et al. Development of vegetable farming: a cause of the emergence of insecticide resistance in populations of *Anopheles gambiae* in urban areas of Benin. *Malaria journal*. 2009; 8:103. [PubMed: 19442297]
- Yang H, Wang Y, Xiao Y, et al. A bi-functional anti-thrombosis protein containing both direct-acting fibrin(ogen)olytic and plasminogen-activating activities. *PloS one*. 2011; 6:e17519. [PubMed: 21423730]
- Zdobnov EM, von Mering C, Letunic I, et al. Comparative genome and proteome analysis of *Anopheles gambiae* and *Drosophila melanogaster*. *Science*. 2002; 298:149–159. [PubMed: 12364792]
- Zheng W, Gianoulis TA, Karczewski KJ, Zhao H, Snyder M. Regulatory variation within and between species. *Annual review of genomics and human genetics*. 2011; 12:327–346.




**Figure 1.** Representative larval breeding habitats of *An. gambiae* (rural; left panel) and *An. coluzzii* (urban; right panel) sampled during this study in Yaoundé, Cameroon.






**Figure 2.** Sampling localities in and around Yaoundé, Cameroon. Four urban localities (pink shading) and four rural localities contain habitats exclusive to *An. gambiae* and *An. coluzzii*, respectively. Areas were shaded pink based on an index of urbanization (reflecting the proportion of built-up surface per spatial sampling unit), extracted from remotely sensed data (Kamdem *et al.* 2012).




**Figure 3.**

Volcano plot displaying gene expression differences ( $\log_2$ -fold change) between untreated larval samples of *An. coluzzii* and *An. gambiae*. Each symbol represents one gene that had detectable expression in either species, in samples of 2<sup>nd</sup> instar larvae (blue cross) or 4<sup>th</sup> instar larvae (red triangle). Relative differences in signal intensity along the X-axis reflect up-regulation in *An. coluzzii* when positive, and up-regulation in *An. gambiae* when negative; vertical lines indicate the  $\log_2$ -fold-change cut-off of  $\pm 1.8$ . The Y-axis displays  $\log_{10}$ -transformed *P*-values associated with Bayes-moderated *t*-tests of differential gene expression; the horizontal line indicates the 0.05 threshold level for significance.



**Figure 4.** Functional enrichment of genes that are over-expressed in field-collected samples of 2<sup>nd</sup> or 4<sup>th</sup> instar larvae of *An. coluzzii* and *An. gambiae*. The numbers of significant genes for each DAVID annotation cluster are indicated above each bar. Enrichment scores are displayed within each bar. GST, glutathione S transferase.

**Figure 5.**

Functional enrichment of genes that are overexpressed in 4<sup>th</sup> instar larvae from laboratory colonies of *An. coluzzii* and *An. gambiae*. A, *An. coluzzii* CAM from Yaoundé, Cameroon compared to *An. gambiae* Pimperena from Mali. B, Comparison of two colonies from Mali: *An. coluzzii* Mali-NIH and *An. gambiae* Pimperena. The numbers of significant genes for each DAVID annotation cluster are indicated above each bar. Enrichment scores are displayed within each bar. CYP450, cytochrome P450; UGT, UDP-glucuronosyltransferase; CE, carboxylesterase; GST, glutathione S transferase; PI, peptidase inhibitor; TEP, thioester-containing protein; Redox, oxidoreductase; CTL, C-type lectin.

**Table 1**

Geographic coordinates of sampling localities in Yaoundé, Cameroon

Description	Locality Name	Longitude; Latitude
Rural ( <i>An. gambiae</i> )	Ngalli II	11°30'36"E; 4°01'12"N
	Nkolkoumou I	11°23'38"E; 3°51'39"N
	Nkolkoumou II	11°24'09"E; 3°51'44"N
	Bibon Biboum	11°25'12"E; 3°47'24"N
Urban ( <i>An. coluzzii</i> )	Mballa II	11°31'12"E; 3°54'00"N
	Mokolo Elobi	11°30'22"E; 3°52'49"N
	Mokolo Messa	11°30'04"E; 3°52'21"N
	Voirie Municipale	11°31'12"E; 3°51'36"N

**Table 2**

Experimental design for microarray-based transcriptional profiling of *An. coluzzii* and *An. gambiae* larvae from Yaoundé, Cameroon

Species	Treatment	Instar	Replicates hybridized (analyzed)
<i>An. coluzzii</i>	None	2 <sup>nd</sup>	4
		4 <sup>th</sup>	4 (3)
	24 h auto-translocation	2 <sup>nd</sup>	4 (3)
		4 <sup>th</sup>	4
	24 h allo-translocation	2 <sup>nd</sup>	4
		4 <sup>th</sup>	4
<i>An. gambiae</i>	None	2 <sup>nd</sup>	4
		4 <sup>th</sup>	4
	24 h auto-translocation	2 <sup>nd</sup>	4 (3)
		4	4
	24 h allo-translocation	2 <sup>nd</sup>	4
		4 <sup>th</sup>	4

Table 3

Immune-related genes<sup>1</sup> (AGAP last five digits) upregulated in field-collected *An. coluzzii* or *An. gambiae* larvae

Immune Gene Family <sup>1</sup>	<i>An. coluzzii</i> 2 <sup>nd</sup> instar	4 <sup>th</sup> instar	<i>An. gambiae</i> 2 <sup>nd</sup> instar	4 <sup>th</sup> instar
Anti-Microbial Peptides			06722*	
Autophagy Genes	04023	00933		
Glucan Binding Proteins	04455	02798	12544*	12544*
Caspases			09214*	02270; 09214; 13184
Clip Domain Serine Proteases	03250*; 03251; 04719; 11789; 11791	03250; 04719	10193*	07411; 10193*; 10196;
C-Type Lectins	00871; 07409*	00940; 02911	10772*	10772*; 11277
Fibrinogen-Related Proteins	02005; 04916; 04917; 04918; 04996*; 04997*; 09184; 09728*; 10761; 10763*; 11197; 11223; 11225; 11239	04916; 04917; 04918; 05848*; 06914; 09728*; 10759; 10762*; 10811*; 11225; 11231; 11239		
IMD Pathway	06473	06473		
JAKSTAT Signal Transduction		08354; 010423		
Leucine-Rich Repeat Proteins	07033*; 07036*	07039*		
Lysozymes	07344*			
MD2-Like Receptors	12352	01956; 02852; 07415		10734
Nimrod	12386			
Peptidoglycan Recognition Proteins	05552			
Peroxidases	03714; 13327			
Prophenoloxidases	04975; 04981	04975*		
Scavenger Receptors	01375*; 03139; 04847; 09670			
Serine Protease Inhibitors				01377
Small RNA Regulatory Pathway	11717*	11537; 12523		09781
Thioester-Containing Proteins	10815	10814		
Toll-Like Receptors	06974*	06974		

<sup>1</sup> Gene assignments based on (Bartholomay et al. 2010; Waterhouse et al. 2007; Waterhouse et al. 2010). Asterisks indicate genes upregulated 24 h after transfer to laboratory culture.

Table 4

Detoxification genes (AGAP last 5 digits) over-expressed in field-collected and colonized *An. coluzzii* larvae

Gene class	Field-collected 2 <sup>nd</sup> instar	Field-collected 4 <sup>th</sup> instar	CAM 4 <sup>th</sup> instar	Mali-NIH 4 <sup>th</sup> instar
CYP450	CYP301A1 (06082), CYP9J5 (12296)	<b>CYP325C2</b> (02205), CYP325B1 (02210), CYP4D16 (13241)	<b>CYP9K1</b> (00818), CYP6AA1 (02862), <b>CYP6P3</b> (02865), <b>CYP6P4</b> (02867), CYP6P1 (02868), CYP6S1 (08204), CYP6M1 (08209), <b>CYP6M2</b> (08212), CYP6M4 (08214), <b>CYP6Z3</b> (08217), <b>CYP6Z2</b> (08218), CYP4H17 (08358), CYP4H27 (08552), CYP9J5 (12296)	<b>CYP6M2</b> (08212), CYP4C36 (09241), CYP6M4 (08214), <b>CYP12F1</b> (08022)
CE	COBE4C (05370), COBE2C (05371), COBE3C (05372), COEAE3D (05758)	COEAE1D (05756)	COEAE1A (01723), COEAE3D (05758), COEJHE5E (05837), COE13O (11507)	---
GST	GSTD10 (04383), <b>GSTE5</b> (09192), <b>GSTE2</b> (09194), <b>GSTE1</b> (09195), <b>GSTE3</b> (09197)	<b>GSTE2</b> (09194), <b>GSTE1</b> (09195)	GSTT2 (00888), <b>GSTD11</b> (04378), <b>GSTD6</b> (04379), <b>GSTD12</b> (04380), <b>GSTD3</b> (04382)	GSTT2 (00888), <b>GSTD11</b> (04378)
UGT	UGT5 (02783)	UGT5 (02783)	UGT (06222), UGT (07028), UGT (07029), UGT (11564)	---

CYP450, cytochrome P450; CE, carboxylesterase; GST, glutathione S transferase; UGT, UDP-glucuronosyltransferase. Genes underlined and bolded have been validated to metabolize insecticide (Daborn *et al.* 2012; David *et al.* 2013; Ranson *et al.* 2001); those in bold have been associated with insecticide resistance (David *et al.* 2013; Djouaka *et al.* 2008; Tene Fossog *et al.* 2013b)