

RESEARCH PAPER

Requirement for A-type cyclin-dependent kinase and cyclins for the terminal division in the stomatal lineage of *Arabidopsis*

Kezhen Yang, Hongzhe Wang, Shan Xue, Xiaoxiao Qu, Junjie Zou and Jie Le*

Key Laboratory of Plant Molecular Physiology, Institute of Botany, Chinese Academy of Sciences, 20 Nanxincun, Beijing 100093, China

* To whom correspondence should be addressed. E-mail: lejie@ibcas.ac.cn

Received 10 October 2013; Revised 20 February 2014; Accepted 7 March 2014

Abstract

The *Arabidopsis* stoma is a specialized epidermal valve made up of a pair of guard cells around a pore whose aperture controls gas exchange between the shoot and atmosphere. Guard cells (GCs) are produced by a symmetric division of guard mother cells (GMCs). The R2R3-MYB transcription factor FOUR LIPS (FLP) and its paralogue MYB88 restrict the division of a GMC to one. Previously, the upstream regions of several core cell cycle genes were identified as the direct targets of FLP/MYB88, including the B-type cyclin-dependent kinase *CDKB1;1* and A2-type cyclin *CYCA2;3*. Here we show that *CDKA;1* is also an immediate direct target of FLP/MYB88 through the binding to cis-regulatory elements in the *CDKA;1* promoter region. *CDKA;1* activity is required not only for normal GMC divisions but also for the excessive cell overproliferation in *flp myb88* mutant GMCs. The impaired defects of GMC division in *cdkb1;1 1;2* mutants could be partially rescued by a stage-specific expression of *CDKA;1*. Although targeted overexpression of *CDKA;1* does not affect stomatal development, ectopic expression of the D3-type cyclin *CYCD3;2* induces GC subdivision, resulting in a stoma with 3–4 GCs instead of the normal two. Co-overexpression of *CDKA;1* with *CYCD3;2*, but not with *CYCA2;3*, confers a synergistic effect with respect to GC subdivision. Thus, in addition to a role in stomatal formative asymmetric divisions at early developmental stages, *CDKA;1* is needed in triggering GMC symmetric divisions at the late stage of stomatal development. However, timely down-regulation of *CDKA;1*–*CYCD3* activity is required for restriction of GC proliferation.

Key words: *Arabidopsis*, cell cycle, stomatal development, symmetric division, transcription factor.

Introduction

In *Arabidopsis thaliana*, stomata consist of a pair of guard cells (GCs) surrounding a pore that permits gas exchange between internal plant tissues and the atmosphere. Stomata are produced through a series of divisions including at least one asymmetric and one symmetric division. The meristemoid mother cell (MMC) undergoes an asymmetric entry division to produce a small triangular-shaped meristemoid and a larger sister cell. Meristemoids usually undergo 1–3 rounds of asymmetric divisions and eventually differentiate into guard mother cells (GMCs). GMCs then divide once symmetrically

to produce a pair of GCs (Bergmann and Sack, 2007). The final developmental stage of stomata is GC differentiation, pore formation, and GC shape control (Nadeau and Sack, 2002).

Three basic-helix–loop–helix (bHLH) transcription factors, SPEECHLESS (SPCH), MUTE, and FAMA, are required for the successive stages of stomatal development (Ohashi-Ito and Bergmann, 2006; MacAlister *et al.*, 2007; Pillitteri *et al.*, 2007). SPCH is essential for MMC formation, stomatal entry divisions, and maintenance of meristemoid stem cell

Abbreviations: CDKA, 1, CYCLIN DEPENDENT KINASE A1; CYCD3, CYCLIN D3; CYCA2, CYCLIN A2; DAPI, 4',6-diamidino-2-phenylindole; FLP, FOUR LIPS; GC, guard cell; GMC, guard mother cell; MMC, meristemoid mother cell; RBR1, RETINOBLASTOMA-RELATED 1; SGC, single guard cell.

© The Author 2014. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

activity (Pillitteri and Torii, 2012; Lau and Bergmann, 2012). MUTE promotes the transition of meristemoids into GMCs (Pillitteri *et al.*, 2007). Loss of function of *FAMA* causes tumours composed of stacked narrow cells, but overexpression of *FAMA* induces ectopic GCs without GMC divisions, suggesting the dual roles of *FAMA* in regulating differentiation and proliferation (Ohashi-Ito and Bergmann, 2006).

Two redundant R2R3 MYB transcription factors, FOUR LIPS (FLP) and MYB88, restrict the symmetrical cell division of the GMC (Lai *et al.*, 2005). Double mutants of *flp myb88* display stacked cells like *fama* mutants but some cells still could acquire the GC fate (Xie *et al.*, 2010). Reducing the activity of CYCLIN DEPENDENT KINASE B1;1 (CDKB1;1) by overexpression of the dominant-negative form *CDKB1;1.N161* or loss of function of both *CDKB1;1* and *CDKB1;2* (*cdkb1;1 1;2*) blocks the GMC symmetric division at the G₂ to M phase transition of the cell cycle, resulting in the formation of single guard cells (SGCs) (Boudolf *et al.*, 2004a; Xie *et al.*, 2010). FLP/MYB88 can bind directly to a *cis*-regulatory element in the *CDKB1;1* promoter and can suppress *CDKB1;1* transcription levels. Although *FAMA* functions at the GMC to GC transition and regulates *CDKB1;1* expression as well, *FAMA* probably acts in a parallel pathway different from FLP/MYB88 (Ohashi-Ito and Bergmann, 2006).

Since CDK activation depends on its association with cyclin partners, co-expression of *CDKB1;1* and *CYCLIN A2;3* (*CYCA2;3*) enhances the kinase activity of CDKB1;1 and triggers ectopic cell divisions (Boudolf *et al.*, 2009). Defective GMC divisions are found in *cyca2* mutants, while the *cdkb1;1 cyca2;234* quadruple mutant displays more SGCs than *cyca2;234* triple mutants, though there is no phenotype in the *cdkb1;1* single mutant, suggesting that *CYCA2* and *CDKB1* genes synergistically promote GMC division. Consistently, a sustained high *CYCA2;3* and *CDKB1;1* expression is found in *flp myb88* epidermal tumours, indicating that FLP/MYB88 repress *CYCA2;3* and *CDKB1;1* transcription in a timely manner after GMC division to prevent cell overproliferation (Xie *et al.*, 2010; Vanneste *et al.*, 2011).

In yeast, there is only a single cyclin-dependent kinase, and its low, intermediate, or high activity is required for DNA replication licensing, DNA replication initiation, and cell mitosis, respectively (Porter, 2008). In *Arabidopsis*, the only one A-type CDK, CDKA;1 (also called CDC2A), which can rescue the fission yeast *cdc2* mutant, is required for cell cycle regulation during pollen, embryo, root, and shoot apical stem development (Inze and De Veylder, 2006; Gutierrez, 2008). Previous chromatin immunoprecipitation (ChIP)-chip experiments with antibodies against FLP/MYB88 revealed that the upstream promoter region of the *CDKA;1* gene might be the target for binding by FLP/MYB88 (Xie *et al.*, 2010), indicating a putative role for *CDKA;1* in stomatal GMC divisions.

In animals, CDK–cyclin complexes control G₁ to S progression in the cell cycle through regulating phosphorylation of *RETINOBLASTOMA* proteins and releasing the E2F/DP complex (Harbour and Dean, 2000). Recently, the partial redundancy among the A- and B1-type CDKs in the *RETINOBLASTOMA-RELATED1* (*RBRI*; the homologue of the human tumour suppressor Retinoblastoma in

Arabidopsis) regulatory network has been intensively investigated in plants. RBR1 is phosphorylated by several cyclin–CDK complexes, such as, CYCD6/CDKB1;1, CYCD3;1/CDKA;1, or CYCB1;1/CDKB1;1 (Cruz-Ramirez *et al.* 2012; Nowack *et al.*, 2012). However, ChIP experiments also revealed that *CDKB1;1* and *CDKB1;2* are the direct transcriptional targets of RBR1 (Nowack *et al.*, 2012). CYCD6 transcript levels are indirectly regulated by RBR1 through interaction with the SCARECROW (SCR) transcription factor, which in complex with SHORT ROOT regulates *CYCD6;1* expression during root stem cell asymmetric division (Cruz-Ramirez *et al.*, 2012). A negative regulatory feedback loop of RBR1 on CDK activity was also found in maize endosperm (Sabelli *et al.*, 2013).

Expression of *CDKA;1* in the stomatal lineage, under control of the *TOO MANY MOUTHS* (*TMM*) promoter, could partially rescue *cdkb1;1 1;2* stomata defects, suggesting functional redundancy between CDKA;1 and CDKB1s, in agreement with the arrested GMC divisions in the *cdka;1* null mutant (Weimer *et al.*, 2012). In addition, stomata in *RBR1* RNA interference (RNAi) lines often consist of four GCs (Borghi *et al.*, 2010). A relationship between two types of CDKs and RBR1 has been proposed for stomatal asymmetric division (Weimer *et al.*, 2012), but whether a similar network is required for the symmetric divisions during the last stage of stomatal development has not been well characterized.

It has been shown recently that FLP and MYB88 conditionally restrict the G₁ to S transition during formation of stomata (Lee *et al.*, 2013). Here it is shown that *CDKA;1*, like *CDKB1;1*, is also a direct target of FLP/MYB88 through binding to the *cis*-regulatory elements in its promoter. CDKA;1 activity is required for both normal GMC division and cell overproliferation in *flp myb88* mutants.

Materials and methods

Plant materials and growth conditions

All genotypes were in a Columbia-0 (Col-0) ecotypic background. *Arabidopsis thaliana* plants were grown on soil or half-strength Murashige and Skoog (MS) medium at 22–24 °C with 16/8 h light/dark cycles.

Plasmid construction and plant transformation

The following constructs and transgenic plants were generated using the primers shown in Supplementary Table S1 available at JXB online: *pSPCH:CDKA;1*, *pSPCH:CYCD3;2*, *pMUTE:CDKA;1*, *pMUTE:CDKA;1.N146*, *pMUTE:CYCD3;2*, *pFAMA:CDKA;1*, *pFAMA:CDKA;1.N146*, *pFAMA:CYCD3;1*, *pFAMA:CYCD3;2*, *pFAMA:CYCD3;3*, and *pFAMA:CYCA2;3*. The promoter fragments of *SPCH*, *MUTE*, *FAMA*, as well as full-length cDNAs of *CDKA;1*, *CDKA;1.N146*, *CYCD3;1*, *CYCD3;2*, *CYCD3;3*, and *CYCA2;3* were obtained by PCR. All resulting DNA fragments were cloned into the pMD19-T vector (TaKaRa) and then subcloned into a pCAMBIA1300 vector (CAMBIA) and transformed into wild-type plants. Transgenic plants were selected on half-strength MS medium containing 25 µg l⁻¹ hygromycin. Plants harbouring multiple constructs were created by crossing and confirmed by PCR. To obtain a viable *cdka;1* null mutant, a *pLAT52:CDKA;1* construct, in which the full-length cDNA of *CDKA;1* is driven by the pollen-specific *LAT52* promoter, was transformed into heterozygous *cdka;1*^{+/-} plants (SALK_106809). The homozygosity of

cdka;1 plants was confirmed by PCR using the primers shown in [Supplementary Table S1](#) at *JXB* online.

β-Glucuronidase (*GUS*) staining

Before staining, young seedlings were incubated in 90% acetone for 2 h at 4 °C. Seedlings were then washed in phosphate buffer and immersed in X-gluc solution (1 mg ml⁻¹ 5-bromo-4-chloro-3-indolyl β-D-glucuronide, 2 mM ferricyanide, and 0.5 mM ferrocyanide in 100 mM phosphate buffer, pH 7.0) overnight at 37 °C in the dark. Seedlings were then cleared and imaged using an Olympus BX51 microscope.

Cell viability assay

True leaves were immersed in a 0.1% neutral red solution for 15 min at room temperature before imaging with an Olympus BX51 microscope.

Microscopy

To obtain differential interference contrast (DIC) images, 2-week-old cotyledons were treated with destaining solution (containing 75% ethanol and 25% acetic acid) for 30 min or overnight at room temperature until the chlorophyll was cleared. After a treatment with basic solution (7% NaOH in 60% ethanol) for 15 min at room temperature, the samples were rehydrated via an ethanol series (40, 20, and 10%) for 15 min at each step and then placed in 5% ethanol and 25% glycerol for 30 min. Materials were mounted in 50% glycerol and imaged using an Olympus BX51 microscope. For fluorescence, samples were stained with 0.5% propidium iodide and fluorescence was imaged using a confocal laser scanning microscope (FV1000-MPE, Olympus).

DAPI staining and measurement of DNA content

For analysis of nuclei in GCs, cotyledons were dissected and fixed in 70% ethanol for 3 h, incubated in 4',6-diamidino-2-phenylindole (DAPI) staining solution for at least 30 min, and excited by UV fluorescence. To score relative DNA levels, the total integrated density of DAPI fluorescence from selected nuclei was analysed using ImageJ (<http://rsb.info.nih.gov/ij/>, last accessed 20 March 2014) with fluorescence from nearby areas subtracted to standardize relative fluorescence levels. GCs resembling those in the wild type as well as those in epidermal cells that were newly divided were used as the references.

Real-time qPCR

Total RNA from 10-day-old seedlings were extracted using TRNzol reagent (<http://www.tiangen.com>, last accessed 20 March 2014). Reverse transcription was performed using a Promega kit (<http://www.promega.com>, last accessed 20 March 2014). Amplification of the *KAT1* gene was used as an internal control. Real-time quantitative PCR (RT-qPCR) experiments were repeated three times independently. The cDNA was amplified using SYBR Premix Ex Taq™ (TaKaRa) with a Corbett RG3000.

Yeast one-hybrid assay

The region between base pairs 859 and 568 upstream of the *CDKA;1* gene was split into a 191 bp upstream fragment and a 100 bp downstream fragment, subjected to PCR, and ligated into the pLacZi2μ vector. *FLP* and *MYB88* cDNA were cloned into pB42AD. Yeast one-hybrid assays were performed as previously described ([Tang et al., 2012](#)).

Protein expression

Full-length cDNA sequences of *FLP* and *MYB88* were amplified by using the primers shown in [Supplementary Table S1](#) at *JXB* online. *FLP* and *MYB88* cDNA fragments were then ligated into pET-28a. Fusion proteins were expressed in the BL21 (DE3)

strain of *Escherichia coli* by induction with 0.1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) at 18 °C for 24 h. His-FLP and His-MYB88 proteins were purified by Ni-NTA agarose (GE Healthcare) following the manufacturer's instructions.

Electrophoretic mobility shift assay (EMSA)

Oligonucleotide probes were synthesized and labelled with biotin at the 3' end (Thermo Scientific) and the probe sequences are shown in [Supplementary Table S1](#) at *JXB* online. EMSA was performed using a Light Shift Chemiluminescent EMSA kit (Thermo Scientific). Briefly, biotin-labelled probes were incubated in 1× binding buffer, 2.5% glycerol, 50 mM KCl, 5 mM MgCl₂, 0.05% NP-40, and 10 mM EDTA with or without proteins at room temperature for 20 min. For probe competition controls, non-labelled probes were added to the reactions.

Yeast two-hybrid assay

Gal4 system vectors were used for yeast two-hybrid assays (Clontech). The full-length coding sequences of *CDKA;1.N146* and *CYCD3;2* were amplified using the primers shown in [Supplementary Table S1](#) at *JXB* online. cDNA fragments from *CYCD3;2* were cloned into a pGBKT7 vector, and then *CDKA;1.N146* cDNA fragments were cloned into a pGADT7 vector. Constructs were then co-transformed into an AH109 yeast strain and selected on SD/-Leu-Trp or SD/-Leu-Trp-His-Ade plates. X-Gal activity was analysed according to the manufacturer's instructions (Clontech).

Bimolecular fluorescence complementation (BiFC) assay

Coding sequences of *CDKA;1.N146* and *CYCD3;2* were amplified using the primers shown in [Supplementary Table S1](#) at *JXB* online. Fragments were cloned into the pSPYNE-35S and pSPYCE-35S vectors ([Walter et al., 2004](#)) to obtain *CYCD3;2-YN* and *CDKA;1.N146-YC*. *CYCD3;2-YN* and *CDKA;1.N146-YC* were co-transformed into *Arabidopsis* protoplasts and imaged using a laser scanning confocal microscope (FV1000-MPE, Olympus).

Pull-down assays

Full coding sequences of *CDKA;1.N146* and *CYCD3;2* were amplified by using the primers shown in [Supplementary Table S1](#) at *JXB* online. The sequences were then ligated into pET-28a and pGEX4T-1 vectors, respectively. Fusion proteins were expressed in the BL21 (DE3) strain of *E. coli* by induction with 0.2 mM IPTG at 17 °C for 4 h. The harvested culture (5000 rpm, 15 min, 4 °C) was re-suspended with ice-cold phosphate-buffered saline (PBS) and then lysed by sonication. The cell lysate was cooled to 4 °C and centrifuged at 10 000 rpm for 15 min. The glutathione *S*-transferase (GST)-*CYCD3;2* supernatant was loaded onto 0.5 ml of glutathione-Sepharose (GE Healthcare) and washed with PBS. The GST-*CYCD3;2* fusion protein on glutathione-Sepharose (GE Healthcare) was then incubated with the His6-*CDKA.N146* supernatant. After 1 h incubation at 4 °C, the agarose was washed twice with PBS and eluted with 10 mM reduced glutathione elution buffer. Elution samples were separated by 12% SDS-PAGE, transferred to a PVDF membrane (Millipore) using a semi-dry blotting system (Bio-Rad), and then incubated with an anti-His6 monoclonal antibody followed by AP-conjugated anti-mouse antibody. The colour reaction was performed using an NBT/BCIP Kit (Invitrogen).

Results

CDKA;1 activity is required for the late stages of stomatal development

Stomata formation is greatly inhibited in *cdka;1* null mutant cotyledons, which demonstrates the essential role of *CDKA;1*

in stomatal formative divisions. The occasional occurrence of arrested GMCs and expression of *CDKA;1* at late stages of the stomatal cell lineage suggests an involvement of *CDKA;1* in the subsequent events as well (Xie *et al.*, 2010; Weimer *et al.*, 2012). Overexpression of a dominant-negative allele of *CDKA;1* (*CDKA;1.N146*) was previously used to reveal the function of *CDK* genes in plant development (Hemerly *et al.*, 1995; De Veylder *et al.*, 2000; Joubès *et al.*, 2004; Gaamouche *et al.*, 2010). *MUTE* and *FAMA* are specifically expressed at the transition phases of meristemoid to GMC and GMC to GC, respectively. Thus, the *CDKA;1.N146* construct was driven by the upstream regulatory regions of *MUTE* and *FAMA* genes. Targeted expression of either *MUTE:CDKA;1.N146* or *FAMA:CDKA;1.N146* did not affect the initiation of stomatal stem cells and morphogenesis of pavement cells (Fig. 1A–C). However, *MUTE:CDKA;1.N146* transgenic lines exhibit large abnormally shaped cells expressing a GC fate marker, E1728 (Gardner *et al.*, 2009), demonstrating that those round cells have acquired GC identity (inset in Fig. 1B).

Loss of *CDKA;1* function impairs the last mitotic division in the male gametophyte, leading to 50% of pollen with two gametes, and no homozygous *cdka;1* seedlings are produced in self-pollinated *cdka;1^{+/+}* plants (Iwakawa *et al.*, 2006). Therefore, viable homozygous *cdka;1* seedlings were recovered from *cdka;1^{+/+}* plants by transformation with *pLAT52:CDKA;1* constructs. As in *MUTE:CDKA;1.N146* seedlings, abnormal undivided round cells were also found in the leaves of homozygous *cdka;1* seedlings. Consistently, expression of the E1728 marker suggests that the undivided *cdka;1* GMCs could eventually also acquire GC cell fate (Fig. 1D).

Similar fluorescent intensity of DAPI-stained nuclei in normal 2-GC stomata and in nuclei of *MUTE:CDKA;1.N146* SGCs indicated that *CDKA;1.N146* prevented DNA synthesis and caused cell cycle arrest (Fig. 1E–H, O), this is in contrast to the a 4C DNA content in SGCs from *cdkb1;1 1;2*, *cyca2;234* mutants or *35S:CDKB1;1.N161* transgenic plants (Xie *et al.*, 2010; Vanneste *et al.*, 2011). Little effect on the symmetric division was observed when the dominant-negative *CDKA;1* was activated later with the *FAMA:CDKA;1.N146* construct (Fig. 1C). Therefore, *CDKA;1* activity is required for the early events, such as the G₁ to S transition, in the GMC cell cycle.

Identification of FLP/MYB88-binding sites in the *CDKA;1* promoter

The weak *flp-1* allele displays a stomatal cluster phenotype that often contains four stacked GCs caused by an extra round of division in a GMC (Lai *et al.*, 2005). The *flp-1 myb88* double mutant displays an enhanced stomatal phenotype with more and larger stomatal clusters (Fig. 1I). To determine whether *CDKA;1* is required for the additional GMC divisions in the *flp-1 myb88* mutant background, *cdka;1 flp-1 myb88* triple mutants were generated. The typical GC stacks of the *flp-1 myb88* mutants were not found in triple mutants, instead SGCs are often present, suggesting that the cell overproliferation caused by mutations of *FLP* and *MYB88*

required *CDKA;1* activity (Fig. 1J). DAPI staining reveals a single nucleus in the round cells from the triple mutant, implying that arrested division is not due to a defective cytokinesis upon mitosis (Fig. 1K–N). Using the transcript level of a GC-specific gene *KATI* as the internal reference for RT-qPCR analysis, it was found that the transcription of *CDKA;1*, like that of *CDKB1;1*, was up-regulated in *flp-1 myb88* plants, indicating that *FLP/MYB88* negatively regulate *CDKA;1* transcription (Fig. 1P).

Previous ChIP-chip experiments with *FLP/MYB88* antisera also revealed that the region between base pairs 859 and 568 upstream of the translational start site of the *CDKA;1* gene might be the target of *FLP/MYB88* (Xie *et al.*, 2010). To identify the *FLP/MYB88* direct binding sites in the *CDKA;1* promoter, the above promoter region was split into 191 bp and 100 bp fragments, named fragment ‘a’ and ‘b’, respectively (Fig. 2A). Yeast one-hybrid assay indicates that the binding sites are present within the fragment ‘a’ (Fig. 2B). Then the sequence in fragment ‘a’ was scanned and four putative *FLP/MYB88 cis*-regulatory binding elements were found (Xie *et al.*, 2010), namely TGCGG, AACCC, TCCCC, and TTCCC, termed elements A–D, respectively (Fig. 2A). Each of these sites was tested in EMSAs using tagged His-*FLP* or His-*MYB88* fusion proteins. Since elements C and D are closely located within the *CDKA;1* promoter, all nucleotides within either the C or D elements were replaced with adenine in EMSA analysis. The EMSA results show *FLP* and *MYB88* can interact with elements A, and B, and with a sequence composed of both the C and D elements (Fig. 2C; Supplementary Fig. S1 at JXB online).

CDKA;1 could rescue defective GMC divisions in *cdkb1* mutants

CDKB1;1 and *FLP* show a similar expression pattern from the late GMC to the newly formed GC, in agreement with the role of *FLP* in regulating *CDKB1;1* transcription (Xie *et al.*, 2010). Targeted expression of *CDKB1;1* under the control of the *FAMA* promoter (*FAMA:CDKB1;1*) is sufficient to complement the SGC phenotype found in *cdkb1;1 1;2* double mutants (Table 1). The *CDKA;1* construct driven by *FAMA* was introduced into *cdkb1;1 1;2* double mutants. Interestingly, expression of this *FAMA:CDKA;1* could reduce the number of SGCs formed in *cdkb1;1 1;2*, indicating that *CDKA;1* can partially substitute for *CDKB1* genes in promoting GMC divisions (Table 1).

A- and D-type cyclins are rate limiting for division after the terminal stomatal division

Previous studies revealed that *CDKB1;1* and *CYCA2;3* form a functional complex regulating the mitosis to endocycle transition (Boudolf *et al.*, 2009). *CYCA2s* and *CDKB1;1* contribute synergistically to the *CDK* activity required for GMC divisions (Vanneste *et al.*, 2011). The dominant-negative *CDKB1;1.N161* competes with the endogenous *CDKs*, leading to SGCs (Fig. 3A, B). Thus experiments were carried out to examine whether an ectopic expression of *CYCA2;3*




Fig. 1. *CDKA;1* is required for excessive GMC divisions in *flp-1 myb88* mutants. (A) Wild-type epidermis. (B, C) The expression of a dominant-negative *CDKA;1.N146* construct driven by the *MUTE* promoter (*MUTE:CDKA;1.N146*), but not the *FAMA* promoter (*FAMA:CDKA;1.N146*), leads to formation of SGCs. Inset: expression of the mature GC marker E1728 in an SGC. (D) SGCs were found in loss-of-function *cdka;1* leaves. Inset: E1728. (E–H) Paired DAPI fluorescence micrographs and DIC images showing normal GCs in wild-type (E, F) and uninucleate SGCs in *MUTE:CDKA;1.N146* (G, H). (I) Stomatal clusters in the *flp-1 myb88* double mutant. (J) The *flp-1 myb88* cluster is repressed in the *cdka;1* background. An arrow indicates an SGC. (K–N) DAPI staining of SGCs in *cdka;1* and the *flp-1 myb88 cdka;1* triple mutant. (O) Quantitative analysis of DAPI fluorescence revealed that *MUTE:CDKA;1.N146* SGCs contain a DNA content comparable with that of normal GCs. (P) Transcript levels of *SPCH*, *CDKA;1*, and *CDKB1;1* in *flp-1 myb88* double mutants. Bars=20 μ m. (This figure is available in colour at JXB online.)

under the control of the *FAMA* promoter has an impact on GMC divisions. Although SGCs were still occasionally found, the defect of GMC division in *35S:CDKB1;1.N161* plants was overcome by expression of *FAMA:CYCA2;3* (Table 1; Fig. 3D). Strikingly, expression of *FAMA:CYCA2;3* in wild-type plants produced stomata with 3–4 guard cells (Fig. 3C), indicating that elevated levels of *CYCA2;3* sustain divisions in the GCs after the GMC symmetric division.

Moreover expression of each member gene of the *CYCD3* family under the control of the *FAMA* promoter induced three- or four-celled stomata as well (Fig. 3E; Supplementary Fig. S2 at JXB online). No SGC was found in *35S:CDKB1;1.N161* plants harbouring *FAMA:CYCD3;2*, indicating that increasing levels of D-type cyclin can reverse the G₂ arrest in these dominant-negative *CDKB1;1.N161* transgenic plants (Fig. 3F). This suggested that sufficient CDK activity still persists in the GC after




Fig. 2. FLP directly binds the *CDKA;1* promoter. (A) Schematic diagram of four putative FLP/MYB88 *cis*-regulatory binding elements in the *CDKA;1* promoter, labelled as elements A–D. (B) Yeast-one-hybrid result shows that FLP and MYB88 can bind to fragment ‘a’ of the *CDKA;1* promoter. (C) EMSA results show that FLP can bind each binding element in *CDKA;1*. Lanes 1, 4, 7, 10, 12, probes only; lanes 2, 5, 8, 11, 13, probes with His-FLP fusion proteins; lanes 3, 6, 9, probes, His-FLP proteins, plus non-labelled competitors. Lanes 1–3, element A; lanes 4–7, element B; lanes 7–9, element C+D; lanes 10–13, nucleotides in element C or D were substituted by nucleotide A. (This figure is available in colour at JXB online.)

the terminal division, but that the levels of A- or D-type cyclins become rate limiting for another round of division (Table 1). RT-qPCR assays revealed that *CDKA;1* transcript levels normalized against the transcripts of the GC-specific *KATI* gene (Xie *et al.*, 2010) were up-regulated in both *FAMA:CYCA2;3* and *FAMA:CYCD3;2* transgenic plants, suggesting that an enhancement in *CDKA;1* expression is associated with *CYCA2;3*- or *CYCD3;2*-promoted cell proliferation (Fig. 3G). However, given that *FAMA:CDKA;1* (Fig. 5A) is not sufficient to induce stomata with extra GCs, the level of cyclins might become rate limiting for divisions in the GC.

Elevating *CYCA2;3* levels in GC causes cellular fate reversion in GCs

FAMA is also required for GMC to GC differentiation to acquire GC fate. Loss of *FAMA* function leads to narrow cell tumours that lack GC fate. *FAMA:CYCD3;2* was crossed into a *fama-1* null mutant. Elevating *CYCD3* levels did not trigger a synergistic effect on cell proliferation in the *fama-1* mutant background (Fig. 4A). In contrast, *FAMA:CYCD3;2* in the *flp-1 myb88* double mutant background, in which some cells of the stomatal clusters could acquire GC fate, produced perpendicular as well as parallel divisions, an additive phenotype consistent with *FAMA:CYCD3;2* inducing GC subdivisions after GMC division (Fig. 4B).

Table 1. Quantification of formation of single guard cells in cotyledon epidermis

Genotype	SGC (%)	SGC	Counts
Col	0	0	680
<i>cdkb1;1 1;2</i>	31	384	1238
<i>FAMA:CDKB1;1 cdkb1;1 1;2</i> line 1	0	0	964
<i>FAMA:CDKB1;1 cdkb1;1 1;2</i> line 2	0	0	1000
<i>FAMA:CDKA;1 cdkb1;1 1;2</i> line 1	22.5	231	1026
<i>FAMA:CDKA;1 cdkb1;1 1;2</i> line 2	17.6	115	650
<i>35S:CDKB1;1.N161</i>	13	169	1308
<i>FAMA:CYCA2;3 35S:CDKB1.N161</i>	2.3	35	1502
<i>FAMA:CYCD3;2 35S:CDKB1.N161</i>	0	0	852

Expression of *CDKB1;1:GUS* is found in GMCs and young GCs, but is low in mature GCs (Fig. 4C). Overexpression of *FAMA:CYCD3;2* induced an enhanced expression of *CDKB1;1:GUS* in subdivided GCs. However, the equal *GUS* expression levels among all subdivided GCs indicate that these divisions are symmetric (Fig. 4F). *TMM:TMM-GFP* (green fluorescent protein) is a stomatal lineage marker expressed in GMCs, young GCs, and stomatal lineage ground cells; but it is absent in mature guard cells (Fig. 4D). The absence of strong *TMM:TMM-GFP* expression in *FAMA:CYCD3;2* subdividing GCs suggests that these extra divisions occurred at the final stage of stomatal development (Fig. 4G).




Fig. 3. Overexpression of *CYCA2;3* and *CYCD3;2* can reduce the number of SGCs in *35S:CDKB1;1.N161*. (A) Wild-type epidermis. (B) SGCs were found in *35S:CDKB1;1.N161* epidermis. (C) *FAMA:CYCA2;3*-induced GC subdivisions, leading to stomata containing 3–4 GCs. (D) *FAMA:CYCA2;3* partially compensates the defects of GMC division in *35S:CDKB1;1.N161*. Inset: an occasionally found SGC (refer to [Supplementary Fig. S4](#) at *JXB* online). (E) *FAMA:CYCD3;2* induces GCs to subdivide, leading to GCs with 3–4 stomata. (F) No SGC was found in *35S:CDKB1;1.N161* plants harbouring *FAMA:CYCD3;2*. (G) The *CDKA;1* transcript levels in *FAMA:CYCD3;2* and *FAMA:CYCA2;3* transgenic plants. Bars=20 μ m. (This figure is available in colour at *JXB* online.)

Moreover, like the wild type, all subdivided GCs eventually express the mature GC marker E1728 (Fig. 4E, H), indicating that *CYCD3;2*-induced extra divisions are indeed symmetric divisions in GC and uncoupled from GC differentiation.

Overexpression of *CYCA2;3* at the stage of *FAMA* expression caused a dramatic increase in the *CYCA2;3* transcription level ([Supplementary Fig. S3A](#) at *JXB* online). Expression of *FAMA:CYCA2;3* also induced a differential increasing expression of *CDKB1;1:GUS* in some subdivided GCs, which was not observed in *FAMA:CYCD3;2* lines (arrow in Fig. 4I versus F). Strikingly, ectopic high expression of *TMM:TMM-GFP* was found in some subdivided cells in *FAMA:CYCA2;3* stomata, implying a cell fate reversion from a GC back to a precursor cell or at least to an immature GC (Fig. 4J). Consistently, the GC marker E1728 is not always expressed evenly in the different subdivided cells within the same stoma, implying that their GC differentiation was disturbed (Fig. 4K). In addition, collapsed cells were often found in *FAMA:CYCA2;3* stomata, which are not found in *FAMA:CYCD3;2*, confirmed by a cell viability assay using

Neutral Red dyes ([Supplementary Figs S3B–E, S4](#) at *JXB* online).

CDKA;1–CYCD3;2 complexes stimulate extra symmetric GC subdivisions

To probe the genetic relationship between *CDKA;1* and *CYCD3;2* during stomatal development, targeted *CYCD3;2* and *CDKA;1* overexpression was initially introduced individually into wild-type plants under the respective control of the *SPCH*, *MUTE*, and *FAMA* promoters. Subsequently, lines with *CYCD3;2* and *CDKA;1* driven by the same promoter were crossed. No obvious stomatal defects were found in plants harbouring both *SPCH:CDKA;1* and *SPCH:CYCD3;2*, or in those containing *MUTE:CDKA;1* and *MUTE:CYCD3;2* ([Supplementary Fig. S5A, B](#) at *JXB* online). Although stomatal development is normal in *FAMA:CDKA;1* transformants (Fig. 5A), co-expression of *FAMA:CDKA;1* with *FAMA:CYCD3;2* greatly enhanced the GC subdivision




Fig. 4. Ectopic expression of *CYCD3;2* or *CYA2;3* induces GC subdivisions. (A) The loss of *FAMA* function phenotype is epistatic to *FAMA:CYCD3;2*, presumably because *fama-1* blocks the acquisition of GC fate. Inset: a *fama-1* tumour. (B) *FAMA:CYCD3;2* and *flp-1 myb88* together show additive phenotypes of stomatal clusters and GC subdivisions. Inset: *flp-1 myb88* stomatal cluster. (C) *CDKB1;1:GUS*, a marker for the stomatal terminal differentiation stage, expressed from the transition stage of GMC to GC in the wild type. Note the weak *GUS* expression in a mature stoma (an arrow). (D) *TMM:TMM-GFP* expression in stomatal lineage cells in the wild type. (E) *E1728*, a mature GC marker expressed in wild-type GCs. (F) Equally enhanced *CDKB1;1:GUS* expression in a subdivided *FAMA:CYCD3;2* stoma. Arrowheads point to the extra cell division planes. (G) No *TMM:TMM-GFP* expression in *FAMA:CYCD3;2*-subdivided GCs. (H) Subdivided GCs induced by *FAMA:CYCD3;2* eventually exhibit a GC fate as shown by the expression of the *E1728* marker. An arrow points to the subdivision plane. (I) Expression of *CDKB1;1:GUS* in a stoma with four GCs of *FAMA:CYA2;3*. An arrow points to the GC showing a differential higher *GUS* staining than its sister cell. (J) Ectopic expression of *TMM:TMM-GFP* could be found in one of the subdivided GCs, indicated by an arrow. (K) *FAMA:CYA2;3*-induced GC subdivision mimics *FAMA:CYCD3;2*, but some of them were unequal divisions. *E1728* GFP fluorescence is from a mature GC but was absent in another GC (arrow). Bars=20 μ m. (This figure is available in colour at *JXB* online.)

phenotype (Fig. 5B; Supplementary Fig. S5C at *JXB* online). In *FAMA:CYCD3;2* plants, most stomata (90%) contained 3–4 GCs, while 5% were normal and 5% had >4 GCs. The co-expression of both *FAMA:CYCD3;2* and *FAMA:CDKA;1* in wild-type plants increased the fraction of stomata containing >4 GCs to 21% and the percentage of normal stomata was reduced to 1% (Fig. 6).

In contrast, the co-expression of a dominant-negative version of *FAMA:CDKA;1.N146* with *FAMA:CYCD3;2* increased the fraction of normal stomata to ~30% and decreased the number of stomata containing >4 GCs (Fig. 6). However, *CDKB1;1* activity is not required for *CYCD3;2*'s functions, since co-expression with either *FAMA:CDKB1;1* or *35S:CDKB1;1.N161* has no impact on




Fig. 5. The synergistic effect of co-expression of *CYCD3;2* and *CDKA;1* in promoting GC subdivisions requires *RBR1*. (A) No obvious stomatal defect was detected in *FAMA:CDKA;1*. (B) Co-expression of *FAMA:CDKA;1* with *FAMA:CYCD3;2* augments GC subdivisions. (C, D) Co-expression of *FAMA:CYCA2;3* with either *FAMA:CYCD3;2* or *FAMA:CDKA;1* mimics stomatal phenotypes in *FAMA:CYCA2;3*. (E) GC subdivisions in *FAMA:amiRNA-RBR1*. (F) *RBR1* transcription is significantly suppressed in a *FAMA:amiRNA-RBR1* line. Bars=20 μ m. (This figure is available in colour at JXB online.)

FAMA:CYCD3;2-induced GC subdivisions (Fig. 6). The above synergistic effects on GC subdivisions were not found between *CYCA2;3* and *CDKA;1*, or between *CYCA2;3* and *CYCD3;2* (Fig. 5C, D; Supplementary Fig. S4 at JXB online).

The potential for *CYCD3;2* to interact directly with *CDKA;1* was demonstrated in yeast two-hybrid assays and BiFC *in vivo* assays (Supplementary Fig. S5D, E at JXB online). Furthermore, a pull-down experiment further proved that *CYCD3;2* forms a protein complex with *CDKA;1* *in vitro* (Supplementary Fig. S5F at JXB online).

Suppression of *RBR1* also induced GC subdivisions

FAMA-driven expression of *CYCA2;3* or *CYCD3;2* induced 3–4 cells radially arranged around the pore. This

stomatal phenotype resembles those stomata found in an inducible RNAi against *RBR1* plants (Borghi *et al.*, 2010). Development of pavement cells and stomatal formation are also disrupted by the loss of *RBR1* function, consistent with its wide expression in the epidermis (Supplementary Fig. S6A at JXB online). GC subdivision is observed in a line of artificial microRNA (*amiRNA*) against *RBR1* under the *FAMA* promoter, *FAMA:amiRNA-RBR1*, in which *RBR1* transcription is significantly suppressed (Fig. 5E, F).

The *RBR1:GUS* expression pattern and RT-qPCR assays reveal that the impact of *FAMA:CYCD3;2* on *RBR1* transcription is limited (Supplementary S6B, C at JXB online). DAPI staining analysis demonstrates that the DNA ploidy levels in the subdivided GCs found in *FAMA:amiRNA-RBR1* and *FAMA:CYCD3;2* epidermis are the same as in normal GCs, indicating that ectopic expression of *CYCD3;2* or suppression of *RBR1* promotes cell proliferation only (Supplementary Fig. S6D–J).

Most of the stomata in *FAMA:CYCD3;2 FAMA:CDKA;1* consist of 3–4 GCs (78%), while 21% of stomata have >4 GCs. Each daughter GC has its own cell outline and contacts with the adjacent GCs, displaying a ‘string of sausages’-like stoma (Type II in Supplementary Fig. S7 at JXB online). In one of the *FAMA:amiRNA-RBR1* lines, 25.7% stomata mimic the ‘string of sausages’-like stomata. Around 30% of stomata in this *FAMA:amiRNA-RBR1* line exhibit ectopic divisions in their GCs. However, these extra divisions happened in original GCs (produced by GMC division) as well, but did not alter the original GC shape (Type III in Supplementary Fig. S7).

Discussion

CDKA;1 activity is required for GMC division

Whereas in yeast a single *CDKA;1* regulates both the G_1 to S and G_2 to M transition of the mitotic cell cycle, the *CDKB* kinases, together with *CDKA;1* kinase complexes, operate at the G_2 to M transition in *Arabidopsis* plants (Menges *et al.*, 2005). *CDKA;1* has pleiotropic functions in plant development, including the maintenance of shoot and root apical meristems (Dissmeyer *et al.*, 2009; Nowack *et al.*, 2012). Mutations in *CDKA;1* induce a failure in the double fertilization and lead to the abortion of homozygous *cdka;1* seeds. Homozygous *cdka;1* mutants recovered using a *PRO_{CDKA;1}:CDKA;1:YFP* construct exhibit defects in embryogenesis and in the maintenance of the root and shoot apical meristems (Nowack *et al.*, 2012). Homozygous *cdka;1* seedlings were recovered via a pollen-specific promoter to drive *CDKA;1* gene expression (*LAT52:CDKA;1*). These *cdka;1* homozygous seedlings showed significant reduction in stomatal formation and undivided GMCs, indicating that *CDKA;1* activity is required for both stomatal formative asymmetric divisions and terminal symmetric divisions, consistent with the expression of *CDKA;1* throughout the stomatal lineage cells (Serna and Fenoll, 1997). Using a dominant-negative *CDKA;1.N146* construct driven by either the *MUTE* or the *FAMA* promoter, it was possible to dissect further the role of *CDKA;1* in stomatal development. Triggering the expression




Fig. 6. Quantification of effects of different CDK forms on *FAMA:CYCD3;2* stomatal phenotypes. Quantification of stomatal phenotypes from 14-day-old cotyledons. A total of 800–1000 stomata from 20–30 cotyledons were scored for each genotype. Black, SGCs; oblique, normal stomata with two GCs; grey, stomata with three GCs; hatched, stomata with four GCs; dotted, stomata with >4 GCs.

of dominant-negative *CDKA;1.N146* with the *MUTE* promoter impaired the GMC symmetric division in forming the two GCs, whereas later expression of *CDKA;1.N146* with the *FAMA* promoter hardly influenced the symmetric division, indicating that the timing of *CDKA;1* activity plays an important role in the early events corresponding to the specification of the GMC. However, even in these GMCs with reduced *CDKA;1* activity, the cell cycle can be uncoupled from GMC to GC differentiation as the undivided GMC arrested early in the cell cycle can eventually acquire mature GC fate, in line with the GC differentiation upon interfering with *CDKB1* activity (Xie et al., 2010).

Abnormal SGCs were first reported in *35S:CDKB1;1.N161*, then in *cdkb1;1 1;2* double mutants, and in *cyca2;234* triple mutants (Boudolf et al., 2004a; Xie et al., 2010; Vanneste et al., 2011). Despite their similar cell size, SGCs in loss-of-function *cdka;1* homozygous mutants or targeted expression of dominant-negative *CDKA;1.N146* lines contain nuclear DNA levels comparable with those of GCs with 2C DNA levels in wild-type stomata, suggesting that *CDKA;1* acts before S-phase at the G_1 to S transition of the cell cycle (Fig. 7). In contrast, SGCs produced in *cdkb1* or *cyca2* mutants display twice the DNA content of wild-type GCs, as a result of cell cycle arrested at the G_2 to M transition (Boudolf et al., 2004a; Xie et al., 2010; Vanneste et al., 2011).

Overexpression of *CDKA;1* under control of *FAMA* at the same stage could partially rescue the GMC division in the *cdkb1;1 1;2* mutant, suggesting that elevated *CDKA;1* kinase activity can, at least partially, substitute for *CDKB1* activity.

Either *CDKA;1* or *CDKB1* kinases can phosphorylate the same substrates, albeit with a different efficiency, for cell cycle progression. Alternatively, elevating the *CDKA;1* level might reduce the threshold for the overall *CDKB1* kinase activity required for G_2 to M transition. In line with this observation, expression of *CDKA;1* driven by the *TMM* promoter displayed a similar rescuing effect (Weimer et al., 2012). Although *CDKA;1* activity is generally more important for the G_1 to S transition and *CDKB1*s are required for G_2 to M progression (Boudolf et al., 2004a), the present results indicate that the mechanism for G_2 to M transition has some degree of flexibility, and that in the absence of *CDKB1*, elevated *CDKA;1* can trigger the G_2 to M transition.

CDKA;1 could be a direct target of *FLP/MYB88*

A model has been proposed for the function of *FLP/MYB88* in stomatal development, in which *FLP/MYB88* enforce cell cycle exit after GMC division by timely suppression of cell cycle genes for further divisions. This is supported by the expression pattern of the promoter reporters, such as *CDKB1;1:GFP* and *CYCA2;3:GUS-GFP* which are expressed in late GMCs and young GCs, similar to the expression pattern of *FLP/MYB88* or *FAMA* (Xie et al., 2010; Vanneste et al., 2011). Using a *CDKA;1* transcriptional reporter included a fusion of yellow fluorescent protein (YFP) with a cyclin B destruction box (DB), *ProCDKA;1:YFP-DB*, *CDKA;1* expression appears to shut off immediately after GMC division (Xie et al., 2010). Up-regulation of *CDKA;1* and *CDKB1;1* transcript levels was found in *flp-1 myb88* mutants. Hence it seems that *CDKA;1* and the earlier reported *CDKB1* genes (Xie et al., 2010) are both involved in the GMC divisions and regulated by *FLP/MYB88* transcription factors (Fig. 7).

FLP and *MYB88* genes encode an atypical two-MYB-repeat transcription factor with binding preferences different from those of other known MYBs. However, *FLP/MYB88* binds to the *CDKB1;1* promoter via the [A/T/G][A/T/G]C[C/G][C/G] consensus sequence, a *cis*-regulatory element overlapping with that of cell cycle transcription factor E2Fa (Boudolf et al., 2004b; Xie et al., 2010). Here four *cis*-regulatory elements in the *CDKA;1* promoter for *FLP/MYB88* binding were identified, suggesting that *CDKA;1*, like *CDKB1;1*, is directly regulated by the *FLP/MYB88* transcription factors, as part of the mechanism to restrict symmetric divisions to a single event.

Elevating cyclin A or D confers ectopic cell cycle activation in GC

Elevating *CDKA;1* in the GMC with the *pFAMA:CDKA;1* construct (bypassing the *FLP/MYB88* control) was not sufficient to confer additional divisions, raising the question of whether cyclin levels were rate limiting for ectopic cell cycle activation. Constitutive *CYCD3;1* expression alters leaf epidermal cell proliferation and stomatal density, but does not induce morphological defects in stomatal development (Dewitte and Murray, 2003; Dewitte et al., 2007; Elsner et al., 2012). However, the targeted expression of *CYCD3;1* in




Fig. 7. Regulatory network of terminal division of stomatal development. When the meristemoid (M) acquires a GMC fate, *CDKA;1* activity is required for G₁ to S transition in the cell cycle. In the loss-of-function *cdka;1* mutant or dominant-negative *CDKA;1.N146* transgenic lines, an undivided GMC still could acquire a GC fate, resulting in SGCs with 2C ploidy. *CDKB1* and *CYCA2* are required for the G₂ to M transition phase. In the *cdkb1;1 1;2* double mutant or dominant-negative *CDKB1;1.N161* lines, the undivided GMCs differentiate into SGCs with 4C ploidy. Mutants of *cyca2;234* also produce SGCs with 4C ploidy. *CDKA;1*, *CDKB1;1*, and *CYCA2* transcription can all be negatively regulated by FLP/MYB88 transcriptional factors through *cis*-regulatory elements. Expression of *CDKA;1* could partially rescue GMC division defects. Overexpression of *CYCA2* or *CYCD3* also reduces the number of SGCs in *CDKB1;1.N161*. In *flp myb88* and *fama* mutants, high activity of *CDKA;1* and *CDKB1* causes excessive GMC divisions. RBR1 is essential to prevent GC subdivision, but the regulatory network of CDK–cyclin–RBR1 and the feedback loop in terminal division remain unclear. (This figure is available in colour at *JXB* online.)

normally endocycling trichomes by the *GLABRA2* promoter triggers mitosis, leading to subdivided trichomes, a phenotype not observed in *35S:CYCD3;1* plants (Schellmann *et al.*, 2002; Dewitte *et al.*, 2003). A similar difference between targeted and constitutive expression of *CYCD3* genes was observed in stomatal development. Whereas constitutive *CYCD3;1* expression does not interfere with the GCs and allows for cell cycle arrest, the *FAMA* promoter probably increases local *CYCD3;2* levels sufficiently to induce GC subdivisions, a gain-of-function phenotype, like *GL2:CYCD3;1* in trichomes.

A similar phenotype was observed when *CYCA2;3* levels were increased during and after the last division. However, in the case of *CYCA2;3*, the formed GC occasionally reverted to a precursor state or caused cell collapse. Indeed, an extremely high transcript level of *CDKA;1* was observed in *FAMA:CYCA2;3* stomata, but it is not clear whether the programmed cell death was stimulated by altered levels of *CDKA;1* or RBR1, a regulation model proposed in trichome and endosperm development (Schnittger *et al.*, 2003; Sabelli *et al.*, 2013). Hence, cyclins can confer ectopic cell cycle activation, suggesting that the maintenance of two-celled stomata requires accurate control of cyclin A and D levels.

Cyclin A and D require different CDKs in order to stimulate GC division

Although the *cdkb1;1* single mutant displays normal stomata, *cdkb1;1 cyca2;234* quadruple mutants show more SGCs than *cyca2;234* triple mutants, suggesting that *CYCA2*s and *CDKB1;1* contribute synergistically to the CDK activity required for GMC division (Vanneste *et al.*, 2011). Co-expressing *CDKB1;1* with *CYCA2;3* could enhance *CYCA2;3*-associated kinase activity, demonstrating that *CYCA2;3* and *CDKB1;1* form a functional complex stimulating cell divisions. Introduction of *CYCA2;3* driven by the *35S* promoter could complement the endoreduplication phenotype of *35S:CDKB1;1.N161* plants, but could not rescue its GMC division defects (Boudolf *et al.*, 2009). Here the targeted expression of *CYCA2;3* has little effect when *CDKB1;1* function is impaired by overexpression of dominant-negative *CDKB1;1.N161*. Although overexpression of either *CYCA2;3* or *CYCD3;2* was associated with up-regulation of the *CDKA;1* transcript, it appears that only *CYCD3;2/CDKA;1*, and not *CYCA2;3/CDKA;1*, is able to stimulate the G₂ to M transition (Healy *et al.*, 2001; Imai *et al.*, 2006). The findings show that the *CYCD3;2* and *CDKA;1*

proteins can interact directly. The extensive subdivision of GCs induced by transformation with *FAMA:CYCD3;2* can in turn be partially repressed by the concomitant expression of the dominant-negative *CDKA;1.N146* construct. In addition, the degree of GC subdivision is profoundly enhanced by co-expression of *CDKA;1*. The gain or loss of function of *CDKB1* genes (via the co-expression of *CDKB1;1* or the dominant-negative *CDKB1;1.N161* construct) does not affect the extent of *FAMA:CYCD3;2*-induced GC subdivision, suggesting that the functions of *CDKA;1* and *CDKB1s* differ with respect to their cooperation with *CYCD3;2*.

RBR1 restricts GC proliferation

Arabidopsis possess only a single *RBR1* gene modulating cell division and endoreduplication (Gutzat *et al.*, 2012). Cell cycle progression is regulated through *RBR1* protein phosphorylation by G₁/S kinases such as *CYCD-CDKA;1* (Nakagami *et al.*, 2002). Indeed the function of *CDKA;1* in stomatal asymmetric division was found to be mediated by *RBR1* (Nowack *et al.*, 2012). Genes required for stomata initiation, such as *TMM* and *SPCH*, were up-regulated in *RBR1* plants, but expression of *MUTE* and *FAMA* was not affected (Borghini *et al.*, 2010). However, similar to overexpression of *CYCD3;2* or *CYCA2;3*, suppression of *RBR1* in GMCs and GCs with the *FAMA:amiRNA-RBR1* caused extra division in GCs but not in GMCs, the subdividing GCs being a phenotype distinct from the narrow parallel arranged small cells in *flp myb88* or *fama* mutants. Consistent with this, overexpressing *FAMA:CYCD3;2* in *fama-1* has no influence on the size of cell tumours. The *RBR1* level is regulated appropriately in wild-type GCs to ensure GCs do not re-enter the cell cycle and maintain GC integrity. Thus, besides a role in stomatal initiation, *RBR1* functions prevent GC overproliferation during the last stage of stomatal formation.

In summary, over-riding the control mechanisms involved in cell cycle arrest in the last phase of GC development by elevating levels of *CYCD/CDKA;1*, *CYCA/CDKB* kinases, or alternatively by suppressing the *RBR1* level sustains cell proliferation in the stomatal lineage, resulting in GC subdivisions. However, since both *CDKA;1* and *CDKB;1* activity are required for proper execution of the terminal cell cycle, timely suppression of *CDKA;1* expression by *FLP/MYB* is likely to be contributing to the final cell cycle arrest in mature stomata (Fig. 7).

Supplementary data

Supplementary data are available at *JXB* online.

Figure S1. EMSA results show that MYB88 can bind the *CDKA;1* promoter.

Figure S2. Expression of *CYCD3;1* or *CYCD3;3* promotes GC subdivisions.

Figure S3. Overexpression of *CYCA2;3* induces GC subdivision and cell collapse.

Figure S4. Quantification of the effect of co-expression of *CYCA2;3* and *CDKB1;1* in promoting GC subdivisions.

Figure S5. *CYCD3;2* directly interacts with *CDKA;1*.

Figure S6. *RBR1* expression and GC subdivision.

Figure S7. Comparison of stomatal phenotypes between *CYCD3;2 CDKA;1* co-expression and *RBR1 RNAi* lines.

Table S1. Primer sequences used in this article.

Acknowledgements

We thank Fred Sack and Eun-kyoung Lee for providing *flp-1 myb88*, *cdkb1;1* *l;2* mutants, *TMM:TMM-GFP*, and *FAMA:amiRNA-RBR1* lines. We also thank Dr Walter Dewitte for his critical reading and discussion. We thank Dr De Ye for providing *LATS2* plasmids. This work was supported by grants from the National Natural Science Foundation of China, 30971652 and 31271463 (to JL), 31071198 (to KY), Hundred Talents Program and KSCX2-YW-N-073 from the Chinese Academy of Sciences (to JL).

References

- Bergmann DC, Sack FD. 2007. Stomatal development. *Annual Review of Plant Biology* **58**, 163–181.
- Borghini L, Gutzat R, Futterer J, Laizet YH, Hennig L, Grissem W. 2010. Arabidopsis RETINOBLASTOMA-RELATED is required for stem cell maintenance, cell differentiation, and lateral organ production. *The Plant Cell* **22**, 1792–1811.
- Boudolf V, Barroco R, Engler Jde A, Verkest A, Beeckman T, Naudts M, Inze D, De Veylder L. 2004a. B1-type cyclin-dependent kinases are essential for the formation of stomatal complexes in *Arabidopsis thaliana*. *The Plant Cell* **16**, 945–955.
- Boudolf V, Lammens T, Boruc J, *et al.* 2009. CDKB1;1 forms a functional complex with *CYCA2;3* to suppress endocycle onset. *Plant Physiology* **150**, 1482–1493.
- Boudolf V, Vlieghe K, Beeckman T, Magyar Z, Acosta JAT, Maes S, Van Der Schueren E, Inze D, De Veylder L. 2004b. The plant-specific cyclin-dependent kinase CDKB1;1 and transcription factor E2Fa-DPa control the balance of mitotically dividing and endoreduplicating cells in *Arabidopsis*. *The Plant Cell* **16**, 2683–2692.
- Cruz-Ramirez A, Diaz-Trivino S, Blilou I, *et al.* 2012. A bistable circuit involving SCARECROW–RETINOBLASTOMA integrates cues to inform asymmetric stem cell division. *Cell* **150**, 1002–1015.
- De Veylder L, Beeckman T, Van Montagu M, Inze D. 2000. Increased leakiness of the tetracycline-inducible Triple-Op promoter in dividing cells renders it unsuitable for high inducible levels of a dominant negative *CDC2aAt* gene. *Journal of Experimental Botany* **51**, 1647–1653.
- Dewitte W, Murray JAH. 2003. The plant cell cycle. *Annual Review of Plant Biology* **54**, 235–264.
- Dewitte W, Scofield S, Alcasabas AA, *et al.* 2007. Arabidopsis *CYCD3* D-type cyclins link cell proliferation and endocycles and are rate-limiting for cytokinin responses. *Proceedings of the National Academy of Sciences, USA* **104**, 14537–14542.
- Dissmeyer N, Weimer AK, Pusch S, *et al.* 2009. Control of cell proliferation, organ growth, and DNA damage response operate independently of dephosphorylation of the Arabidopsis Cdk1 homolog *CDKA;1*. *The Plant Cell* **21**, 3641–3654.
- Elsner J, Michalski M, Kwiatkowska D. 2012. Spatiotemporal variation of leaf epidermal cell growth: a quantitative analysis of *Arabidopsis thaliana* wild-type and triple cyclinD3 mutant plants. *Annals of Botany* **109**, 897–910.
- Gaamouche T, Manes C-LdO, Kwiatkowska D, *et al.* 2010. Cyclin-dependent kinase activity maintains the shoot apical meristem cells in an undifferentiated state. *The Plant Journal* **64**, 26–37.
- Gardner MJ, Baker AJ, Assie J-M, Poethig RS, Haseloff JP, Webb AAR. 2009. GAL4 GFP enhancer trap lines for analysis of stomatal guard cell development and gene expression. *Journal of Experimental Botany* **60**, 213–226.
- Gutierrez C. 2008. Coordination of cell division and differentiation. In: Verma DPS, Hong Z, eds. *Cell division control in plants*. Heidelberg: Springer-Verlag, 377–393.

- Gutzat R, Borghi L, Grisse W.** 2012. Emerging roles of RETINOBLASTOMA-RELATED proteins in evolution and plant development. *Trends in Plant Science* **17**, 139–148.
- Harbour JW, Dean DC.** 2000. Chromatin remodeling and Rb activity. *Current Opinion in Cell Biology* **12**, 685–689.
- Healy JMS, Menges M, Doonan JH, Murray JAH.** 2001. The Arabidopsis D-type cyclins CycD2 and CycD3 both interact *in vivo* with the PSTAIRE cyclin-dependent kinase Cdc2a but are differentially controlled. *Journal of Biological Chemistry* **276**, 7041–7047.
- Hemerly A, de Almeida Engler J, Bergounioux C, Van Montagu M, Engler G, Inze D, Ferreira P.** 1995. Dominant negative mutants of the Cdc2 kinase uncouple cell division from iterative plant development. *EMBO Journal* **14**, 3925–3936.
- Imai KK, Ohashi Y, Tsuge T, Yoshizumi T, Matsui M, Oka A, Aoyama T.** 2006. The A-type cyclin CYCA2;3 is a key regulator of ploidy levels in Arabidopsis endoreduplication. *The Plant Cell* **18**, 382–396.
- Inze D, De Veylder L.** 2006. Cell cycle regulation in plant development. *Annual Review of Genetics* **40**, 77–105.
- Iwakawa H, Shinmyo A, and Sekine M.** 2006. Arabidopsis CDKA1;1, a cdc2 homologue, controls proliferation of generative cells in male gametogenesis. *The Plant Journal* **45**, 819–831.
- Joubès J, De Schutter K, Verkest A, Inzé D, De Veylder L.** 2004. Conditional, recombinase-mediated expression of genes in plant cell cultures. *The Plant Journal* **37**, 889–896.
- Lai L, Nadeau JA, Lucas J, Lee E-K, Nakagawa T, Zhao L, Geisler MJ, Sack FD.** 2005. The Arabidopsis R2R3 MYB proteins FOUR LIPS and MYB88 restrict divisions late in the stomatal cell lineage. *The Plant Cell* **17**, 2754–2767.
- Lau OS, Bergmann DC.** 2012. Stomatal development: a plant's perspective on cell polarity, cell fate transitions and intercellular communication. *Development* **139**, 3683–3692.
- Lee E, Liu X, Eglit Y, Sack F.** 2013. FOUR LIPS and MYB88 conditionally restrict the G1/S transition during stomatal formation. *Journal of Experimental Botany* **64**, 5207–5219.
- MacAlister CA, Ohashi-Ito K, Bergmann DC.** 2007. Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. *Nature* **445**, 537–540.
- Menges M, De Jager SM, Grisse W, Murray JAH.** 2005. Global analysis of the core cell cycle regulators of Arabidopsis identifies novel genes, reveals multiple and highly specific profiles of expression and provides a coherent model for plant cell cycle control. *The Plant Journal* **41**, 546–566.
- Nadeau JA, Sack FD.** 2002. Stomatal development in Arabidopsis. *The Arabidopsis Book*, e0066.
- Nakagami H, Kawamura K, Sugisaka K, Sekine M, Shinmyo A.** 2002. Phosphorylation of retinoblastoma-related protein by the cyclin D/cyclin-dependent kinase complex is activated at the G1/S-phase transition in tobacco. *The Plant Cell* **14**, 1847–1857.
- Nowack MK, Harashima H, Dissmeyer N, Zhao XA, Bouyer D, Weimer AK, De Winter F, Yang F, Schnittger A.** 2012. Genetic framework of cyclin-dependent kinase function in Arabidopsis. *Developmental Cell* **22**, 1030–1040.
- Ohashi-Ito K, Bergmann DC.** 2006. Arabidopsis FAMA controls the final proliferation/differentiation switch during stomatal development. *The Plant Cell* **18**, 2493–2505.
- Pillitteri LJ, Sloan DB, Bogenschutz NL, Torii KU.** 2007. Termination of asymmetric cell division and differentiation of stomata. *Nature* **445**, 501–505.
- Pillitteri LJ, Torii KU.** 2012. Mechanisms of stomatal development. *Annual Review of Plant Biology* **63**, 591–614.
- Porter A.** 2008. Preventing DNA over-replication: a Cdk perspective. *Cell Division* **3**, 3.
- Sabelli PA, Liu Y, Dante RA, et al.** 2013. Control of cell proliferation, endoreduplication, cell size, and cell death by the retinoblastoma-related pathway in maize endosperm. *Proceedings of the National Academy of Sciences, USA* **110**, E1827–E1836.
- Schellmann S, Schnittger A, Kirik V, Wada T, Okada K, Beermann A, Thumfahrt J, Jurgens G, Hulskamp M.** 2002. TRIPTYCHON and CAPRICE mediate lateral inhibition during trichome and root hair patterning in Arabidopsis. *EMBO Journal* **21**, 5036–5046.
- Schnittger A, Weinl C, Bouyer D, Schobinger U, Hulskamp M.** 2003. Misexpression of the cyclin-dependent kinase inhibitor ICK1/KRP1 in single-celled Arabidopsis trichomes reduces endoreduplication and cell size and induces cell death. *The Plant Cell* **15**, 303–315.
- Serna L, Fenoll C.** 1997. Tracing the ontogeny of stomatal clusters in Arabidopsis with molecular markers. *The Plant Journal* **12**, 747–755.
- Tang W, Wang W, Chen D, Ji Q, Jing Y, Wang H, Lin R.** 2012. Transposase-derived proteins FHY3/FAR1 interact with PHYTOCHROME-INTERACTING FACTOR1 to regulate chlorophyll biosynthesis by modulating HEMB1 during deetiolation in Arabidopsis. *The Plant Cell* **24**, 1984–2000.
- Vanneste S, Coppens F, Lee E, et al.** 2011. Developmental regulation of CYCA2s contributes to tissue-specific proliferation in Arabidopsis. *EMBO Journal* **30**, 3430–3441.
- Walter M, Chaban C, Schütze K, et al.** 2004. Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *The Plant Journal* **40**, 428–438.
- Weimer AK, Nowack MK, Bouyer D, Zhao X, Harashima H, Naseer S, De Winter F, Dissmeyer N, Geldner N, Schnittger A.** 2012. Retinoblastoma related1 regulates asymmetric cell divisions in Arabidopsis. *The Plant Cell* **24**, 4083–4095.
- Xie Z, Lee E, Lucas JR, Morohashi K, Li D, Murray JAH, Sack FD, Grotewold E.** 2010. Regulation of cell proliferation in the stomatal lineage by the Arabidopsis MYB FOUR LIPS via direct targeting of core cell cycle genes. *The Plant Cell* **22**, 2306–2321.