



Correction

Correction: Post-Heparin LPL Activity Measurement Using VLDL As a Substrate: A New Robust Method for Routine Assessment of Plasma Triglyceride Lipolysis Defects

The *PLOS ONE* Staff

Notice of Republication

This article was republished on May 14, 2014, due to multiple instances of " \pm " being published as "+" The publisher apologizes for this error. Additionally, some symbols were missing from Figure 3. Please download the PDF again to view the corrected article and Figure 3. The originally published, uncorrected article and the republished, corrected article are provided here for reference.

Supporting Information

File S1. Originally published, uncorrected article.

(PDF)

File S2. Republished, corrected article.

(PDF)

Reference

1. Di Filippo M, Marçais C, Charrière S, Marmontel O, Broyer M, et al. (2014) Post-Heparin LPL Activity Measurement Using VLDL As a Substrate: A New Robust Method for Routine Assessment of Plasma Triglyceride Lipolysis Defects. *PLoS ONE* 9(5): e96482. doi:10.1371/journal.pone.0096482

Citation: The *PLOS ONE* Staff (2014) Correction: Post-Heparin LPL Activity Measurement Using VLDL As a Substrate: A New Robust Method for Routine Assessment of Plasma Triglyceride Lipolysis Defects. *PLoS ONE* 9(6): e99721. doi:10.1371/journal.pone.0099721

Published: June 2, 2014

Copyright: © 2014 The *PLOS ONE* Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.