

Published in final edited form as:

Inorg Chem. 2013 May 6; 52(9): 5236–5245. doi:10.1021/ic400171z.

Sequential oxidations of thiolates and the cobalt metallocenter in a synthetic metallopeptide: Implications for the biosynthesis of nitrile hydratase

Arnab Dutta^{†,§}, Marco Flores[†], Souvik Roy^{†,§}, Jennifer C. Schmitt[‡], G. Alexander Hamilton[†], Hilairy E. Hartnett^{†,∞}, Jason Shearer[‡], and Anne K. Jones^{†,§,*}

[†]Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ 85287

[§]Center for Bio-Inspired Solar Fuel Production, Arizona State University, Tempe, AZ 85287

[∞]School of Earth and Space Exploration; Arizona State University, Tempe, AZ 85287

[‡]Department of Chemistry, University of Nevada, Reno, Nevada 89557

Abstract

Cobalt nitrile hydratases (Co-NHase) contain a catalytic cobalt (III) ion coordinated in an N₂S₃ first coordination sphere comprised of two amidate nitrogens and three cysteine-derived sulfur donors: a thiolate (-SR), a sulfenate (-S(R)O⁻), and a sulfinato (-S(R)O₂⁻). The sequence of biosynthetic reactions that leads to the post-translational oxidations of the metal and the sulfur ligands is unknown, but the process is believed to be initiated directly by oxygen. Herein we utilize cobalt bound in an N₂S₂ first coordination sphere by a seven amino acid peptide known as SODA (ACDLPCG) to model this oxidation process. Upon exposure to oxygen, Co-SODA is oxidized in two steps. In the first fast step (seconds), magnetic susceptibility measurements demonstrated that the metallocenter remains paramagnetic, i.e. Co²⁺, and sulfur K-edge X-ray absorption spectroscopy (XAS) is used to show that one of the thiolates is oxidized to sulfinato. In a second process on a longer timescale (hours), magnetic susceptibility measurements and Co K-edge XAS show that the metal is oxidized to Co³⁺. Unlike other model complexes, additional slow oxidation of the second thiolate in Co-SODA is not observed, and a catalytically active complex is never formed. The likely reason is the absence of the axial thiolate ligand. In essence, the reactivity of Co-SODA can be described as intermediate between previously described models which either quickly convert to final product or are stable in air, and it offers a first glimpse into a possible oxidation pathway for nitrile hydratase biosynthesis.

Introduction

Post-translational modification of cysteinyl thiolates (RS⁻) to such products as disulfide bonds (RS-SR'), sulfenates (RSO⁻) or sulfinates (RSO₂⁻) is relatively common in biology and has been found to play roles in signal transduction, protein structure, and enzyme

*Corresponding Author: jonesak@asu.edu Tel: 480-965-0356. Fax: 480-965-2747.

Supporting Information Additional mass spectral, NMR, and EXAFS data. Partial molecular orbital diagrams and hypothetical oxidation scheme. This material is available free of charge via the Internet at <http://pubs.acs.org>.

functionality.^{1–3} Cysteine residues are also commonly used in metalloenzymes to ligate metals or metalloclusters. Nonetheless, the hydrolytic enzymes nitrile hydratase (NHase) and thiocyanate hydrolase (SCNase) are the only known examples of metalloenzymes requiring sulfenate and sulfinato ligands for activity.^{4–12} The active site of NHase consists of a low spin Fe(III) or Co(III) ion in an N_2S_3 coordination environment provided by two amide nitrogens from the protein backbone and three cysteines found in a highly conserved $CXXC(SO_2H)SC(SOH)$ motif (Figure 1). It is interesting to note that each of the three coordinating cysteines is present in a different oxidation state: one thiolate, one sulfenate and one sulfinato. The oxidized cysteines are found in equatorial coordination sites along with the amides, and the thiolate ligand occupies an axial position *trans*- to the presumed substrate-binding site. A number of roles have been proposed for the oxidized cysteines including modulating the Lewis acidity of the metal site, serving as a base during the catalytic cycle, and modifying substrate/product binding affinity.¹³

Conversion of nitriles to amides is an industrially important reaction.^{14, 15} To help elucidate the mechanistic details of the enzymatic catalysis of these transformations, a number of small-molecule mimics of NHase have been synthesized.^{16–22} However, few of these have even modest nitrile hydrolysis activity, and most of those that utilize oxidized sulfur contain bis-sulfinato ligands since controlling the extent of oxidation of the metal-bound thiolates has proven difficult.²³ This has been especially true for reactions of thiolates with oxygen, as opposed to O-atom transfer reagents, which tend to yield sulfinato complexes.^{18, 23–34} Nonetheless, oxygen is believed to be the oxidant *in vivo* for modification of the sulfur atoms of NHase.^{5, 12, 35–37} This stands in marked contrast to a wealth of existing synthetic inorganic literature describing synthesis of metal bound sulfinato species in which reactions with molecular oxygen are seldom reported. Generally, Co(III) and other metal sulfinato complexes have been prepared by one of three routes: direct reaction of a sulfinato salt and Co(III), SO_2 insertion into a Co(III)-C bond, or oxidation of a Co(III) thiolate via hydrogen peroxide, a much stronger oxidant than molecular oxygen.^{38–44} Biosynthetic studies have shown that apo-NHase, *i.e.*, the metal-free protein with cysteine thiolates, can be activated by incubation with Co^{2+} and appropriate activation mediating proteins via a novel mechanism that has been referred to as “self-subunit swapping”.^{35–37} First, the cobalt binding α -subunit is post-translationally modified and loaded with cobalt while the protein is part of a trimeric α_2 complex with NhlE, a maturation mediator. Then, the holo- α -subunit is exchanged with an apo- α -subunit from the tetrameric apo- $\alpha_2\beta_2$ NHase to form the functional enzyme. Surprisingly, cobalt cannot be directly incorporated into apo- $\alpha_2\beta_2$ NHase, and this swapping of apo- and holo- proteins is an obligatory part of the maturation. However, although this symphony of protein subunits is now well documented, the chemical steps allowing for specific cysteine modification, cobalt incorporation and oxidation, and exchange of the two forms of the α -subunit are unknown.

Recently, Shearer and coworkers demonstrated the first functional metallopeptide mimic of Co-NHase.²² The peptide coordinates Co in an N_2S_3 coordination sphere, and, as for the enzyme, two of the cysteines must be oxidized to observe modest activity with acrylonitrile. In this work, we utilize a related seven amino acid peptide known as SODA ($ACDLPCG$) to coordinate Co(II) in a less sulfur-rich N_2S_2 environment. We show that, upon exposure of

the Co(II)-metallopeptide to air, the first oxidation occurs at the ligands without a change in the oxidation state of the metal (Scheme 1). This ligand-oxidized, sulfur bound Co(II) system is then further oxidized to Co(III) to produce a fully oxidized species (Scheme 1). We propose that these differentially oxidized states are relevant to understanding the biosynthesis of Co-NHase, and possible mechanisms for metal incorporation into the enzyme and the roles of the distinctively oxidized thiolates present in the first coordination sphere during this process are discussed (Scheme S1).

Experimental Section

All inorganic syntheses were performed under a nitrogen atmosphere using a double-manifold Schlenk vacuum line. Unless otherwise specified, all the chemicals and solvents, of highest available grades, were obtained from Sigma-Aldrich (Milwaukee, WI) and were used as received. Aqueous solutions were prepared using purified water (resistivity = 18.2 M Ω -cm).

Physical Measurements

Optical, FTIR, and CD spectroscopy

UV-visible measurements were executed on a Hewlett-Packard 8453 spectrophotometer using quartz cuvettes with a 1 cm pathlength. NMR spectra were recorded at 400 MHz (^1H) using Varian Liquid-State NMR instruments in D₂O (99.9%, Cambridge Isotopes Laboratories Inc.). FTIR (Fourier-transform infrared) spectra of the samples were recorded on a Thermo Nicolet Avatar-360 spectrometer in KBr pellets. A KBr pellet without sample was used as a reference for the background. For the samples, each spectrum is an average of 512 scans at 1 cm⁻¹ resolution. The KBr pellets for the reduced samples were prepared in an anaerobic glove box. Circular dichroism (CD) spectroscopy was performed on a Jasco-815 spectropolarimeter using a rectangular quartz cell with a pathlength of 0.1 cm.

EPR spectroscopy

Electron paramagnetic resonance (EPR) spectra were recorded between 5K and 7K using a Bruker ELEXSYS E580 CW X-band spectrometer (Bruker, Silberstreifen, Germany) equipped with an Oxford model 900 EPL liquid helium cryostat (Oxford instruments, Oxfordshire, UK). The magnetic field modulation frequency was 100 kHz; the amplitude was 1 mT; the microwave power was between 1–4 mW and the microwave frequency was 9.42 GHz; the sweep time was 168 s. Samples for the EPR experiments were prepared in a 1:1 mixture of aqueous 50 mM N-ethyl morpholine (NEM) buffer (pH 7.5) and glycerol (BDH, West Chester, PA) before transfer to a quartz EPR tube. EPR spectra were simulated using EasySpin (v 3.1.1), a computational package developed by Stoll and Schweiger⁴⁵ and based on Matlab (The MathWorks, Natick, MA, USA). The model used for the EPR simulations is based on a single high-spin Co²⁺ ion ($S = 3/2$). The variable parameters were: the g values (g_x , g_y and g_z), the line widths (B_x , B_y and B_z), and the zero-field splitting parameters D and E . The fitting procedure was similar to the one previously described by Flores *et al.*⁴⁶

Magnetic susceptibility

The molecular magnetic susceptibility (χ_M), effective magnetic moment (μ_{eff}) for the partially oxidized Co-SODA, and the time course measurement for the oxidation of reduced Co-SODA after exposure to air were measured via the Evans NMR method by monitoring shifts in the water solvent peaks at either 400 or 500 MHz (^1H) using a Varian Liquid State NMR instrument.^{47, 48}

Mass spectrometry

MALDI-MS (matrix assisted laser desorption/ionization mass spectrometry) characterization of apo-SODA was performed using α -cyano-4-hydroxycinnamic acid as matrix on a Voyager DE STR. ESI-MS (electrospray ionization mass spectrometry) was accomplished using a Thermo Quantum Discovery Max triple-quadrupole mass spectrometer. Measurements were conducted in positive (+) and negative (-) ionization modes, using a methanol/water (50:50 by volume) mobile phase at a flow rate of $10 \mu\text{L min}^{-1}$ and the following ionization conditions: spray voltage, 4000 (+, -); capillary temperature, 270°C ; sheath gas pressures, 25 (+) and 15 (-); auxiliary gas pressure, 2 (+, -). Isotope Pattern Calculator v4.0, developed by Junhua Yan, was used for the calculation of the molecular mass with isotopic abundances.

Co K-edge and S K-edge X-ray Absorption Data Collection and Analysis

Solutions of Co-SODA were prepared as 1:1 CoSODA (50 mM NEM buffer, pH 7.5):glycerol samples, injected in aluminum sample holders between windows made from Kapton tape (3M; Minneapolis, MN, catalog no. 1205) and quickly frozen in liquid nitrogen. Co K-edge data were collected on beamline X3b at the National Synchrotron Light Source (Brookhaven National Laboratories; Upton, NY). The samples were maintained at 20 K throughout data collection in an evacuated He Displex cryostat and collected in fluorescence mode on a Canberra 31 element solid-state Ge detector. Total count rates for all channels were maintained under 30 kHz, and a deadtime correction was not applied. The data were collected in 10 eV steps below the edge (7609 – 7689 eV), 0.5 eV steps in the edge region (7689 – 7759 eV), 2 eV steps in the near edge region (7759 – 8009 eV) and 5 eV steps in the far edge region (8009 eV – 14 keV). The spectra represent the averaged sum of three independent data scans. The sulfur K-edge data were collected as solution samples at room temperature on beamline X19a at the National Synchrotron Light Source (Brookhaven National Laboratories; Upton, NY). Solutions of CoSODA (50 mM NEM buffer, pH 7.5) were injected into Lucite sample holders between two polyester film windows ($2.5 \mu\text{m}$ thickness; SPI Supplies; West Chester, PA, catalogue no. 01865-AB) adhered to the sample holders using an epoxy resin. Data were collected in a sample chamber that was continuously purged with He(g) and data were recorded in fluorescence mode on a Canberra passivated implanted planar silicon (PIPS) detector. Data were collected from 200 eV below to 300 eV above the edge to obtain a good baseline for data analysis. In the pre-edge region, data were collected in 5 eV steps (2270 – 2465 eV); in the edge region, data were collected in 0.1 eV steps (2465 – 2475 eV); in the far edge region, data were collected in 2 eV steps (2475 – 2765 eV). All data were calibrated to the white line of Na_2SO_4 . The spectra presented represent the summed average of 5 scans.

All Co and S K-edge spectra were processed and analyzed as previously described.^{22, 49} Briefly, S K-edge spectra were obtained by averaging the data sets, applying a baseline, and fitting the pre-edge region to a polynomial function. This baseline was then subtracted from the whole spectrum. The region above the edge jump was fit to a two knot cubic spline, and the data normalized to the edge height. The pre-edge and rising-edge features were modeled as pseudo-Voigt line shapes (a 1:1 sum of Gaussian and Lorentzian line shapes). Co K-edge spectra were processed and analyzed using the XAS data analysis suit *EXAFS123*⁵⁰ which operates in Igor Pro⁵¹ and requires phase and amplitude functions generated within FEFF 8.20.⁵² All refinements are based on Fourier-filtered $k^3(\chi)$ data over the energy range of $k = 2.0 - 12.0 \text{ \AA}^{-1}$ and back-transformed over the range of $r = 1.0 - 3.0 \text{ \AA}$.

Peptide synthesis and purification

The seven amino acid peptide referred to as SODA (ACDLPCG) was synthesized, purified and quantitated as described previously.⁵³ The reaction yield was 60% and the measured peptide mass (678 m/z, MALDI-MS (positive ion) $[M]^+$) agreed well with the calculated peptide mass, 677.8 amu.

Synthesis and purification of NHasem1 (CCDLPCGVYDPA) and the subsequent preparation of Co-NHasem1 were performed as described by Shearer *et al.*²²

Co-SODA preparation in reduced and oxidized forms

Metallopeptide samples were prepared analogously to the procedure described by Neupane *et al.* for the synthesis of the corresponding Ni-metallopeptide.⁵⁴ One equivalent of an aqueous solution of Co(II) (ten times concentrated compared to SODA solution) was added dropwise to one equivalent of SODA in 50 mM NEM buffer (pH= 7.5) solution. The concentration of the SODA solution was determined using Ellman's test.⁵⁵ Both solutions were degassed and saturated with nitrogen prior to the reaction. The reaction mixture immediately turned green upon addition of the cobalt solution. If exposed to oxygen, the reduced Co-SODA solution rapidly changes to brown.

Reduced Co-SODA—The electronic absorption spectrum (in buffered aqueous solution, pH 7.5) had the following features: λ_{max} nm ($\epsilon \text{ M}^{-1}\text{cm}^{-1}$): 725 (210), 681 (230), 609 (225), 350 (shoulder); 303 (2250). The ESI MS (negative ion mode) yielded a $[M-H]^-$ peak with an m/z of 731.01 that corresponded well with the calculated mass of 731.1 amu for this species.

Partially oxidized Co-SODA—The features of the electronic absorption spectrum (in buffered aqueous solution, pH 7.5) are as follows: λ_{max} nm ($\epsilon \text{ M}^{-1}\text{cm}^{-1}$): 600 (shoulder), 460 (shoulder), 355 (1100). ESI MS (positive ion mode) yielded an $[M+O+3H]^+$ molecular ion peak at 752.04 m/z; the calculated mass was 752.17 amu. The measured effective magnetic moment was $\mu_{\text{eff}} = 4.38 \text{ B.M.}$

Fully oxidized Co-SODA—The electronic absorption spectrum features (in buffered aqueous solution, pH 7.5) included: λ_{max} nm ($\epsilon \text{ M}^{-1}\text{cm}^{-1}$): 570 (shoulder), 450 (shoulder), 340 (1500). The ESI-MS (positive ion mode) yielded an $[M+O+3H]^+$ molecular ion peak at 752.06 m/z, and the calculated mass was 752.17 amu.

Results and Discussion

Coordination of Co(II) by SODA

The seven amino acid peptide SODA (ACDLPCG) is derived from the nickel enzyme superoxide dismutase and is known to coordinate a Ni(II) ion via the two cysteines together with the amino terminus and a backbone amide nitrogen.^{56–58} Thus it seemed likely that a similar coordination set could accommodate Co(II), and the interactions between Co(II) and SODA were investigated via optical spectroscopy under both anaerobic and aerobic conditions. First, an anaerobic aqueous Co(II) solution was added dropwise to a buffered SODA solution (pH 7.5) resulting in a green solution. As shown in Figure 2, the UV-vis absorbance spectrum of this reduced, cobalt bound metalloprotein includes three distinct ligand field (LF) bands in the visible region at 725 nm ($\epsilon = 210 \text{ M}^{-1}\text{cm}^{-1}$), 681 nm ($\epsilon = 230 \text{ M}^{-1}\text{cm}^{-1}$) and 609 nm ($\epsilon = 225 \text{ M}^{-1}\text{cm}^{-1}$). These bands have the appropriate energies and extinction coefficients for the ${}^4\text{T}_1(\text{P}) \leftarrow {}^4\text{A}_2(\text{F})$ transitions expected to arise from a tetrahedral Co(II) (d^7) center (Table 1, Figure 2), and are split, as expected, due to spin-orbit coupling.^{59–61} The stoichiometry of the interaction between Co(II) and SODA was determined by monitoring the absorbances at these three wavelengths as a function of added Co(II) as shown in the inset of Figure 2. Titrations at all three wavelengths are consistent with a 1:1 interaction between Co(II) and SODA. Further evidence for 1:1 binding was provided by mass spectrometry. A $[\text{Co-SODA}]^-$ molecular ion peak with the expected isotopic pattern was observed in a negative-mode ESI-MS experiment (Figure S1). In addition to the LF bands, two stronger bands at 350 nm ($\epsilon = 1650 \text{ M}^{-1}\text{cm}^{-1}$) and 303 nm ($\epsilon = 2250 \text{ M}^{-1}\text{cm}^{-1}$) were observed in the far UV region of the optical spectrum for reduced Co-SODA (Figure 3). Based on comparison to other Co-substituted proteins such as Co-rubredoxin, Co-metallothionein, and Co liver alcohol dehydrogenase, these higher energy bands likely arise from cysteinyl sulfur to Co(II) charge transfer (CT).^{62–64} Comparison of these proteins has demonstrated that each Co-thiolate bond in a metalloprotein or peptide typically contributes $\epsilon = 900\text{--}1200 \text{ M}^{-1}\text{cm}^{-1}$ in such a ligand to metal charge transfer (LMCT) transition. Since $\epsilon_{303}(\text{Co-SODA})=2250 \text{ M}^{-1}\text{cm}^{-1}$, it is likely that Co-SODA includes two Co-thiolate bonds.⁶⁵

The CD spectrum of reduced Co-SODA consists of four transitions: a negative band at 280 nm, a positive band at 312 nm, a positive band at 404 nm, and a negative band at 544 nm (Figure 4, Table 1). The two highest energy bands are likely related to the $\text{S} \rightarrow \text{Co}$ LMCT transitions, which are CD active because the first coordination sphere of the cobalt is non-centrosymmetric. The neighboring chiral amino acid based transitions also induce chirality into the metal centered chromophores resulting in the two lower energy features and providing additional evidence that the peptide is involved in coordination of the metal.^{54, 66}

To evaluate the possibility that the other two ligands to the tetrahedral Co(II) in Co-SODA were nitrogenous, FTIR spectra in the region $400\text{--}1000 \text{ cm}^{-1}$ were collected. Three characteristic $\tilde{\nu}(\text{Co-N})$ stretching frequencies were observed at 452, 475 and 602 cm^{-1} for Co-SODA which are absent in the spectrum of the SODA apo-peptide (Figure 5). The two lower energy bands are consistent with symmetric and asymmetric Co-N stretching vibrations for two *cis*-positioned M-N bonds, for M=metal. The highest energy band likely

arises from out of plane bending of C=O for a metal-bound amide, which is affected by the M-N bond strength since the two bonds are shared in a ring structure (Scheme 1).^{67, 68} Thus, it is very likely that the remaining two ligands of the primary coordination sphere are the amino terminus and a backbone amide, as seen already in the Ni-SODA peptide.

Aerobic oxidation of Co-SODA

The effects of molecular oxygen on the reduced Co-SODA sample were investigated by exposing it to air. The green, reduced Co-SODA was found to be oxygen sensitive. This was indicated by a color change to reddish-brown within a few minutes when exposed to air. The UV-vis spectrum recorded after 30 minutes of exposure to air consists of a nearly featureless, increase in absorption in the visible region with broad shoulders at 600 and 460 nm (Figure 3, Table 1). In the far UV region, a sharp band is seen at 355 nm. The prominent LF bands observed for the reduced Co-SODA bleached immediately upon air oxidation, suggesting changes in the tetrahedral coordination geometry around the cobalt center. Additional subtle changes in the optical spectra were observed on longer time scales (24–72 hours). The bands of the initially formed oxidized species were subtly shifted to 570 nm, 450 nm, and 340 nm, respectively (Figure 3, Table 1). After this second oxidized species was formed, no additional spectral changes were observed. These modifications of the optical spectra upon exposure to air suggest a stepwise oxidation of the reduced Co-SODA by oxygen; a partially oxidized intermediate forms first followed later by the fully oxidized product.

CD spectra were also recorded after exposure of the reduced Co-SODA to oxygen. After exposure of the sample to air for 30 minutes, a new spectrum with bands at 267 nm (negative), 306 nm (positive), 420 nm (positive) and 536 nm (negative) arose, suggesting changes in the interactions between the peptide and the cobalt ion. After 24 hours of exposure to air, the CD spectrum was further changed. In particular, both band positions and elliptical polarization (positive or negative) were changed, and a new positive feature appeared at 470 nm (Figure 4). This suggests further rearrangements in the metal-ligand electronic environment after extended interactions with oxygen. Thus CD spectroscopy provides evidence complementary to UV-vis absorption spectroscopy that two distinct species are sequentially formed by air oxidation of Co-SODA.

Magnetic susceptibility changes upon exposure of Co-SODA to oxygen

There are two distinct sites in Co-SODA at which oxidation could easily occur: the cysteinyl thiolates and the metal. Oxidation of the metal from Co(II) to Co(III) would result in a change from a paramagnetic (d^7 , $S=3/2$) starting material to a diamagnetic (d^6 , $S=0$) product. Thus we used the Evans NMR method to determine the effect of air oxidation on the magnetic susceptibility of the sample. In this experiment, the shift of a solvent NMR resonance in a sample containing the complex of interest is monitored relative to a pure solvent sample. The frequency difference between the solvent signal in the presence and the absence of the metal complex can be directly correlated to the magnetic susceptibility of the sample. As shown in Figure 6, the reduced sample is paramagnetic (0 minute data), resulting in two distinct ^1H NMR resonances for the pure H_2O and the H_2O in the presence of Co-SODA (For a solution of ~ 1.9 mM metalloprotein in the 500 MHz NMR instrument, a shift

of 30 Hz was observed.). Figure 6 shows analogous H₂O ¹H NMR spectra collected after the Co-SODA sample was exposed to air for varying lengths of time. The paramagnetism of the Co-SODA sample continued to be observed for several hours after exposure to air. The molecular magnetic susceptibility (χ_M) of the partially oxidized sample was evaluated using the Evans NMR method.^{47, 48} The calculated effective magnetic moment (μ_{eff}) was 4.38 Bohr magnetons. Based on the relation: $\mu_{\text{eff}} = [n(n+2)]$, in which n= number of unpaired electrons, it is likely that the partially oxidized sample contains at least three unpaired electrons (*i.e.*, an S=3/2 system), corresponding to Co(II). Finally, after 24 hours, the solvent signal collapsed into a single peak indicating that the system became diamagnetic. The persistence of paramagnetism for several hours suggests that the initial oxidation does not change the oxidation state of the cobalt. Instead, only on longer time scales is a diamagnetic, Co(III) species formed.

To explore the hypothesis that the initial site of oxidation of the metalloprotein is the cysteine ligands, attempts were made to detect sulfinate or sulfenate groups via FTIR (see Figure S5). Although upon oxidation of the peptide extensive changes in the 1200–900 cm⁻¹ region of the spectrum were observed, the bands could not be unambiguously assigned, leaving interpretation relatively meaningless.

EPR spectroscopy

NMR experiments demonstrated paramagnetism in the reduced and partially oxidized Co-SODA samples. EPR spectra were recorded for these paramagnetic samples to gain insight into their electron spin distributions and corresponding changes upon oxidation. Figure 7A shows the EPR spectrum of the reduced Co-SODA sample. The spectrum was simulated using a spin Hamiltonian (see materials and methods) for a high spin Co(II) (d⁷ system, S=3/2) system.^{69, 70} The good agreement between the experimental and simulated data ($\sigma = 0.94\%$) suggests, as expected, that the spectrum arises from a high spin Co(II) center. The parameters obtained from simulation, g values and zero field splitting (ZFS) constraints for the reduced Co-SODA species, are shown in Table 2. The small E/D value indicates the presence of an effectively axial field around the paramagnetic Co(II) center. The EPR spectrum of the partially oxidized sample (Figure 7B, Table 2) was also simulated with an S=3/2 spin Hamiltonian ($\sigma = 1.20\%$). Although the spectra are quite similar, the g-values for the two species were not identical. This is most clearly seen from their rhombicity parameters [$(g_x - g_y/g_z) \times 100\%$], which have values of 18.102 and 12.170 for the reduced and partially oxidized samples, respectively. The decrease in rhombicity after the partial oxidation of the complex is consistent with a change in the coordination geometry of the cobalt with no change in the oxidation state of the metal.

X-ray absorbance spectroscopy

Cobalt X-ray absorbance spectra (XAS) were obtained to confirm the oxidation state and characterize the first coordination sphere of the cobalt in the partially oxidized and completely oxidized samples. In the XAS spectrum of the fully oxidized sample (Figure 8A), a weak (0.06(1) eV refined area relative to edge height) pre-edge feature at ~7719 eV was observed that arises from the dipole forbidden Co(1s→3d) transition. This transition is ordinarily observed only for cobalt centers in a non-centrosymmetric environment. The edge

energy of ~ 7723 eV corresponds to the presence of a Co(III) center in the fully oxidized metalloprotein sample, consistent with the diamagnetism observed via NMR. The extended X-ray absorbance fine structure (EXAFS) region of the XAS data is best modeled as a pseudo-octahedrally coordinated cobalt center (Figure 8D) in an $S_2(N/O)_4$ coordination environment (Figure 8B,C). The two sulfur scatterers are 2.24 \AA from the Co(III) ion. The N/O ligands could be resolved into three shells: two N/O ligands at 2.13 \AA , one N/O ligand at 1.97 \AA , and one nitrogen ligand at 1.84 \AA from the cobalt center. The last one is likely to be a deprotonated amide nitrogen ligand from the peptide backbone. The scatterer at 1.97 \AA could be the amino terminus of the peptide, and the ligands at longer distances are likely solvent molecules or other loosely bound amides.

As shown in Figure 9 changes in the oxidation state of the Co and its first coordination sphere upon oxidation of Co-SODA were investigated by comparison of the XANES regions of the spectra for the fully oxidized, partially oxidized, and reduced (literally unoxidized) metalloproteins. The edge energy of the partially oxidized sample is $2.9(1)$ eV lower in energy than that of the fully oxidized sample. These edge energies are consistent with a change from Co(II) in the partially oxidized sample to Co(III) in the fully oxidized sample. The area under the pre-edge feature for the partially oxidized sample is $0.04(1)$ eV refined area/edge height, which is also consistent with a six coordinate geometry in the partially oxidized sample as observed in the fully oxidized sample. An EXAFS spectrum from a partially oxidized sample obtained via an experiment analogous to that shown in Figure 8 is shown in Figure S4. However, since the EXAFS data for the partially oxidized sample were of relatively poor quality at high k , the spectrum was only refined over the range $2\text{--}11 \text{ \AA}^{-1}$ and BT. Like the completely oxidized sample, the partially oxidized sample is best fit to a model including two sulfur ligands and four N/O ligands in the first coordination sphere (Figure S4). The Co K-edge energy for the reduced Co-SODA (7720.1 eV) is very similar to that of the partially oxidized sample, suggesting that both are in the Co(II) oxidation state (Figure 9). However, the pre-edge feature is most prominent in the spectrum from the reduced species ($0.20(1)$ eV refined area/edge height (Figure 9) suggesting that the reduced sample features cobalt in the least centrosymmetric environment. Similarly, analysis of the EXAFS region of the spectrum from the reduced sample indicated that it consisted of Co in a four coordinate N_2S_2 site (Figures S6 and S7). Although the EXAFS data is relatively noisy so that it should be interpreted with caution, this observation is consistent with the assignment of the reduced species as a tetrahedral $Co-N_2S_2$ species predicted from the UV-vis and FTIR data above.

Although FTIR could, in principle, be used to detect the SO stretches associated with oxidation of thiolates, as shown in Figure S5 in this case this approach was inconclusive in determining the initial site of oxidation in the peptide because a number of other vibrational modes of similar energies ($1200\text{--}900 \text{ cm}^{-1}$) were also impacted by oxidation of the peptide. Definitely assigning all of the observed changes simply proved untenable. On the other hand, S K-edge X-ray absorbance spectroscopy (XAS) is very sensitive to sulfur oxidation state, and a number of natural proteins and model complexes related to nitrile hydratase have been previously studied using this technique.^{71, 72} Sulfur K-edge X-ray absorbance spectra of the reduced, partially oxidized and fully oxidized metalloproteins described herein were obtained to unambiguously determine the oxidation states of the cysteinate ligands in each

sample. As shown in Figure 10, the reduced sample contained, as expected, only unoxygenated cysteinate sulfurs; a pre-edge peak at 2470.9(1) eV was observed corresponding to the S(1s) \rightarrow Co(3d) transition. At higher energy (2472.6(1) eV), the S(1s) \rightarrow (C-S(σ^*)) transition is observed. In contrast, the S K-edge results suggest the completely oxidized Co(III) complex contains one unmodified cysteinate and one that has been oxidized to a sulfinate (SO_2^-) ligand. The S K-edge spectrum for this sample includes a relatively small pre-edge feature at 2477.8(3) eV together with a strong peak at 2480.8(3) eV. These correspond to $\text{SO}_2(1s) \rightarrow \text{Co}(3d)$ and $\text{SO}_2(1s) \rightarrow \text{C-S}(\sigma^*)$ transitions, respectively. In addition the S(1s) \rightarrow Co(3d) transition is observed at 2470.1(1) eV and the S(1s) \rightarrow (C-S(σ^*)) transition at 2473.3(2) eV indicating the presence of a reduced thiolate sulfur ligand. The spectrum of the partially oxidized sample is remarkably similar to that of the completely oxidized metalloprotein and is also consistent with the sulfur being present as thiolate and sulfinate species. These data provide clear spectroscopic evidence that the first step in the oxidation process is modification of one of the cysteines to form sulfinate.

Concluding Remarks

As depicted in Scheme 1, the data presented herein suggest the tetrahedral high spin d^7 Co(II)-SODA is air oxidized in a two-step process. The immediate changes in the UV-vis and CD spectra upon exposure of Co-SODA to air suggest oxidation occurs rapidly and results in formation of a pseudo-octahedral species. However, perhaps surprisingly, NMR and EPR spectra indicate that the first oxidation product formed is still paramagnetic with three unpaired electrons. This indicates Co cannot be the first species oxidized. Instead, the S K-edge XAS spectrum of this intermediate shows that the first oxidation occurs at a single cysteine ligand to form a sulfinate. Then, it is only on longer time scales that Co oxidizes to the 3+ oxidation state becoming a diamagnetic low spin d^6 system. EXAFS data suggest this fully oxidized metalloprotein contains a pseudo-octahedrally coordinate Co(III) ion ligated by an unmodified cysteinate as well as an oxidized sulfinate ligand. The remainder of the coordination sphere is four N/O donors present in two pairs at different distances. Two of those are likely to be the peptide ligands from the reduced peptide and the additional ones are expected to be solvent molecules. Thus air oxidation of Co-SODA causes two sequential types of changes. First oxidation of one of the cysteines occurs; then oxidation of the metal follows.

Oxidation of cysteinyl thiolates can result in a number of products including disulfide bonds, sulfinate and sulfenate. Thus the question of how each cysteine in the active site of nitrile hydratase achieves a unique oxidation state comes to the fore, and Co-SODA, in which one cysteinate is modified and the other remains reduced, is an interesting model. For these metalloproteins/peptides, formation of disulfide bonds is likely of little concern due to the relative inertness of Co(II) to substitution reactions. On the other hand, in the case of Co-SODA, a peptide originally designed to bind not Co but Ni, oxidation of one of the cysteines likely provides more stable bonding. This lowers the energies of the occupied Co d orbitals and increases the driving force for oxidation of the metal to form a low spin d^6 system. Furthermore, binding of S-O ligands likely strains the tetrahedral geometry of Co(II)-SODA towards a more favorable octahedral environment, again providing ideal conditions for oxidation of the metal. We can speculate that once the metal is oxidized in Co-SODA there

is relatively little energetic gain from oxidation of the second thiolate so that it remains unmodified.

Comparison of Co SODA to other model complexes for nitrile hydratase suggests reasons for the unique reactivity of this metalloprotein. Angelosante and coworkers reported the synthesis of an unsymmetrical, square planar S=1 Co(II) complex in which the metal is coordinated in an N₂S₂ environment consisting of two thiolates and two amidates.⁷³ Interestingly, this complex mimics the asymmetric equatorial coordination environment of nitrile hydratase that consists of two five-membered rings and a single six-membered ring. However, in contrast to Co-SODA, the reported complex does not undergo air oxidation of either the thiolates or the metal.⁷³ In essence, the square planar complex appears to be too stable to promote the oxidation. On the other hand, Shearer and coworkers reported the reactivity of an N₂S₃ coordinated Co-metalloprotein known as Co-NHaseM1.⁷⁴ Like Co-SODA, reduced Co-NHaseM1 binds Co(II) in a tetrahedral environment consisting of two thiolates and two amidates. Upon exposure to air, both the metal and two of the three thiolates are oxidized. As shown in Fig. S3, we undertook NMR experiments with Co-NHaseM1 to compare its oxidation pathway to that of Co-SODA. Unlike Co-SODA, upon oxidation Co-NHaseM1 rapidly forms a diamagnetic Co(III) state. Furthermore, both equatorial thiolates are likely oxidized. The increased rate of metal oxidation in Co-NHaseM1 is likely a result, in large part, of the additional axial thiolate ligand and the replacement of an amine with an amide ligand. As shown in Scheme S2, the strongly σ -donating axial thiolate likely interacts with the metal d_{z²} orbital, changing its character from non-bonding to antibonding – significantly raising its energy. Similarly, the amide of Co-NHaseM1 provides stronger π -bonding than the amine of Co-SODA further increasing the ligand field stabilization energy of the N₂S₃ peptide and stabilizing the low spin d⁶ oxidized state. Thus oxidation of Co-NHaseM1 is considerably more favorable energetically than oxidation of Co-SODA. Essentially, Co-SODA is intermediate between the relatively inert complex of Angelosante and coworkers and the reactive metalloprotein Co-NHaseM1. Furthermore, it is likely the tetrahedral geometry of the reduced Co-SODA species and the lack of an axial thiolate that allow it to occupy this unique niche of reactivity. Nonetheless, more detailed computational investigation will be necessary to confirm this hypothesis.

Finally, we end by speculating as to the applicability of Co-SODA to understanding the details of metal incorporation and protein modification in the natural nitrile hydratase enzyme. Cobalt is generally transported into microbes as Co(II) ion, and it is likely that Co is incorporated into the apo-nitrile hydratase α subunit in this reduced oxidation state.^{75, 76} Biochemical evidence has arisen that, at least *in vitro*, post-translational modification of the nitrile hydratase protein and oxidation of the metal are autocatalytic.⁵ The reactivity of Co-SODA in air lends further credence to such a model. By analogy to Co-SODA, we hypothesize that interaction of the reduced NHase protein with air first results in oxidation of one or both of the equatorial sulfur ligands. Initial binding of the metal in a geometry that is more tetrahedral than square planar geometry would likely promote the specific protein modifications. The modified bonding with the oxidized ligands could then provide the driving force for a subtle change in geometry and oxidation of the metal to the active form of the enzyme. At the moment, these ideas are hypothetical, but further investigation of both

the enzyme and relevant metalloprotein models is expected to clarify the remaining unknowns of this unique biosynthetic process. Moreover, we anticipate that designed metalloproteins will prove to be an exciting tool for exploring metallocenter biosynthesis in a biologically relevant, *in vitro*, chemical context.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported through the Center for Bio-Inspired Solar Fuel Production an Energy Frontier Research Center funded by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences under Award Number DE-SC0001016. J.M.S. acknowledges the NSF for financial support (CHE-0844234). H.E.H. acknowledges NSF for financial support (EAR-0846188). XAS data collection at the National Synchrotron Light Source (NSLS) at Brookhaven National Laboratory was supported by the U.S. Department of Energy, Division of Materials Sciences and Division of Chemical Sciences. Beamline X3B at NSLS is supported by the NIH. This publication was made possible in part by the Center for Synchrotron Biosciences grant, P30-EB-009998, from the National Institute of Biomedical Imaging and Bioengineering (NIBIB). A.D. also thanks the Science Foundation of Arizona for a graduate fellowship. We are grateful to Dr. Ulrich Haussermann for technical assistance preparing anaerobic KBr pellets.

REFERENCES

1. Paulsen CE, Carroll KS. *ACS Chem. Biol.* 2010; 5:47–62. [PubMed: 19957967]
2. Claiborne A, Yeh JI, Mallett TC, Luba J, Crane EJ, Charrier V, Parsonage D. *Biochemistry.* 1999; 38:15407–15416. [PubMed: 10569923]
3. Dickinson BC, Chang CJ. *Nat. Chem. Biol.* 2011; 7:504–511. [PubMed: 21769097]
4. Nagashima S, Nakasako M, Dohmae N, Tsujimura M, Takio K, Odaka M, Yohda M, Kamiya N, Endo I. *Nat. Struct. Mol. Biol.* 1998; 5:347–351.
5. Arakawa T, Kawano Y, Katayama Y, Nakayama H, Dohmae N, Yohda M, Odaka M. *J. Am. Chem. Soc.* 2009; 131:14838–14843. [PubMed: 19785438]
6. Arakawa T, Kawano Y, Kataoka S, Katayama Y, Kamiya N, Yohda M, Odaka M. *J. Mol. Biol.* 2007; 366:1497–1509. [PubMed: 17222425]
7. Katayama Y, Hashimoto K, Nakayama H, Mino H, Nojiri M, Ono T, Nyunoya H, Yohda M, Takio K, Odaka M. *J. Am. Chem. Soc.* 2006; 128:728–729. [PubMed: 16417356]
8. Huang WJ, Jia J, Cummings J, Nelson M, Schneider G, Lindqvist Y. *Structure.* 1997; 5:691–699. [PubMed: 9195885]
9. Miyanaga A, Fushinobu S, Ito K, Wakagi T. *Biochem. Biophys. Res. Commun.* 2001; 288:1169–1174. [PubMed: 11700034]
10. Nakasako M, Odaka M, Yohda M, Dohmae N, Takio K, Kamiya N, Endo I. *Biochemistry.* 1999; 38:9887–9898. [PubMed: 10433695]
11. Song L, Wang M, Shi J, Xue Z, Wang M-X, Qian S. *Biochem. Biophys. Res. Commun.* 2007; 362:319–324. [PubMed: 17716629]
12. Murakami T, Nojiri M, Nakayama H, Dohmae N, Takio K, Odaka M, Endo I, Nagamune T, Yohda M. *Protein Sci.* 2000; 9:1024–1030. [PubMed: 10850812]
13. Endo I, Odaka M, Yohda M. *Trends Biotechnol.* 1999; 17:244–248. [PubMed: 10354562]
14. Zhou Z, Hashimoto Y, Kobayashi M. *Actinomycetologica.* 2005; 19:18–26.
15. Kobayashi M, Nagasawa T, Yamada H. *Trends Biotechnol.* 1992; 10:402–408. [PubMed: 1368882]
16. Noveron JC, Olmstead MM, Mascharak PK. *J. Am. Chem. Soc.* 1999; 121:3553–3554.
17. Kovacs JA. *Chem. Rev.* 2004; 104:825–848. [PubMed: 14871143]

18. Kung I, Schweitzer D, Shearer J, Taylor WD, Jackson HL, Lovell S, Kovacs JA. *J. Am. Chem. Soc.* 2000; 122:8299–8300.
19. Shearer J, Kung IY, Lovell S, Kaminsky W, Kovacs JA. *J. Am. Chem. Soc.* 2001; 123:463–468. [PubMed: 11456548]
20. Galvez C, Ho DG, Azod A, Selke M. *J. Am. Chem. Soc.* 2001; 123:3381–3382. [PubMed: 11457080]
21. Rat M, Alves de Sousa R, Vaissermann J, Leduc P, Mansuy D, Artaud I. *J. Inorg. Biochem.* 2001; 84:207–213. [PubMed: 11374583]
22. Shearer J, Callan PE, Amie J. *Inorg. Chem.* 2010; 49:9064–9077. [PubMed: 20831172]
23. Masitas CA, Kumar M, Mashuta MS, Kozlowski PM, Grapperhaus CA. *Inorg. Chem.* 2010; 49:10875–10881. [PubMed: 20973591]
24. Farmer PJ, Solouki T, Mills DK, Soma T, Russell DH, Reibenspies JH, Darensbourg MY. *J. Am. Chem. Soc.* 1992; 114:4601–4605.
25. Mirza SA, Pressler MA, Kumar M, Day RO, Maroney MJ. *Inorg. Chem.* 1993; 32:977–987.
26. Farmer PJ, Verpeaux J-N, Amatore C, Darensbourg MY, Musie G. *J. Am. Chem. Soc.* 1994; 116:9355–9356.
27. Noveron JC, Olmstead MM, Mascharak PK. *J. Am. Chem. Soc.* 2001; 123:3247–3259. [PubMed: 11457060]
28. Kaasjager VE, Bouwman E, Gorter S, Reedijk J, Grapperhaus CA, Reibenspies JH, Smeets JJ, Darensbourg MY, Derecskei-Kovacs A, Thomson LM. *Inorg. Chem.* 2002; 41:1837–1844. [PubMed: 11925177]
29. Lee C-M, Hsieh C-H, Dutta A, Lee G-H, Liaw W-F. *J. Am. Chem. Soc.* 2003; 125:11492–11493. [PubMed: 13129340]
30. Lee C-M, Chen C-H, Chen H-W, Hsu J-L, Lee G-H, Liaw W-F. *Inorg. Chem.* 2005; 44:6670–6679. [PubMed: 16156625]
31. Chohan BS, Maroney MJ. *Inorg. Chem.* 2006; 45:1906–1908. [PubMed: 16499349]
32. Grapperhaus CA, Darensbourg MY, Sumner LW, Russell DH. *J. Am. Chem. Soc.* 1996; 118:1791–1792.
33. Galardon E, Giorgi M, Artaud I. *Chem. Commun.* 2004:286–287.
34. Bourles E, de Sousa RA, Galardon E, Giorgi M, Artaud I. *Angew. Chem. Int. Edit.* 2005; 44:6162–6165.
35. Zhou Z, Hashimoto Y, Shiraki K, Kobayashi M. *Proc. Natl. Acad. Sci. USA.* 2008; 105:14849–14854. [PubMed: 18809911]
36. Zhou ZM, Hashimoto Y, Kobayashi M. *J. Biol. Chem.* 2009; 284:14930–14938. [PubMed: 19346246]
37. Zhou Z, Hashimoto Y, Cui T, Washizawa Y, Mino H, Kobayashi M. *Biochemistry.* 2010; 49:9638–9648. [PubMed: 20886813]
38. Sloan CP, Krueger JH. *Inorg. Chem.* 1975; 14:1481–1485.
39. Lange BA, Libson K, Deutsch E, Elder RC. *Inorg. Chem.* 1976; 15:2985–2989.
40. Higashi LS, Lundeen M, Hilti E, Seff K. *Inorg. Chem.* 1977; 16:310–313.
41. Lundeen M, Firor RL, Seff K. *Inorg. Chem.* 1978; 17:701–706.
42. Elder RC, Heeg MJ, Payne MD, Trkula M, Deutsch E. *Inorg. Chem.* 1978; 17:431–440.
43. Kubota M, Rothrock RK, Kernan MR, Haven RB. *Inorg. Chem.* 1982; 21:2491–2493.
44. Bertz SH, Dabagh G. *Inorg. Chem.* 1985; 24:3488–3490.
45. Stoll S, Schweiger A. *J. Magn. Reson.* 2006; 178:42–55. [PubMed: 16188474]
46. Flores M, Isaacson R, Abresch E, Calvo R, Lubitz W, Feher G. *Biophys. J.* 2007; 92:671–682. [PubMed: 17071655]
47. Evans DF. *J. Chem. Soc.* 1959:2003–2005.
48. Grant DH. *J. Chem. Educ.* 1995; 72:39.
49. Shearer J, Dehestani A, Abanda F. *Inorg. Chem.* 2008; 47:2649–2660. [PubMed: 18330983]
50. Shearer, J.; Scarrow, RS. EXAFS123. University of Nevada; Reno: 2011.

51. IgorPro. Wavemetrics, Inc.; Lake Oswego, OR: 2011.
52. Ankudinov AL, Ravel B, Rehr JJ, Conradson SD. *Phys. Rev. B.* 1998; 58:7565–7576.
53. Dutta A, Hamilton GA, Hartnett HE, Jones AK. *Inorg. Chem.* 2012; 51:9580–9588. [PubMed: 22924594]
54. Neupane KP, Gearty K, Francis A, Shearer J. *J. Am. Chem. Soc.* 2007; 129:14605–14618. [PubMed: 17985883]
55. Ellman GL. *Arch. Biochem. Biophys.* 1958; 74:443–450. [PubMed: 13534673]
56. Choudhury SB, Lee J-W, Davidson G, Yim Y-I, Bose K, Sharma ML, Kang S-O, Cabelli DE, Maroney MJ. *Biochemistry.* 1999; 38:3744–3752. [PubMed: 10090763]
57. Wuerges J, Lee J-W, Yim Y-I, Yim H-S, Kang S-O, Carugo KD. *Proc. Natl. Acad. Sci. USA.* 2004; 101:8569–8574. [PubMed: 15173586]
58. Barondeau DP, Kassmann CJ, Bruns CK, Tainer JA, Getzoff ED. *Biochemistry.* 2004; 43:8038–8047. [PubMed: 15209499]
59. Lever, ABP. *Inorganic Electronic Spectroscopy.* 3rd ed.. Elsevier; 1986.
60. Petros AK, Shaner SE, Costello AL, Tierney DL, Gibney BR. *Inorg. Chem.* 2004; 43:4793–4795. [PubMed: 15285646]
61. Petros AK, Reddi AR, Kennedy ML, Hyslop AG, Gibney BR. *Inorg. Chem.* 2006; 45:9941–9958. [PubMed: 17140191]
62. Drum DE, Vallee BL. *Biochem. Biophys. Res. Commun.* 1970; 41:33–39. [PubMed: 4318855]
63. May SW, Kuo J-Y. *Biochemistry.* 1978; 17:3333–3338. [PubMed: 687587]
64. Vasak M, Kaegi JHR, Holmquist B, Vallee BL. *Biochemistry.* 1981; 20:6659–6664. [PubMed: 6272845]
65. VanZile ML, Cospier NJ, Scott RA, Giedroc DP. *Biochemistry.* 2000; 39:11818–11829. [PubMed: 10995250]
66. Nakashini, K.; Berova, N.; Woody, RW. *Circular Dichroism: Principles and Applications.* 1st ed.. Wiley-VCH publishers; Darmstadt: 1994.
67. Condrate RA, Nakamoto K. *J. Chem. Phys.* 1965; 42:2590–2598. [PubMed: 14294946]
68. Kincaid JR, Nakamoto K. *Spectrochim. Acta, Part A.* 1976; 32:277–283.
69. Rizzi AC, Brondino CD, Calvo R, Baggio R, Garland MT, Rapp RE. *Inorg. Chem.* 2003; 42:4409–4416. [PubMed: 12844314]
70. Pietrzyk P, Srebro M, Rado M, Sojka Z, Michalak A. *J. Phys. Chem. A.* 2011; 115:2316–2324. [PubMed: 21351791]
71. Dey A, Chow M, Taniguchi K, Lugo Mas P, Davin S, Maeda M, Kovacs JA, Odaka M, Hodgson KO, Hedman B, Solomon EI. *J. Am. Chem. Soc.* 2006; 128:533–541. [PubMed: 16402841]
72. Rompel A, Cinco RM, Latimer MJ, McDermott AE, Guiles RD, Quintanilha A, Krauss RM, Sauer K, Yachandra VK, Klein MP. *Proc. Natl. Acad. Sci. USA.* 1998; 95:6122–6127. [PubMed: 9600928]
73. Angelosante J, Schopp L, Lewis B, Vitalo A, Titus D, Swanson R, Stanley A, Abolins B, Frome M, Cooper L, Tierney D, Moore C, Rheingold A, Daley CA. *J. Biol. Inorg. Chem.* 2011; 16:937–947. [PubMed: 21638158]
74. Shigehiro S, Nakasako M, Dohmae N, Tsujimura M, Tokoi K, Odaka M, Yohda M, Kamiya N, Endo I. *Nat. Struct. Biol.* 1998; 5:347–351. [PubMed: 9586994]
75. Komeda H, Kobayashi M, Shimizu S. *Proc. Natl. Acad. Sci. USA.* 1997; 94:36–41. [PubMed: 8990157]
76. Okamoto S, Eltis LD. *Metallomics.* 2011; 3:963–970. [PubMed: 21804980]

Synopsis

Air oxidation of the cobalt-bound seven amino acid peptide ACDLPCG (Co-SODA) is characterized as a model for maturation of nitrile hydratase. Co-SODA is oxidized in two steps. In the first step, the metallocenter remains unchanged while one of the thiolates is oxidized to sulfinate. In the second step, the cobalt center is oxidized to Co^{3+} .




Figure 1. Heterodimeric cobalt nitrile hydratase and its metal active site (PDB code: 1UGP, figure prepared using PyMOL software [The PyMOL Molecular Graphics System, Version 0.99rc6, Schrödinger, LLC]). The green color corresponds to the α chain and the blue color to the β chain.




Figure 2.

UV-vis spectrum of Co-SODA (solid line), partially oxidized Co-SODA (dotted line, 30 minutes of air oxidation), completely oxidized Co-SODA (dashed line, 24 hours of air exposure) All solutions are in aqueous 50 mM NEM buffer at pH 7.5. The inset shows an enlargement of the region between 500 and 1000 nm featuring bands associated with ligand field (LF) transitions for reduced Co-SODA.




Figure 3.

Spectral changes in the UV-vis spectrum of SODA upon stepwise addition under anaerobic conditions of an aqueous solution (50 mM NEM buffer, pH 7.7) of Co(II). The inset shows the binding curves extracted at three different wavelengths from the spectra (squares 681 nm, circles 709 nm, and triangles 725 nm). The ratios of Co(II):SODA for which spectra are shown are 0:1, 0.13:1, 0.26:1, 0.39:1, 0.52:1, 0.65:1, 0.78:1, 0.91:1, 1.04:1, 1.17:1, and 1.3:1.




Figure 4. CD spectra for reduced Co-SODA (solid black line), partially oxidized Co-SODA (dotted line), completely oxidized Co-SODA (dashed line) and the SODA apo-peptide (solid grey line). All metalloprotein samples were in aqueous 50 mM NEM buffer at pH 7.5.




Figure 5. FTIR absorption spectra from the SODA apo-peptide (solid line at the bottom), reduced Co-SODA (solid line in the middle), and the partially oxidized Co-SODA (dash dotted line) in the 400–700 cm⁻¹ region.




Figure 6.

^1H NMR resonances as a function of time exposed to air for H_2O in an aqueous sample of reduced Co-SODA [in an NMR tube (O.D. \sim 3mm)]. The sample was placed within another NMR tube (O.D. \sim 5 mm) containing pure H_2O , and the resonances are referenced such that the frequency of the signal of the blank H_2O (measured independently) is zero.




Figure 7. Continuous wave (CW) EPR spectra of (A) reduced Co-SODA (in aqueous 50 mM NEM buffer at pH 7.5) at 5 K and (B) partially oxidized Co-SODA sample (in aqueous 50 mM NEM buffer at pH 7.5) at 7 K. The experimental data and the simulated data are shown as solid black line and dotted lines, respectively.




Figure 8.

(A) XANES region of the Co K-edge X-ray absorption spectrum of fully oxidized Co-SODA. (B) FF k^3 data and (C) FT k^3 data (experimental data, simulated data, and the difference spectrum are represented as the red solid line, blue dashed line, and green dotted line, respectively). (D) Primary coordination geometry around cobalt in oxidized Co-SODA and metal-ligand bond distances from EXAFS. Refinement parameters: Co-S, $n = 2$ (restrained), $r = 2.241(13) \text{ \AA}$, $\sigma^2 = 0.0009(2) \text{ \AA}^2$; Co-N, $n = 2$ (restrained), $r = 2.130(9) \text{ \AA}$, $\sigma^2 = 0.0012(4) \text{ \AA}^2$; Co-N, $n = 1$ (restrained), $r = 1.968(7) \text{ \AA}$, $\sigma^2 = 0.0011(8) \text{ \AA}^2$; Co-N, $n = 1$ (restrained), $r = 1.844(6) \text{ \AA}$, $\sigma^2 = 0.0011(4) \text{ \AA}^2$; Co-C, $n = 5$ (restrained), $r = 2.50(2) \text{ \AA}$, $\sigma^2 = 0.0016(2) \text{ \AA}^2$; $E_0 = 7725.1 \text{ eV}$.




Figure 9. Comparison of the XANES region of the X-ray absorption spectrum of reduced (green), partially oxidized (red), and fully oxidized Co-SODA (blue).





Figure 10. S K-edge X-ray absorption spectra of fully reduced (green), partially oxidized (red) and fully oxidized (blue) CoSODA.

**Scheme 1.**

Hypothesized sequence for oxidation of Co(II)-SODA in air.

Table 1

Optical and CD data for various Co-SODA species

Sample	Electronic absorption	CD absorption
	λ (nm) (ϵ ($M^{-1} cm^{-1}$))	λ (nm) (ϵ ($M^{-1} cm^{-1}$))
Reduced Co-SODA	725 (210)	404 (0.20)
	681 (230)	544 (-0.15)
	609 (225)	312 (0.54)
	350 (shoulder, 1700)	280 (-2.58)
	303 (2250)	
Partially oxidized Co-SODA	600 (shoulder, 140)	536 (-0.8)
	460 (shoulder, 430)	420 (0.37)
	355 (1100)	306 (2.18)
		267 (-3.92)
Fully oxidized Co-SODA	570 (shoulder, 195)	536 (0.7)
	450 (shoulder, 480)	470 (0.96)
	340 (1500)	414 (shoulder, 0.36)
		328 (-5.88)
		286 (-3.87)

Table 2

Simulation parameters for the EPR spectra of reduced and partially oxidized Co-SODA

Parameter ^a	Reduced Co-SODA	Partially oxidized Co-SODA
$g_x (\pm 0.001)$	2.527	2.476
g_y	2.111	2.199
g_z	2.298	2.276
B_x (MHz)	3480	3695
B_y	5544	6096
B_z	2824	2491
D (MHz)	5×10^8	5×10^8
E/D	0.02	0.02
$[(g_x - g_y/g_z) \times 100\%]$	18.102	12.170

^aThe simulation parameters were the following: the principal g values (g_x , g_y and g_z), the line widths (B_x , B_y and B_z) and the zero-field splitting parameters, D and E .

Table 3

Summary of K-edge energies from Co and S XAS.

Sample	Co K-edge energy (eV)		S K-edge energy (eV)			
	Co (1s→3d)	Co edge	S(1s)→Co(3d)	S(1s)→C-S (σ^*)	SO ₂ (1s)→Co(3d)	SO ₂ (1s)→C-S(σ^*)
Reduced Co-SODA	7719.3(2)	7720.1(3)	2470.9(2)	2472.6(2)	N/A	N/A
Partially oxidized Co-SODA	7719.2(1)	7720.1(4)	2470.4(2)	2473.2(2)	2477.6(7)	2480(1)
Fully oxidized Co-SODA	7719.4(1)	7723.2(3)	2470.1(2)	2473.2(2)	2477.8(6)	2480(1)