

A Molecular Survey of *Campylobacter jejuni* and *Campylobacter Coli* Virulence and Diversity

Mahdi Ghorbanalizadgan¹, Bita Bakhshi^{*1}, Anoshirvan Kazemnejad Lili²,
Shahin Najar-Peerayeh¹ and Bahram Nikmanesh³

¹Dept. of Bacteriology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran; ²Dept. of Biostatistics, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran; ³Children's Hospital Medical Center, Tehran university of Medical Sciences, Tehran, Iran

Received 21 December 2013; revised 16 February 2014; accepted 17 February 2014

ABSTRACT

Background: The aim of this study was to determine the prevalence of virulence-associated genes and enterobacterial repetitive intergenic consensus PCR (ERIC-PCR) analysis of *Campylobacter* spp. isolated from children with diarrhea in Iran. **Methods:** A total of 200 stool specimens were obtained from children under 5 years during July 2012 to July 2013. Detection of *C. jejuni* and *C. coli* was performed by standard biochemical and molecular methods. The presence of virulence-associated genes and genetic diversity of isolates was examined using PCR and ERIC-PCR analyses. **Results:** A total of 12 (6%) *Campylobacter* spp. were isolated from patients including 10 (4.5%) *C. jejuni* and 2 (1.5%) *C. coli*. The *flaA*, *cadF* and *ciaB* genes were present in 100% of isolates, while no plasmid of *virB11* gene was present in their genome. The prevalence of invasion-associated marker was 100% among *C. coli* and was not detected in *C. jejuni* isolates. The distribution of both *pldA* and the genes associated with cytolethal distending toxin (CDT) was 58.3% in *C. jejuni* isolates. Seven distinct ERIC-PCR profiles were distinguished in three clusters using ERIC-PCR analysis. Genotyping analysis showed a relative correlation with geographic location of patients and virulence gene content of isolates. **Conclusion:** To our knowledge, this is the first molecular survey of *Campylobacter* spp. in Iran concerning genotyping and virulence gene content of both *C. jejuni* and *C. coli*. ERIC-PCR revealed appropriate discriminatory power for clustering *C. jejuni* isolates with identical virulence gene content. However, more studies are needed to clearly understand the pathogenesis properties of specific genotypes. *Iran. Biomed. J.* 18 (3): 158-164, 2014

Keywords: *Campylobacter jejuni*, *Campylobacter coli*, Ddiarrhea, Virulence factors

INTRODUCTION

Campylobacter spp. especially *C. jejuni* and *C. coli* are considered as potential etiological agents that caused many undiagnosed cases of acute diarrhea in children in developing countries including Iran [1-3]. The purpose of our experiments was to determine the rate and molecular survey of *C. jejuni* and *C. coli* virulence and also their diversity that leads to the practical applications to elucidate *Campylobacter* colonization and the control of this organism. Recurrent exposure to these organisms might raise level of specific immunity correlated with age in developing countries. Therefore, children younger than 5 years of age are mainly affected by these organisms [4].

Above organisms are fastidious and need nutrient-

rich-based medium and microaerobic atmosphere [5]. This reason may be the main cause that *Campylobacter* spp. are not applied in routine diagnostic programs of clinical laboratories in most developing countries. The pathogenicity of *Campylobacter* species is dependent on their ability to bind to the human intestinal cells and *CadF* protein. This protein, which is encoded by *cadF* gene, is responsible for *Campylobacter* binding to extracellular matrix of human intestinal cells [6]. Another gene, *flaA*, encodes a flagella protein which mediates motility, colonization, and invasion of gastrointestinal tract and it is essential for establishing human infection.

The *ciaB* (an invasion protein), *virB11* (the IV secretory system), and *pldA* (an outer membrane phospholipase A) genes encode proteins associated with increased bacterial invasion on cultured epithelial

cells; however, their exact roles in invasion have remained to be elucidated [7]. Cytolethal distending toxin (CDT) is encoded by three linked genes, including *cdtA*, *cdtB*, and *cdtC*. In epidemiology of infectious diseases, bacterial typing is of great value in source tracking studies. To analyze the genetic relatedness of *C. jejuni*, several molecular typing methods based on PCR have been developed. enterobacterial repetitive intergenic consensus PCR (ERIC-PCR) has been shown high discriminatory power, good legibility, and ease of use in most epidemiological investigations.

In this study, we aimed to determine the prevalence of virulence-associated genes and ERIC-PCR analysis of *C. jejuni* and *C. coli* isolated from children with diarrhea. CDT is one of the most characterized virulence factors in *Campylobacter* spp. Pathogenesis induces cell cycle arrest in G2 phase and promotes DNA damage together with apoptotic death in human monocytic cells; therefore, its presence is supposed to be associated with the severity of the disease.

MATERIALS AND METHODS

Study design and data collection procedure. The study was started after obtaining ethical approval from Research and Publication ethics office of Tarbiat Modares University (Tehran, Iran). A total of 200 stool specimens were collected from children with acute diarrhea attended to two major Children's Hospital Medical Center in Tehran from July 2012 to July 2013. All children under five years of age were included in this study. Children with persistent diarrhea or previous treatment with antibiotics in the last 5 days were excluded from the study. Acute diarrhea was defined as diarrhea which takes about 14 days or less. Demographic data were collected by a co-worker resident in hospital. All samples were transported to the laboratory in a modified Cary-Blair transport medium (5.00 g/L sodium chloride, 1.50 g/L sodium thioglycollate, 1.10 g/L disodium phosphate, and 0.09 g/L calcium chloride, pH 8.4 ± 0.2 at 25°C) with reduced agar content (1.6 g/L). Identification of *C. jejuni* and *C. coli* was performed by the standard culture, Gram staining, and conventional biochemical tests and confirmed by molecular methods [8, 9].

Confirmation of presumptive *Campylobacter* species by duplex-PCR. Samples were incubated at 42°C for 48-72 h in microaerophilic conditions onto modified charcoal cefoperazone deoxycholate agar medium (10 g/L meat extract, 10 g/L peptone, 5 g/L sodium chloride, 4 g/L bacteriological charcoal, 3 g/L casein hydrolysate, 1 g/L sodium deoxycholate, 0.25

g/L iron (II) sulfate, 0.25 g/L sodium pyruvate, and 15 g/L agar, pH 7.4 ± 0.2 at 25°C) plus campylobacter CCDA selective supplement (cefoperazone 3,200 mg/L). DNA templates were extracted by boiling method [10]. Confirmation of *Campylobacter* spp. was performed by PCR amplification of *cadF* gene. A duplex-PCR was applied for simultaneous detection of *hipO* and *asp* genes, specific to *C. jejuni* and *C. coli*, respectively [11]. PCR was performed in a 25-μl reaction mixture, containing 10 ng DNA template, 2.5 μl PCR buffer 10×, 200 μM dNTP, 5 mM MgCl₂, 0.1 μM each primer, 1 unit of Taq DNA polymerase, and deionized water. The primer sequences and their designation are shown in Table 1. The *C. jejuni* ATCC 29428 and *C. coli* ATCC 43478 were used as reference strains.

Detection of virulence/invasion-associated genes.

The presence of virulence/invasion-associated genes, including invasion-associated marker (*iam*), *pldA*, and *ciaB*, (responsible for *Campylobacter* invasion and attachment), *virB11* (involved in *Campylobacter* virulence), and CDT were investigated using specific primers which specifically amplify within the coding region of each gene. The distribution of *flaA* gene (responsible for *Campylobacter* attachment) was examined by primers specifically designed according to the *flaA* locus sequence of *C. jejuni* strain (GenBank accession no. AF050186.1). Due to the species-allele-specification of CDT sequence, two separate primer pairs were used which were selected according to *cdt* locus sequence of *C. jejuni* and *C. coli* standard strains.

Enterobacterial repetitive intergenic consensus PCR. ERIC-PCR assay was performed according to the method introduced by Versalovic and colleagues [12]. The primer sequences, their designation, and amplification conditions are depicted in Table 1. ERIC-PCR amplification reactions were performed in a 25-μl reaction mixture, containing 10 ng genomic DNA, 2.5 μl reaction buffer 10×, 200 μM dNTP, 5 mM MgCl₂, 0.2 μM each primer, and 1 unit Taq DNA polymerase. The reaction was placed in a DNA thermal cycler (Mini Bio-Rad, USA). ERIC-PCR patterns were analyzed based on the Dice similarity coefficient using GelClust software [13].

RESULTS

The study population was made up of 200 children with acute diarrhea, including 110 (55%) male and 90 (45%) female with the mean age of 27.4 months (2.3 years). Among 200 stool samples, *Campylobacter* spp., including 10 (4.5%) *C. jejuni* and 2 (1.5%) *C. coli* were isolated from 12 (6%) samples.

Table 1. Primers, PCR conditions, and respective references

Primers	Sequence (5'→3')	Target	PCR condition			Amplicon (bp)	References
			Denaturin	Annealin	Extension		
cadFU cadFR	TTGAAGGTAATTTAGATATG CTAATACCTAAAGTTGAAAC	<i>cadF</i>	94°C, 30 s	43°C, 30 s	72°C, 30 s	400	[11]
hipOU hipOR	GAAGAGGTTTGGGTGGTG AGCTAGCTTCGCATAATAACTTG	<i>hipO</i>	94°C, 30 s	53°C, 30 s	72°C, 30 s	735	[11]
aspU aspR	GGTATGATTTCTACAAAGCGAG ATAAAAAGACTATCGTCGCGTG	<i>asp</i>	94°C, 30 s	53°C, 30 s	72°C, 30 s	500	[11]
flaAU flaAR	TTTCGTATTAACACAAATGGTGC CTGTAGTAATCTTAAAACATTTTG	<i>flaA</i>	94°C, 45 s	46°C, 45 s	72°C, 60 s	1743	this study
cdtjAU cdtjAR	AGGACTTGAACCTACTTTTC AGGTGGAGTAGTAAAAACC	<i>Cj-cdtA</i>	94°C, 30 s	55°C, 30 s	72°C, 30 s	631	[31]
cdtj BU cdtjBR	ATCTTTTAAACCTTGCTTTTGC GCAAGCATTAAAATCGCAGC	<i>Cj-cdtB</i>	94°C, 30 s	56°C, 30 s	72°C, 30 s	714	[31]
cdtjCU cdtjCR	TTTAGCCTTTGCAACTCCTA AAGGGGTAGCAGCTGTAA	<i>Cj-cdtC</i>	94°C, 30 s	55°C, 30 s	72°C, 30 s	524	[31]
cdtCAU cdtCAR	ATTGCCAAGGCTAAAATCTC GATAAAGTCTCCAAAACCTGC	<i>Cc-cdtA</i>	94°C, 30 s	55°C, 30 s	72°C, 30 s	329	[31]
cdtCBU cdtCBR	TTTAATGTATTATTTGCCGC TCATTGCCTATGCGTATG	<i>Cc-cdtB</i>	94°C, 30 s	56°C, 30 s	72°C, 30 s	413	[31]
cdtCCU cdtCCR	TAGGGATATGCACGCAAAAG GCTTAATACAGTTACGATAG	<i>Cc-cdtC</i>	94°C, 30 s	55°C, 30 s	72°C, 30 s	313	[31]
ciaBU ciaBR	TGCTAGCCATACTTAGGCGTTTT TTGATAATAGCGGACAATTTGAAA	<i>ciaB</i>	94°C, 30 s	54°C, 30 s	72°C, 30 s	610	this study
pldAU pldAR	AAGCTTATGCGTTTTT TATAAAGCTTTCTCC	<i>PldA</i>	94°C, 30 s	46°C, 30 s	72°C, 30 s	913	[32]
iamAU iamAR	GCGCAAAATATTATCACCC TTCACGACTACTATGCGG	<i>iam</i>	94°C, 30 s	47°C, 30 s	72°C, 30 s	518	[33]
virB11U virB11R	GAACAGGAAGTGAAAAACTAGC TCCCGCATTGGGCTATATG	<i>virB11</i>	94°C, 30 s	52°C, 30 s	72°C, 120 s	708	[33]
ERICF ERICR	ATGTAAGCTCCTGGGGATTCA AAGTAAGTGACTGGGTGAGCG	ERIC	94°C, 30 s	52°C, 30 s	72°C, 300 s	variable	[34]




Fig. 1. PCR and Duplex-PCR for species identification of *Campylobacter* spp. and *C. jejuni*/*C. coli* isolates, respectively; Lane1, *cadF* gene; lane 2, *asp* gene; lane 3, *hipO* gene, and M, 1 kb DNA size marker.

Duplex-PCR assay. The *cadF* gene was positive in 12 (100%) suspected cultures, and an amplification band of 400 bp was obtained for *cadF*⁺ isolates indicative of *Campylobacter* spp. Duplex-PCR for species identification indicated the presence of 10 (4.5%) *C. jejuni* and 2 (1.5%) *C. coli* isolates, respectively (Fig. 1).

Prevalence of virulence-associated genes and cytotoxin distending toxin. All of our *Campylobacter* isolates harbored *flaA* and *ciaB* genes. CDT encoding genes (*cdtA*, *cdtB*, and *cdtC*) involved in CDT production, and *pldA* was found in 7 (58.3%) of *C. jejuni* isolates corresponding to amplification bands of 631, 714, 524, and 913 bp, respectively. However, no CDT encoding genes were found among *C. coli* isolates. The plasmid *virB11* gene was also absent among our *Campylobacter* spp. isolates. The prevalence of *iam* was 100% among *C. coli*, while no amplification was obtained for *C. jejuni* isolates (Table 2).

Enterobacterial repetitive intergenic consensus PCR. In total, 10 out of 12 *Campylobacter* isolates under study produced 7 distinguishable banding profiles by ERIC-PCR genomic fingerprinting, which were corresponded to 10 *C. jejuni* isolates. Dendrogram of ERIC-PCR was created with UPGMA algorithm, which revealed 3 major clusters with 4, 4, and 2 members. No obvious banding pattern was obtained for *C. coli* isolates even after multiple attempts (Fig. 2).

DISCUSSION

Campylobacteriosis is one of the most common bacterial causes of food-borne illness and leading cause of bacterial diarrheal disease in the world [9]. In the present study, the prevalence of *Campylobacter* spp. among diarrheic children was about 6%. In a few previous studies performed in Iran, the prevalence of *Campylobacter* spp. was reported to be 8%, 10.8%, and 8.7% in 2007, 2009, and 2011, respectively [3, 14, 15]. This observation shows the trend of *Campylobacter* infections to be almost unchanged through years in this country.

The prevalence of *Campylobacter* species among diarrheic children was reported to be 5.4% in Turkey [14], 7% in India [15] and 11.1% in Lebanon [16]. As reported by WHO [17], the incidence of *Campylobacteriosis* was 9.3 per 1,000 people in Europe America [17]. All isolates in the present study, either *C. Jejuni* or *C. coli*, harbored *cadF*, *flaA*, and *ciaB* genes. These genes are essential virulence factors involved in *Campylobacter* adhesion and colonization to human intestinal epithelial cells during human infection. The ubiquitous existence of the highly conserved *cadF* gene in 100% of *Campylobacter* spp. was previously reported by Konkel and coworkers [18] and was subsequently used by other investigators for successful detection of *Campylobacter* spp. [19, 20].

The prevalence of virulence-associated genes (*ciaB* and *flaA*) was reported to be 80-100% in different studies concerning *Campylobacter* spp. infections in children with moderate to severe diarrhea [21-23]. Similarly, these genes were also detected within all of our isolates. The *ciaB* and *flaA* are both involved in maximal invasion of human intestinal cells. This result can justify broadly existence of these genes in clinical *Campylobacter* spp.

The *virB11* gene was not found in any of *Campylobacter* isolates under study. This finding is in agreement with the studies by other investigators who did not find *virB11* gene among *Campylobacter* isolates of children from Brazil and Bangladesh [23, 24]. However, a few other studies indicated the prevalence of *virB11* to be 10.7-22.7% among clinical isolates [21, 25]. This finding emphasizes the low prevalence of type IV secretion system apparatus in

Table 2. Prevalence of virulence and toxin genes in *C. jejuni* and *C. coli* isolates under study

Species (No.)	No. of PCR positive (%)										
	<i>cadF</i>	<i>hipO</i>	<i>asp</i>	<i>virB11</i>	<i>ciaB</i>	<i>iam</i>	<i>pldA</i>	<i>flaA</i>	<i>cdtA</i>	<i>cdtB</i>	<i>cdtC</i>
<i>C. jejuni</i> (10)	10 (100)	10 (100)	0 (0)	0 (0)	10 (100)	0 (0)	7 (58.3)	10 (100)	7 (58.3)	7 (58.3)	7 (58.3)
<i>C. coli</i> (2)	2 (100)	0 (0)	2 (100)	0 (0)	2 (100)	2 (100)	0 (0)	2 (100)	0 (0)	0 (0)	0 (0)
Total (12)	12 (100)	10 (83.4)	2 (16.6)	0 (0)	12 (100)	2 (16.6)	7 (58.3)	12 (100)	7 (58.3)	7 (58.3)	7 (58.3)




Fig. 2. Dendrogram generated from the ERIC-PCR profiles of *C. jejuni* isolates from humans in relation to their Profile of virulence-associated genes. Strains 1, 2, 3, and 4 share 80% overall similarity, but strains 7, 8, 9, and 10 share 70% overall similarity. ■ positive □ Negative

Campylobacter spp., which can probably be due to plasmid basis of the gene.

In this study, the *iam* gene that codes for *iam* was detected in 100% of *C. coli* isolates, while none of *C. jejuni* isolates harbored the gene. Similar results have been also reported regarding the high prevalence of *iam* gene among *C. coli* as well as its absence or low distribution among *C. jejuni* isolates of children with diarrhea in Brazil [26]. However, several studies demonstrated no substantial difference in its occurrence among the two species. This result shows that *iam* frequency is controversial [21, 27, 28].

The distribution of both *pldA* and the genes associated with CDT production (*cdtA*, *cdtB*, and *cdtC*) was 58.3% in *C. jejuni* isolates, while none of the genes were detected among *C. coli* isolates. The CDT toxin induces cell cycle arrest in G2 phase and promotes DNA damage; therefore, its presence is supposed to be associated with the severity of the disease in *C. jejuni*. However, variations which may occur within *cdt* gene sequences and may affect their detection through amplification methods should not be ignored during interpretation of negative results in *C. coli*. Moreover, a direct correlation was observed in

this study between detection of *pldA* gene and the presence of white and red blood cells in stool of patients, which may be due to contribution of *pldA* gene product in pathological changes in intestinal epithelium. In agreement with our results, Rizal and colleagues [21] reported the presence of *cdt* and *pldA* genes among 50% and 55% of *C. jejuni* isolates, respectively, while none of the genes were detected within their *C. coli* isolates. Seven distinct ERIC-PCR profiles were distinguished from 10 *C. jejuni* isolates which were located in three clusters by ERIC-PCR. Cluster analysis showed that all isolates (no. 4) within cluster I were isolated from patients who lived in the south of Tehran. Except one isolate, all others within cluster II (no. 3) have also isolated from the patients in a similar geographical location in Tehran (east and center). Moreover, all of the isolates in cluster I and II revealed identical virulence gene content. Two isolates within cluster III were isolated from west of Tehran, but no clear correlation could be determined between ERIC-PCR profile and virulence genes content of isolates within this cluster. Sahilah and colleagues [29] reported that no specific relationship could be extracted between ERIC-PCR analysis and virulence gene

content of their *C. jejuni* isolates. However, Wardak *et al.* [30] showed that ERIC-PCR could clearly divide *C. jejuni* and *C. coli* into two clusters.

ERIC-PCR was unable to type our *C. coli* isolates which raises the question that to how extent is the typeability power of ERIC-PCR for *C. coli* strains. This emphasizes that more studies are needed to clearly understand the ability and role of this typing method in *C. coli* epidemiological studies. However, it is noteworthy that ERIC-PCR analysis has been proved as a well-documented molecular tool in epidemiological studies of *C. jejuni* strains.

To our knowledge, this is the first molecular survey of *C. jejuni* and *C. Coli* genotypes in Iran. Nevertheless, further studies are needed to more clearly understand the correlation between virulence-associated genes and specific genotypes of *C. jejuni* and *C. coli* clinical isolates.

ACKNOWLEDGMENTS

This work was supported by a grant from Research council of Tarbiat Modares University, and it is a part of Ph.D. thesis of Mahdi Ghorbanalizadgan in medical bacteriology branch. We thank M. Akbari and L. Kashi for their contributions to this study.

REFERENCES

- Sangaré L, Nikiéma A, Zimmermann S, Sanou I, Congo-Ouédraogo M, Diabaté A, et al. *Campylobacter* Spp. Epidemiology and Antimicrobial Susceptibility in a Developing Country, Burkina Faso (West Africa). *Afr J Clin Exp Microbiol.* 2012 May; 13(2):10-117.
- Silva J, Leite D, Fernandes M, Mena C, Gibbs PA, and Teixeira P. *Campylobacter* spp. as a Foodborne Pathogen: A Review. *Front Microbiol.* 2011 Sep; 2:200.
- Jafari F, Garcia-Gil L, Salmanzadeh-Ahrabi S, Shokrzadeh L, Aslani MM, Pourhoseingholi MA, et al. Diagnosis and prevalence of enteropathogenic bacteria in children less than 5 years of age with acute diarrhea in Tehran children's hospitals. *J Infect.* 2009 Jan; 58(1): 21-7.
- Koenraad P, Rombouts F, and Notermans S. Epidemiological aspects of thermophilic *Campylobacter* in water-related environments: a review. *Water Environ Res.* 1997 Jan; 69(1):52-63.
- Ica T, Caner V, Istanbulu O, Nguyen HD, Ahmed B, Call DR, et al. Characterization of mono- and mixed-culture *Campylobacter jejuni* biofilms. *Appl Environ Microbiol.* 2011 Dec; 78(4): 1033-1038.
- Nayak R, Stewart TM, and Nawaz MS. PCR identification of *Campylobacter coli* and *Campylobacter jejuni* by partial sequencing of virulence genes. *Mol Cell Probes.* 2005 Jun; 19(3):187-193.
- Konkel ME, Kim BJ, Rivera-Amill V, and Garvis SG. Bacterial secreted proteins are required for the internalization of *Campylobacter jejuni* into cultured mammalian cells. *Mol Microbiol.* 1999 May; 32(4): 691-701.
- Isenberg HD. Clinical microbiology procedures handbook. 2nd ed. ASM Press, USA; 2007.
- Ripabelli G, Tamburro M, Minelli F, Leone A, Sammarco ML. Prevalence of virulence-associated genes and cytotoxic distending toxin production in *Campylobacter* spp. isolated in Italy. *Comp Immunol Microbiol Infect Dis.* 2010 Jul; 33(4):355-64.
- Mohran ZS, Arthur RR, Oyofa BA, Peruski LF, Wasfy MO, Ismail TF, et al. Differentiation of *Campylobacter* isolates on the basis of sensitivity to boiling in water as measured by PCR-detectable DNA. *Appl Environ Microbiol.* 1998 Jan; 64(1): 363-5.
- Al Amri A, Senok AC, Ismaeel AY, Al-Mahmeed AE, Botta GA. Multiplex PCR for direct identification of *Campylobacter* spp. in human and chicken stools. *J Med Microbiol.* 2007 Oct; 56(10):1350-5.
- Versalovic J, Koeuth T, Lupski JR. Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res.* 1991 Dec; 19(24): 6823-31.
- Khakabimamaghani S, Najafi A, Ranjbar R, Raam M. GelClust: a software tool for gel electrophoresis images analysis and dendrogram generation. *Comput Methods Programs Biomed.* 2013 Aug; 111(2):512-8.
- Kayman T, Abay S, Hizlisoy H. Identification of *Campylobacter* spp. isolates with phenotypic methods and multiplex polymerase chain reaction and their antibiotic susceptibilities. *Mikrobiyol Bul.* 2013 Apr; 47(2):230-9.
- Mukherjee P, Ramamurthy T, Bhattacharya MK, Rajendran K, Mukhopadhyay AK. *Campylobacter jejuni* in hospitalized patients with diarrhea, Kolkata, India. *Emerg Infectious Dis.* 2013 Jul; 19(7): 1155-6.
- Dabboussi F, Alam S, Mallat H, Hlais S, Hamze M. Preliminary study on the prevalence of *Campylobacter* in childhood diarrhoea in north Lebanon. *East Mediterr Health J.* 2012 Dec; 18(12): 1225-8.
- WHO and FAO. The global view of campylobacteriosis: report of an expert consultation. www.who.int/foodsafety/publications/foodborne_disease/global_view_campylobacteriosis/en/2012.html
- Konkel ME, Gray SA, Kim BJ, Garvis SG, Yoon J. Identification of the enteropathogens *Campylobacter jejuni* and *Campylobacter coli* based on the *cadF* virulence gene and its product. *J Clin Microbiol.* 1999 Mar; 37(3): 510-7.
- Wieczorek K, Denis E, Lynch O, Osek J. Molecular characterization and antibiotic resistance profiling of *Campylobacter* isolated from cattle in Polish slaughterhouses. *Food Microbiol.* 2013 May; 34(1):130-6.
- Parahitayawa N, Jin L, Leung W, Yam W, and Samaranayake L. Microbiology of odontogenic bacteremia: beyond endocarditis. *Clin Microbiol Rev.* 2009 Apr; 22(1): 46-64.
- Rizal A, Kumar A, and Vidyarthi AS. Prevalence of pathogenic genes in *Campylobacter jejuni* isolates from poultry and human. *Internet J Food Safety.* 2010 Jan;

- 12:29-34.
22. Hamidian M, Sanaei M, Azimi-Rad M, Tajbakhsh M, Dabiri H, and Zali M-R. fla-typing, RAPD analysis, isolation rate and antimicrobial resistance profile of *Campylobacter jejuni* and *Campylobacter coli* of human origin collected from hospitals in Tehran, Iran. *Ann Microbiol.* 2011 Sep; 61(2): 315-21.
 23. Quetz Jda S, Lima IF, Havt A, Prata MM, Cavalcante PA, Medeiros PH, et al. *Campylobacter jejuni* infection and virulence-associated genes in children with moderate to severe diarrhoea admitted to emergency rooms in northeastern Brazil. *J Med Microbiol.* 2012 Apr; 61(Pt 4):507-13.
 24. Talukder KA, Aslam M, Islam Z, Azmi IJ, Dutta DK, Hossain S, et al. Prevalence of virulence genes and cytotoxin production in *Campylobacter jejuni* isolates from diarrheal patients in Bangladesh. *J Clin Microbiol.* 2008 Apr; 46(4): 1485-88.
 25. Datta S, Niwa H, Itoh K. Prevalence of 11 pathogenic genes of *Campylobacter jejuni* by PCR in strains isolated from humans, poultry meat and broiler and bovine faeces. *J Med Microbiol.* 2003 Apr; 52(4):345-8.
 26. Andrzejewska M, Klawe J, Szczepańska B, Śpica D. Occurrence of virulence genes among *Campylobacter jejuni* and *Campylobacter coli* isolates from domestic animals and children. *Pol J Vet Sci.* 2011; 14(2): 207-11.
 27. Rozynek E, Dzierzanowska-Fangrat K, Jozwiak P, Popowski J, Korsak D, Dzierzanowska D. Prevalence of potential virulence markers in Polish *Campylobacter jejuni* and *Campylobacter coli* isolates obtained from hospitalized children and from chicken carcasses. *J Med Microbiol.* 2005 Jul; 54(7): 615-9.
 28. Carvalho AC, Ruiz-Palacios GM, Ramos-Cervantes P, Cervantes L-E, Jiang X, Pickering LK. Molecular characterization of invasive and noninvasive *Campylobacter jejuni* and *Campylobacter coli* isolates. *J Clin Microbiol.* 2001 Apr; 39(4): 1353-9.
 29. Sahilah AM, Tuan Suraya TS, Noraida I, Ahmad Azuhairi A, Chai LC, Son R. Detection of virulence genes and enterobacterial repetitive intergenic consensus-PCR (ERIC-PCR) analysis among raw vegetables isolates of *Campylobacter jejuni*. *Int Food Res J.* 2010 Apr; 17: 681-90.
 30. Wardak S, Jagielski M. Evaluation of genotypic and phenotypic methods for the differentiation of *Campylobacter jejuni* and *Campylobacter coli* clinical isolates from Poland. PFGE, ERIC-PCR, PCR-flaA-RFLP and MLST. *Med Dosw Mikrobiol.* 2009; 61(1):63-77.
 31. Asakura M, Samosornsuk W, Taguchi M, Kobayashi K, Misawa N, Kusumoto M, et al. Comparative analysis of cytotoxin production (cdt) genes among *Campylobacter jejuni*, *C. coli*, and *C. fetus* strains. *Microb Pathog.* 2007 May-Jun; 42(5):174-83.
 32. Biswas D, Hannon SJ, Townsend HG, Potter A, Allan BJ. Genes coding for virulence determinants of *Campylobacter jejuni* in human clinical and cattle isolates from Alberta, Canada, and their potential role in colonization of poultry. *Int Microbiol.* 2011 Mar; 14(1): 25-32.
 33. Müller J, Schulze F, Müller W, Hänel I. PCR detection of virulence-associated genes in *Campylobacter jejuni* strains with differential ability to invade Caco-2 cells and to colonize the chick gut. *Vet Microbiol.* 2006 Mar; 113(1-2):123-9.
 34. Zulkifli Y, Alitheen N, Son R, Raha AR, Samuel L, Yeap SK, Nishibuchi M. Random amplified polymorphic DNA-PCR and ERIC PCR analysis on *Vibrio parahaemolyticus* isolated from cockles in Padang, Indonesia. *Int Food Res J.* 2009 Apr; 16(2): 141-50.