



Published in final edited form as:

Pathog Dis. 2014 June ; 71(1): 81–89. doi:10.1111/2049-632X.12117.

Elevated Suppressor of Cytokine Signaling-1 (SOCS-1): A Mechanism for Dysregulated Osteoclastogenesis in HIV Transgenic Rats

Mark K. Lafferty^a, Lori Fantry^b, Joseph Bryant^{c,d}, Odell Jones^d, Dima Hammoud^f, M. Neale Weitzmann^e, George K. Lewis^a, Alfredo Garzino-Demo^a, and William Reid^{a,f}

^aDivision of Basic Science and Vaccine Research, Institute of Human Virology, and Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore MD 21201

^bDepartment of Medicine and Institute of Human Virology, University of Maryland School of Medicine, Baltimore MD. 21201

^cDepartment of Pathology and Institute of Human Virology, University of Maryland School of Medicine, Baltimore, MD 21201

^dAnimal Models Division, Institute of Human Virology, University of Maryland School of Medicine, Baltimore MD 21201

^eDivision of Endocrinology and Metabolism and Lipids, Emory University School of Medicine, Atlanta, GA 30322; Atlanta Veterans Affairs Medical Center, Decatur, GA 30033 and Winship Cancer Institute, Emory University School of Medicine 30322

^fRadiology and Imaging Sciences, National Institutes of Health/Clinical Center, Bethesda, MD, USA

Abstract

Accelerated bone loss leading to osteopenia, osteoporosis, and bone fracture is a major health problem that is increasingly common in human immunodeficiency virus (HIV) infected patients. The underlying pathogenesis is unclear but occurs in both treatment naïve and individuals receiving antiretroviral therapies. We developed an HIV-1 transgenic rat that exhibits many key features of HIV disease including HIV-1 induced changes in bone mineral density (BMD). A key determinant in the rate of bone loss is the differentiation of osteoclasts, the cells responsible for bone resorption. We found HIV-1 transgenic osteoclast precursors (OCP) express higher levels of suppressor of cytokine signaling-1 (SOCS-1) and TNF receptor associated factor 6 (TRAF6) and are resistant to interferon-gamma (IFN- γ) mediated suppression of osteoclast differentiation. Our data suggest that dysregulated SOCS-1 expression by HIV-1 transgenic OCP promotes

Correspondence to William Reid, DVM, Radiology and Imaging Sciences, National Institutes of Health/Clinical Center, Bethesda, MD, USA. william.reid@nih.gov.

Authorship

W.R., N.W. and M.K.L. conceived the study. W.R. and M.K.L. designed the experiments, performed the experiments, analyzed the data and wrote the manuscript. All coauthors interpreted the data and reviewed the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

osteoclastogenesis leading to the accelerated bone loss observed in this animal model. We propose that elevated SOCS-1 expression in OCP antagonizes the inhibitory effects of IFN- γ and enhances receptor activator of NF- κ B ligand (RANKL) signaling which drives osteoclast differentiation and activation. Understanding the molecular mechanisms of HIV-associated BMD changes has the potential to detect and treat bone metabolism disturbances early and improve the quality of life in patients.

Keywords

SOCS-1; HIV-1; Osteoclast; Osteoporosis

Introduction

The introduction of antiretroviral drugs has dramatically improved both survival and quality of life of HIV patients. However, improved survival has resulted in the emergence of long-term complications associated with HIV-1 disease. In many respects HIV infection recapitulates conditions of accelerated senescence (Deeks, 2009). Two manifestations of natural senescence are skeletal degradation and decreases in immune competence. The two may be closely related; recent studies suggest that immune cells regulate bone homeostasis and cause changes in bone mineral density (BMD) during inflammatory conditions (Gao et al., 2007; Li et al., 2007; Weitzmann and Pacifici, 2007). Recent studies of HIV-1 infected patients show abnormalities in BMD and in bone turnover markers. The prevalence of bone loss in HIV patients range from 22–50% for osteopenia and from 3–20% for osteoporosis (Amiel et al., 2004; Carr et al., 2001; Madeddu et al., 2004; Tebas et al., 2000; Yin et al., 2005). Of concern are recent studies that suggest fractures occur more commonly in HIV-positive men and women, especially in older patients, contributing to HIV-associated morbidity and mortality (Dolan et al., 2006; Ofotokun and Weitzmann; Prior et al., 2007; Stone et al.; Triant et al., 2008; Womack et al., 2011; Young et al., 2011).

The etiology of HIV-1 bone loss is likely multifactorial. Known risk factors for accelerated bone loss are common in HIV-1 infected patients such as low body weight, low vitamin D levels, hypogonadism and alcohol abuse (McComsey et al.; Yin et al.). A meta-analysis of several cross-sectional studies report that the odds ratio for osteoporosis in HIV patients is 3.7 compared to age-matched controls (Brown and Qaqish, 2007). In addition, HIV disease and antiretroviral therapies are risk factors for accelerated bone loss. Cross-sectional studies have shown a correlation between antiretroviral therapy itself and bone loss (Dolan et al., 2006; Mondy et al., 2003; Nolan et al., 2001). However, longitudinal studies of patients on antiretroviral therapy show BMD stabilizes over time (Bolland et al., 2007; Dolan et al., 2006). Further, HIV-1 infection is associated with bone loss in both treated and treatment naïve patients and the HIV-1 proteins gp120 and Vpr *in vitro* increase expression of receptor activator of NF- κ B ligand (RANKL), the key osteoclastogenic cytokine (Brown and Qaqish, 2007; Fakruddin and Laurence, 2003, 2005; Gibellini et al., 2007; Madeddu et al., 2004; McComsey et al., 2010; Paton et al., 1997).

The adult skeleton continuously undergoes bone remodeling to shape and repair damaged and worn bone (Manolagas and Jilka, 1995). Osteoblasts and osteoclasts are the primary cells responsible for bone formation and bone resorption, respectively. The breakdown of bone by osteoclasts is a critical function in bone homeostasis but is also implicated in the pathogenesis of various bone diseases including postmenopausal osteoporosis and inflammatory conditions such as periodontitis (Teitelbaum, 2000). Osteoclasts are large multinucleated hematopoietic cells of the myeloid lineage that develop from precursors following stimulation with macrophage/monocyte-colony forming factor (M-CSF) and RANKL (Boyle et al., 2003), which bind to their receptors c-Fms (also called CSF-1R) and RANK, respectively. M-CSF supports survival and proliferation of myeloid progenitors and promotes generation of osteoclast precursors (OCP) that express RANK (Arai et al., 1999). RANKL, a member of the TNF superfamily of cytokines, provides the critical signal that drives development of OCP and activation of mature osteoclasts (Arai et al., 1999; Kong et al., 1999b; Lacey et al., 1998; Yasuda et al., 1998b). RANKL binding RANK induces recruitment of the adaptor protein TNF receptor associated factor 6 (TRAF6) and activation of the transcription factors nuclear factor κ B (NF- κ B), activation protein 1 (AP-1) and nuclear factor of activated T cells and cytoplasmic 1 (NFATc1), which transactivate osteoclastogenic genes (Takayanagi et al., 2002; Takayanagi et al., 2000; Wong et al., 1998). RANKL is expressed by osteoclasts, chondrocytes, osteocytes, osteoblasts, stromal cells, T cells, and B cells in either a membrane bound or soluble form (Kong et al., 1999b; Lacey et al., 1998; Nakashima et al., 2011; Takayanagi et al., 2000; Vikulina et al., 2010; Xiong et al., 2011). Expression is upregulated by vitamin D₃, prostaglandin E₂, parathyroid hormone, TNF- α , IL-1, IL-6, IL-11, and IL-17 (Kong et al., 1999b; Kotake et al., 1999; Nakashima and Takayanagi, 2008; Vikulina et al., 2010; Wada et al., 2006; Wong et al., 1997).

Osteoclastogenesis is inhibited by IFN- γ and osteoprotegerin (OPG), a soluble decoy receptor of RANKL that blocks osteoclast formation *in vitro* and bone resorption *in vivo* (Simonet et al., 1997; Teitelbaum, 2000; Yasuda et al., 1998b). IFN- γ strongly suppresses osteoclastogenesis *in vitro*, which may be attributable to multiple inhibitory mechanisms. IFN- γ induces apoptosis, suppresses expression of RANK by OCP, down-regulates cathepsin K expression, and blocks RANKL-RANK downstream signaling events (Gao et al., 2007; Kamolmatyakul et al., 2001; Takahashi et al., 1986; Takayanagi et al., 2002; Takayanagi et al., 2005; van't Hof and Ralston, 1997; Wong et al., 1998). IFN- γ inhibits RANK signaling by accelerating the proteasome-mediated degradation of the key adaptor molecule TRAF6 (Takayanagi et al., 2000). Upon binding to its receptor, IFN- γ activates the Janus kinases Jak1 and Jak2 and phosphorylates the transcription factor signal transducer and activator of transcription (STAT)-1, which results in the induction of IFN-responsive gene transcription (Dalpke et al., 2003). IFN- γ and STAT-1 induce expression of SOCS-1, a potent feedback inhibitor of IFN- γ signaling that also cross-inhibits signaling by type 1 IFN receptors and the IL-4 receptor in many lineages of immune cells (Hu and Ivashkiv, 2009).

We have reported that HIV-1 transgenic (Tg) rats have both reduced type 1 cytokine production (IFN- γ and IL-2) and type 1 cytokine responses and a concomitant increase in IL-10 production, which are also observed in patients during progression to AIDS (Clerici

and Shearer, 1993; Reid et al., 2004; Yadav et al., 2009; Yadav et al., 2006). We have shown that IL-10 induces over-expression of SOCS-1 in HIV-1 Tg rat CD4⁺ T cells and dendritic cells, thereby disrupting the IL-12- IFN- γ signaling axis (Yadav et al., 2009). We showed that SOCS-1 is likewise elevated in CD4⁺ T cells from HIV-1 infected patients and is correlated with defective IFN- γ signaling (Reid et al., 2004; Reid et al., 2001a; Yadav et al., 2009; Yadav et al., 2006). It was recently reported that the HIV-1 Tg rat undergoes severe osteoclastic bone resorption and shows an imbalanced ratio of RANKL to OPG in B cells (Vikulina et al.). Here, we demonstrate that along with dysregulated induction of SOCS-1 by OCP, the OCP are resistant to suppression of osteoclast differentiation by IFN- γ . Therefore, we propose that elevated SOCS-1 expression by OCP abrogates IFN- γ mediated control of osteoclastogenesis in the HIV-1 Tg rat and hypothesize that overproduction of SOCS-1 during HIV-1 infection is an important mechanism by which osteoclastogenesis is augmented, leading to an increase in bone loss. This study will help to understand the pathogenesis of HIV-1 induced bone loss in infected patients.

Material and methods

HIV-1 Tg and non-Tg rats

The construction of the HIV-1 transgene and production of the Tg rats have been described (Reid et al., 2001a). Mature (12–15 months) pathogen free Tg rats and age-matched Fisher 344/NHsd non-Tg rats were used in our analysis and were housed under pathogen free conditions in micro-isolator cages on HEPA filtered ventilated racks. The University of Maryland School of Medicine Animal Care and Use Committee approved the experimental protocol.

OCP and splenic mononuclear cell isolation and flow cytometry

Osteoclast precursors were isolated from the bone marrow and RBC were removed by osmotic lyses. Splenic mononuclear cells were isolated using Histopaque-1083 (Sigma-Aldrich). OPC were stained with anti CD11b-FITC (Antigenex America) and isolated using positive selection (Miltenyi Biotec) under conditions described by the manufacture. The positive population was stained for RANK by staining with anti RANK-PE (Imgenex). Flow cytometry was as described previously (Reid et al., 2004) and data were analyzed by FlowJo software. RANK surface expression levels were quantified using QuantiBRITE PE beads (BD Biosciences).

Biochemical indices of bone resorption

Serum C-terminal telopeptide of collagen, a marker of bone resorption, and serum osteocalcin, a specific marker for bone formation, were measured in rats 12–14 months of age using RATlaps and Rat-MID ELISAs respectively (Immunodiagnostic Systems). Samples were measured in triplicate and averaged for each rat.

Real-time PCR

Relative levels of specific mRNA were quantified by real-time RT-PCR analysis using the IQ5 Multicolor Realtime PCR Detection System (Bio-Rad Laboratories). Isolated OCP from control and Tg rats were stimulated at 1.0×10^6 cells/ml at indicated times with IFN- γ or 5

hours with 50 ng/ml sRANKL. Total cellular RNA was prepared using an RNeasy mini kit (Qiagen). First-strand cDNA was synthesized using iScript cDNA Synthesis Kit (Bio-Rad) and analyzed using IQ SYBR Green PCR kit (Bio-Rad Laboratories). Rat specific primer sets for RANKL, OPG, SOCS-1, 18s, tartrate resistant acid phosphatase 5, and cathepsin K were synthesized (Bio-polymer core, University of Maryland, Baltimore or Realltimeprimers.com): RANKL forward primer: 5'- TTT GCT CAC CTC ACCATC AA; reverse primer: 5' – TCC GTT GCT TAA CGT CAT GT; OPG forward primer: 5'- TCC GGA AAC AGA GAA GCA AC; reverse primer: 5' – TGT CCA CCA GAA CACTCA GC; SOCS-1 forward primer: 5'- AGC CAT CCT CGT CCT CGT C; reverse primer: 5'- GCG GAA GGT GCG GAA GTG; 18s forward primer: 5'-GCC TTT CTT CAT TGT CCA GA; reverse primer: 5'- AAA CTT TGG ACG CAG TCT TG; tartrate resistant acid phosphatase 5 forward primer: 5'- CAA CTT CAT GGA CCC TTC TG; reverse primer: 5'- ACC CAT TAG GGG ATA AGC AG and cathepsin K forward primer: 5'- CTT GGC TCG GAA TAA GAA CA; reverse primer: 5'- GAG GCC ACA ACT CTC AGA AA. Samples were run in triplicate and the yield of PCR product was normalized to 18S ribosomal RNA. To control for DNA contamination, equal amounts of RNA were used without reverse transcriptase.

In vitro osteoclastogenesis

Bone marrow cells were collected from the femurs and tibias of 12–15 month old rats. These cells were suspended in a culture dish with α MEM containing 10% FBS for 24 hours at 37°C. Non-adherent cells were collected without contaminating RBC and washed in α MEM. Cells were cultured in 24 well plates (1.0×10^6 cells/ml) in the presence of 20 ng/ml mouse M-CSF (R&D Systems) for 3 days. Change media and culture adherent cells in 20 ng/ml M-CSF and 100 ng/ml rat RANKL (Antigenix America) for an additional 5–7 days with or without rat IFN- γ (BD Pharmingen). The cells were fixed and stained for tartrate-resistant acid phosphatase 5 (TRAP) using a leukocyte acid phosphatase kit (Sigma-Aldrich) according to manufacturer's instructions. The number of TRAP-positive multinuclear cells (>3 nuclei/cell) were determined by counting. RNA was also isolated from adherent cells as described above.

Immunoblotting

Osteoclasts were cultured with or without IFN- γ (10 ng/ml) for 2 hours. Cells were lysed with RIPA buffer (Sigma-Aldrich) containing 0.1mM PMSF, 1 \times EDTA-free protease inhibitor cocktail (Calbiochem), 1 \times Phosphatase Inhibitor Cocktail 2 and 3 (Sigma-Aldrich). Total protein concentration was determined by DC Protein Assay (Bio-Rad Laboratories), and equal amounts of total protein were subjected to sodium dodecyl sulfate–polyacrylamide gel electrophoresis analysis. The primary antibodies included anti-SOCS-1 (Cell Signaling Technology), TRAF6 (Santa Cruz Biotechnology) and anti- β actin (Thermo Scientific).

Cell proliferation assay

Cell proliferation was determined by [3 H]-thymidine uptake.

Statistics

Statistical significance was determined using GraphPad InStat (GraphPad Software). Simple comparisons were made using unpaired or paired 2 tailed Student's t-test for parametric data or Mann-Whitney test for unpaired non-parametric data. Multiple comparisons were made using one-way ANOVA with Tukey Kramer post test. $P < 0.05$ was considered statistically significant. All data is presented as Mean \pm SEM.

Results

HIV-1 Tg rats express increased serum markers of bone resorption

We previously reported that HIV-1 Tg rats have low number of T cells with Th1 effector/memory phenotype, reduced production of IFN- γ and high levels of SOCS-1 expressed in lymphoid and myeloid cells, which led us to hypothesize that IFN- γ signaling is compromised in this animal model, interfering with suppression of osteoclastogenesis (Hayashi et al., 2002; Ohishi et al., 2005; Takayanagi et al., 2000). Consistent with these findings, osteoclastogenic abnormalities include an uncoupling of serum biochemical indices of bone resorption and formation (C-terminal telopeptide of collagen (CTx) and osteocalcin (OCN), respectively) and an abnormal ratio of RANKL to the decoy receptor OPG, a key determinant of the rate of osteoclastogenesis and bone resorption (Haskelberg et al., 2011; Yasuda et al., 1998a; Yasuda et al., 1998b). Therefore, we measured serum levels of CTx, and OCN. Serum CTx (a bone resorption marker) was significantly increased in HIV-1 Tg rats relative to age-matched non-Tg controls (13 ± 1.5 ng/ml and 8.2 ± 0.8 ng/ml, respectively, t-test; $p = 0.018$; $n = 5$) (Figure 1A), while differences in serum OCN (a bone formation marker) levels in HIV- Tg rats relative to age-matched non-Tg controls did not reach significance (Figure 1B).

HIV-1 Tg rat PBMC express an increased ratio of RANKL to OPG

An increase in the ratio of RANKL to OPG accelerates the rate of osteoclastic bone resorption. To further assess osteoclastic bone resorption we measured levels of RANKL and OPG mRNA from HIV-1 Tg and non-Tg control PBMC using real time RT-PCR. Higher levels of RANKL mRNA was measured in HIV-1 Tg rats compared to controls (4.4 ± 0.09 and 1.1 ± 0.38 , respectively; Mann-Whitney; $p = 0.02$). In contrast, the relative expression of OPG mRNA did not differ significantly. This increase in the ratio of RANKL to OPG mRNA levels (Figure 2) suggests enhanced osteoclastogenesis as the basis for the accelerated bone resorption identified in the HIV-1 Tg rat. Similar results were also observed in mononuclear cells isolated from the spleen. HIV-1 Tg splenic mononuclear cells express elevated levels of SOCS-1 (Mann-Whitney; $p = 0.018$; $n = 3$) and RANKL (Mann-Whitney; $p = 0.0005$; $n = 3$) mRNA (Figure S1A and B, respectively) compared to control. Similar levels of OPG mRNA (Figure S1C) were observed resulting in an increase in the ratio of RANKL to OPG (Mann-Whitney; $p = 0.0005$; $n = 3$) (Figure S1D).

HIV-1 Tg rats express increased SOCS-1 mRNA and protein

We hypothesized that compromised IFN- γ signaling mediated by SOCS-1 prevents effective suppression of osteoclast differentiation. Therefore, we analyzed SOCS-1 expression in

HIV-1 Tg and control OCP. HIV-1 Tg and non-Tg control OCP were treated with IFN- γ for 2 hours. Figure 3A shows that HIV-1 Tg OCP had approximately 2.0 fold greater basal levels of SOCS-1 mRNA relative to non-Tg controls and a highly significant 14.7 fold increase (ANOVA; $p=0.008$) following IFN- γ stimulation. Treatment with IFN- γ induced higher SOCS-1 protein expression in HIV-1 Tg OCP compared to non-Tg control OCP (Figure 3B). In the absence of IFN- γ treatment, HIV-1 Tg and non-Tg control OCP express similar levels of the RANK receptor and no significant difference in proliferation was observed (Supplemental Figure S2A–C).

HIV-1 Tg rats are resistant to IFN- γ mediated suppression of osteoclast differentiation

We tested whether the elevated SOCS-1 expression correlated with lack of suppression of osteoclast differentiation following treatment with exogenous IFN- γ . As shown in Figure 4A, significantly more HIV-1 Tg OCP differentiated into tartrate-resistant acid phosphatase 5 (TRAP)⁺ multi-nucleated cells, in the presence of 500 and 1000 pg/ml IFN- γ (Mann-Whitney; $p=0.008$ and 0.032 , $n=5$, respectively) than non-Tg controls. Resistance to IFN- γ suppression by HIV-1 Tg OCP was confirmed by measuring expression of mRNA for the osteoclast-specific gene TRAP and the predominant protease in bone-resorption, cathepsin K, by real time RT-PCR. As shown in Figure 4B, there was a significant 3.7 and 2.8 fold increase of TRAP (t-test; $p=0.048$; $n=4$) and cathepsin K (t-test; $p=0.048$; $n=3$) mRNA detected in developing osteoclasts from HIV-1 Tg rats relative to non-Tg controls following treatment with 1000 pg/ml of IFN- γ , respectively. Our findings that SOCS-1 is elevated in HIV-1 Tg OCP (Figure 3A and B) and that HIV-1 Tg OCP are resistant to IFN- γ suppression of the RANKL induced bone-resorbing enzymes, TRAP and cathepsin K (Figure 4A and B), suggest that increased SOCS-1 expression attenuates anti-osteoclastogenesis mediated by IFN- γ . Therefore, we tested whether IFN- γ mediated degradation of TRAF6, an adaptor protein critical in RANKL signaling, is disrupted in HIV-1 Tg OCP. HIV-1 Tg and non-Tg control OCP were treated with RANKL and IFN- γ for 24 hours. RANKL treated HIV-1 Tg OCP express higher levels of TRAF6 protein compared to non-Tg control OCP as determined by Western blot (Figure 4C). Further, RANKL treated HIV-1 Tg OCP express higher levels of TRAF6 following treatment with 500 pg/ml of IFN- γ compared to non-Tg controls (Figure 4C). These data suggest that increased SOCS-1 expression by HIV-1 Tg OCP is associated with attenuated IFN- γ inhibition of osteoclastogenesis.

RANKL induction of SOCS-1 in non-Tg control osteoclast precursors

IFN- γ inhibits osteoclastogenesis of OCP by suppressing the expression of c-fms and/or RANK signaling through TRAF6. RANKL induces SOCS-1 in OCP (Hayashi et al., 2002); therefore, we measured SOCS-1 mRNA levels in non-Tg control rat OCP following 5 hours of treatment with 50 ng/ml of sRANKL. SOCS-1 mRNA increased 2.8 fold (paired t-test; $p=0.0159$; $n=5$) relative to unstimulated controls (Figure 5). These data suggest that the increased RANKL expression by HIV-1 Tg PBMC (Figure 2) plays a role in increased SOCS-1 expression by HIV-1 Tg OCP and the reduced ability of IFN- γ to attenuate osteoclast differentiation (Figure 4).

Discussion

In the past, few HIV infected patients lived long enough to experience the morbidity and mortality of bone loss. Measurement of BMD as a routine test in elderly HIV-infected patients has not previously been recommended. A detailed history and physical to assess individual risk for osteopenia/osteoporosis is now recommended. Abnormal clinical laboratory values obtained during the course of HIV treatment (i.e. an elevated alkaline phosphatase or low testosterone) suggest the need to test for changes in BMD. No studies, however, have thus far addressed the underlying mechanism(s) for abnormal BMD reported in HIV-1 disease; consequently the pathogenesis remains poorly understood. In this study we have identified a potential pathological mechanism of HIV-1 induced bone loss mediated by SOCS-1 enhancement of osteoclastogenesis. The differentiation of osteoclasts is dependent on signals from RANK, stimulated by its ligand RANKL (Kong et al., 1999a; Vikulina et al.; Wong et al., 1997). Upregulation of RANKL by inflammatory cytokines such as TNF- α contributes to osteoclastogenesis (Lam et al., 2000; Zhang et al., 2001). TNF- α has not, however, been implicated in increased RANKL expression in HIV-1 Tg rats (Vikulina et al.). The HIV-1 proteins Vpr and gp120 enhance expression of RANKL (Fakruddin and Laurence, 2003, 2005). We have previously shown elevated serum levels of gp120 in the HIV-1 Tg rat (Reid et al., 2001b); therefore increased RANKL expression may be a consequence of the expression of this HIV-1 transgene protein. Relevantly, we have demonstrated that the HIV Tg rat expresses elevated levels of SOCS-1 and that IFN- γ treatment results in increased levels of TRAF6 and impaired suppression of osteoclastogenesis. Along with reduced production of IFN- γ by CD4+ T cells (Reid et al., 2004; Yadav et al., 2009) and increased RANKL expression, these results suggest that SOCS-1 amplifies the osteoclastogenic activity of RANKL thereby enhancing bone loss in the HIV-1 Tg rat. In this model, IL-10 induction of SOCS-1 by CD4+ T cells and dendritic cells inhibits both LPS and IFN- γ signaling. Additionally, we reported that increased SOCS-1 expression by HIV-1 infected patients altered IFN- γ signaling by CD4 T cells (Yadav et al., 2009).

Here, consistent with previous findings, we demonstrate that similar to HIV-1 infected patients, the HIV-1 Tg rat undergoes pathological bone resorption. From a mechanistic viewpoint, we demonstrate that HIV-1 Tg rats have increased levels of the serum bone resorption protein CTx and higher expression of RANKL mRNA resulting in an increased ratio of RANKL to OPG in PBMC and mononuclear cells isolated from the spleen. However, we did not observe a concurrent decrease in OPG mRNA, as previously reported in bone marrow and total splenic cells (Vikulina et al., 2010). We now report that HIV-1 Tg rat OCP express higher levels of SOCS-1 and TRAF6 which in conjunction with elevated RANKL expression enhances osteoclastogenesis and resistance to suppression of osteoclast differentiation by IFN- γ signaling. The molecular mechanism of suppression of RANK signaling by IFN- γ has not been clarified; Takayanagi *et al.* have demonstrated that IFN- γ induces the ubiquitination and degradation of TRAF6 by a STAT1-dependent mechanism in murine cells (Takayanagi et al., 2000) while Ji *et al.* demonstrated inhibition of RANK expression and osteoclastogenesis by TLR-4 and IFN- γ signaling synergy, likely through a down-regulation of c-Fms expression (Ji et al., 2009). In conclusion, our data suggest a link

between high levels of SOCS-1 by OCP and enhanced RANK signaling and resistance to IFN- γ induced suppression of osteoclastogenesis. Understanding the mechanisms of HIV-1 induced bone loss and the role played by over-expression of SOCS-1 will be critical for early detection of changes in BMD and in developing effective therapy. We speculate that elevated SOCS-1 levels may be predictive for reduced BMD and an increased likelihood of HIV-1 induced fragility fractures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. Marvin Reitz for his critical reading of this manuscript. We also thank Dr. Bryan Taylor, manager of the Core facility at the Institute of Human Virology for his technical assistance. This work was supported by National Institute of Allergy and Infectious Diseases, NIH grants R01A1063171 and R01A1087181 and National Institute of Neurological Disorders And Stroke, NIH grant R01NS066842.

References

- Amiel C, Ostertag A, Slama L, Baudoin C, N'Guyen T, Lajeunie E, Neit-Ngeilh L, Rozenbaum W, De Vernejoul MC. BMD is reduced in HIV-infected men irrespective of treatment. *J Bone Miner Res.* 2004; 19:402–409. [PubMed: 15040828]
- Arai F, Miyamoto T, Ohneda O, Inada T, Sudo T, Brasel K, Miyata T, Anderson DM, Suda T. Commitment and differentiation of osteoclast precursor cells by the sequential expression of c-Fms and receptor activator of nuclear factor kappaB (RANK) receptors. *J Exp Med.* 1999; 190:1741–1754. [PubMed: 10601350]
- Bolland MJ, Grey AB, Horne AM, Briggs SE, Thomas MG, Ellis-Pegler RB, Woodhouse AF, Gamble GD, Reid IR. Bone mineral density remains stable in HAART-treated HIV-infected men over 2 years. *Clinical endocrinology.* 2007; 67:270–275. [PubMed: 17547686]
- Boyle WJ, Simonet WS, Lacey DL. Osteoclast differentiation and activation. *Nature.* 2003; 423:337–342. [PubMed: 12748652]
- Brown TT, Qaqish RB. Response to Berg et al. 'Antiretroviral therapy and the prevalence of osteopenia and osteoporosis: a meta-analytic review'. *Aids.* 2007; 21:1830–1831. [PubMed: 17690589]
- Carr A, Miller J, Eisman JA, Cooper DA. Osteopenia in HIV-infected men: association with asymptomatic lactic acidemia and lower weight pre-antiretroviral therapy. *AIDS.* 2001; 15:703–709. [PubMed: 11371684]
- Clerici M, Shearer GM. A TH1-->TH2 switch is a critical step in the etiology of HIV infection. *Immunology today.* 1993; 14:107–111. [PubMed: 8096699]
- Dalpe AH, Eckerle S, Frey M, Heeg K. Triggering of Toll-like receptors modulates IFN-gamma signaling: involvement of serine 727 STAT1 phosphorylation and suppressors of cytokine signaling. *EurJImmunol.* 2003; 33:1776–1787.
- Deeks SG. Immune dysfunction, inflammation, and accelerated aging in patients on antiretroviral therapy. *Top HIV Med.* 2009; 17:118–123. [PubMed: 19890183]
- Dolan SE, Kanter JR, Grinspoon S. Longitudinal analysis of bone density in human immunodeficiency virus-infected women. *J Clin Endocrinol Metab.* 2006; 91:2938–2945. [PubMed: 16735489]
- Fakruddin JM, Laurence J. HIV envelope gp120-mediated regulation of osteoclastogenesis via receptor activator of nuclear factor kappa B ligand (RANKL) secretion and its modulation by certain HIV protease inhibitors through interferongamma/RANKL cross-talk. *J Biol Chem.* 2003; 278:48251–48258. [PubMed: 12975380]

- Fakruddin JM, Laurence J. HIV-1 Vpr enhances production of receptor of activated NF-kappaB ligand (RANKL) via potentiation of glucocorticoid receptor activity. *Arch Virol*. 2005; 150:67–78. [PubMed: 15449141]
- Gao Y, Grassi F, Ryan MR, Terauchi M, Page K, Yang X, Weitzmann MN, Pacifici R. IFN-gamma stimulates osteoclast formation and bone loss in vivo via antigen-driven T cell activation. *J Clin Invest*. 2007; 117:122–132. [PubMed: 17173138]
- Gibellini D, Borderi M, De Crignis E, Cicola R, Vescini F, Caudarella R, Chiodo F, Re MC. RANKL/OPG/TRAIL plasma levels and bone mass loss evaluation in antiretroviral naive HIV-1-positive men. *Journal of medical virology*. 2007; 79:1446–1454. [PubMed: 17705184]
- Haskelberg H, Carr A, Emery S. Bone turnover markers in HIV disease. *AIDS reviews*. 2011; 13:240–250. [PubMed: 21975360]
- Hayashi T, Kaneda T, Toyama Y, Kumegawa M, Hakeda Y. Regulation of receptor activator of NF-kappa B ligand-induced osteoclastogenesis by endogenous interferon-beta (INF-beta) and suppressors of cytokine signaling (SOCS). The possible counteracting role of SOCSs- in IFN-beta-inhibited osteoclast formation. *J Biol Chem*. 2002; 277:27880–27886. [PubMed: 12023971]
- Hu X, Ivashkiv LB. Cross-regulation of signaling pathways by interferongamma: implications for immune responses and autoimmune diseases. *Immunity*. 2009; 31:539–550. [PubMed: 19833085]
- Ji JD, Park-Min KH, Shen Z, Fajardo RJ, Goldring SR, McHugh KP, Ivashkiv LB. Inhibition of RANK expression and osteoclastogenesis by TLRs and IFN-gamma in human osteoclast precursors. *J Immunol*. 2009; 183:7223–7233. [PubMed: 19890054]
- Kamolmatyakul S, Chen W, Li YP. Interferon-gamma down-regulates gene expression of cathepsin K in osteoclasts and inhibits osteoclast formation. *J Dent Res*. 2001; 80:351–355. [PubMed: 11269728]
- Kong YY, Feige U, Sarosi I, Bolon B, Tafuri A, Morony S, Capparelli C, Li J, Elliott R, McCabe S, et al. Activated T cells regulate bone loss and joint destruction in adjuvant arthritis through osteoprotegerin ligand. *Nature*. 1999a; 402:304–309. [PubMed: 10580503]
- Kong YY, Yoshida H, Sarosi I, Tan HL, Timms E, Capparelli C, Morony S, Oliveira-dos-Santos AJ, Van G, Itie A, et al. OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature*. 1999b; 397:315–323. [PubMed: 9950424]
- Kotake S, Udagawa N, Takahashi N, Matsuzaki K, Itoh K, Ishiyama S, Saito S, Inoue K, Kamatani N, Gillespie MT, et al. IL-17 in synovial fluids from patients with rheumatoid arthritis is a potent stimulator of osteoclastogenesis. *J Clin Invest*. 1999; 103:1345–1352. [PubMed: 10225978]
- Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, et al. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell*. 1998; 93:165–176. [PubMed: 9568710]
- Lam J, Takeshita S, Barker JE, Kanagawa O, Ross FP, Teitelbaum SL. TNF-alpha induces osteoclastogenesis by direct stimulation of macrophages exposed to permissive levels of RANK ligand. *J Clin Invest*. 2000; 106:1481–1488. [PubMed: 11120755]
- Li Y, Toraldo G, Li A, Yang X, Zhang H, Qian WP, Weitzmann MN. B cells and T cells are critical for the preservation of bone homeostasis and attainment of peak bone mass in vivo. *Blood*. 2007; 109:3839–3848. [PubMed: 17202317]
- Madeddu G, Spanu A, Solinas P, Calia GM, Lovigu C, Chessa F, Mannazzu M, Falchi A, Mura MS. Bone mass loss and vitamin D metabolism impairment in HIV patients receiving highly active antiretroviral therapy. *Q J Nucl Med Mol Imaging*. 2004; 48:39–48. [PubMed: 15195003]
- Manolagas SC, Jilka RL. Bone marrow, cytokines, and bone remodeling. Emerging insights into the pathophysiology of osteoporosis. *N Engl J Med*. 1995; 332:305–311. [PubMed: 7816067]
- McComsey GA, Tebas P, Shane E, Yin MT, Overton ET, Huang JS, Aldrovandi GM, Cardoso SW, Santana JL, Brown TT. Bone disease in HIV infection: a practical review and recommendations for HIV care providers. *Clin Infect Dis*. 51:937–946. [PubMed: 20839968]
- McComsey GA, Tebas P, Shane E, Yin MT, Overton ET, Huang JS, Aldrovandi GM, Cardoso SW, Santana JL, Brown TT. Bone disease in HIV infection: a practical review and recommendations for HIV care providers. *Clin Infect Dis*. 2010; 51:937–946. [PubMed: 20839968]

- Mondy K, Yarasheski K, Powderly WG, Whyte M, Claxton S, DeMarco D, Hoffmann M, Tebas P. Longitudinal evolution of bone mineral density and bone markers in human immunodeficiency virus-infected individuals. *Clin Infect Dis*. 2003; 36:482–490. [PubMed: 12567307]
- Nakashima T, Hayashi M, Fukunaga T, Kurata K, Oh-Hora M, Feng JQ, Bonewald LF, Kodama T, Wutz A, Wagner EF, et al. Evidence for osteocyte regulation of bone homeostasis through RANKL expression. *Nat Med*. 2011; 17:1231–1234. [PubMed: 21909105]
- Nakashima T, Takayanagi H. The dynamic interplay between osteoclasts and the immune system. *Arch Biochem Biophys*. 2008; 473:166–171. [PubMed: 18410741]
- Nolan D, Upton R, McKinnon E, John M, James I, Adler B, Roff G, Vasikaran S, Mallal S. Stable or increasing bone mineral density in HIV-infected patients treated with nelfinavir or indinavir. *AIDS*. 2001; 15:1275–1280. [PubMed: 11426072]
- Ototokun I, Weitzmann MN. HIV-1 infection and antiretroviral therapies: risk factors for osteoporosis and bone fracture. *Curr Opin Endocrinol Diabetes Obes*. 17:523–529. [PubMed: 20844427]
- Ohishi M, Matsumura Y, Aki D, Mashima R, Taniguchi K, Kobayashi T, Kukita T, Iwamoto Y, Yoshimura A. Suppressors of cytokine signaling-1 and -3 regulate osteoclastogenesis in the presence of inflammatory cytokines. *J Immunol*. 2005; 174:3024–3031. [PubMed: 15728516]
- Paton NI, Macallan DC, Griffin GE, Pazianas M. Bone mineral density in patients with human immunodeficiency virus infection. *Calcif Tissue Int*. 1997; 61:30–32. [PubMed: 9192508]
- Prior J, Burdge D, Maan E, Milner R, Hankins C, Klein M, Walmsley S. Fragility fractures and bone mineral density in HIV positive women: a case-control population-based study. *Osteoporos Int*. 2007; 18:1345–1353. [PubMed: 17665239]
- Reid W, Abdelwahab S, Sadowska M, Huso D, Neal A, Ahearn A, Bryant J, Gallo RC, Lewis GK, Reitz M. HIV-1 transgenic rats develop T cell abnormalities. *Virology*. 2004; 321:111–119. [PubMed: 15033570]
- Reid W, Sadowska M, Denaro F, Rao S, Foulke J Jr. Hayes N, Jones O, Doodnauth D, Davis H, Sill A, et al. An HIV-1 transgenic rat that develops HIV-related pathology and immunologic dysfunction. *Proc Natl Acad Sci USA*. 2001a; 98:9271–9276. [PubMed: 11481487]
- Reid W, Sadowska M, Denaro F, Rao S, Foulke J Jr. Hayes N, Jones O, Doodnauth D, Davis H, Sill A, et al. An HIV-1 transgenic rat that develops HIV-related pathology and immunologic dysfunction. *Proc Natl Acad Sci U S A*. 2001b; 98:9271–9276. [PubMed: 11481487]
- Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, Luthy R, Nguyen HQ, Wooden S, Bennett L, Boone T, et al. Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell*. 1997; 89:309–319. [PubMed: 9108485]
- Stone B, Dockrell D, Bowman C, McCloskey E. HIV and bone disease. *Arch Biochem Biophys*. 503:66–77. [PubMed: 20682280]
- Takahashi N, Mundy GR, Roodman GD. Recombinant human interferon-gamma inhibits formation of human osteoclast-like cells. *J Immunol*. 1986; 137:3544–3549. [PubMed: 3097126]
- Takayanagi H, Kim S, Koga T, Nishina H, Isshiki M, Yoshida H, Saiura A, Isobe M, Yokochi T, Inoue J, et al. Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev Cell*. 2002; 3:889–901. [PubMed: 12479813]
- Takayanagi H, Ogasawara K, Hida S, Chiba T, Murata S, Sato K, Takaoka A, Yokochi T, Oda H, Tanaka K, et al. T-cell-mediated regulation of osteoclastogenesis by signalling cross-talk between RANKL and IFN-gamma. *Nature*. 2000; 408:600–605. [PubMed: 11117749]
- Takayanagi H, Sato K, Takaoka A, Taniguchi T. Interplay between interferon and other cytokine systems in bone metabolism. *Immunol Rev*. 2005; 208:181–193. [PubMed: 16313349]
- Tebas P, Powderly WG, Claxton S, Marin D, Tantisiriwat W, Teitelbaum SL, Yarasheski KE. Accelerated bone mineral loss in HIV-infected patients receiving potent antiretroviral therapy. *AIDS*. 2000; 14:F63–F67. [PubMed: 10770534]
- Teitelbaum SL. Bone resorption by osteoclasts. *Science*. 2000; 289:1504–1508. [PubMed: 10968780]
- Triant VA, Brown TT, Lee H, Grinspoon SK. Fracture prevalence among human immunodeficiency virus (HIV)-infected versus non-HIV-infected patients in a large U.S. healthcare system. *J Clin Endocrinol Metab*. 2008; 93:3499–3504. [PubMed: 18593764]

- van't Hof RJ, Ralston SH. Cytokine-induced nitric oxide inhibits bone resorption by inducing apoptosis of osteoclast progenitors and suppressing osteoclast activity. *J Bone Miner Res.* 1997; 12:1797–1804. [PubMed: 9383684]
- Vikulina T, Fan X, Yamaguchi M, Roser-Page S, Zayzafoon M, Guidot DM, Ofotokun I, Weitzmann MN. Alterations in the immuno-skeletal interface drive bone destruction in HIV-1 transgenic rats. *Proc Natl Acad Sci U S A.*
- Vikulina T, Fan X, Yamaguchi M, Roser-Page S, Zayzafoon M, Guidot DM, Ofotokun I, Weitzmann MN. Alterations in the immuno-skeletal interface drive bone destruction in HIV-1 transgenic rats. *Proc Natl Acad Sci U S A.* 2010; 107:13848–13853. [PubMed: 20643942]
- Wada T, Nakashima T, Hiroshi N, Penninger JM. RANKL-RANK signaling in osteoclastogenesis and bone disease. *Trends Mol Med.* 2006; 12:17–25. [PubMed: 16356770]
- Weitzmann MN, Pacifici R. T cells: unexpected players in the bone loss induced by estrogen deficiency and in basal bone homeostasis. *Ann N Y Acad Sci.* 2007; 1116:360–375. [PubMed: 18083938]
- Womack JA, Goulet JL, Gibert C, Brandt C, Chang CC, Gulanski B, Fraenkel L, Mattocks K, Rimland D, Rodriguez-Barradas MC, et al. Increased risk of fragility fractures among HIV infected compared to uninfected male veterans. *PLoS One.* 2011; 6:e17217. [PubMed: 21359191]
- Wong BR, Josien R, Lee SY, Vologodskaja M, Steinman RM, Choi Y. The TRAF family of signal transducers mediates NF-kappaB activation by the TRANCE receptor. *J Biol Chem.* 1998; 273:28355–28359. [PubMed: 9774460]
- Wong BR, Rho J, Arron J, Robinson E, Orlinick J, Chao M, Kalachikov S, Cayani E, Bartlett FS 3rd, Frankel WN, et al. TRANCE is a novel ligand of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. *J Biol Chem.* 1997; 272:25190–25194. [PubMed: 9312132]
- Xiong J, Onal M, Jilka RL, Weinstein RS, Manolagas SC, O'Brien CA. Matrix-embedded cells control osteoclast formation. *Nat Med.* 2011; 17:1235–1241. [PubMed: 21909103]
- Yadav A, Fitzgerald P, Sajadi MM, Gilliam B, Lafferty MK, Redfield R, Reid W. Increased expression of suppressor of cytokine signaling-1 (SOCS-1): A mechanism for dysregulated T helper-1 responses in HIV-1 disease. *Virology.* 2009; 385:126–133. [PubMed: 19135225]
- Yadav A, Pati S, Nyugen A, Barabitskaja O, Mondal P, Anderson M, Gallo RC, Huso DL, Reid W. HIV-1 transgenic rat CD4+ T cells develop decreased CD28 responsiveness and suboptimal Lck tyrosine dephosphorylation following activation. *Virology.* 2006; 353:357–365. [PubMed: 16828835]
- Yasuda H, Shima N, Nakagawa N, Mochizuki SI, Yano K, Fujise N, Sato Y, Goto M, Yamaguchi K, Kuriyama M, et al. Identity of osteoclastogenesis inhibitory factor (OCIF) and osteoprotegerin (OPG): a mechanism by which OPG/OCIF inhibits osteoclastogenesis in vitro. *Endocrinology.* 1998a; 139:1329–1337. [PubMed: 9492069]
- Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinosaki M, Mochizuki S, Tomoyasu A, Yano K, Goto M, Murakami A, et al. Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci U S A.* 1998b; 95:3597–3602. [PubMed: 9520411]
- Yin M, Dobkin J, Brudney K, Becker C, Zadel JL, Manandhar M, Adesso V, Shane E. Bone mass and mineral metabolism in HIV+ postmenopausal women. *Osteoporos Int.* 2005; 16:1345–1352. [PubMed: 15754081]
- Yin MT, McMahon DJ, Ferris DC, Zhang CA, Shu A, Staron R, Colon I, Laurence J, Dobkin JF, Hammer SM, et al. Low bone mass and high bone turnover in postmenopausal human immunodeficiency virus-infected women. *J Clin Endocrinol Metab.* 95:620–629. [PubMed: 19965927]
- Young B, Dao CN, Buchacz K, Baker R, Brooks JT. Increased rates of bone fracture among HIV-infected persons in the HIV Outpatient Study (HOPS) compared with the US general population, 2000–2006. *Clin Infect Dis.* 2011; 52:1061–1068. [PubMed: 21398272]
- Zhang YH, Heulsmann A, Tondravi MM, Mukherjee A, Abu-Amer Y. Tumor necrosis factor-alpha (TNF) stimulates RANKL-induced osteoclastogenesis via coupling of TNF type 1 receptor and RANK signaling pathways. *J Biol Chem.* 2001; 276:563–568. [PubMed: 11032840]




Figure 1.

Bone resorption in HIV-1 Tg rats. (A) C-terminal telopeptide (CTx) and (B) osteocalcin, markers of bone resorption and bone formation respectively, were quantified in the serum of non-Tg and HIV-1 Tg rats by ELISA. Samples were analyzed in triplicate.




Figure 2.

Increased RANKL and OPG ratio in HIV-1 Tg rat. RANKL and OPG mRNA was quantified using real time RT-PCR from PBMCs for HIV-1 Tg (n=4) and non-Tg (n=6) as described in the *Material and methods* section and the ratio determined for relative expression. Samples were analyzed in triplicate and data normalized to the 18s ribosomal RNA.





Figure 3.

SOCS-1 mRNA and protein expression are elevated in HIV-1 Tg rats. (A) OCP (1.0×10^6 /ml) from non-Tg and HIV-1 Tg rats were stimulated with for 2 hours with 10ng/ml of IFN- γ and levels of SOCS-1 mRNA were determined by real-time quantitative RT-PCR. Samples were analyzed in triplicate and data normalized to the expression of 18s ribosomal RNA. (B) OCP from non-Tg (n=3) and HIV-1 Tg (n=3) rats were stimulated for 2 hours with 10ng/ml of IFN- γ . SOCS-1 and β actin were detected by Western blotting.

**Figure 4.**

SOCS-1 over-expression confers resistance to suppression of osteoclast differentiation by IFN- γ . (A) OCP (1.0×10^6 /ml in a 24 well plate) were cultured with 20 ng/ml of M-CSF and 100 ng/ml RANKL with various concentrations of IFN- γ for 7–8 days. Osteoclast numbers (TRAP⁺ multinuclear cells (MNC)) were determined from duplicate samples. TRAP⁺ MNC cells are represented as a percent of osteoclast produced without added IFN- γ . Total RNA was isolated from developing osteoclast cultured for 5-days with IFN- γ (1000 pg/ml), and RT-PCR performed for (B) TRAP and Cathepsin K. Shown are expression levels of TRAP and Cathepsin K in HIV-1 Tg cells compared to non-Tg. Samples were analyzed in triplicate and data normalized to the 18s ribosomal RNA. (C) Non-Tg and HIV-1 Tg OCP were cultured in a 20 ng/ml of M-CSF and stimulated with 100 ng/ml RANKL and IFN- γ for 24 hours. TRAF6 and β actin were detected by Western blotting.




Figure 5.

SOCS-1 mRNA induction by RANKL in non-Tg control OCP. OCP from non-Tg controls stimulated with soluble (s) RANKL (50 ng/ml) for 5 h. Shown is the expression of SOCS-1 mRNA in control OCP. Levels of SOCS-1 mRNA were determined by real time RT-PCR as described in *Material and methods*. Samples were analyzed in triplicate and data normalized to the expression of 18s ribosomal RNA.