

## Archaeobiotics

### Proposed therapeutic use of archaea to prevent trimethylaminuria and cardiovascular disease

Jean-François Brugère<sup>1\*</sup>, Guillaume Borrel<sup>1,2</sup>, Nadia Gaci<sup>1</sup>, William Tottey<sup>1</sup>, Paul W O'Toole<sup>2</sup>, and Corinne Malpuech-Brugère<sup>3</sup>

<sup>1</sup>EA 4678 CIDAM; Clermont Université; Université d'Auvergne; Clermont-Ferrand, France; <sup>2</sup>Department of Microbiology and Alimentary Pharmabiotic Centre; University College Cork; Cork, Ireland; <sup>3</sup>Clermont Université; Université d'Auvergne – INRA; UMR 1019 UNH; Clermont-Ferrand, France

**T**rimethylamine (TMA) is produced by gut bacteria from dietary ingredients. In individuals with a hereditary defect in flavin-containing monooxygenase 3, bacterial TMA production is believed to contribute to the symptoms of trimethylaminuria (TMAU; fish-odor syndrome). Intestinal microbiota TMA metabolism may also modulate atherosclerosis risk by affecting trimethylamine oxide (TMAO) production levels. We propose that reducing TMA formation in the gut by converting it to an inert molecule could be used to prevent or limit these human diseases, while avoiding the major drawbacks of other clinical interventions. Reducing TMA levels by microbiological interventions could also be helpful in some vaginoses. Particular members of a recently discovered group of methanogens, that are variably present in the human gut, are unusual in being apparently restricted to utilizing only methyl compounds including TMA as substrates. We confirmed experimentally that one of these strains tested, *Methanomassiliicoccus luminyensis* B10, is able to deplete TMA, by reducing it with H<sub>2</sub> for methanogenesis. We therefore suggest that members of this archaeal lineage could be used as treatments for metabolic disorders.

#### The Gut Origin of TMA in Human

The tertiary amine trimethylamine (TMA) is a volatile compound which has a characteristic fishy odor, and is formed by bacterial reduction of trimethylamine

oxide (TMAO). This alkylamine is detected as unpleasant by the human olfactory system at even very low levels, thereby preventing humans from ingesting rotting fish. Ironically, this molecule is also endogenous in humans, being synthesized in the gut and sometimes in the vagina by the endogenous microbiota. The microbiome of the gastrointestinal tract is better characterized than that of other body sites, so only this niche will be discussed here, although relevant metabolic processes in the vagina are probably similar.

In the gut, TMA is formed by microbial conversion of dietary ingredients<sup>1</sup> and is further absorbed before being oxidized into TMAO in the liver (see Fig. 1). Such TMA precursors include TMAO (in abundance in seafood), and choline (eggs, soybean, cauliflower), which is likely the most important TMA source. Recent data also show that L-carnitine<sup>2</sup> found in red meat (and in some energy drinks and dietary supplements) is converted into TMA. The metabolic pathways leading to TMA are thought to be exclusively microbial in humans: the mechanism of conversion of choline by an anaerobic mammalian gut *Desulfovibrio* has recently been identified and relies on the *cut* (choline utilization) operon, in which *cutC* encodes a glyceryl radical enzyme with a choline trimethylamine-lyase activity.<sup>3</sup> Choline is an important factor for human health, with an adequate intake for adults of 425 mg and 550 mg per day being recommended for women and men respectively in the US.<sup>4</sup> Choline depletion resulting from a choline-deficient diet

**Keywords:** trimethylaminuria (TMAU), cardiovascular disease (CVD), TMA, TMAO, thermoplasmata-related methanogens, archaeobiotics, choline, L-carnitine, vaginosis, *Methanomassiliicoccus* spp

\*Corresponding author: Jean-François Brugère; Email: jf.brugere@udamail.fr

Submitted: 04/15/2013; Revised: 10/01/2013;

Accepted: 10/08/2013;

Published Online: 10/31/2013

<http://dx.doi.org/10.4161/gmic.26749>

causes clinical consequences such as non-alcoholic fatty-liver disease (NAFLD) and muscle damage.<sup>5,6</sup> Dietary lecithin (phosphatidylcholine, PC) also feeds into plasma TMAO levels by gut conversion into choline (and then to TMA), as shown by mouse studies.<sup>7</sup> Thus, limiting intake of choline or its precursors to reduce TMA production is not feasible. However such a reduction of TMA levels is highly desirable because TMA is related to at least two disorders.

### TMA is Implicated in Disease

Trimethylaminuria (TMAU) is caused by a metabolic anomaly whereby sufferers emit a pervasive fishy odor caused by the excretion of TMA in their breath, sweat and urine. As TMAU is still under-recognized and often goes undiagnosed, those affected often suffer from psychological problems and social stress.<sup>8</sup> Although no large prevalence studies are currently available, people with compromised ability to N-oxidize trimethylamine into the odorless TMAO are probably common.<sup>9</sup> TMAU can be classified into a primary genetic form and an acquired (or secondary) form. The genetic form is now well characterized with approximately 15% of single nucleotide polymorphisms (SNPs) in the *FMO3* gene (encoding a flavin-containing monooxygenase) associated with either lack of or diminished enzymatic oxidation in the liver of TMA into TMAO.<sup>9</sup> In the healthy individual, more than 95% of TMA is oxidized into TMAO in this way, while genetic defects in *FMO3* result in accumulation of TMA in blood, and then in breath, sweat and urine (Fig. 1, pathway A). Acquired TMAU, sometimes transient, includes conditions featuring elevated urinary levels of TMA due to dietary and hormonal factors in combination with enzymatic activity and metabolism in the gut. Important examples include transient menstrual TMAU, overload of dietary precursors of TMA (see above), and impaired hepatic function.<sup>10</sup> Treatment relies mainly on an empirical dietary approach, or several complementary therapeutic adjuncts, including soaps which limit the volatility of TMA from the skin, together with brief courses of antimicrobials (neomycin, metronidazole)

which are not always effective.<sup>10</sup> This last approach limits the bacterial load, and in extenso, gut conversion of TMAO, choline and derivatives into TMA.

Gut microbiota metabolism of phosphatidylcholine (PC) and L-carnitine can promote atherosclerosis.<sup>2,7,11</sup> This is because their plasma metabolites (choline, betaine and TMAO) are risk factors for cardiovascular disease (CVD) in humans. Experimentally, dietary choline supplementation has been shown in a seminal paper to significantly enhance aortic lesion area in mice genetically prone to atherosclerosis, as did supplementation of the diet with TMAO.<sup>7</sup> Furthermore, gut microbiota depletion by antibiotics led to inhibition of the dietary choline-induced cardiovascular effects. Mice with depleted gut microbiota did not experience increased TMAO plasma levels<sup>7</sup> (Fig. 1, pathway B) nor did humans.<sup>11</sup> Human gut microbiota is also required to form TMAO from L-carnitine, and plasma TMAO is likely the primary driver of cardiovascular risks rather than L-carnitine itself.<sup>2</sup>


A fishy odor frequently occurs in bacterial vaginosis.<sup>12</sup> It is important to stress that, in this instance, TMA does not originate from urine, but rather from in situ production of TMA by the vaginal microbiota through a process similar to that which occurs in the gut.

### Hypothesis: That TMA Could Be Depleted as it is Synthesized

We were prompted by recent microbiology studies to ask if TMA could be depleted in vivo by bioconversion into a molecule with no undesirable properties. The underlying biochemistry was described more than 35 y ago in the rumen of cows, where choline is metabolized via TMA and onwards into methane.<sup>13</sup> Rumen microbes like the methanogenic archaea *Methanosarcina barkeri* metabolize methyl compounds, including TMA, to methane for growth.<sup>14</sup> A distinct archaeal group (the Rumen Cluster C; RCC) was subsequently identified as being putatively able to convert methyl compounds including TMA and methylamines into methane. Half the human subjects examined to date also produce methane, usually by harboring the methanogen *Methanobrevibacter*

*smithii*, or less frequently, *Methanosphaera stadtmanae* (reviewed in ref. 15). It is also noteworthy that *M. smithii* has been identified in vaginal samples.<sup>15</sup> We thus asked if methylotrophic methanogens would be able to survive and deplete TMA in the human gut (or in the vagina), a notion that would be greatly supported if methanogens that naturally occur in the human gut could metabolize TMA.

The trigger for proposing this hypothesis now is that, until recently, only the methanogens *M. smithii* and *M. stadtmanae* were recognized as residents of the human gut. They belong to the order Methanobacteriales, one of the six known orders of methanogens. However in 2008, the existence of a seventh order inhabiting the human gut (hereafter referred to as the Mx-lineage) was proposed,<sup>16</sup> particularly common in older subjects,<sup>17</sup> and *Methanomassiliicoccus luminyensis*, the first and unique isolated member of this order metabolizes methanol with hydrogen for methane production.<sup>18</sup> Genomic sequences of three of these unusual methanogens are currently available, that of *M. luminyensis*,<sup>19</sup> and genome sequences that we recently determined for “*Candidatus Methanomethylophilus alvus*”<sup>20</sup> and “*Candidatus Methanomassiliicoccus intestinalis*”<sup>21</sup> grown in an enrichment culture from human stool. “*Ca. Methanomethylophilus alvus*” belongs to the RCC cluster that was recently highlighted for its putative involvement in TMA consumption in the rumen,<sup>23</sup> while *M. luminyensis* and “*Ca. Methanomassiliicoccus intestinalis*” are phylogenetically closely related to the RCC cluster.<sup>24</sup> These three genomes have all of the genes necessary for reduction of methanol using hydrogen, but also for the reduction of tri- di- and monomethylamine with hydrogen (Fig. 2), while the two other possible pathways for methanogenesis (CO<sub>2</sub> reduction with H<sub>2</sub> and acetoclastic methanogenesis) are genetically incomplete. Whereas the capacity of these strains to reduce methanol with hydrogen was previously demonstrated, their ability to grow on TMA with hydrogen has not been investigated. We therefore tested this hypothesis on the sole strain of this group available in pure culture, *M. luminyensis* B10, and thus confirmed for the first time



**Figure 1.** Origin and fate of TMA in the human gut, and the Archaeobiotics concept. Gut microbiota synthesis of TMA is realized from TMAO, choline, PC and L-carnitine. The TMA is then absorbed and goes to the liver, routes (A) or (B). In the case of route (A), a partial or total defect in FMO3-oxidation into TMAO leads to increased level and diffusion of TMA in breath, urine and sweat. When FMO3 liver oxidation is functional (B), the increase of TMAO in blood is associated with atherosclerosis.<sup>2,7,11</sup> Therefore, converting TMA directly in the gut using Archaeobiotics belonging to the seventh methanogenic order, naturally-occurring in the gut, route (C) should be envisaged. Interestingly, these archaea are only able to perform methanogenesis using methyl compounds (see Fig. 2), because the two other pathways are absent (CO<sub>2</sub> reduction with H<sub>2</sub> and aceticlastic pathway): this would increase the efficiency of TMA conversion.


that this strain is able to grow on TMA with hydrogen (Fig. 3). *M. luminyensis* B10 was also able to grow on the by-products of TMA catabolism, dimethylamine and then monomethylamine, both being used with hydrogen for methanogenesis (data not shown). These results are in agreement with the annotated genetic complement of this strain and strongly suggest the same metabolic capacity in “*Ca. M. alvus*” and “*Ca. M. intestinalis*” because they harbor orthologs of all the relevant *M. luminyensis* genes involved in these pathways. Collectively this suggests to us that natural methanogenic inhabitants of

the human gut will be able to metabolize TMA, and could deplete this metabolite as it is formed by bacterial elements of the microbiota.

### Proposal: The Archaeobiotics Concept

Drawing together these ecological and biochemical strands, we now propose the Archaeobiotics Concept viz. the intestinal application of specific archaeal strains for the treatment of human diseases including cardiovascular disease and TMAU (Fig. 1). The fishy-odor aspect of some

vaginal conditions could be addressed using pessary delivery. Focusing on CVD and TMAU, there are several advantages. Methane, the TMA-metabolite produced by these archaea is usually considered biologically inert in humans. The pathway and relevant coding sequences for catalyzing TMA conversion to methane by “*Ca. Methanomethylophilus alvus*” and “*Ca. Methanomassiliicoccus intestinalis*” based on annotation of the genomes we recently sequenced is shown in Figure 2; a similar pathway is inferred from the *M. luminyensis* genome. As for other methanogenic archaea, it includes a



**Figure 2.** Metabolic pathway of TMA conversion deduced from genomic data from species of the 7th order of methanogens. The pathway of TMA conversion involves a trimethylamine:corrinoid methyltransferase (MttB) which contains a pyrrolysine, an unusual amino-acid. The synthesis of this amino-acid and its transfer on a tRNA<sup>Pyl</sup> to form a pyrrolysyl-tRNA<sup>Pyl</sup>, and its insertion during the translation of *mttB* are presented in the green part. The pathway of trimethylamine (TMA) conversion, involving the pyrrolysine containing MttB, is presented in the orange part. Steps in the pathway between named compounds are indicated by blue arrows. The purple arrow indicates the transcription and translation of *mttB*. The dotted arrows indicate steps that are not detailed for the clarity of the scheme. Recognized enzyme names (see KEGG map 00680) are provided, flanked by the corresponding protein accession numbers in the “*Ca. M. alvus*” (blue) and “*Ca. M. intestinalis*” (red) genomes. Only accession numbers of enzymes specifically involved in the depletion of TMA are presented, some being shared with dimethylamine (DMA) and monomethylamine (MMA) depletion (not shown here).

methyltransferase MttB with a predicted pyrrolysine residue,<sup>22</sup> which is associated with its synthetic machinery. The fact that these strains were isolated from humans might facilitate regulatory compliance. *M. luminyensis* like *M. smithii* is susceptible to bacitracin, metronidazole, ordinazole and squalamine,<sup>25</sup> providing the reassurance of contingency eradication. An alternative to administering live cells might be therapeutic administration of the relevant enzymes (Fig. 2), all of which are also encoded by *M. luminyensis*.

Paralleling the experience with the development of probiotic bacterial strains for human consumption, it has been shown that particular archaeal methanogen strains are likely to be more suitable for clinical use

in this application. For example, we established that some strains seem to be evolutionarily more adapted to the human gut ecosystem, harboring genes predicted to confer resistance to bile salts (bile salt hydrolases occur in “*Ca. Methanomethylophilus alvus*” species, but not in *M. luminyensis* nor in “*Ca. Methanomassiliicoccus intestinalis*”). Consideration of factors other than TMA depletion alone is thus appropriate (see also below).


### Challenges for Therapeutic Usage for Archaea to Deplete TMA

Technological hurdles to overcome will include establishing the best way to deliver

oxygen-sensitive microorganisms into the human gut, and whether or not this administration will lead to cell levels that are sufficient to allow clinically relevant TMA depletion. Continuous administrations may be necessary if subjects do not become colonized. There are additional biological challenges; will these strains be able to effectively deplete TMA in the gut? Are the ensuing risk reductions for cardiovascular disease achieved, as they were by antibiotic therapy to limit TMA production?<sup>11</sup> It is also important to concede that archaeal abundance is not uniformly associated with increased health in mammals; for example, *Bacteroides thetaiotaomicron*-*M. smithii* co-colonization of germ-free mice increased host adiposity.<sup>26</sup> Archaea in general appear to be more abundant in

**Figure 3.** Growth of *Methanomassiliicoccus luminyensis* strain B10 on trimethylamine (TMA) and hydrogen ( $H_2$ ). (A) The growth of *M. luminyensis* B10 on methanol (M, 50 mM) and  $H_2$  (previously described by Dridi et al.<sup>19</sup>) was used as a positive control for growth and maximal cell density. (B) We now show that *M. luminyensis* B10 also grows on TMA (15 mM) in the presence of  $H_2$  (filled diamonds) but not on TMA in the absence of  $H_2$  (open diamonds). Accordingly, TMA is depleted in presence of  $H_2$  (filled triangles) and not in the absence of  $H_2$  (open triangles). The composition of the gas phase was measured at the end of the experiment (C) and revealed the production of methane ( $CH_4$ ) and the depletion of  $H_2$  in presence of either M or TMA. No  $CH_4$  was produced in the absence of  $H_2$  and  $CH_4$  was not produced nor  $H_2$  depleted in the absence of the methylated compound substrate (M or TMA). *M. luminyensis* B10 was obtained from DSMZ (DSM No. 25720) and cultivated in DSMZ medium 119, with rumen fluid replacing the sludge fluid. When  $H_2$  was present, initial atmosphere composition was:  $N_2/H_2/CO_2$  (55:35:10), and in absence of  $H_2$ :  $N_2/CO_2$  (75:25). Growth in each condition was performed in triplicate. Trimethylamine concentration was measured as described by Krätzer et al.<sup>30</sup> OD, Optical Density.

native Africans than in African Americans, likely indicating dietary influences on the microbiota.<sup>27,28</sup> Diet could conceivably be controlled or modulated to promote methanogen levels in recipients of archaeobiotics. We cannot currently explain why the Mx-lineage of Archaea appears from the literature to be more abundant in older subjects, so we are determining fecal archaeal levels in the ELDERMET cohort subjects, that are well phenotyped for diet, health and bacterial microbiota.<sup>29</sup> A possible factor is the trophic interaction with other microbiota elements, and that a more favorable ecology emerges in some older subjects. This raises the provocative issue of what the effects of introducing a methanogenic archaea might be on the microbiota of the recipient. The Mx-methanogens are very likely not acetoclastic but would compete with other hydrogenotrophic archaea and bacteria, so the downstream effects on microbiome, metabolome and host physiology would need detailed exploration in vitro, in animal models, and ultimately in humans. Although the application of archaeobiotics as proposed here faces significant challenges, we contend that the anticipated benefit of their use to mitigate the life-time risk of CVD and to treat TMAU warrants a concerted research effort in this area by the scientific community.



## Disclosure of Potential Conflict of Interests

No potential conflict of interest was disclosed.

## Acknowledgments

This work was supported by two PhD. Scholarship supports, one from the French “Ministère de l’Enseignement Supérieur et de la Recherche” to NG,

the other of the European Union (UE) and the Auvergne Council to WT (FEDER). PWOT was supported by Science Foundation Ireland through a Principal Investigator award and by an FHRI award to the ELDERMET project by the Dept. Agriculture, Fisheries and Marine of the Government of Ireland. We thank F Shanahan and EMM Quigley

for helpful critique of the manuscript. In Memoriam to Albert Jacquard (b. 12-23-1925, † 09-11-2013), geneticist and great humanist.

## References

- Zhang AQ, Mitchell SC, Smith RL. Dietary precursors of trimethylamine in man: a pilot study. *Food Chem Toxicol* 1999; 37:515-20; PMID:10456680; [http://dx.doi.org/10.1016/S0278-6915\(99\)00028-9](http://dx.doi.org/10.1016/S0278-6915(99)00028-9)
- Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, Britt EB, Fu X, Wu Y, Li L, et al. Intestinal microbiota metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. *Nat Med* 2013; 19:576-85; PMID:23563705; <http://dx.doi.org/10.1038/nm.3145>
- Craciun S, Balskus EP. Microbial conversion of choline to trimethylamine requires a glycol radical enzyme. *Proc Natl Acad Sci U S A* 2012; 109:21307-12; PMID:23151509; <http://dx.doi.org/10.1073/pnas.1215689109>
- Institute of Medicine (U.S.). Standing Committee on the Scientific Evaluation of Dietary Reference Intakes., Institute of Medicine (U.S.). Panel on Folate Other B Vitamins and Choline., Institute of Medicine (U.S.). Subcommittee on Upper Reference Levels of Nutrients. Dietary reference intakes for thiamin, riboflavin, niacin, vitamin B<sub>6</sub>, folate, vitamin B<sub>12</sub>, pantothenic acid, biotin, and choline. Washington, D.C.: National Academy Press, 1998.
- Buchman AL, Dubin MD, Moukarzel AA, Jenden DJ, Roch M, Rice KM, Gornbein J, Ament ME. Choline deficiency: a cause of hepatic steatosis during parenteral nutrition that can be reversed with intravenous choline supplementation. *Hepatology* 1995; 22:1399-403; PMID:7590654
- Spencer MD, Hamp TJ, Reid RW, Fischer LM, Zeisel SH, Fodor AA. Association between composition of the human gastrointestinal microbiome and development of fatty liver with choline deficiency. *Gastroenterology* 2011; 140:976-86; PMID:21129376; <http://dx.doi.org/10.1053/j.gastro.2010.11.049>
- Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, Feldstein AE, Britt EB, Fu X, Chung YM, et al. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* 2011; 472:57-63; PMID:21475195; <http://dx.doi.org/10.1038/nature09922>
- Ayesh R, Mitchell SC, Zhang A, Smith RL. The fish odour syndrome: biochemical, familial, and clinical aspects. *BMJ* 1993; 307:655-7; PMID:8401051; <http://dx.doi.org/10.1136/bmj.307.6905.655>
- D’Angelo R, Esposito T, Calabrò M, Rinaldi C, Robledo R, Varriale B, Sidoti A. FMO3 allelic variants in Sicilian and Sardinian populations: trimethylaminuria and absence of fish-like body odor. *Gene* 2013; 515:410-5; PMID:23266626; <http://dx.doi.org/10.1016/j.gene.2012.12.047>
- Mackay RJ, McEntyre CJ, Henderson C, Lever M, George PM. Trimethylaminuria: causes and diagnosis of a socially distressing condition. *Clin Biochem Rev* 2011; 32:33-43; PMID:21451776
- Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, Wu Y, Hazen SL. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med* 2013; 368:1575-84; PMID:23614584; <http://dx.doi.org/10.1056/NEJMoa1109400>
- Wolrath H, Ståhlbom B, Hallén A, Forsum U. Trimethylamine and trimethylamine oxide levels in normal women and women with bacterial vaginosis reflect a local metabolism in vaginal secretion as compared to urine. *APMIS* 2005; 113:513-6; PMID:16086821; [http://dx.doi.org/10.1111/j.1600-0463.2005.apm\\_175.x](http://dx.doi.org/10.1111/j.1600-0463.2005.apm_175.x)
- Neill AR, Grime DW, Dawson RM. Conversion of choline methyl groups through trimethylamine into methane in the rumen. *Biochem J* 1978; 170:529-35; PMID:646798
- Hippe H, Caspari D, Fiebig K, Gottschalk G. Utilization of trimethylamine and other N-methyl compounds for growth and methane formation by *Methanosarcina barkeri*. *Proc Natl Acad Sci U S A* 1979; 76:494-8; PMID:284366; <http://dx.doi.org/10.1073/pnas.76.1.494>
- Conway de Macario E, Macario AJ. Methanogenic archaea in health and disease: a novel paradigm of microbial pathogenesis. *Int J Med Microbiol* 2009; 299:99-108; PMID:18757236; <http://dx.doi.org/10.1016/j.ijmm.2008.06.011>
- Mihajlovski A, Alric M, Brugère JF. A putative new order of methanogenic Archaea inhabiting the human gut, as revealed by molecular analyses of the *mcrA* gene. *Res Microbiol* 2008; 159:516-21; PMID:18644435; <http://dx.doi.org/10.1016/j.resmic.2008.06.007>
- Mihajlovski A, Doré J, Levenez F, Alric M, Brugère J-F. Molecular evaluation of the human gut methanogenic archaeal microbiota reveals an age-associated increase of the diversity. *Environ Microbiol Rep* 2010; 2:272-80; PMID:23766078; <http://dx.doi.org/10.1111/j.1758-2229.2009.00116.x>
- Dridi B, Fardeau ML, Ollivier B, Raoult D, Drancourt M. *Methanomassiliicoccus luminyensis* gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces. *Int J Syst Evol Microbiol* 2012; 62:1902-7; PMID:22859731; <http://dx.doi.org/10.1099/ijs.0.033712-0>
- Gorlas A, Robert C, Gimenez G, Drancourt M, Raoult D. Complete genome sequence of *Methanomassiliicoccus luminyensis*, the largest genome of a human-associated Archaea species. *J Bacteriol* 2012; 194:4745; PMID:22887657; <http://dx.doi.org/10.1128/JB.00956-12>
- Borrel G, Harris HM, Tottey W, Mihajlovski A, Parisot N, Peyretailade E, Peyret P, Gribaldo S, O’Toole PW, Brugère JF. Genome sequence of “*Candidatus Methanomethylphilus alvus*” Mx1201, a methanogenic archaeon from the human gut belonging to a seventh order of methanogens. *J Bacteriol* 2012; 194:6944-5; PMID:23209209; <http://dx.doi.org/10.1128/JB.01867-12>
- Borrel G, Harris HM, Parisot N, Gaci N, Tottey W, Mihajlovski A, et al. Genome Sequence of “*Candidatus Methanomassiliicoccus intestinalis*” Isoire-Mx1, a Third Thermoplasmatales-Related Methanogenic Archaeon from Human Faeces. *Genome Announc* 2013; 1.
- Krzycki JA. The direct genetic encoding of pyrrolysine. *Curr Opin Microbiol* 2005; 8:706-12; PMID:16256420; <http://dx.doi.org/10.1016/j.mib.2005.10.009>
- Poulsen M, Schwab C, Jensen BB, Engberg RM, Spang A, Canibe N, Højberg O, Milinovich G, Fragner L, Schleper C, et al. Methylophilic methanogenic Thermoplasmata implicated in reduced methane emissions from bovine rumen. *Nat Commun* 2013; 4:1428; PMID:23385573; <http://dx.doi.org/10.1038/ncomms2432>
- Borrel G, O’Toole PW, Harris HM, Peyret P, Brugère JF, Gribaldo S. Phylogenomic data support a seventh order of methylophilic methanogens and provide insights into the evolution of methanogenesis. *Genome Biol Evol* 2013; 5:1769-80; PMID:23985970; <http://dx.doi.org/10.1093/gbe/evt128>
- Dridi B, Fardeau ML, Ollivier B, Raoult D, Drancourt M. The antimicrobial resistance pattern of cultured human methanogens reflects the unique phylogenetic position of archaea. *J Antimicrob Chemother* 2011; 66:2038-44; PMID:21680581; <http://dx.doi.org/10.1093/jac/dkr251>
- Samuel BS, Gordon JI. A humanized gnotobiotic mouse model of host-archaeal-bacterial mutualism. *Proc Natl Acad Sci U S A* 2006; 103:10011-6; PMID:16782812; <http://dx.doi.org/10.1073/pnas.0602187103>
- O’Keefe SJ, Chung D, Mahmoud N, Sepulveda AR, Manafé M, Arch J, Adada H, van der Merwe T. Why do African Americans get more colon cancer than Native Americans? *J Nutr* 2007; 137(Suppl):175S-82S; PMID:17182822
- Nava GM, Carbonero F, Ou J, Benefiel AC, O’Keefe SJ, Gaskins HR. Hydrogenotrophic microbiota distinguish native Africans from African and European Americans. *Environ Microbiol Rep* 2012; 4:307-15; PMID:23760794; <http://dx.doi.org/10.1111/j.1758-2229.2012.00334.x>
- Claesson MJ, Jeffery IB, Conde S, Power SE, O’Connor EM, Cusack S, Harris HM, Coakley M, Lakshminarayanan B, O’Sullivan O, et al. Gut microbiota composition correlates with diet and health in the elderly. *Nature* 2012; 488:178-84; PMID:22797518; <http://dx.doi.org/10.1038/nature11319>
- Krätzer C, Carini P, Hovey R, Deppenmeier U. Transcriptional profiling of methyltransferase genes during growth of *Methanosarcina mazei* on trimethylamine. *J Bacteriol* 2009; 191:5108-15; PMID:19525341; <http://dx.doi.org/10.1128/JB.00420-09>