



Published in final edited form as:

J Immunol. 2014 June 15; 192(12): 5490–5498. doi:10.4049/jimmunol.1400722.

Antigen-dependent versus -independent activation of iNKT cells during infection

Keli L. Holzapfel^{*}, Aaron J. Tyznik[†], Mitchell Kronenberg[†], and Kristin A. Hogquist^{*}

^{*}Department of Laboratory Medicine and Pathology, Center for Immunology, University of Minnesota, Minneapolis, MN 55455, USA

[†]Division of Developmental Immunology, La Jolla Institute for Allergy and Immunology, La Jolla, CA 92037, USA

Abstract

CD1d-reactive invariant natural killer T cells (iNKT) play a vital role in determining the characteristics of immune responses to infectious agents. Previous reports suggest that iNKT cell activation during infection can be: 1) solely driven by cytokines from innate immune cells; 2) require microbial-antigen; or 3) require self-antigen. In this study, we examined the role of antigen receptor stimulation in iNKT cells during several bacterial and viral infections. To test for antigen receptor signaling, Nur77^{gfp} BAC transgenic mice, which upregulate GFP in response to antigen receptor but not inflammatory signals, were analyzed. iNKT cells in the reporter mice infected with mouse cytomegalovirus (MCMV) produced IFN γ but did not upregulate GFP, consistent with their reported CD1d-independent activation. However, two bacteria known to produce lipid antigens for iNKT cells induced GFP expression and cytokine production. In contrast, although *Salmonella typhimurium* (*S. typhimurium*) was proposed to induce the presentation of a self-lipid, iNKT cells produced IFN γ , but did not upregulate GFP after infection *in vivo*. Even in CD1d-deficient hosts, iNKT cells were still able to produce IFN γ after *S. typhimurium* infection. Furthermore, while it has been proposed that endogenous lipid presentation is a result of Toll-like receptor (TLR) stimulation of antigen presenting cells, injection of different TLR agonists led to iNKT cell IFN γ but not increased GFP expression. These data indicate that robust iNKT cell responses to bacteria as well as viruses can be obtained in the absence of antigenic stimulation.

Introduction

CD1d-reactive invariant Natural Killer T cells (iNKT cells) are a distinct lineage of $\alpha\beta$ T lymphocytes with an invariant T cell antigen receptor (TCR) composed in mice of the α -chain variable region 14 (V α 14) and the α -chain joining region 18 (J α 18) paired with a restricted subset of TCR- β chains (1). As a result of this TCR expression, iNKT cells are able to recognize several types of glycolipid antigens in the context of CD1d, a non-classical major histocompatibility class I (MHC-I)-like antigen presenting molecule. Upon glycolipid presentation, and most famously with the strong agonist α -galactosylceramide (α GalCer), iNKT cells are able to rapidly produce cytokines such as IFN- γ and IL-4 (2). This rapid

cytokine production contributes to the recruitment and activation of other cell types during an immune response, which has been shown to influence a variety of diseases, including cancer (3), autoimmunity (4, 5), and pathogenic infections (6). Therefore, insight into their activation is crucial for the general understanding of how iNKT cells contribute to immune responses.

One area that remains to be fully elucidated is the extent to which antigen recognition by the iNKT TCR contributes to the activation of iNKT cells during various infections. Previous reports have shown that iNKT cells can be activated through the TCR by certain infectious agents that produce glycolipid antigens (7-11). For example, iNKT cells are activated by glycosylated diacylglycerol antigens from *Streptococcus pneumoniae* (*S. pneumoniae*) in a CD1d-dependent manner (10). Furthermore, iNKT deficient mice infected with *S. pneumoniae* have significantly reduced survival compared to infected wild type mice (9). In addition, *Sphingomonas paucimobilis* (*S. paucimobilis*), which is a Gram-negative bacteria that does not express LPS, produces glycosphingolipids (GSLs) with an α -linked glucuronic or galacturonic acid that are able to activate iNKT cells (7, 8, 12). More specifically, it was shown that these GSLs activate iNKT cell proliferation and cytokine secretion, which was dependent on CD1d (11).

However, iNKT cells are able to rapidly produce cytokines during a variety of infections where the pathogen is not known to produce an exogenous stimulating lipid (11, 13), raising the question of how iNKT cells are activated to rapidly produce cytokines. It was previously proposed that during these types of infections, iNKT cell activation occurs partly in response to the presentation of an endogenous lipid capable of stimulating the iNKT TCR (11, 13, 14). During these types of infections, iNKT cell responses are induced, at least in part, as a result of stimulation of TLRs on antigen presenting cells (APCs), which not only causes the secretion of cytokines, such as IL-12 or type I IFN, that participate in the iNKT cell activation process, but which also causes increased synthesis and presentation by the APCs of a stimulating endogenous lipid (14-16). This process is sometimes referred to as indirect activation, because rather than providing the antigen, the bacteria are inducing the synthesis of both a stimulatory antigen and cytokines by host APCs. Although the full spectrum of endogenous stimulatory lipids remains to be identified, and their relative importance remains to be conclusively assessed, two GSLs, isoglobotrihexosylceramide (iGb3) and β -D-glucopyranosylceramide (β GlcCer) are known to be self-lipids (14, 17, 18).

Yet it is clear that in some situations T lymphocytes, such as $\gamma\delta$ T cells and CD4 and CD8 memory T cells, become activated independently of antigen receptor stimulation, sometimes referred to as non-cognate activation (19-23). Therefore, it is possible that certain microorganisms are able to activate iNKT cells independently of their TCR (24, 25). This antigen-independent model has been proposed to explain iNKT cell activation during infection with mouse cytomegalovirus (MCMV) (25). Upon infection with MCMV, iNKT cells demonstrated an activated phenotype and produced IFN γ . This activation resulted from APC secretion of IL-12 and IL-18, and there was little or no effect on IFN γ production when CD1d was blocked or deleted (25, 26).

Further understanding of the indirect activation mode, whereby iNKT cells are activated by endogenous lipids, is important as it may be occurring in diseases such as type I diabetes and asthma (27-29). Greater mechanistic understanding might also provide a profound way to control iNKT cells during infections. However, one challenge to greater understanding is that it remains difficult to distinguish TCR-mediated from non-TCR mediated activation. The activation parameters typically used in the study of iNKT cells are CD69 upregulation, cytokine production, and reduced recovery of iNKT cells using tetramers, presumed to be secondary to TCR internalization. All of these features, however, can also be a consequence of non-TCR mediated activation. In this study, we examined the role of antigen receptor stimulation in iNKT cell activation during infections *in vivo* using Nur77^{gfp} BAC transgenic mice, which upregulate GFP in response to antigen receptor but not inflammatory signals (30). Such mice were infected and examined for cytokine production and GFP expression as indicators of iNKT cell activation and iNKT TCR stimulation, respectively.

Materials and Methods

Mice

Nur77^{gfp} B6 mice were previously described (30). B6 (C57BL/6NCr) and B6.SJL (B6-LY.5/Cr) mice were obtained from the National Cancer Institute. CD1d^{-/-} B6 (B6.129S6-Cd1d1/Cd1d2tm1Spb/J) mice and V α 14-transgenic B6 (C57BL/6-Tg(Cd4-TcraDN32D3)1Alben/J) mice were obtained from The Jackson Laboratory. Fabry mice deficient for the enzyme α -galactosidase A (B6;129-Gla^{tm1Kul}/J) were obtained from F. Winau (Harvard Medical School). KN2 mice on the B6 background were obtained from M. Mohrs (Trudeau Institute). KN2 Nur77^{gfp} B6 mice were generated by the crossing of KN2 and Nur77^{gfp} mice. Nur77^{gfp} V α 14 TCR transgenic B6 mice were generated by the crossing of V α 14 TCR transgenic B6 and Nur77^{gfp} B6 mice. All animal experimentation was approved by and performed according to the guidelines from the Institutional Animal Care and Use Committees at the University of Minnesota or at the La Jolla Institute for Allergy & Immunology.

Flow cytometry, antibodies, and intracellular cytokine staining

Single-cell suspensions were prepared from spleens, and hepatic mononuclear cells were separated by Percoll gradient centrifugation. All cells were resuspended in staining buffer (PBS with 5% FBS). Antibodies to surface markers were added, and cells were stained on ice for 20 min. Antibodies used were from eBioscience, BD, or BioLegend. For iNKT cell staining, CD1d monomers loaded with biotinylated PBS57 or unloaded monomers were obtained from the tetramer facility of the US National Institutes of Health. Cells were analyzed on an LSRII (Becton Dickinson) and data was processed with FlowJo software (TreeStar). A dump strategy (CD11b, F4/80, and CD8) was used to eliminate non-specific events.

For intracellular detection of cytokines, cells were stained for surface markers as described above, and then fixed and permeabilized using the Cytotfix/CytoPerm buffer (BD Biosciences) according to the manufacturer's protocol. Anti-IFN γ (eBioscience) was added

and the cells were stained on ice for 30 min. Cells were washed twice and then resuspended in staining buffer before analysis on an LSRII.

iNKT cell purification

For iNKT cell adoptive transfer experiments, a single-cell suspension was prepared from Nur77^{gfp} V α 14-transgenic mice spleens, and from hepatic mononuclear cells separated by Percoll gradient centrifugation. This single-cell suspension was depleted of B cells and CD8 T cells using anti-CD19 and anti-CD8 α conjugated to magnetic beads and MACS columns (Miltenyi Biotec), according to manufacturer's protocol. The percentage of iNKT cells in the enriched fraction was ~30%, and the equivalent of $\sim 3 \times 10^6$ iNKT cells was adoptively transferred per mouse.

In vivo Infections

Mouse cytomegalovirus (MCMV-K181) was obtained from C. Benedict (La Jolla Institute for Allergy & Immunology). Mice were infected intra-peritoneally with 1×10^5 PFU diluted in 300 μ l PBS. *S. paucimobilis* was obtained from M. Kronenberg (La Jolla Institute for Allergy & Immunology). *S. paucimobilis* cultured in Tryptic Soy Broth (BD) at 37°C were collected at a mid-log phase and washed with PBS. Mice were inoculated intravenously with $\sim 1 \times 10^9$ colony forming units diluted in 200 μ l PBS. *Streptococcus pneumoniae* URF918 (clinical isolate, serotype 3) was obtained from M. Kronenberg (La Jolla Institute for Allergy & Immunology). *S. pneumoniae* cultured in Todd-Hewitt broth (BD) at 37°C were collected at a mid-log phase and then washed with PBS. Mice were inoculated intravenously with $\sim 1 \times 10^7$ colony-forming units diluted in 200 μ l PBS. *Salmonella typhimurium* (SL1344) was obtained from S. McSorley (University of California, Davis). *S. typhimurium* cultured in Todd-Hewitt broth (BD) at 37°C were collected at a mid-log phase and then washed with PBS. Mice were inoculated intravenously with $\sim 1 \times 10^6$ colony-forming units diluted in 200 μ l PBS. As a control for all infections, 2 μ g of α GalCer diluted in 200 μ l of PBS was injected intravenously, and spleen and liver were harvested 2-4 hours later.

TLR agonists

Mice were injected intravenously in a final volume of 200 μ l with 50 μ g of *E. coli* derived LPS diluted in PBS, or with 10 μ g of ODN 1826 diluted in endotoxin free water. Spleen and liver were analyzed at the indicated timepoints after injection.

Lipid-pulsed bone marrow DCs

Bone marrow cells from femurs of mice were cultured for 7 days (5×10^6 cells/well) at 37°C in 6-well cell culture dishes with complete RPMI medium in the presence of recombinant murine GM-CSF (50 ng/ml, PeproTech) and IL-4 (10 ng/ml, PeproTech). On day six, cells were pulsed with either 100 ng – 1 μ g/ml of α GalCer (KRN7000, Avanti Polar Lipids), 1 μ g/ml OCH (Alexis Biochemicals), 1 μ g/ml of β GlcCer (C24:1 Glucosyl(β) Ceramide (d18:1/24:1(15Z)), Avanti Polar Lipids), 1 μ g/ml of iGb3 provided by D. Zhou (MD Anderson) or 1 μ l/ml solvent (2:1 methanol:chloroform) for 12-15 hours at 37 C. BMDCs were intravenously injected into Nur77^{gfp} mice (0.5 – 1.0×10^6 cells/mouse) and endogenous splenic iNKT cells were analyzed 16 h later.

Mixed-bone marrow chimeras

Total bone marrow cells were prepared from the femurs of Nur77^{gfp} B6.SJL (CD45.1⁺CD45.1⁺), B6 (CD45.2⁺CD45.2⁺), or Fabry B6 (CD45.2⁺CD45.2⁺) donor mice, and samples were depleted of mature T cells with anti-Thy1.2 (30-H12; Biolegend) and complement. Recipient mice (CD45.2⁺ CD45.1⁺) were lethally irradiated (1,000 rads) and received 1×10^7 adult bone marrow cells. Chimeras were analyzed 8 weeks after transplantation.

Statistical analysis

Prism software (GraphPad) was used for statistical analysis. Unpaired two-tailed *t* tests were used for data analysis.

Results

iNKT cells from the Nur77^{gfp} reporter mouse upregulate GFP in response to antigenic lipids

Nur77^{gfp} BAC transgenic mice encode GFP in the locus of Nur77 (*Nr4a1*), an immediate early gene downstream of the TCR. Previous work showed that T lymphocytes from these mice expressed GFP only after antigen receptor stimulation, and not after inflammatory signals, cytokines, or stimuli that promote cell survival or trafficking (30). iNKT cells in this reporter mouse exhibited a very low level of GFP in the periphery at steady state, suggesting that iNKT cells are not continuously receiving stimulation through their TCR (30). However, when injected with potent TCR agonist α GalCer, iNKT cells greatly increased expression of GFP (30). Therefore, we used the Nur77^{gfp} mouse to examine iNKT TCR activation *in vivo*.

Using OT-I TCR transgenic CD8 T lymphocytes, the Nur77 reporter was shown to respond to very weak TCR stimuli, even those that support positive selection and are incapable of inducing proliferation of mature T cells. Therefore we sought to test the sensitivity of GFP upregulation after iNKT cell TCR stimulation. Most antigenic lipids have lower affinities for the iNKT TCR compared to α GalCer (31, 32), and we tested OCH, β -D-glucopyranosylceramide (β GlcCer), and isoglobotrihexosylceramide (iGb3), in addition to α GalCer. OCH is an analog of α GalCer, and is reported to also strongly activate iNKT cells, albeit with a slightly weaker affinity for the iNKT TCR than α GalCer (31, 33). iGb3, and more recently β GlcCer, were identified as putative endogenous self-lipids capable of activating iNKT cells through the TCR (17, 18). iGb3 is a weak activator of iNKT cells with a much lower affinity for the iNKT TCR than α GalCer, in the low μ M range compared to as low as 1 nM for complexes of α GalCer bound to CD1d (34, 35). β GlcCer, when loaded into CD1d tetramers, is at least capable of binding to a proportion of iNKT cells, suggesting that it is a stronger ligand than iGb3, although an affinity measurement per se has not been reported (17).

We tested the ability of these lipids to activate the iNKT TCR by pulsing bone marrow derived dendritic cells (BMDC), and then intravenously injecting them into Nur77^{gfp} mice. After 18 hours, iNKT cells were isolated and examined directly *ex vivo* by flow cytometry

for GFP upregulation. All four lipids induced an upregulation of GFP as compared to iNKT cells from a control mouse injected with unpulsed BMDC (Fig. 1). α GalCer consistently induced the strongest increase, although OCH induced only slightly less. Of the putative self-lipids, β GlcCer induced a higher upregulation of GFP than iGb3. CD69 was also upregulated, showing the same order of expression (α GalCer>OCH> β Glc>iGb3) (Fig. 1), although its use as a marker of iNKT TCR stimulation during infections is limited by the fact that it can also be upregulated by inflammatory stimuli (30, 36). These results suggest that in the Nur77^{gfp} mouse, GFP expression is a sensitive tool for assessing iNKT TCR stimulation.

MCMV infection does not stimulate iNKT cells through their TCR

MCMV does not contain a stimulatory lipid that activates the iNKT TCR; however, MCMV has been shown to activate iNKT cells during infection *in vivo*, as demonstrated by increased CD25 and CD69 expression, and by IFN- γ production detected by intracellular cytokine staining as early as 24 hours after infection (25). This activation is thought to be predominantly a result of IL-12 secretion, and IL-18 and/or type I IFN secretion from APCs that were activated through TLR9 (26, 37). Importantly, there was little to no effect on iNKT cell IFN γ production in experiments where CD1d was blocked and/or deleted (25). Since CD1d-mediated antigen is not considered a key-player in the MCMV-induced activation of iNKT cells, we tested this putatively cytokine-mediated model of iNKT cell activation *in vivo* using the Nur77^{gfp} mouse. Similar to previous reports, iNKT cells from Nur77^{gfp} mice infected with MCMV showed a statistically significant increase in the percentage of hepatic iNKT cells that produce IFN γ 36 hours post infection (Fig. 2B). Interestingly, the level of GFP expression remained the same at all time points examined, suggesting that iNKT cells did not receive TCR stimulation during MCMV infection (Fig. 2A), and confirming that other inflammatory stimuli produced during infections do not cause GFP upregulation (30).

S. pneumoniae and *S. paucimobilis* stimulate iNKT cells through their TCR

We next infected Nur77^{gfp} mice with microbes that are known to produce lipid antigens that activate iNKT cells (7, 10). *S. pneumoniae* is an extracellular Gram-positive pathogen, which produces glycosylated diacylglycerol antigens (9, 10). These antigens are highly abundant in the bacteria, they can stimulate the iNKT TCR, and iNKT cells were shown to be important for clearance and host survival (10). *S. paucimobilis* is a Gram-negative bacteria that does not produce LPS, but does produce a GSL with an α -linked glucuronic acid, which also has been shown to directly stimulate the iNKT TCR when presented by CD1d (7, 8). *S. paucimobilis* also produces another GSL with a tetrasaccharide head group, that is either weakly antigenic (38) or not antigenic (39).

We infected Nur77^{gfp} mice with *S. pneumoniae* systemically by intravenous injection, harvested iNKT cells 6 or 20 hours later, and analyzed cytokine by intracellular staining directly *ex vivo* without re-stimulation (Fig. 3A). As expected, iNKT cells produced IFN γ (Fig. 3B) and upregulated CD69 (not shown). Consistent with a TCR-dependent activation mode, the expression of GFP increased in iNKT cells as well (Fig. 3A). This increase was statistically significant compared to the uninfected control mice, but was well below that

seen after injection of α GalCer, indicating that although lipids are presented during microbial infections that are able to activate iNKT cells through their TCR, they are weaker agonists than α GalCer. This result is consistent with previous reports that lipid antigens from *S. pneumoniae* presented by CD1d have an affinity for the iNKT TCR in the low μ M range, compared to the low nM range for α GalCer complexes with CD1d (10, 40). Notably, the GFP level was higher in the subset of iNKT cells producing IFN γ , suggesting that a TCR signal activated cytokine production.

Similarly, iNKT cells upregulated GFP expression 18 hours after infection with *S. paucimobilis* (Fig. 4A). This increase was statistically significant compared to the uninfected controls, but again was well below that seen after injection of α GalCer (Fig. 4A), which is consistent with previous reports (7). For these experiments we used Nur77^{efp}/KN2 mice that have a human CD2 gene engineered into one allele of the IL-4 locus, because it was shown that *Sphingomas* GSLs also induce IL-4 production by iNKT cells (7). *S. paucimobilis* infection resulted in the production of IL-4 by iNKT cells, as judged by increased hCD2 staining, which interestingly was primarily produced by cells that had upregulated GFP (Fig. 4B), again supporting the idea that a TCR signal activates IL-4 in iNKT cells. Overall, these data indicate that *S. paucimobilis* and *S. pneumoniae* infections, in which the bacteria have glycolipid antigens for the iNKT TCR, activate iNKT cells through TCR stimulation.

***S. typhimurium* infection does not stimulate iNKT cells through their TCR**

Having analyzed a cytokine mediated model of activation and a two direct activation models, we next analyzed a third proposed model of iNKT cell activation. In this case, we infected mice with a microbe that has not been shown to produce a microbial antigen for iNKT TCR stimulation, but one that putatively causes APCs to present an endogenous self-lipid that stimulates the iNKT TCR, namely *Salmonella typhimurium* (11, 13). *S. typhimurium* is a Gram-negative LPS-positive pathogen in which a microbial lipid antigen has not been detected, but which causes an activation of iNKT cells that has been shown to be partially CD1d dependent, but which is also dependent on MyD88 expression by the APCs (11) and on their ability to produce IL-12 when activated (13). Unexpectedly, when Nur77^{efp} mice were infected with *S. typhimurium*, the iNKT cells did not upregulate GFP expression (Fig. 5A). Despite this, there was increased expression of CD69 (data not shown) and an increase in the percentage of iNKT cells producing IFN γ (Fig. 5B). The iNKT cells producing IFN γ did not have a higher mean expression of GFP (Fig. 5B), unlike what was observed for cytokine producing iNKT cells in *S. pneumoniae* or *S. paucimobilis* infection (Fig. 3B and 4B). To ensure that we did not miss a relevant time point *in vivo*, we evaluated GFP expression at earlier (4 hours) and later (72 hours) times, but did not observe a GFP increase (Fig. 5A). However, at all of the tested time points, there was a statistically significant increase in the percentage of iNKT cells producing IFN γ (Fig. 5B). These data suggest that iNKT cells were not receiving stimulation through their TCR during *S. typhimurium* infection, despite being activated to produce IFN γ .

Given that this result contradicts previous reports (13, 15), we tested if iNKT cells could produce IFN γ in the absence of CD1d expressing APCs during *S. typhimurium* infection. This was done by enriching for iNKT cells from the liver and spleen of V α 14 TCR

transgenic crossed to Nur77^{gfp} transgenic mice, and by adoptively transferring these cells into either CD1d-deficient or WT mice. Host mice were infected with *S. typhimurium* 24 hours after receiving the iNKT cells, and 20 hours after infection the transferred iNKT cells were analyzed. The transferred iNKT cells did not upregulate GFP expression regardless of whether they were recovered from CD1d-deficient or WT control hosts (Fig. 6), consistent with the result obtained previously (Fig. 5). Despite this, iNKT cells from both types of infected hosts produced IFN γ , and the percentage of cytokine producing iNKT cells was not significantly different. These data suggest that iNKT cells were activated independently of CD1d, and thus independently of endogenous self-lipid antigen, during *S. typhimurium* infection.

TLR stimulation does not cause TCR-dependent activation of iNKT cells

The presentation of endogenous self-lipids by CD1d leading to the stimulation of iNKT cells has been proposed to occur as a result of TLR stimulation of APCs (14-16). Specifically, stimulation of TLR4 or TLR9 was suggested to lead to increased synthesis and/or presentation of endogenous self-lipids (11, 14-16). In order to test this, we injected Nur77^{gfp} mice with either LPS to stimulate TLR4, or CpG oligodeoxynucleotides (ODNs) to stimulate TLR9, and examined iNKT cell activation 4, 24, or 36 hours later. There was no increase in GFP in iNKT cells at any time point after TLR activation, despite robust production of IFN γ and elevated expression of CD69 at 4 hours for LPS and 36 hours for CpG (Fig. 7 and data not shown).

This contradicts the conclusions from a previous report, in which it was found that TLR stimulation led to the inhibition of the enzyme α -galactosidase A (α -Gal A) (14). α -Gal A was proposed to convert a stimulatory lipid to a non-stimulatory lipid, and thus the inhibition of this enzyme would result in the increased presentation of the stimulatory lipid for iNKT cell activation (14). Therefore, we used the Nur77^{gfp} mice with α -Galactosidase A deficient (α -Gal A^{-/-}) mice to test this notion. Initially, we looked at GFP expression in Nur77^{gfp}: α -Gal A^{-/-} mixed bone marrow chimeras to examine iNKT cell activation in the steady state *in vivo*. However, we saw no GFP expression in the recovered Nur77^{gfp} iNKT cells (Fig. S1). Next, we analyzed a model of acute activation of iNKT cells by examining GFP expression after adoptive transfer of iNKT cells from Nur77^{gfp} V α 14 TCR transgenic mice into α -Gal A^{-/-} or WT hosts. The Nur77^{gfp} V α 14 TCR transgenic iNKT cells did not have increased GFP when recovered from α -Gal A^{-/-} hosts (Fig. S2). Lastly, we performed a direct comparison with the reported data, but analyzed GFP expression as a measure of iNKT TCR stimulation in addition to the decrease in the α GalCer-loaded CD1d-tetramer⁺ TCR β ⁺ population used in the previous report as a measure of activation (14). Similar to the prior study, we transferred α -Gal A^{-/-} BMDC, β GlcCer pulsed α -Gal A^{-/-} BMDC or WT BMDC into Nur77^{gfp} recipients. iNKT cells from animals injected with α -Gal A^{-/-} BMDC failed to show increased GFP expression in comparison to animals injected with WT BMDC, while increased GFP expression was seen in iNKT cells from animals injected with β GlcCer pulsed α -Gal A^{-/-} BMDC (Fig. S3).

Activation of iNKT cells can lead to a decreased recovery when using tetramer, due to TCR internalization (41) and/or activation induced cell death (42, 43). This reduced recovery has

been used as a measure of iNKT cell activation *in vivo*. However, mice from α -Gal A^{-/-} BMDC injected groups exhibited variability in the percentage of α GalCer-loaded CD1d-tetramer⁺ TCR β ⁺ populations recovered (Fig. S4). Overall, our data are not consistent with the hypothesis that TLR stimulation of APCs leads to the presentation of endogenous self-lipid for the stimulation of iNKT TCRs.

Discussion

iNKT cells are able to rapidly produce cytokines not only in response to bacterial, viral, and fungal infections, but also in response to cancers, various inflammatory conditions, and autoimmune diseases (44). The invariant nature of the iNKT cell TCR α chain must restrict the diversity of lipids that are capable of stimulating iNKT cells, but despite this, diverse structures have been reported for both the microbial and self-lipids recognized by these cells including GSLs, diacylglycerols, phospholipids and cholesterol containing compounds (45). Also, several TCR independent modes of activation of iNKT cells have been reported, including the cytokine-mediated activation described above, but also stimulation mediated by engagement of activating NK receptors such as NKG2D (46) and activation of iNKT cells in the liver by engagement of β -adrenergic receptors (47). This ability to respond in such diverse contexts raises the question as to the importance of TCR-dependent pathways in those situations in which iNKT cells play a role in the immune response, and also, when the TCR is involved, the relative weight of recognition of self versus foreign glycolipids.

Here we used the Nur77^{gfp} mouse as a sensitive tool to examine the role of iNKT cell TCR stimulation during various infections. We showed that this reporter mouse can detect iNKT cell responses to self glycolipid antigens presented by CD1d that have an affinity for the iNKT cell TCR of approximately 5 μ M or weaker, nearly 10³-fold weaker than the affinity for α GalCer presented by CD1d. Using this system, we showed that iNKT cells are stimulated through their TCR when infected with pathogens known to have antigenic lipids, such as in *S. pneumoniae* infection and *S. paucimobilis* infection.

Using infection with MCMV, we showed that iNKT cells became activated in the absence of TCR stimulation, consistent with other results showing that cytokines from APCs were sufficient to activate a protective response by iNKT cells after this infection (25, 37). Further evidence showing that iNKT cells can undergo TCR-independent stimulation was recently reported. Using a transgenic mouse in which the iNKT TCR could be conditionally ablated, the authors showed that TCR-negative iNKT cells could still mount a response to LPS with similar amounts of IFN γ produced compared to iNKT cells that retained TCR expression (48).

More surprising was our findings that iNKT cell TCR stimulation could not be detected after either *S. typhimurium* infection or TLR stimulation of APCs, as these observations run counter to a currently popular model (45). It could be argued that the reporter mouse is not sufficiently sensitive, and that while we have tested some of the weakest reported TCR agonists, that *S. typhimurium* induces the presentation of a self-antigen that has a very low abundance, and/or TCR affinity, that it cannot give an activation signal in the Nur77^{gfp} mice. We consider this unlikely, however, because when transferred to CD1d^{-/-} deficient

hosts, we found that the iNKT cell IFN γ response to *S. typhimurium* was not impaired. Regarding the reported TCR-dependence of the iNKT cell response to *S. typhimurium*, much of the data relied on *in vitro* studies. For example, *in vitro* experiments showing that CD1d^{-/-} BMDC were ineffective at inducing IFN γ production by tetramer sorted iNKT cells in response to *S. typhimurium* are difficult to reconcile with our data. *In vivo* studies used CD1d-blocking antibodies to demonstrate TCR dependent IFN γ production. It is possible that antibodies to CD1d could have had a direct effect on APC, independent of iNKT cells, that led to a reduction in inflammatory cytokine production, and a consequent reduction in IFN γ production by iNKT cells (49).

Similarly, we did not find that TLR activation of APCs induced a detectable TCR signal in iNKT cells, although it was previously reported to increase the synthesis of enzymes responsible for the synthesis of stimulatory self-antigens (13, 16, 17, 50). Although again, other *in vivo* experiments showed that iNKT cells with and without a TCR produced similar levels of IFN γ in response to LPS (48). One possible explanation is that a stronger stimulation of the innate immune system achieved *in vivo* may somehow inhibit TCR dependent activation of iNKT cells. TLR stimulation was also reported to inhibit the synthesis of an enzyme, α -Gal A, that would degrade a putative lipid self-antigen (14). Regarding the role of α -Gal A, using the Nur77^{efp} reporter mouse, we did not find evidence that APCs in mice deficient for this enzyme induced a TCR signal in iNKT cells *in vivo*. In the α -Gal A deficient model, the previous data supporting TCR-dependent activation of iNKT cells relied upon loss of the α GalCer-loaded CD1d-tetramer⁺ TCR β ⁺ population and on the increased expression of the surface marker CD69 (14). In our hands, however, population loss was not a consistent and reliable measure, and CD69 expression is not strictly TCR dependent, as it is also influenced by inflammatory stimuli.

In the cases of infection with *S. pneumoniae* and *S. paucimobilis*, in which there was a detectable TCR signal in the Nur77^{efp} reporter mouse, the question remains as to whether that signal was due to the presence of a foreign antigen or the induction of the synthesis of a self-antigen, although these alternatives are not mutually exclusive, and both types of antigens could be present (17). The evidence provided here is only correlative with the need for a foreign antigen to induce a TCR-mediated signal in the infections studied, although if self-antigen were dominant during each of the three bacterial infections, it is puzzling that a signal from the reporter mouse could only be obtained using bacteria with a foreign antigen. There is no evidence that the iNKT cells from the *S. typhimurium* infected mice were less activated; in fact, by intracellular cytokine staining they were at least as activated as the populations from *S. pneumoniae* or *S. paucimobilis* infected mice. It has not been possible to resolve the issue of foreign versus self-antigen recognition so far, however, as it has not been feasible to remove either the microbial antigen or the self-antigen by genetic deletion, without causing unacceptable secondary effects. For example, *S. pneumoniae* mutants that cannot synthesize the glycosylated diacylglycerol antigen have proved not to be viable (M. Kronenberg, unpublished). Regarding the self-antigen, problems include the diversity of self-antigens reported, with both GSLs and phospholipids, and secondary effects on the health of the mice and the functioning of the endosomal system in APCs when basic components of glycolipid synthesis pathways are eliminated (51, 52).

Although *S. typhimurium* infection did not stimulate the iNKT TCR as a result of endogenous lipid antigen presentation in our experiments, we are not proposing that this pathway for iNKT cell activation is unimportant. It is known that iNKT cells do receive TCR stimulation from endogenous lipid during iNKT cell thymic selection (53, 54). For example, in the Nur77^{gfp} mice, intrathymic stage 0 iNKT cell precursors expressed a high level of GFP (30). Furthermore, thymic NKT2 cells from BALB/c mice, which produce IL-4 at steady state, showed an increased expression of GFP compared to NKT1 or NKT17 cells (55). Additionally, other types of infections remain to be tested for iNKT cell TCR stimulation as was done in this report. For example, infection with the helminth parasite *Schistosoma mansoni*, in which iNKT cell activation has been shown to be dependent on CD1d after schistosome egg encounter, remains one of a number of possible examples of iNKT cell TCR stimulation mediated by an endogenous lipid, as a foreign lipid from *S. mansoni* has not been reported (56).

In conclusion, we have demonstrated that the Nur77^{gfp} mouse provides a sensitive tool for directly measuring iNKT cell TCR stimulation that detects endogenous and foreign lipid agonists even when they vary greatly in potency. Additionally, our data provided insight into the role of iNKT cell TCR stimulation in iNKT cell activation during infections, differentiating between infections that are dependent on TCR stimulation and infections that are not. We showed that TCR-independent activation applies to bacterial as well as viral infections, and furthermore, that the ability to detect TCR-mediated activation *in vivo* correlated with the presence of foreign antigen. This approach may be useful to identify those situations in which a self-antigen mediated activation of iNKT cells is most relevant, for example, in the context of sterile inflammatory conditions such as ischemia reperfusion injury (57).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to thank Jane Ding for excellent technical support, and You Jeong Lee for helpful discussion and critical reading of the manuscript. Markus Mohrs (Trudeau Institute) provided KN2 mice. Florian Winau (Harvard) provided Fabry mice. Stephen McSorley (University of California, Davis) provided *Salmonella typhimurium* (SL1344). Chris A. Benedict (University of California, San Diego) provided mouse cytomegalovirus (MCMV-K181). The authors declare no competing financial interests.

This research was supported by NIH grants R37-AI39560 (to K.A.H.) and T32 HD060536 training grant (to K.L.H.). Mitchell Kronenberg and Aaron Tzysnik's research was supported by R01 AI69296, R37 AI71922 and R21 AI076864 (M.K.), and F32 AI80087 (to A.J.T.).

Abbreviations used in this paper

iNKT	invariant Natural Killer T cell
MCMV	mouse cytomegalovirus
<i>S. typhimurium</i>	<i>Salmonella typhimurium</i>

αGalCer	α -Galactosylceramide
<i>S. pneumoniae</i>	<i>Streptococcus pneumoniae</i>
<i>S. paucimobilis</i>	<i>Sphingomonas paucimobilis</i>
GSL	glycosphingolipid
iGb3	isoglobotrihexosylceramide
βGlcCer	β -D-glucopyranosylceramide
BMDC	bone marrow derived dendritic cell
α-Gal A	α -Galactosidase A

References

1. Lantz O, Bendelac A. An invariant T cell receptor alpha chain is used by a unique subset of major histocompatibility complex class I-specific CD4+ and CD4-8- T cells in mice and humans. *J. Exp. Med.* 1994; 180:1097–1106. [PubMed: 7520467]
2. Kawano T, Cui J, Koezuka Y, Toura I, Kaneko Y, Motoki K, Ueno H, Nakagawa R, Sato H, Kondo E, Koseki H, Taniguchi M. CD1d-restricted and TCR-mediated activation of valpha14 NKT cells by glycosylceramides. *Science.* 1997; 278:1626–1629.
3. Cui J, Shin T, Kawano T, Sato H, Kondo E, Toura I, Kaneko Y, Koseki H, Kanno M, Taniguchi M. Requirement for Valpha14 NKT cells in IL-12-mediated rejection of tumors. *Science.* 1997; 278:1623–1626. [PubMed: 9374462]
4. Hong S, Wilson MT, Serizawa I, Wu L, Singh N, Naidenko OV, Miura T, Haba T, Scherer DC, Wei J, Kronenberg M, Koezuka Y, Van Kaer L. The natural killer T-cell ligand alpha-galactosylceramide prevents autoimmune diabetes in non-obese diabetic mice. *Nat. Med.* 2001; 7:1052–1056. [PubMed: 11533710]
5. Shi FD, Van Kaer L. Reciprocal regulation between natural killer cells and autoreactive T cells. *Nat. Rev. Immunol.* 2006; 6:751–760. [PubMed: 16998508]
6. Tupin E, Kinjo Y, Kronenberg M. The unique role of natural killer T cells in the response to microorganisms. *Nat. Rev. Microbiol.* 2007; 5:405–417. [PubMed: 17487145]
7. Kinjo Y, Wu D, Kim G, Xing GW, Poles MA, Ho DD, Tsuji M, Kawahara K, Wong CH, Kronenberg M. Recognition of bacterial glycosphingolipids by natural killer T cells. *Nature.* 2005; 434:520–525. [PubMed: 15791257]
8. Sriram V, Du W, Gervay-Hague J, Brutkiewicz RR. Cell wall glycosphingolipids of *Sphingomonas paucimobilis* are CD1d-specific ligands for NKT cells. *Eur. J. Immunol.* 2005; 35:1692–1701. [PubMed: 15915536]
9. Kawakami K, Yamamoto N, Kinjo Y, Miyagi K, Nakasone C, Uezu K, Kinjo T, Nakayama T, Taniguchi M, Saito A. Critical role of Valpha14+ natural killer T cells in the innate phase of host protection against *Streptococcus pneumoniae* infection. *Eur. J. Immunol.* 2003; 33:3322–3330. [PubMed: 14635040]
10. Kinjo Y, Illarionov P, Vela JL, Pei B, Girardi E, Li X, Li Y, Imamura M, Kaneko Y, Okawara A, Miyazaki Y, Gomez-Velasco A, Rogers P, Dahesh S, Uchiyama S, Khurana A, Kawahara K, Yesilkaya H, Andrew PW, Wong CH, Kawakami K, Nizet V, Besra GS, Tsuji M, Zajonc DM, Kronenberg M. Invariant natural killer T cells recognize glycolipids from pathogenic Gram-positive bacteria. *Nat. Immunol.* 2011; 12:966–974. [PubMed: 21892173]
11. Mattner J, Debord KL, Ismail N, Goff RD, Cantu C 3rd, Zhou D, Saint-Mezard P, Wang V, Gao Y, Yin N, Hoebe K, Schneewind O, Walker D, Beutler B, Teyton L, Savage PB, Bendelac A. Exogenous and endogenous glycolipid antigens activate NKT cells during microbial infections. *Nature.* 2005; 434:525–529. [PubMed: 15791258]

12. Kawahara K, Seydel U, Matsuura M, Danbara H, Rietschel ET, Zahringer U. Chemical structure of glycosphingolipids isolated from *Sphingomonas paucimobilis*. *FEBS Lett.* 1991; 292:107–110. [PubMed: 1959589]
13. Brigl M, Bry L, Kent SC, Gumperz JE, Brenner MB. Mechanism of CD1d-restricted natural killer T cell activation during microbial infection. *Nat. Immunol.* 2003; 4:1230–1237. [PubMed: 14578883]
14. Darmoise A, Teneberg S, Bouzonville L, Brady RO, Beck M, Kaufmann SH, Winau F. Lysosomal alpha-galactosidase controls the generation of self lipid antigens for natural killer T cells. *Immunity.* 2010; 33:216–228. [PubMed: 20727792]
15. Brigl M, Tatituri RV, Watts GF, Bhowruth V, Leadbetter EA, Barton N, Cohen NR, Hsu FF, Besra GS, Brenner MB. Innate and cytokine-driven signals, rather than microbial antigens, dominate in natural killer T cell activation during microbial infection. *J. Exp. Med.* 2011; 208:1163–1177. [PubMed: 21555485]
16. Paget C, Malleveay T, Speak AO, Torres D, Fontaine J, Sheehan KC, Capron M, Ryffel B, Faveeuw C, Leite de Moraes M, Platt F, Trottein F. Activation of invariant NKT cells by toll-like receptor 9-stimulated dendritic cells requires type I interferon and charged glycosphingolipids. *Immunity.* 2007; 27:597–609. [PubMed: 17950005]
17. Brennan PJ, Tatituri RV, Brigl M, Kim EY, Tuli A, Sanderson JP, Gadola SD, Hsu FF, Besra GS, Brenner MB. Invariant natural killer T cells recognize lipid self antigen induced by microbial danger signals. *Nat. Immunol.* 2011; 12:1202–1211. [PubMed: 22037601]
18. Zhou D, Mattner J, Cantu C 3rd, Schrantz N, Yin N, Gao Y, Sagiv Y, Hudspeth K, Wu YP, Yamashita T, Teneberg S, Wang D, Proia RL, Levery SB, Savage PB, Teyton L, Bendelac A. Lysosomal glycosphingolipid recognition by NKT cells. *Science.* 2004; 306:1786–1789. [PubMed: 15539565]
19. Dieli F, Caccamo N, Meraviglia S, Ivanyi J, Sireci G, Bonanno CT, Ferlazzo V, La Mendola C, Salerno A. Reciprocal stimulation of gammadelta T cells and dendritic cells during the anti-mycobacterial immune response. *Eur. J. Immunol.* 2004; 34:3227–3235. [PubMed: 15376192]
20. Sutton CE, Lalor SJ, Sweeney CM, Breerton CF, Lavelle EC, Mills KH. Interleukin-1 and IL-23 induce innate IL-17 production from gammadelta T cells, amplifying Th17 responses and autoimmunity. *Immunity.* 2009; 31:331–341. [PubMed: 19682929]
21. O'Donnell H, Pham OH, Li LX, Atif SM, Lee SJ, Ravesloot MM, Stolfi JL, Nuccio SP, Broz P, Monack DM, Baumler AJ, McSorley SJ. Toll-like Receptor and Inflammasome Signals Converge to Amplify the Innate Bactericidal Capacity of T Helper 1 Cells. *Immunity.* 2014; 40:213–224. [PubMed: 24508233]
22. Berg RE, Cordes CJ, Forman J. Contribution of CD8+ T cells to innate immunity: IFN-gamma secretion induced by IL-12 and IL-18. *Eur. J. Immunol.* 2002; 32:2807–2816. [PubMed: 12355433]
23. Soudja SM, Ruiz AL, Marie JC, Lauvau G. Inflammatory monocytes activate memory CD8(+) T and innate NK lymphocytes independent of cognate antigen during microbial pathogen invasion. *Immunity.* 2012; 37:549–562. [PubMed: 22940097]
24. Nagarajan NA, Kronenberg M. Invariant NKT cells amplify the innate immune response to lipopolysaccharide. *J. Immunol.* 2007; 178:2706–2713. [PubMed: 17312112]
25. Tyznik AJ, Tupin E, Nagarajan NA, Her MJ, Benedict CA, Kronenberg M. Cutting edge: the mechanism of invariant NKT cell responses to viral danger signals. *J. Immunol.* 2008; 181:4452–4456. [PubMed: 18802047]
26. Wesley JD, Tessmer MS, Chaukos D, Brossay L. NK cell-like behavior of V alpha 14i NK T cells during MCMV infection. *Plos Pathog.* 2008; 4
27. Van Kaer L. NKT cells: T lymphocytes with innate effector functions. *Curr. Opin. Immunol.* 2007; 19:354–364. [PubMed: 17428648]
28. Cardell SL. The natural killer T lymphocyte: a player in the complex regulation of autoimmune diabetes in non-obese diabetic mice. *Clin. Exp. Immunol.* 2006; 143:194–202. [PubMed: 16412042]
29. Matangkasombut P, Pichavant M, Dekruyff RH, Umetsu DT. Natural killer T cells and the regulation of asthma. *Mucosal Immunol.* 2009; 2:383–392. [PubMed: 19587638]

30. Moran AE, Holzapfel KL, Xing Y, Cunningham NR, Maltzman JS, Punt J, Hogquist KA. T cell receptor signal strength in Treg and iNKT cell development demonstrated by a novel fluorescent reporter mouse. *J. Exp. Med.* 2011; 208:1279–1289. [PubMed: 21606508]
31. Sullivan BA, Nagarajan NA, Wingender G, Wang J, Scott I, Tsuji M, Franck RW, Porcelli SA, Zajonc DM, Kronenberg M. Mechanisms for glycolipid antigen-driven cytokine polarization by Valpha14i NKT cells. *J. Immunol.* 2010; 184:141–153. [PubMed: 19949076]
32. Borg NA, Wun KS, Kjer-Nielsen L, Wilce MC, Pellicci DG, Koh R, Besra GS, Bharadwaj M, Godfrey DI, McCluskey J, Rossjohn J. CD1d-lipid- antigen recognition by the semi-invariant NKT T-cell receptor. *Nature.* 2007; 448:44–49. [PubMed: 17581592]
33. Wun KS, Cameron G, Patel O, Pang SS, Pellicci DG, Sullivan LC, Keshipeddy S, Young MH, Uldrich AP, Thakur MS, Richardson SK, Howell AR, Illarionov PA, Brooks AG, Besra GS, McCluskey J, Gapin L, Porcelli SA, Godfrey DI, Rossjohn J. A Molecular Basis for the Exquisite CD1d-Restricted Antigen Specificity and Functional Responses of Natural Killer T Cells. *Immunity.* 2011; 34:327–339. [PubMed: 21376639]
34. Zajonc DM, Savage PB, Bendelac A, Wilson IA, Teyton L. Crystal structures of mouse CD1d-iGb3 complex and its cognate Valpha14 T cell receptor suggest a model for dual recognition of foreign and self glycolipids. *J. Mol. Biol.* 2008; 377:1104–1116. [PubMed: 18295796]
35. Pellicci DG, Clarke AJ, Patel O, Mallevaey T, Beddoe T, Le Nours J, Uldrich AP, McCluskey J, Besra GS, Porcelli SA, Gapin L, Godfrey DI, Rossjohn J. Recognition of beta-linked self glycolipids mediated by natural killer T cell antigen receptors. *Nat. Immunol.* 2011; 12:827–U823. [PubMed: 21804559]
36. Shioh LR, Rosen DB, Brdickova N, Xu Y, An J, Lanier LL, Cyster JG, Matloubian M. CD69 acts downstream of interferon-alpha/beta to inhibit S1P1 and lymphocyte egress from lymphoid organs. *Nature.* 2006; 440:540–544. [PubMed: 16525420]
37. Tyznik AJ, Verma S, Wang Q, Kronenberg M, Benedict CA. Distinct Requirements for Activation of NKT and NK Cells during Viral Infection. *J. Immunol.* 2014 in press.
38. Tyznik AJ, Farber E, Girardi E, Birkholz A, Li Y, Chitale S, So R, Arora P, Khurana A, Wang J, Porcelli SA, Zajonc DM, Kronenberg M, Howell AR. Glycolipids that elicit IFN-gamma-biased responses from natural killer T cells. *Chem. Biol.* 2011; 18:1620–1630. [PubMed: 22195564]
39. Long XT, Deng S, Mattner JC, Zang Z, Zhou D, McNary N, Goff RD, Teyton L, Bendelac A, Savage PB. Synthesis and evaluation of stimulatory properties of Sphingomonadaceae glycolipids. *Nat. Chem. Biol.* 2007; 3:559–564. [PubMed: 17660835]
40. Girardi E, Yu ED, Li Y, Tarumoto N, Pei B, Wang J, Illarionov P, Kinjo Y, Kronenberg M, Zajonc DM. Unique interplay between sugar and lipid in determining the antigenic potency of bacterial antigens for NKT cells. *PLoS Biol.* 2011; 9:e1001189. [PubMed: 22069376]
41. Wilson MT, Johansson C, Olivares-Villagomez D, Singh AK, Stanic AK, Wang CR, Joyce S, Wick MJ, Van Kaer L. The response of natural killer T cells to glycolipid antigens is characterized by surface receptor down-modulation and expansion. *Proc. Natl. Acad. Sci. USA.* 2003; 100:10913–10918. [PubMed: 12960397]
42. Eberl G, MacDonald HR. Selective induction of NK cell proliferation and cytotoxicity by activated NKT cells. *Eur. J. Immunol.* 2000; 30:985–992. [PubMed: 10760785]
43. Matsuda JL, Naidenko OV, Gapin L, Nakayama T, Taniguchi M, Wang CR, Koezuka Y, Kronenberg M. Tracking the response of natural killer T cells to a glycolipid antigen using CD1d tetramers. *J. Exp. Medicine.* 2000; 192:741–753.
44. Bendelac A, Savage PB, Teyton L. The biology of NKT cells. *Annu. Rev. Immunol.* 2007; 25:297–336. [PubMed: 17150027]
45. Brennan PJ, Brigl M, Brenner MB. Invariant natural killer T cells: an innate activation scheme linked to diverse effector functions. *Nat. Rev. Immunol.* 2013; 13:101–117. [PubMed: 23334244]
46. Kuylentsterna C, Bjorkstrom NK, Andersson SK, Sahlstrom P, Bosnjak L, Paquin-Proulx D, Malmberg KJ, Ljunggren HG, Moll M, Sandberg JK. NKG2D performs two functions in invariant NKT cells: Direct TCR-independent activation of NK-like cytotoxicity and co-stimulation of activation by CD1d. *Eur. J. Immunol.* 2011; 41:1913–1923. [PubMed: 21590763]
47. Wong CHY, Jenne CN, Lee WY, Leger C, Kubes P. Functional Innervation of Hepatic iNKT Cells Is Immunosuppressive Following Stroke. *Science.* 2011; 334:101–105. [PubMed: 21921158]

48. Vahl JC, Heger K, Knies N, Hein MY, Boon L, Yagita H, Polic B, Schmidt-Supprian M. NKT cell-TCR expression activates conventional T cells in vivo, but is largely dispensable for mature NKT cell biology. *PLoS Biol.* 2013; 11:e1001589. [PubMed: 23853545]
49. Colgan SP, Hershberg RM, Furuta GT, Blumberg RS. Ligation of intestinal epithelial CD1d induces bioactive IL-10: critical role of the cytoplasmic tail in autocrine signaling. *Proc. Natl. Acad. Sci. USA.* 1999; 96:13938–13943. [PubMed: 10570177]
50. Salio M, Speak AO, Shepherd D, Polzella P, Illarionov PA, Veerapen N, Besra GS, Platt FM, Cerundolo V. Modulation of human natural killer T cell ligands on TLR-mediated antigen-presenting cell activation. *Proc. Natl. Acad. Sci. USA.* 2007; 104:20490–20495. [PubMed: 18077358]
51. Gadola SD, Silk JD, Jeans A, Illarionov PA, Salio M, Besra GS, Dwek R, Butters TD, Platt FM, Cerundolo V. Impaired selection of invariant natural killer T cells in diverse mouse models of glycosphingolipid lysosomal storage diseases. *J. Exp. Med.* 2006; 203:2293–2303. [PubMed: 16982810]
52. Schumann J, Facciotti F, Panza L, Michieletti M, Compostella F, Collmann A, Mori L, De Libero G. Differential alteration of lipid antigen presentation to NKT cells due to imbalances in lipid metabolism. *Eur. J. Immunol.* 2007; 37:1431–1441. [PubMed: 17492806]
53. Bendelac A, Bonneville M, Kearney JF. Autoreactivity by design: innate B and T lymphocytes. *Nat. Rev. Immunol.* 2001; 1:177–186. [PubMed: 11905826]
54. Gopin L. iNKT cell autoreactivity: what is 'self' and how is it recognized? *Nat. Rev. Immunol.* 2010; 10:272–277. [PubMed: 20224567]
55. Lee YJ, Holzapfel KL, Zhu J, Jameson SC, Hogquist KA. Steady-state production of IL-4 modulates immunity in mouse strains and is determined by lineage diversity of iNKT cells. *Nat. Immunol.* 2013; 14:1146–1154. [PubMed: 24097110]
56. Mallevaey T, Zanetta JP, Faveeuw C, Fontaine J, Maes E, Platt F, Capron M, de-Moraes ML, Trottein F. Activation of invariant NKT cells by the helminth parasite *schistosoma mansoni*. *J. Immunol.* 2006; 176:2476–2485. [PubMed: 16456008]
57. Wallace KL, Marshall MA, Ramos SI, Lannigan JA, Field JJ, Strieter RM, Linden J. NKT cells mediate pulmonary inflammation and dysfunction in murine sickle cell disease through production of IFN-gamma and CXCR3 chemokines. *Blood.* 2009; 114:667–676. [PubMed: 19433855]




Fig. 1. iNKT cells upregulated GFP in response to various antigenic lipids, including self-lipids
 Expression of GFP (top) and CD69 (bottom) by splenic iNKT cells from B6 Nur77^{flp} mice 18 h after injection with bone marrow-derived unpulsed DCs (cntrl) or BMDCs pulsed with αGalCer, OCH, or the self-lipids βGlcCer or iGb3 as described in Materials and Methods. Histograms are representative of eight independent experiments. Average relative mean fluorescence intensity (MFI) of GFP (normalized to that in CD4 T cells in each experiment) is indicated.




Fig. 2. MCMV infection induced iNKT cell cytokine production, but not GFP expression
 Nur77^{gfp} B6 mice were infected with MCMV *i.p.* and tet⁺ liver iNKT cells were analyzed by flow cytometry 36 h later for GFP expression (A) and IFN γ production (B). Results show uninfected (gray), MCMV infected (red) and α GalCer treated animals 3 h after injection (black). Data are representative of two independent experiments, with at least 3 animals per time point. The histogram in (B) shows GFP expression of the IFN γ ⁻ iNKT cell population (black) versus the IFN γ ⁺ iNKT cell population (red) from an animal 36 h post infection. In graphs, each dot represents one mouse (Uninfected n = 6; 12 h post infection n = 3; 24 h post infection n = 3; 36 h post infection n = 6; 3 h post α GalCer injection n = 3). **P = 0.0026, percentage uninfected IFN γ ⁺ iNKT cells versus percentage 36 h post infection IFN γ ⁺ iNKT cells (unpaired two-tailed t-test).




Fig. 3. *S. pneumoniae* infection induced cytokine production and GFP expression
 Nur77^{gfp} B6 mice were infected with *S. pneumoniae* *i.v.* and tet⁺ iNKT cells were analyzed by flow cytometry for GFP expression (A) and IFN γ production (B). Results shown are from uninfected mice (gray), mice 6 hr post *S. pneumoniae* infection (red) and α GalCer treated animals 2.5 h after injection (black). The histogram in (B) shows GFP expression in the IFN γ ⁻ iNKT cell population (black) versus the IFN γ ⁺ iNKT cell population (red) from 20 h post infection. Data show representative dot plots of two independent experiments with 6 animals per time point. For percentage of GFP⁺ iNKT cells graph (A), *** P = 0.0006, uninfected versus 6 h post infection; and *** P = 0.0007, uninfected versus 20 h post infection. For percentage of IFN- γ ⁺ iNKT cells graph (B), * P = 0.0478, uninfected versus 6 h post infection; * P = 0.0483, uninfected versus 20 h post infection (all using unpaired two-tailed t-test).




Fig. 4. *S. paucimobilis* infection induced cytokine production and GFP expression

IL-4 reporter (KN2) Nur77^{gfp} B6 mice were infected with *S. paucimobilis* *i.v.* and tet⁺ iNKT cells were analyzed by flow cytometry 18 h post infection for GFP (A) and human CD2 expression (B). Histograms in (A) show GFP expression for uninfected (gray), 18 h infected (red) and 3 h αGalCer injected (black) animals. The histogram in (B) shows GFP expression on the hCD2⁻iNKT cell population (black) versus the hCD2⁺ iNKT cell population (red) from an animal 18 h post infection. Data show representative dot plots of two independent experiments with 4 to 6 animals per time point. ***P* = 0.0082, uninfected versus 18 h post infection (unpaired two-tailed t-test).




Fig. 5. *S. typhimurium* infection induced cytokine production, but not GFP expression
 Nur77^{gfp} B6 mice were infected with *S. typhimurium* *i.v.* and tet⁺ iNKT cells were analyzed by flow cytometry directly *ex vivo* for GFP expression (A) and IFN- γ production (B). The histogram shows uninfected mice (gray), 20 h *S. typhimurium* infected (red) and 3 h α GalCer injected (black) animals. The histogram in (B) show GFP expression of the IFN γ ⁻ iNKT cell population (black) versus the IFN γ ⁺ iNKT cell population (red) from an animal 20 h post infection. Data show representative dot plots and histogram of four independent experiments with 4 to 6 animals per time point. For percentage of IFN γ ⁺ iNKT cells graph (B), **P = 0.0059, uninfected versus 4 h post infection; ***P < 0.0001, uninfected versus 20 h post infection; ***P < 0.0001, uninfected versus 72 h post infection (all with unpaired two-tailed t-test).




Fig. 6. iNKT cell cytokine production is CD1d independent during *S. typhimurium* infection
iNKT cells were purified from Nur77^{gfp} Vα14 TCR transgenic B6 mice and transferred into either WT hosts (iNKT → WT) or CD1d^{-/-} hosts (iNKT → CD1d^{-/-}). Mice were then infected with *S. typhimurium* *i.v.* and the transferred tet⁺ iNKT cells were isolated and analyzed by flow cytometry for intracellular IFNγ 20 h post infection. Dot plots (left) show representative data of iNKT cells isolated from an uninfected WT host (top, left), an uninfected CD1d^{-/-} host (bottom, left), an infected WT host (top, right) or infected CD1d^{-/-} host (bottom, right) from two independent experiments with 3 to 6 animals per group. **P* = 0.0127, uninfected in WT host versus 20 h post infection in WT host; **P* = 0.0381, uninfected in CD1d^{-/-} host versus 20 h post infection in CD1d^{-/-} host (unpaired two-tailed t-test).




Fig. 7. iNKT cell cytokine production is CD1d independent during TLR stimulation

Nur77^{gfp} B6 mice were injected with LPS (top panel) or CpG (bottom panel) *i.v.* and tet⁺ iNKT cells were analyzed by flow cytometry 4 or 36 h later, respectively, for GFP expression, CD69 expression, and IFN γ production. Representative histograms show GFP expression (left) or CD69 expression (middle) for uninjected (gray), LPS or CPG injected (red), or 2-3 h α GalCer injected animals (black) from two independent experiments with 3 animals per group. For percentage of IFN γ ⁺ iNKT cells graph (right, top), $**P = 0.0026$, uninjected versus 4 h post LPS injection. For percentage of IFN γ ⁺ iNKT cells graph (right, bottom), $**P = 0.0092$, uninjected versus 36 h post CpG injection (unpaired two-tailed t-test).