

BMJ Open Marital status and survival after oesophageal cancer surgery: a population-based nationwide cohort study in Sweden

Nele Brusselaers,¹ Fredrik Mattsson,¹ Asif Johar,² Anna Wikman,² Pernilla Lagergren,² Jesper Lagergren,^{1,3} Rickard Ljung^{1,4}

To cite: Brusselaers N, Mattsson F, Johar A, *et al*. Marital status and survival after oesophageal cancer surgery: a population-based nationwide cohort study in Sweden. *BMJ Open* 2014;**4**: e005418. doi:10.1136/bmjopen-2014-005418

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2014-005418>).

Received 7 April 2014
Revised 13 May 2014
Accepted 16 May 2014



¹Unit of Upper Gastrointestinal Surgery, Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden

²Surgical Care Science, Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden

³Division of Cancer Studies, King's College London, London, UK

⁴Unit of Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

Correspondence to
Dr Nele Brusselaers;
nele.brusselaers@ki.se

ABSTRACT

Objectives: A beneficial effect of being married on survival has been shown for several cancer types, but is unclear for oesophageal cancer. The objective of this study was to clarify the potential influence of the marital status on the overall and disease-specific survival after curatively intended treatment of oesophageal cancer using a nationwide population-based design, taking into account the known major prognostic variables.

Design: Prospective, population-based cohort.

Setting: All Swedish hospitals performing surgery for oesophageal cancer during 2001–2005.

Participants: This study included 90% of all patients with oesophageal or junctional cancer who underwent surgical resection in Sweden in 2001–2005, with follow-up until death or the end of the study period (2012).

Primary and secondary outcome measures: Cox regression was used to estimate associations between the marital status and the 5-year overall and disease-specific mortality, expressed as HRs with 95% CIs, with adjustment for sex, age, tumour stage, histological type, complications, comorbidities and annual surgeon volume.

Results: Of all 606 included patients (80.4% men), 55.1% were married, 9.2% were remarried, 22.6% were previously married and 13% were never married. Compared with the married patients, the never married (HR 1.02, 95% CI 0.77 to 1.35), previously married (HR 0.90, 95% CI 0.71 to 1.15) and remarried patients (HR 0.79, 95% CI 0.55 to 1.13) had no increased overall 5-year mortality. The corresponding HRs for disease-specific survival, and after excluding the initial 90 days of surgery, were similar to the HRs for the overall survival.

Conclusions: This study showed no evidence of a better 5-year survival in married patients compared with non-married patients undergoing surgery for oesophageal cancer.

INTRODUCTION

The potential influence of the marital status on health and life expectancy has long been a matter of discussion.^{1 2} Being married has

Strengths and limitations of this study

- Patients with oesophageal cancer have a poor prognosis, even those patients eligible for curative surgery (oesophagectomy).
- Being married has been shown to be a marker of better survival after cancer in general and of specific types, but this has not been studied yet for oesophageal cancer.
- This nationwide population-based study did not find any evidence that the marital status influences long-term survival after oesophagectomy for oesophageal cancer, even after taking into account the known major prognostic variables.
- The absence of a clear association between marital status and long-term survival might be a false-negative result related to the relatively small sample size. Yet, it is also possible that the beneficial effect of being married is non-existent for oesophageal cancer.
- The main strength of this study is the population-based design with almost complete coverage, and complete follow-up of all patients operated in Sweden for oesophageal cancer in 2001–2005. Moreover, complete data on a large variety of variables enabled adjustment for known prognostic factors. Yet, among limitations are that the influence of other confounders or residual confounding by the variables adjusted for can never be ruled out. The statistical power was however limited to detect weak associations.

been shown to be a marker of better survival after cancer in general and of specific types,³ and the survival discrepancy between married and non-married patients with cancer has been claimed to be increasing over time.⁴ Having a partner is believed to lead to a healthier lifestyle, for example, regarding tobacco and alcohol use, physical activity and social support, and can also influence the choice and adherence to treatment, as well as the time elapsing between early

symptoms and a confirmed cancer diagnosis.² However, little is known about such an influence on oesophageal cancer. Only a few studies have investigated the potential role of socioeconomic factors in the prognosis of oesophageal cancer.^{5 6} A previous study by our group showed limited evidence of an association between lower education and worse long-term survival in operated patients with oesophageal cancer (only a significant difference in patients with tumour stage IV).⁷ A recent American study showed that married patients were more likely to be diagnosed with localised disease, and received treatment with curative intent more frequently than non-married patients, also showing a better overall survival.⁸ However, patients with oesophageal cancer have an overall very poor prognosis, mainly due to advanced tumour at the time of diagnosis, and only a minority is eligible for curatively intended treatment.⁹ Surgery (oesophagectomy) plays a main role in the curatively intended therapy for most oesophageal cancer.⁹ The objective of this study was to clarify the potential influence of the marital status on the overall and disease-specific survival after curatively intended treatment of oesophageal cancer using a nationwide population-based design, taking into account the known major prognostic variables.

METHODS

Participants

This was a population-based prospective cohort study, which included 90% of all patients with oesophageal or gastro-oesophageal junctional cancer treated with surgery and the treatment of choice for potentially curable patients in Sweden during the study period, from 2 April 2001 to 31 December 2005. Follow-up for survival ended in 31 August 2012. The median duration of follow-up was 619 days. From the full cohort of 616 patients, 10 patients were excluded because of missing values for the marital status, leaving 606 patients for final analysis.

Informed consent was obtained from each patient before inclusion in the study.

Data sources

The organisation of the comprehensive nationwide network of clinicians that participated in the data collection has been described elsewhere, and most parts of the design of this clinical cohort study have been presented in detail.^{10 11} In brief, information about the tumour stage, histology and localisation, surgical procedures and complications was prospectively collected for all patients through scrutiny of relevant medical records from 174 hospital departments in Sweden.¹¹ In September 2012, the cohort was linked to the Patient Register, the Causes of Death Register and the Longitudinal Integration Database for Health Insurance and Labour Market Studies (LISA). The Patient Register includes all in-hospital care and outpatient specialist

care in Sweden, including codes for diagnoses, surgical procedures and comorbidity with a high level of validity.¹² The Causes of Death Register contains information on the date and cause of death for all deceased Swedish residents since 1952.¹³ The LISA came into use in 1990 and includes a large variety of variables, including socioeconomic status, marital status and work history.¹⁴

Study exposure and outcome

Four mutually exclusive categories for marital status during the calendar year before surgery were married (never divorced or widowed), remarried (after widowhood or divorce), previously married (widowed or divorced) and never married. An additional analysis was performed grouping all patients into two groups: the currently married (married and remarried) and the not currently married patients (previously married and never married).

The main outcome measure was the overall mortality (including all causes of death) up to 5 years after oesophagectomy for oesophageal cancer. Other outcomes were mortality within 5 years after surgery: (1) after exclusion of deaths within first 90 days of surgery (defined as conditional mortality), (2) with oesophageal cancer as an underlying or contributing cause of death (disease-specific mortality or death related to oesophageal cancer) and (3) the combination of conditional and disease-specific mortality.

Statistical analyses

The association between marital status and mortality was analysed by means of multivariable Cox regression and expressed as HRs with 95% CIs. The married group was used as the reference category. Three regression models were employed: (1) without any adjustments (crude model), (2) adjusted for sex, age and tumour stage (basic model) and (3) further adjusted for histological tumour type, major complications, comorbidities and annual surgeon volume for oesophagectomy for cancer (fully adjusted model).¹⁵ Owing to the limited effect on survival, as shown in a previous study,⁷ the educational level was not included in the analyses. The age was categorised as <60, 60–74 or ≥75 years, and the tumour stage as 0–I, II, III or IV according to the sixth edition of the UICC (Union Internationale Contre le Cancer).¹⁶ The histological type was subdivided into adenocarcinoma and squamous cell carcinoma. Numbers of major complications (within 30 days of surgery) and comorbidities were categorised as none, one or more than one, as described previously.^{11 17} The surgeon volume was categorised into two equally sized groups based on the median annual number of oesophagectomies per surgeon (<8 or ≥8).¹⁵ To assess effect modification, stratified survival analyses were performed for the tumour stage and histological type using the same regression models, without adjustment for the stratifying variable.

RESULTS

Patients

Characteristics of the 606 study patients are described in [table 1](#). At the time of surgery, 334 patients were married (55.1%), 56 were remarried (9.2%), 137 were married previously (22.6%) and 79 patients were never married (13%). Compared with the married group, the remarried and never married groups were younger ([table 1](#)). The largest proportion of male patients was found in the never married group (87.3%) and the smallest in the previously married group (75.2%). Tumours were as likely to be of an advanced stage (stage IV) in all four groups (range 10.2–12%), but fewer early tumours (stage I) were found in the never married group (12.7%) compared with the married group (20.7%). Squamous cell carcinoma was found more frequently in the previously married (29.9%) and never married groups (30.4%), compared with the married (21.3%) and remarried group (23.2%). Of all patients, 53.6–62.8% had comorbidity at the time of surgery, which was least in the

remarried group. Married patients were least likely to have been operated on by surgeons with a low annual volume, compared with previously married patients (49.4% vs 58.4%). There were postoperative complications in 35.6–41.8% of the patients. In total, 455 (75.1%) patients died within 5 years of surgery, of whom 429 (94.3%) had oesophageal cancer as an underlying or contributing cause of death.

Marital status and mortality

The proportions of the overall and conditional 5-year mortality were highest in the never married group ([table 1](#)). The absolute 5-year survival rates for married, remarried, previously married and never married patients were 25.7%, 37.5%, 22.6% and 16.5%, respectively. Short-term mortality was slightly lower in the never married group (6.3% vs 8.1–8.9% in the other groups; [table 1](#)). The HRs for mortality after oesophagectomy according to the marital status are presented in [table 2](#). Compared with the married group, no increased HRs of mortality were found

Table 1 Demographic, treatment and tumour characteristics and mortality after oesophagectomy for cancer, categorised by the marital status at the time of diagnosis

	Marital status				Total N (%)
	Married N (%)	Remarried N (%)	Previously married* N (%)	Never married N (%)	
Total	334 (55.1)	56 (9.2)	137 (22.6)	79 (13.0)	606 (100.0)
Age (years)					
<60	63 (18.9)	18 (32.1)	31 (22.6)	33 (41.8)	145 (23.9)
60–74	199 (59.6)	34 (60.7)	69 (50.4)	37 (46.8)	339 (55.9)
≥75	72 (21.6)	4 (7.1)	37 (27.0)	9 (11.4)	122 (20.1)
Sex					
Male	270 (80.8)	45 (80.4)	103 (75.2)	69 (87.3)	487 (80.4)
Female	64 (19.2)	11 (19.6)	34 (24.8)	10 (12.7)	119 (19.6)
Tumour stage					
0–I	69 (20.7)	12 (21.4)	23 (16.8)	10 (12.7)	114 (18.8)
II	100 (29.9)	12 (21.4)	42 (30.7)	24 (30.4)	178 (29.4)
III	125 (37.4)	26 (46.4)	58 (42.3)	36 (45.6)	245 (40.4)
IV	40 (12.0)	6 (10.7)	14 (10.2)	9 (11.4)	69 (11.4)
Histology					
Adenocarcinoma	263 (78.7)	43 (76.8)	96 (70.1)	55 (69.6)	457 (75.4)
Squamous cell carcinoma	71 (21.3)	13 (23.2)	41 (29.9)	24 (30.4)	149 (24.6)
Comorbidity					
None	126 (37.7)	26 (46.4)	51 (37.2)	33 (41.8)	236 (38.9)
One	117 (35.0)	21 (37.5)	49 (35.8)	25 (31.7)	212 (35.0)
More than one	91 (27.3)	9 (16.1)	37 (27.0)	21 (26.6)	158 (26.1)
Surgical volume (operations/year)					
Low (<8)	165 (49.4)	30 (53.6)	80 (58.4)	43 (54.4)	318 (52.5)
High (≥8)	169 (50.6)	26 (46.4)	57 (41.6)	36 (45.6)	288 (47.5)
Postoperative complications					
None	215 (64.4)	35 (62.5)	85 (62.0)	46 (58.2)	381 (62.9)
One	85 (25.5)	14 (25.0)	28 (20.4)	17 (21.5)	144 (23.8)
More than one	34 (10.2)	7 (12.5)	24 (17.5)	16 (20.3)	81 (13.4)
Mortality, 90 days					
Within 90 days	27 (8.1)	5 (8.9)	11 (8.0)	5 (6.3)	48 (7.9)
Overall within 5 years	248 (74.3)	35 (62.5)	106 (77.4)	66 (83.5)	455 (75.1)
Conditional within 5 years†	221 (72.0)	30 (58.8)	95 (75.4)	61 (82.4)	407 (72.9)

*Previously married was defined as patients who have been married before but are living alone (after the death of the partner or divorce).

†Conditional mortality: excluding first 90 days after surgery.

Table 2 Multivariable Cox regression models analysing the association between the marital status at the time of diagnosis and mortality after oesophagectomy for cancer, expressed as HRs with 95% CIs

	Marital status			
	Married (reference)	Remarried HR (95% CI)	Previously married* HR (95% CI)	Never married HR (95% CI)
Overall 5-year mortality				
Model 1	1	0.77 (0.54 to 1.10)	1.09 (0.87 to 1.37)	1.14 (0.87 to 1.49)
Model 2	1	0.79 (0.55 to 1.13)	0.95 (0.75 to 1.20)	1.03 (0.78 to 1.36)
Model 3	1	0.79 (0.55 to 1.13)	0.90 (0.71 to 1.15)	1.02 (0.77 to 1.35)
Conditional† overall 5-year mortality				
Model 1	1	0.73 (0.50 to 1.07)	1.11 (0.87 to 1.41)	1.18 (0.89 to 1.57)
Model 2	1	0.73 (0.50 to 1.08)	0.96 (0.75 to 1.23)	1.04 (0.78 to 1.38)
Model 3	1	0.74 (0.50 to 1.08)	0.93 (0.72 to 1.20)	1.05 (0.78 to 1.41)
Disease-specific‡ 5-year mortality				
Model 1	1	0.79 (0.56 to 1.13)	1.11 (0.88 to 1.40)	1.19 (0.89 to 1.57)
Model 2	1	0.80 (0.56 to 1.15)	0.96 (0.76 to 1.22)	1.03 (0.77 to 1.34)
Model 3	1	0.80 (0.56 to 1.15)	0.92 (0.72 to 1.17)	1.04 (0.78 to 1.39)
Conditional† disease-specific‡ 5-year mortality				
Model 1	1	0.75 (0.51 to 1.10)	1.12 (0.87 to 1.43)	1.21 (0.90 to 1.63)
Model 2	1	0.74 (0.50 to 1.09)	0.97 (0.76 to 1.25)	1.02 (0.76 to 1.38)
Model 3	1	0.74 (0.51 to 1.10)	0.94 (0.73 to 1.22)	1.05 (0.78 to 1.42)

Values are expressed as HRs.

*Previously married was defined as patients who have been married before but are living alone (after the death of the partner or divorce).

†Conditional mortality: excluding first 90 days after surgery.

‡Disease-specific mortality: oesophageal or junctional cancer as the underlying or contributing cause. Model 1: unadjusted. Model 2: adjusted for sex, age, tumour stage. Model 3: adjusted for sex, age, tumour stage, histology, major complications, comorbidity and surgeon volume.

in the other marital status groups. The fully adjusted HR for the overall 5-year mortality was similar in never married patients compared with married patients (HR 1.02, 95% CI 0.77 to 1.35), and the corresponding HRs were very similar for disease-specific survival and survival after excluding the initial 90 days after surgery (table 2). The fully adjusted HR for the various definitions of 5-year mortality was lower in the remarried group (HR ranging from 0.74 to 0.80) and in the previously married group (HR ranging from 0.90 to 0.94), but no statistically significant associations were identified (table 2). The results for disease-specific survival, and for overall or disease-specific survival after excluding the initial 90 days of surgery, were all similar to the HRs for the overall survival (table 2). The fully adjusted HR for the overall mortality in the not currently married group is 0.98 (95% CI 0.80 to 1.19) compared with the currently married group.

The fully adjusted subgroup analyses for all-cause and disease-specific mortality by tumour stage and histological type showed similar results to the main analyses, and no statistically significant associations were identified (data not shown).

DISCUSSION

This study did not show any improved survival among married patients with oesophageal cancer undergoing oesophagectomy compared with other marital status groups after adjustment for several prognostic variables.

The main strength of this study is the population-based design with almost complete coverage, and complete

follow-up of all patients operated in Sweden for oesophageal cancer in 2001–2005. Moreover, complete data on a large variety of variables enabled adjustment for known prognostic factors. Yet, among limitations are that the influence of other confounders or residual confounding by the variables adjusted for can never be ruled out. The statistical power was limited to detect weak associations, but large effects of the marital status on survival in this cohort of patients with cancer are unlikely. Changes in the marital status after surgery were considered of limited impact on the results, since these only occurred in 3.3% of all patients. Moreover, we did not have data on cohabiting without being married, or the extent of social networks, which could have led to an underestimation of potential associations. Despite the multicentre design, differences between hospitals and treatment are unlikely to have influenced any associations between marital status and survival. Socioeconomic variables such as ethnicity and income level may have an impact on survival as well, yet the underlying mechanisms are different. Socioeconomic inequality is a measure for lifetime differences, whereas the marital status reflects social support at the time of the disease.

The absence of a clear association between marital status and long-term survival, such as that found for some other cancer types and oesophageal cancer in the large American study mentioned earlier (including all diagnosed patients),^{3 8} might be a false-negative result related to the relatively small sample size. Yet, it is also possible that the beneficial effect of being married is non-existent for patients with oesophageal cancer receiving surgery,

which has a limited chance of survival despite its curative intent.⁹ Although being married might not influence the survival, social support might be beneficial for other reasons, including the health-related quality of life in these patients. To conclude, this study showed no evidence of improved survival of married patients compared with non-married patients after having undergone surgery for oesophageal cancer.

Contributors All the authors have made substantial contributions to conception and design of the article. JL and PL were responsible for the acquisition of the data. NB, FM and AJ performed the statistical analyses and interpretation of the data. NB has drafted the manuscript, and all the authors revised it critically for important intellectual content. All the authors have given final approval of the version to be published.

Funding This work was supported by the Swedish Research Council (SIMSAM) and the Swedish Cancer Society (Contract number: 13 0537).

Competing interests None.

Patient consent Obtained.

Ethics approval The Regional Ethical Review Board in Stockholm, Sweden.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- Berkson J. Mortality and marital status. Reflections on the derivation of etiology from statistics. *Am J Public Health Nations Health* 1962;52:1318–29.
- Rendall MS, Weden MM, Favreault MM, *et al.* The protective effect of marriage for survival: a review and update. *Demography* 2011;48:481–506.
- Bailey J. Effects of marital status on cancer incidence and survival rates. *Am Fam Physician* 2009;80:1052; author reply 52.
- Kravdal H, Syse A. Changes over time in the effect of marital status on cancer survival. *BMC Public Health* 2011;11:804.
- Thrift AP, Nagle CM, Fahey PP, *et al.* Predictors of survival among patients diagnosed with adenocarcinoma of the esophagus and gastroesophageal junction. *Cancer Causes Control* 2012;23:555–64.
- Gossage JA, Forshaw MJ, Khan AA, *et al.* The effect of economic deprivation on oesophageal and gastric cancer in a UK cancer network. *Int J Clin Pract* 2009;63:859–64.
- Brusselsaers N, Ljung R, Mattsson F, *et al.* Education level and survival after oesophageal cancer surgery: a prospective population-based cohort study. *BMJ* 2013;3:e003754.
- Aizer AA, Chen MH, McCarthy EP, *et al.* Marital status and survival in patients with cancer. *J Clin Oncol* 2013;31:3869–76.
- Lagergren J, Lagergren P. Recent developments in esophageal adenocarcinoma. *CA Cancer J Clin* 2013;63:232–48.
- Lagergren J, Bergström R, Lindgren A, *et al.* Symptomatic gastroesophageal reflux as a risk factor for esophageal adenocarcinoma. *N Engl J Med* 1999;340:825–31.
- Viklund P, Lindblad M, Lu M, *et al.* Risk factors for complications after esophageal cancer resection: a prospective population-based study in Sweden. *Ann Surg* 2006;243:204–11.
- Lagergren K, Derogar M. Validation of oesophageal cancer surgery data in the Swedish Patient Registry. *Acta Oncol* 2012;51:65–8.
- Johansson LA, Westerling R. Comparing Swedish hospital discharge records with death certificates: implications for mortality statistics. *Int J Epidemiol* 2000;29:495–502.
- Statistics-Sweden. Longitudinell integrationsdatabas för Sjukförsäkrings- och Arbetsmarknadsstudier (LISA) 1990–2009. 2011.
- Brusselsaers N, Mattsson F, Lagergren J. Hospital and surgeon volume in relation to long-term survival after oesophagectomy: systematic review and meta-analysis. *Gut* 2013. Published Online First: 22 Nov 2013. doi:10.1136/gutjnl-2013-306074
- Sobin L, Wittekind C. *International Union Against Cancer (UICC): classification of malignant tumours*. 6th edn, 2002.
- Backemar L, Djärv T, Wikman A, *et al.* The role of diabetes and other co-morbidities on survival after esophageal cancer surgery in a population-based study. *Am J Surg* 2013;206:539–43.