Hindawi Publishing Corporation Mediators of Inflammation Volume 2014, Article ID 513027, 2 pages http://dx.doi.org/10.1155/2014/513027

## **Erratum**

## Erratum to "Palmitic Acid Induces Production of Proinflammatory Cytokines Interleukin-6, Interleukin-1 $\beta$ , and Tumor Necrosis Factor- $\alpha$ via a NF- $\kappa$ B-Dependent Mechanism in HaCaT Keratinocytes"

## Bing-rong Zhou, Jia-an Zhang, Qian Zhang, Felicia Permatasari, Yang Xu, Di Wu, Zhi-qiang Yin, and Dan Luo

Department of Dermatology, The First Affiliated Hospital of Nanjing Medical University, Nanjing 210029, China

Correspondence should be addressed to Dan Luo; daniluo2013@njmu.edu.cn

Received 15 March 2014; Accepted 27 April 2014; Published 21 May 2014

Copyright © 2014 Bing-rong Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the paper, the 3 bottom photos in Figure 2(a) were the same as the 3 bottom photos in Figure 2(b), and here we provide the right form of Figure 2(a). We apologize for this oversight and for any confusion that it has caused.

2 Mediators of Inflammation




FIGURE 2: PA-induced nuclear translocation of NF- $\kappa$ B p65 subunit in HaCaT cells immediately after the treatment and 24 hours after the PA depletion. Keratinocytes were untreated or treated with PA (75, 100, 125, and 150  $\mu$ mol/L) for 24 hours under serum-free conditions. Immunostaining was performed with specific mouse anti-p65 antibody followed by Cy3-conjugated mouse anti-rabbit immunoglobulins (red) immediately after the treatment (a) and 24 hours after the PA depletion (b). Images are representative of three independent experiments.