

Identification of genes preventing transgenerational transmission of stress-induced epigenetic states

Mayumi Iwasaki^{a,b,1,2} and Jerzy Paszkowski^{a,b,1,2}

^aDepartment of Plant Biology, University of Geneva, Sciences III, CH-1211 Geneva 4, Switzerland; and ^bThe Sainsbury Laboratory, University of Cambridge, Cambridge CB2 1LR, United Kingdom

Edited* by Jeffery L. Dangl, Howard Hughes Medical Institute and The University of North Carolina, Chapel Hill, NC, and approved April 16, 2014 (received for review February 5, 2014)

Examples of transgenerational transmission of environmentally induced epigenetic traits remain rare and disputed. Abiotic stress can release the transcription of epigenetically suppressed transposons and, noticeably, this activation is only transient. Therefore, it is likely that mechanisms countering the mitotic and meiotic inheritance of stress-triggered chromatin changes must exist but are undefined. To reveal these mechanisms, we screened for *Arabidopsis* mutants impaired in the resetting of stress-induced loss of epigenetic silencing and found that two chromatin regulators, *Decrease in DNA methylation1* (*DDM1*) and *Morpheus' Molecule1* (*MOM1*), act redundantly to restore prestress state and thus erase "epigenetic stress memory". In *ddm1* mutants, stress hyperactivates heterochromatic transcription and transcription persists longer than in the wild type. However, this newly acquired state is not transmitted to the progeny. Strikingly, although stress-induced transcription in *mom1* mutants is as rapidly silenced as in wild type, in *ddm1 mom1* double mutants, transcriptional signatures of stress are able to persist and are found in the progeny of plants stressed as small seedlings. Our results reveal an important, previously unidentified function of *DDM1* and *MOM1* in rapid resetting of stress induced epigenetic states, and therefore also in preventing their mitotic propagation and transgenerational inheritance.

chromatin regulation | DNA methylation

Although environmentally induced traits and their transgenerational transmission in plants have been described repeatedly, trait stability and the involvement of epigenetic mechanisms in their generation remain controversial (1–3). In contrast, it has been well documented that environmental challenges such as elevated temperature can transcriptionally activate chromosomal loci normally silenced by repressive chromatin (4–7). However, this release of epigenetic silencing, unaccompanied by changes in DNA methylation or histone modifications, is only transient (4, 5). Such noncanonical release of transcriptional gene silencing (TGS) is similar to alterations in epigenetic regulation observed in the *mom1* mutant, where release of TGS occurs without major changes in epigenetic marks (8–13). Although, molecular mechanisms used by *MOM1* in TGS regulation are not well understood, genetic studies have linked *MOM1* activity to small interfering RNAs (14) and RNA processing (15). In addition, structure/function studies have suggested that a conserved domain of *MOM1* forms a homodimer, which is possibly required as a binding platform for additional silencing factors (13, 16).

The rapid resilencing of heterochromatic transcription induced by heat stress seems to involve changes in nucleosome occupancy and resilencing is delayed in mutants with impaired chromatin assembly (5). These observations suggest that suppressive chromatin has certain plasticity in response to stress, but also a robust buffering system that resets its prestress state. This counters the persistence of stress-induced epigenetic alterations during subsequent development and thus their transmission to the progeny.

Results and Discussion

To identify factors involved in the erasure of "epigenetic stress memory," we performed a genetic screen using *Arabidopsis* line *LUC25* carrying a transcriptionally silenced transgene encoding firefly luciferase (*LUC*) (14), which as an endogenous chromosomal TGS target loci can be transiently activated after heat stress. First, we introduced the *mom1* mutation into *LUC25* (*mom1 LUC25*). The *mom1* mutation partially releases silencing of *LUC25*, producing weak luciferase signals in roots but not aerial parts, where the *LUC* transgene remains silenced. Importantly, the *LUC* transgene in *mom1 LUC25* is strongly activated by heat stress, similar to *LUC25* (Fig. S1A). We presumed that the introduction of the *mom1* mutation would enhance stress-induced luciferase signals, increasing clarity and thus the efficiency of the mutant screen. Moreover, although the *mom1* mutation does not directly influence the kinetics of stress-induced TGS release, *MOM1* is involved in buffering epigenetic states of chromatin (11). We hypothesized, therefore, that any deficiency in such buffering would facilitate phenotypic detection of additional epigenetic regulators involved in the rapid restoration of TGS after stress and, thus, in the erasure of epigenetic stress memory.

M2 seedlings of mutagenized *mom1 LUC25* were germinated for 5 d and individuals showing enhanced luciferase signals before stress treatment were removed, because these plants release TGS constitutively. The remaining seedlings were subjected to heat stress and plants showing significantly stronger and/or longer-lasting luciferase signals were selected and grown to maturity (Fig. S1B). We examined their M3 progeny to determine whether selected

Significance

Much attention has been drawn to research, which suggests that environmental factors, including stress and maternal care, alter the way the genetic code is executed. These epigenetic changes in gene regulation are thought to be stable enough to be heritable and thus may influence subsequent generations. Such prospects are as intriguing as they are troubling, since it is possible to imagine that accumulation of stress memories over several generations could make life decisions difficult. Therefore, we have questioned whether mechanisms exist that could prevent the inheritance of stress-induced epigenetic changes and have discovered such mechanisms using forward genetics in *Arabidopsis*. Interestingly, one of the critical activities erasing stress memories is conserved between plants and mammals.

Author contributions: M.I. and J.P. designed research; M.I. performed research; M.I. analyzed data; and M.I. and J.P. wrote the paper.

The authors declare no conflict of interest.

*This Direct Submission article had a prearranged editor.

¹Present address: The Sainsbury Laboratory, University of Cambridge, Bateman Street, Cambridge CB2 1LR, United Kingdom.

²To whom correspondence may be addressed E-mail: jerzy.paszkowski@slcu.cam.ac.uk or mayumi.iwasaki@slcu.cam.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1402275111/-DCSupplemental.

phenotypes were heritable. Interestingly, several plants selected in the M2 produced progeny uniformly showing high luciferase signals *prior* to heat stress in the M3. Because such “constitutive” phenotypes had been discarded in the previous M2 generation, we concluded that their appearance in the M3 reflects transgenerational transmission of heat stress-induced TGS release that occurred in the previous plant generation (Fig. 1A). In other words, we may have recovered mutant plants severely impaired in the erasure of epigenetic stress memory. Focusing on four independent lines with these characteristics, we identified causal mutations by a combination of genetic mapping and whole genome sequencing. Two independent mutations resided in a gene encoding nucleosome remodeler DDM1 (17–20) (Fig. 1B). The DDM1 protein, conserved between plants and mammals, is required for maintenance of DNA methylation, thus TGS (21, 22). It has been suggested that DDM1 alters accessibility of H1-containing heterochromatin to DNA methyltransferases (23).

Recovery of *ddm1* mutants in our screen was both surprising and disturbing. Surprising, because *ddm1* mutants are known to

release epigenetic silencing independently of stress and, therefore, should have been eliminated from the screen in the M2. Disturbing, because *ddm1* mutants have a transgenerationally progressive effect on the loss of DNA methylation and silencing release (18, 19). Thus, luciferase activity observed in the M3 could simply reflect this *ddm1* property rather than transgenerational memory of heat stress. To address these reservations, we first analyzed DNA methylation of the *LUC* promoter. Cytosine methylation patterns were only slightly altered, however, surprisingly high levels of methylation remained in the M3 of both mutant lines (Fig. 1C and D), which is unusual for *ddm1* mutants strictly associated with hypomethylation-mediated TGS release (17, 20, 24, 25). This result supported the possibility that the heat stress-activated state of the *LUC* transgene, which is independent of DNA methylation (4, 5), can in fact be maintained and transmitted to the next generation in the *ddm1* background, and therefore, would define a previously unrecognized and potentially crucial activity of DDM1 in reversing TGS after it has been destabilized by environmental changes/stress.

The only way to test this hypothesis was by the recreation and analysis of the *ddm1* mutant line with a naïve and, thus, still-silenced *LUC* transgene (Fig. 2A). To obtain such a line, we crossed the commonly used allele of *ddm1* (*ddm1-2*) (18) into *LUC25* and subjected the F2-segregating progeny to temperature stress. Importantly, under control growth conditions, luciferase signals remained at the *LUC25* level and no differences between segregating F2 individuals were recorded (Fig. 2B). This suggested that the *LUC* transgene remained silent in the first generation of *ddm1* mutants and explains the initially unexpected presence of *ddm1* among plants subjected to heat stress during the mutant screen. Notably, heat stress applied to segregating F2 seedlings revealed individuals with very strong luciferase signals (Fig. 2B) in proportion close to expected segregation ratio for plants homozygous *ddm1* mutation containing *LUC* plants (18.75%). The genotyping of the segregating population at *DDM1* and *LUC* loci confirmed that, all of these plants were found to be homozygous for the *ddm1-2* mutation and contained *LUC* transgene in homozygous or hemizygous state, which had no influence on the intensity of LUC signals. Furthermore, we performed an analogous genetic experiment introducing the *ddm1-2* allele into another line (L5) (26) carrying a silent transgenic locus for the glucuronidase marker gene (*GUS*). As with the *LUC* transgene, we observed heat-stress-dependent hyperactivation of *GUS* expression in *ddm1-2* mutants (Fig. S2). These results demonstrated that DDM1 down-regulates stress-induced heterochromatic transcription. Moreover, this novel *DDM1* activity appears to be independent of changes in DNA methylation, with which so far *DDM1* was very tightly associated.

However, we found that in *ddm1-2* mutants the stress-hyperactivated *LUC* transgene was resiled within a few days (Fig. 2B) and there was no difference in LUC signals in the progeny (F3 generation) between stressed and nonstressed *ddm1-2 LUC25* plants (Fig. 2D).

Considering the genetic screen was performed in the *mom1 LUC25* background, we repeated the genetic reconstruction experiment including the *ddm1-2* mutation. *ddm1-2* was crossed with *mom1 LUC25* and stress-induced LUC activation in the segregating F2 populations was examined as before, as well as its inheritance in the F3 (Fig. S3). LUC phenotyping and subsequent genotyping of F2 plants showed that the *mom1* mutation alone did not affect stress-induced expression of *LUC*, confirming the previous observations with the *mom1 LUC25* line. Although, the stress-induced LUC activation levels in the *ddm1-2 mom1* double mutants were similar to those in the *ddm1-2* single mutants (Fig. S4A), however, the LUC signals remained high only in *ddm1-2 mom1* double mutants (Fig. 2C). Furthermore, in the next (F3) generation, progeny from stressed *ddm1-2 mom1 LUC25* plants showed significantly higher LUC signals than nonstressed *ddm1-2*




Fig. 1. Identification and characterization of mutants showing transgenerational transmission of the heat-stress release of TGS. (A) Bioluminescence images of the progeny (M3 and M4) of two mutant candidates (9.2.1 and 62.2.1) and of the controls (*LUC25*, and *mom1 LUC25*), all grown under control conditions at 21 °C. The two mutant lines are M3 progeny of heat stressed M2 parents recovered from the mutant screen. The green and red signals are luciferase luminescence and auto-fluorescence of chlorophyll, respectively. (B) Two new mutant alleles of the *DDM1* gene (9.2.1 and 62.2.1) were identified in the screen. The *ddm1-2* allele was used for the experiments presented in Fig. 2. (C) DNA methylation distribution at the ubiquitin3 promoter of the *LUC* transgene in 9.2.1 (M3), 62.2.1 (M3), and *LUC 25* (WT). Colored and open circles represent methylated and unmethylated cytosines respectively, with red representing CG sites, blue CHG and green CHH (H can be A, T, or C). (D) Percentage of cytosine methylation in the ubiquitin3 promoter. Black bars, 9.2.1; gray bars, 62.2.1; white bars, *LUC25*.




Fig. 2. Inheritance of stress-induced transcriptional activation of *LUC*. (A) Crossing scheme for the recreation of the *ddm1-2* mutant line with the naïve *LUC* transgene. P0: *ddm1-2* was crossed with WT *LUC25*, F1: heterozygous for *ddm1* and carrying the hemizygous naïve *LUC* transgene, F2: *ddm1* homozygous mutants are segregated in the progeny. F2 seedlings were separated into two subpopulations, one of which was subjected to heat stress. Bioluminescence images were captured (B), and each plant was genotyped at the *DDM1* and *LUC* loci. (B) Bioluminescence images of segregating progeny of a hybrid between *ddm1-2* and *LUC25* (A). The F2 seedlings were expected to include 18.75% of individuals homozygous for the *ddm1-2* mutation and carrying the *LUC* transgene, these were predicted to display enhanced luminescence. Rows of *LUC25* and *mom1 LUC25* plants are shown as a control. Note, *LUC* signals in *mom1* after heat stress are restricted to roots. (C) Bioluminescence images of segregating progeny of a cross between *ddm1-2 mom1 LUC25* (for details, see Fig. S3). The F2 progeny is expected to include 4.69% of individuals homozygous for both *ddm1-2* and *mom1* and carrying the *LUC* transgene. White arrowheads point toward *ddm1-2 mom1* double mutant plants, as revealed by the genotyping of the population at *DDM1* and *MOM1* loci. Quantification of *LUC* signals is displayed in Fig. S4. (D) Bioluminescence images of *ddm1 LUC25* and *ddm1 mom1 LUC25* F3 progenies (as depicted in A) of F2 heat stressed parents at the seedlings stage (+stress) or nonstressed controls (–stress). Two independent F3 populations (upper or lower row) derived from two *ddm1 LUC25* or *ddm1 mom1 LUC25* F2 plants (B and C).

mom1 LUC25 controls (Fig. 2D and Fig. S4B), indicating that the stress-induced active state of the *LUC* transgene initiated at the small seedling stage could persist throughout plant development and be transgenerationally transmitted, however only in *ddm1 mom1* double mutants.

To further investigate stability and possible transgenerational inheritance in *ddm1 mom1* double mutants of stress-triggered TGS release, we examined transcriptional changes at endogenous chromosomal loci. For this, we first performed RNA-seq analyses on four independent populations of the progeny of stressed and control-treated *ddm1-2 mom1* double mutant plants. This genome-wide approach should uncover chromosomal loci stably activated following the stress subjected to the parental plants. The MDS plot analysis of the RNA-seq data revealed well-clustered biological repetitions for each individual line (Fig. 3A), indicating

the robustness of the RNA-seq data. Noticeably, the largest difference between biological repetitions was seen for *ddm1 mom1* double mutant seedlings derived from ancestors not subjected to stress. This may simply reflect an intrinsic property of *ddm1 mom1* double mutants, which exhibit variation in heterochromatin silencing among individual plants (27). Most importantly, the samples were clearly clustered according to whether seedlings of the previous generation were stressed or not (Fig. 3A). Therefore, the genome-wide transcriptional profiles supported and extended our previous conclusion based on transgenic loci that temperature stress-activated transcription occurs genome wide, and that newly acquired transcriptional signatures can be transgenerationally inherited in *ddm1 mom1* double mutants.

Of the loci affected in the progeny of stressed *ddm1-2 mom1*, compared with the progeny of control-treated *ddm1-2 mom1*, 340




Fig. 3. Genome-wide analysis of transcriptional changes in the progeny of heat-stressed *ddm1 mom1*. (A) Multidimensional scaling (MDS) plot (R/Bioconductor) showing the overall similarity of RNA expression patterns between samples using RNA-seq data of two biological repetitions. These were performed with two independent populations (circled) of *ddm1 mom1* progeny plants obtained from stressed or nonstressed parents at the seedling stage (*mom1* transcriptome was used as an additional control). (B) Pie charts showing the difference in functional distribution of up- and down-regulated loci between heat-stressed or non-stressed double mutants. (C) Relative levels of mRNA of selected targets as determined by quantitative RT-PCR. Values were normalized to 18s ribosomal RNA. One sample of stressed *ddm1 mom1* was set to 1. White and gray bars indicate the progeny of control plants and heat-stressed plants, respectively. Pooled F3 plants (~20 individuals), progeny of two independent F2 plants for each category were used for these analyses. Gene annotations of each target: AT1G43880 - ATLANTY1 (TE), AT2G05564 - VANDAL2 (TE), AT5G29560 - caleosin-related family protein, AT5G34790 - VANDAL20 (TE), AT2G12345 - ATHILA3 (TE), AT5G48850 - ATSD1, SULFUR DEFICIENCY-INDUCED 1. Error bars indicate SD of results from three repeated experiments. *P < 0.01, Student *t* test.

loci were up-regulated more than twofold ($P < 0.01$) and 484 down-regulated less than twofold ($P < 0.01$) (Fig. 3B). Approximately 60% of the up-regulated transcripts and 20% of the down-regulated transcripts are derived from transposable elements (TEs) (Fig. 3B). These results are consistent with our previous demonstration that heterochromatic regions enriched in TEs are predominantly transcriptionally activated by temperature stress, and that euchromatic regions are either activated or repressed by this treatment (4). Such a transcriptional signature of the genome-wide stress-induced alteration of transcription appears to be inherited by the progeny of *ddm1 mom1* double mutant plants.

Due to economic constraints, we refrained from genome-wide analyses of plant populations constituting various experimental controls. Instead, using the transcriptional profiling data described above we selected several genomic loci displaying heritable stress-induced alteration of transcription in *ddm1 mom1* double mutants, and examined by quantitative RT-PCR persistence of their transcriptional activation in WT and single *mom1* or *ddm1* mutant plants relative to stress treatments in the preceding generation. No significant differences in transcript levels were found between the progenies of stressed and control WT plants or single mutants, which is in contrast to *ddm1 mom1* double mutants (Fig. 3C). This result supports our previous conclusion, derived from the properties of transgenic loci, that DDM1 and MOM1 both act redundantly in resetting chromatin status destabilized by heat stress, which prevents transgenerational propagation of transcriptional stress memory (Fig. 4).

A closer look at the results of the genome-wide transcriptional analyses reveals that DDM1 and MOM1 are not the only factors reverting the properties of chromatin affected by stress. We found previously that ~3,000 loci are activated under stress conditions analogous to those used here (4) and, thus, only a small fraction (340) remain transgenerationally active in the *ddm1 mom1* double mutants. This result suggests that the prevention of transgenerational transmission of stress memory extends far beyond the activities of DDM1 and MOM1 and, thus, that the unequivocal demonstration of transgenerational transmission of environmentally-induced epigenetic traits remains a significant challenge.

Materials and Methods

Plant Materials. The *LUC25* line was described (14). *mom1 LUC25* was obtained by crossing *mom1-6* and *LUC25*. *mom1-6* seeds were obtained from INRA-Versailles, Genomic Resource Center (FLAG_340E12) and *ddm1-2* seeds were provided by E. Richards (Boyce Thompson Institute for Plant Research, Cornell University, Ithaca, NY). *LUC25* and *mom1-6* are in the Wassilewskija (WS) background and *ddm1-2* in the Columbia (Col-0) background.

Mutagenesis and Screening. *mom1 LUC25* seeds (20,000) were mutagenized in 0.3% EMS for 15 h. After washing with water, seeds were germinated on soil to give 78 M2 pools, each derived from ~150 M1 plants. For each pool, 1,000 seeds were plated on 1/2 MS medium (0.8% agar, 1% sucrose) and screened for mutants by spraying with a luciferin (Biosynth) solution (31.5 mg per 100 mL of water) and examining treated seedlings using an Aequoria dark box with a mounted ORCAII CCD camera (Hamamatsu). Luciferase luminescence and chlorophyll auto-fluorescence image overlays were created using the Wasabi software package (Hamamatsu). Isolated mutants were crossed with WT in the Col-0 background to generate mapping populations.

DNA Methylation Analysis. Bisulphite sequencing was performed as described (14). Sequencing data were analyzed with Kismeth (<http://katahdin.mssm.edu/kismeth>) (28). Primers used for bisulphite sequencing are listed in Table S1.

Histochemical GUS Staining. Staining was performed on cotyledons of 12-d-old plants as described (13).




Fig. 4. Model summarizing a putative epigenetic mechanism preventing the transmission of “stress memory” to progeny. Schematic illustration on a possible chromatin states upon heat stress. Although heterochromatic loci are transcriptionally activated by temperature stress in WT and *mom1* mutants, they are rapidly resiled after stress is removed. Stress-induced transcription is hyper-activated and persists longer in *ddm1* mutants than in WT and *mom1*; however, the altered transcriptional status is not transmitted to the progeny. In contrast, stress-induced transcriptional activation in *ddm1 mom1* double mutants is transgenerationally inherited. The transcriptional activation in *ddm1 mom1* could persist due to altered positioning of nucleosome or other modifications of chromatin properties.

RAN-seq Analyses. Total RNA samples were isolated from 7-d-old seedlings using TRI reagent (Sigma). The libraries were prepared using a TruSeq RNA Sample Prep Kit (Illumina) and sequenced using HiSeq. 2500 (Illumina) with single-end 50 base reads. The trimmed reads were mapped with the TopHat v1.3.3 software to the TAIR10 annotations. The normalization and differential expression analysis were performed with R/Bioconductor package edgeR v2.6.12 (29).

Real-Time Quantitative PCR Analysis. Total RNA was isolated from 7-d-old seedlings using TRI reagent (Sigma). After RQ1 DNase treatment (Promega), cDNA was synthesized with the SuperScript VILO cDNA synthesis kit (Life Technologies). Real-time quantitative PCR analyses were performed using Power SYBR Green PCR Master Mix (Life Technologies) in ABI7900FT (Life Technologies). PCR conditions were 95 °C for 10 min followed by 40 cycles of 95 °C for 15 s and 60 °C for 60 s. Primers are listed in Table S1.

ACKNOWLEDGMENTS. We thank Taisuke Nishimura for preparing EMS-mutagenized *mom1 LUC25* seeds, Christian Megies for technical assistance, Mylène Docquier and Céline Delucinge for help with RNA-seq analysis, Detlef Weigel and Beth Rowan for help with whole genome sequencing analysis, and Patrick King for editing of the manuscript. We also thank all members of the J.P. laboratory for constructive discussions. This work was supported by grants from the Swiss National Science Foundation (31003A-125005), the European Commission through the AENEAS collaborative project (FP7 226477), the Gatsby Charitable Foundation, and the European Research Council.

- Paszkowski J, Grossniklaus U (2011) Selected aspects of transgenerational epigenetic inheritance and resetting in plants. *Curr Opin Plant Biol* 14(2):195–203.
- Boyko A, Kovalchuk I (2011) Genome instability and epigenetic modification—heritable responses to environmental stress? *Curr Opin Plant Biol* 14(3):260–266.

- Mirouze M, Paszkowski J (2011) Epigenetic contribution to stress adaptation in plants. *Curr Opin Plant Biol* 14(3):267–274.
- Tittel-Elmer M, et al. (2010) Stress-induced activation of heterochromatic transcription. *PLoS Genet* 6(10):e1001175.

5. Pecinka A, et al. (2010) Epigenetic regulation of repetitive elements is attenuated by prolonged heat stress in Arabidopsis. *Plant Cell* 22(9):3118–3129.
6. Lang-Mladek C, et al. (2010) Transgenerational inheritance and resetting of stress-induced loss of epigenetic gene silencing in Arabidopsis. *Mol Plant* 3(3): 594–602.
7. Ito H, et al. (2011) An siRNA pathway prevents transgenerational retrotransposition in plants subjected to stress. *Nature* 472(7341):115–119.
8. Amedeo P, Habu Y, Afsar K, Mittelsten Scheid O, Paszkowski J (2000) Disruption of the plant gene MOM releases transcriptional silencing of methylated genes. *Nature* 405(6783):203–206.
9. Steimer A, et al. (2000) Endogenous targets of transcriptional gene silencing in Arabidopsis. *Plant Cell* 12(7):1165–1178.
10. Probst AV, Fransz PF, Paszkowski J, Mittelsten Scheid O (2003) Two means of transcriptional reactivation within heterochromatin. *Plant J* 33(4):743–749.
11. Habu Y, et al. (2006) Epigenetic regulation of transcription in intermediate heterochromatin. *EMBO Rep* 7(12):1279–1284.
12. Vaillant I, Schubert I, Tourmente S, Mathieu O (2006) MOM1 mediates DNA-methylation-independent silencing of repetitive sequences in Arabidopsis. *EMBO Rep* 7(12): 1273–1278.
13. Nishimura T, et al. (2012) Structural basis of transcriptional gene silencing mediated by Arabidopsis MOM1. *PLoS Genet* 8(2):e1002484.
14. Yokthongwattana C, et al. (2010) MOM1 and Pol-IV/V interactions regulate the intensity and specificity of transcriptional gene silencing. *EMBO J* 29(2):340–351.
15. Zhou Y, Zhang J, Lin H, Guo G, Guo Y (2010) MORPHEUS' MOLECULE1 is required to prevent aberrant RNA transcriptional read-through in Arabidopsis. *Plant Physiol* 154(3):1272–1280.
16. Caikovski M, et al. (2008) Divergent evolution of CHD3 proteins resulted in MOM1 refining epigenetic control in vascular plants. *PLoS Genet* 4(8):e1000165.
17. Vongs A, Kakutani T, Martienssen RA, Richards EJ (1993) Arabidopsis thaliana DNA methylation mutants. *Science* 260(5116):1926–1928.
18. Kakutani T, Jeddeloh JA, Flowers SK, Munakata K, Richards EJ (1996) Developmental abnormalities and epimutations associated with DNA hypomethylation mutations. *Proc Natl Acad Sci USA* 93(22):12406–12411.
19. Jeddeloh JA, Bender J, Richards EJ (1998) The DNA methylation locus DDM1 is required for maintenance of gene silencing in Arabidopsis. *Genes Dev* 12(11):1714–1725.
20. Jeddeloh JA, Stokes TL, Richards EJ (1999) Maintenance of genomic methylation requires a SWI2/SNF2-like protein. *Nat Genet* 22(1):94–97.
21. Bourc'his D, Bestor TH (2002) Helicase homologues maintain cytosine methylation in plants and mammals. *Bioessays* 24(4):297–299.
22. Tao Y, et al. (2011) Lsh, chromatin remodeling family member, modulates genome-wide cytosine methylation patterns at nonrepeat sequences. *Proc Natl Acad Sci USA* 108(14):5626–5631.
23. Zemach A, et al. (2013) The Arabidopsis nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing heterochromatin. *Cell* 153(1):193–205.
24. Kakutani T, Munakata K, Richards EJ, Hirochika H (1999) Meiotically and mitotically stable inheritance of DNA hypomethylation induced by ddm1 mutation of Arabidopsis thaliana. *Genetics* 151(2):831–838.
25. Lippman Z, et al. (2004) Role of transposable elements in heterochromatin and epigenetic control. *Nature* 430(6998):471–476.
26. Morel JB, Mourrain P, Béclin C, Vaucheret H (2000) DNA methylation and chromatin structure affect transcriptional and post-transcriptional transgene silencing in Arabidopsis. *Curr Biol* 10(24):1591–1594.
27. Mittelsten Scheid O, Probst AV, Afsar K, Paszkowski J (2002) Two regulatory levels of transcriptional gene silencing in Arabidopsis. *Proc Natl Acad Sci USA* 99(21):13659–13662.
28. Grunman E, et al. (2008) Kismeth: Analyzer of plant methylation states through bisulfite sequencing. *BMC Bioinformatics* 9:371.
29. Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1): 139–140.