


Parahydrogen Induced Polarization of 1-¹³C-Phospholactate-*d*₂ for Biomedical Imaging with >30,000,000-fold NMR Signal Enhancement in Water

Roman V. Shchepin,^{*,†} Aaron M. Coffey,^{†,‡} Kevin W. Waddell,^{†,§} and Eduard Y. Chekmenev^{*,†,‡,||,¶}

[†]Vanderbilt University Institute of Imaging Science (VUIIS), Department of Radiology, [‡]Department of Biomedical Engineering and Biochemistry, [§]Department of Physics and Astronomy, ^{||}Department of Biochemistry, [¶]Vanderbilt-Ingram Cancer Center (VICC), Vanderbilt University, Nashville, Tennessee 37232, United States

Supporting Information

ABSTRACT: The synthetic protocol for preparation of 1-¹³C-phosphoenolpyruvate-*d*₂, precursor for parahydrogen-induced polarization (PHIP) of 1-¹³C-phospholactate-*d*₂, is reported. ¹³C nuclear spin polarization of 1-¹³C-phospholactate-*d*₂ was increased by >30,000,000-fold (5.75 mT) in water. The reported ¹³C polarization level approaching unity (>15.6%), long lifetime of ¹³C hyperpolarized 1-¹³C-phospholactate-*d*₂ (58 ± 4 s versus 36 ± 2 s for nondeuterated form at 47.5 mT), and large production quantities (52 μmoles in 3 mL) in aqueous medium make this compound useful as a potential contrast agent for the molecular imaging of metabolism and other applications.



Hyperpolarized (HP) ¹³C magnetic resonance imaging (MRI) has evolved from proof-of-principle studies^{1,2} to clinical trials over the past decade.³ The high sensitivity of HP MRI was the main driving force behind its application for metabolic imaging. Indeed, HP MRI provides several orders of magnitude signal enhancement over the conventional MRI. Furthermore, HP MRI permits fast imaging of HP contrast agents and their metabolites within the time scale of the hyperpolarization decay, e.g., 1–3 min *in vivo*.^{4–6} Monitoring the metabolic signatures of HP contrast agents⁷ allows one to interrogate disease aggressiveness, progression, and response to treatment.

While dissolution dynamic nuclear polarization (DNP) is the most widely used hyperpolarization method at present,¹ parahydrogen-induced polarization (PHIP)⁸ is an alternative approach, which enables HP contrast agent preparation in less than 1 min.^{4,5} PHIP is a chemical method relying on ultrafast molecular hydrogenation of an unsaturated precursor with parahydrogen enabled by Rh(I) catalysts, Figure 1.^{6,9} The use of hyperpolarization techniques for bioimaging requires a relatively long lifetime of the HP state *in vivo*, which historically has been demonstrated with isolated ¹³C sites in biomolecules, e.g., ¹³C carboxyl sites without directly attached protons with *in vivo* ¹³C *T*₁ > 20 s.³ Isotopic enrichment with ~99% ¹³C is mandatory to boost the sensitivity of the contrast agent by increasing the payload of HP ¹³C magnetization per μmole of injected agent. In the PHIP process, the long-lived ¹³C sites *J*-coupled to nascent parahydrogens are hyperpolarized using polarization transfer from the parahydrogens to ¹³C, Figure 1. In addition to the mandatory ¹³C^{10,11} labeling of carboxyl carbon adjacent to the double or triple carbon–carbon bond

required for molecular addition of parahydrogen,^{10,11} the spin system of the hydrogenated molecule often benefits from simplification to three spins via deuteration: two parahydrogen protons and a ¹³C nucleus, Figure 1. This deuteration of the precursor molecules also lengthens the ¹³C relaxation times *T*₁ of HP states.^{12–14}

Despite the synthetic challenges related to deuterium and ¹³C enrichments, several contrast agents such as 2-hydroxyethyl propionate-1-¹³C,2,3,3-*d*₃ (HEP), 1-¹³C-succinate,^{12–14} tetrafluoropropyl propionate-1-¹³C,2,3,3-*d*₃ (TFPP),^{15,16} ¹⁵N-propargylcholine,¹⁷ and 1-¹³C-phospholactate¹⁸ have been demonstrated. HEP,¹⁹ 1-¹³C-succinate-*d*₂,^{12–14} and TFPP^{15,16} have been used for *in vivo* metabolic imaging of vasculature, cancer, and plaque lipid content, respectively.

The recently introduced concept of –OH protection of the C=C–OH moiety^{18,20} expanded the number of potential PHIP-based HP contrast agents. HP 1-¹³C-phospholactate (PLAC)^{18,21} is particularly attractive, Figure 1, because it can be rapidly dephosphorylated to HP 1-¹³C-lactate by blood phosphatases.²¹ Therefore, bioavailable 1-¹³C-lactate can be prepared by the PHIP method. HP lactate^{22–25} elevation in cancer cells enables *in vivo* metabolic imaging using injection of HP 1-¹³C-lactate.^{26,27}

A low polarization level of ~1% has been the main obstacle for use of HP PLAC, Figure 1. Here, a significantly greater polarization level and larger-scale production for HP PLAC is

Received: March 15, 2014

Accepted: April 14, 2014

Published: April 14, 2014




Figure 1. (a) Rh(I) water-soluble catalyst preparation outside of PHIP polarizer and subsequent catalytic cycle inside the PHIP polarizer leading to ¹H polarized 1-¹³C-phospholactate-d₂ (1-¹³C-PLAC-d₂). (b) Molecular addition of parahydrogen and polarization transfer leading to ¹³C HP 1-¹³C-phospholactate (1-¹³C-PLAC).¹⁸ (c) Molecular addition of parahydrogen and polarization transfer leading to ¹³C HP 1-¹³C-phospholactate-d₂ (1-¹³C-PLAC-d₂).

demonstrated by using fully deuterated PHIP precursor, 1-¹³C-phosphoenolpyruvate-d₂ (1-¹³C-PEP-d₂), and higher hydrogenation reaction temperature, respectively. Finally, a complete synthetic methodology for preparation of 1-¹³C-PEP-d₂ is reported.

RESULTS AND DISCUSSION

Commercial phosphoenolpyruvate (PEP) has been previously synthesized by the Hirschbein procedure,²⁸ where pyruvic acid is the starting material. It has also been shown that protons of pyruvic acid can be exchanged by deuterons of D₂O.^{29,30} However, the product, pyruvic-d₃ acid, was either used *in situ*³⁰ or prepared with low yield.²⁹ For example, the actual yield of the pure deuterated product was not reported, and the yield of a crude material was <50%.²⁹ Moreover, only modest isotopic purity of 70% of pyruvic-d₃ acid was achieved. Therefore, a more efficient exchange procedure was required for preparation of 1-¹³C-PEP-d₂. The exchange procedure shown in Figure 2a




Figure 2. Step-wise optimization of deuterium exchange of pyruvic acid, synthesis of PEP from sodium pyruvate, and preparation of 1-¹³C-phosphoenolpyruvate-d₂ (1-¹³C-PEP-d₂, 6). (a) Deuterium exchange: (a) (i) D₂O (450 mL), 100 °C, 5 h; (ii) 0.95 eq. NaHCO₃; (iii) Rec. D₂O/EtOH; sodium pyruvate-d₃ (2, 54% yield, ratio C₃D₃O₃⁻ to C₃D₂HO₃⁻ = 1:0.28). (b) Potassium phosphoenolpyruvate synthesis based on sodium pyruvate: (b) 0.95 eq. H₂SO₄, 0.95 eq. Br₂ (dry), CCl₄; (c) (i) P(OMe)₃, THF; (ii) H₂O; (iv) KOH to pH ~ 2.7; (v) Rec. H₂O/EtOH; potassium phosphoenolpyruvate (3, 52% over two steps); (c) preparation for potassium 1-¹³C-phosphoenolpyruvate-d₃ (1-¹³C-PEP-d₃): (a) (i) D₂O (450 mL), 100 °C, 5 h; (ii) 0.95 eq. NaHCO₃; (iii) Rec. D₂O/EtOH; sodium 1-¹³C-pyruvate-d₃ (4, 75% yield, C₂¹³CD₃O₃⁻ to C₂¹³CD₂HO₃⁻ = 1:0.25). (b') 0.95 eq. D₂SO₄, 0.95 eq. Br₂ (dry), CCl₄; (c') (i) P(OMe)₃, THF; (ii) D₂O; (iii) KOH to pH ~ 2.7; (iv) Rec. H₂O/EtOH (6, 43% over 2 steps, C₂¹³CH₂D₂O₆P⁻ to C₂¹³CH₃DO₆P⁻ to C₂¹³CH₄O₆P⁻ = 1:0.10:0.05).

was developed after several model experiments with unlabeled pyruvic acid. In this synthetic protocol, pyruvic acid is converted to pyruvic-d₃ acid by exchange with D₂O followed by its conversion to the sodium salt 1 and subsequent recrystallization. The use of a mild base, e.g., sodium bicarbonate, is essential, because stronger bases, e.g., NaOH, cause pyruvate to polymerize (Supporting Information (SI), Figure S7).

While the Hirschbein procedure²⁸ is the best PEP preparation protocol to date, it requires pyruvic acid as the starting material. Therefore, it cannot be directly applied to sodium pyruvate. Our attempt to use the previously reported Dotson procedure³¹ resulted in relatively low yields with variable reproducibility. Here, a new PEP synthesis using sodium pyruvate is reported representing the combination of modified Hirschbein and Dotson protocols, Figure 2b. Optimization of all steps in Figures 2a and 2b allowed one to prepare the main synthetic target 1-¹³C-PEP-d₂ shown in Figure 2c. To summarize, deuteration of 1-¹³C-pyruvic acid resulted in sodium 1-¹³C-pyruvate-d₃ 4 with 75% yield and ~80% isotopic purity. It was converted to 1-¹³C-bromopyruvic-d₂ acid 5, which was reacted with P(OMe)₃ to form the desired 1-¹³C-PEP-d₂ 6 with 43% yield over two steps and ~87% deuterium isotopic purity. This procedure is efficient, because it allows for a high percent deuterium isotopic labeling and high yields of the new isotopomer.

The preliminary PHIP studies with nondeuterated and deuterated 1-¹³C-PEP were conducted using previously described setup and sample preparation¹⁸ in 47.5 mT PHIP polarizer,³² where the hydrogenation high-pressure reactor was maintained at 35 °C. The pilot experiments were conducted with PHIP precursor concentration ~6 mM, [Rh(I) catalyst] ~2.5 mM at pH ~12, and in 99.8% D₂O. The effective ¹³C polarization %P was 1.86 ± 0.55% for 1-¹³C-PLAC-d₂ and %P was 0.13 ± 0.015% for 1-¹³C-PLAC. The effective %P was low

primarily because of an incomplete hydrogenation (data not shown) due to slow hydrogenation reaction. However, the ratio $\%P(1\text{-}^{13}\text{C-PLAC-}d_2)/\%P(1\text{-}^{13}\text{C-PLAC})$ was 14.3 ± 4.5 , representing a significant improvement due to deuteration of PHIP precursor. ^{13}C T_1 increased from 36 ± 2 s (PLAC) to 58 ± 4 s ($1\text{-}^{13}\text{C-PLAC-}d_2$).

The previously described PHIP polarizer design utilized a 47.5 mT permanent magnet with a relatively narrow bore of 89 mm, which cannot be heated above 50 °C. Kadlecik and co-workers³³ demonstrated that the PHIP hydrogenation reaction with similar Rh(I) catalyst and precursors can be significantly accelerated by a reaction temperature increase. Therefore, further studies with high PHIP precursor concentration were conducted using a variable temperature electromagnet PHIP setup operating at 5.75 mT and utilizing the identical chemical reactor. All samples at both polarizing fields were hydrogenated using ultrahigh purity H_2 (~7 atm partial pressure using normal H_2 gas, i.e., 75% ortho-/25% para-, or higher percent para-enrichment) for ~5 s in a 60 mL volume reactor.

The 5.75 mT PHIP polarizer setup enabled experiments at medium (55 °C) and high (68 °C) reaction temperatures for ~5 s-long reaction with ~7 atm of normal H_2 gas pressure, Figure 3. pH dependence of PEP to PLAC conversion via




Figure 3. (a) Diagram of PEP to PLAC conversion using automated PHIP polarizer. (b) pH optimization of PEP (5 mM) in phosphate buffer (25 mM) using Rh(I) catalyst (5.3 mM) at 55 ± 2 °C. (c) pH optimization of PEP (30 mM) in phosphate buffer (30 mM) using Rh(I) catalyst (5.3 mM) in the pH range of interest. Rh(I) catalyst performs several hydrogenation cycles using 30 mM PEP, Figure 1. The reaction temperatures were 68 ± 1 °C (red trace) and 55 ± 1 °C (blue trace). (d) Final pH (of solutions exiting PHIP polarizer) as a function of the starting pH (solutions entering PHIP polarizer) of solutions used in (c). All hydrogenations were performed at ~7 atm of H_2 gas partial pressure during an ~5 s reaction time.

molecular hydrogenation was evaluated next (Figure 3a). Figure 3b,c shows the percent conversion of 5 and 30 mM PEP, respectively, as measured using high-resolution proton NMR spectroscopy. Briefly, D_2O solution of PEP (5 mM) in phosphate buffer (25 mM) was prepared and divided into 25 mL fractions, which were adjusted to seven pH values: 3.9, 5.95, 7.14, 8.07, 9.93, 11.12, and 12.11. An equimolar solution of PEP and Rh (I) catalyst (5.3 mM) was prepared by a previously described procedure from each fraction.³⁴ The solutions collected from the polarizer outputs were analyzed by ^1H NMR spectroscopy using PEP/PLAC vinyl/methyl proton intensities' ratio. Percent conversion (Figure 3b) was calculated as an average of two injections at each pH. A high level (>80%) of hydrogenation was observed in all cases except at pH 12 and above.

While hydrogenation of low concentrated PEP (5 mM) can be useful for *in vitro* applications, significantly higher

concentration of an injectable, aqueous, HP contrast agent is typically required for *in vivo* imaging.^{14,16,23,35} Moreover, previous studies with $1\text{-}^{13}\text{C-succinates}$ ^{12,13} indicated that ^{13}C compounds with exchangeable carboxyl protons near pH ~5–6 may lose polarization rapidly at neutral pH due to significantly shortened T_1 . Therefore, PEP (30 mM) molecular hydrogenation with phosphate buffer (30 mM) was evaluated in the upper-middle pH range (10.0, 10.5, 11.0, and 11.5), away from neutral pH. $[\text{PEP}]/[\text{Rh(I)}]$ ratio was >5.5:1. Polarizer output solutions were analyzed by ^1H NMR spectroscopy. The intensity ratio of PEP vinyl protons and PLAC proton adjacent to the phosphate group of $(\text{CH}_3\text{CH}(\text{OPO}_3\text{HK})\text{CO}_2\text{H})$ yielded % conversion (an average of 2 injections per each pH value, Figure 3c) ranging from 41% to 74%. Higher reaction temperature translated into a 3–6% increase of conversion %. Note that the hydrogenation reaction may continue after exiting PHIP polarizer, but at a significantly lower rate, because the ejected solution temperature returned to ~25 °C and H_2 pressure $\ll 1$ atm.

The initial PEP and produced PLAC have different pK_a values, and hydrogenation also modifies the Rh(I) catalyst by removing NBD from its complex. As a result, pH of solutions passing through PHIP polarizer changes. Figure 3d clearly demonstrates the observed 1.4 ± 0.3 pH shift between starting solution (i.e., entering PHIP polarizer) and final solution (i.e., exiting PHIP polarizer). This effect must be considered since neutral pH, e.g., 7.0–7.5, is required for the *in vivo* injection of HP contrast agents, and pH of solutions exiting PHIP polarizer, e.g., pH 8.5–10.5, must be adjusted to the neutral pH range.

The above evaluation of PEP hydrogenation was used in HP studies of the fully labeled $1\text{-}^{13}\text{C-PEP-}d_2$ precursor **6** (Figure 4a), and ~26 mM solution of $1\text{-}^{13}\text{C-PEP-}d_2$ **6** ($[\text{PEP}]/[\text{Rh(I)}] > 6.5:1$) with 30 mM of phosphate buffer in D_2O was prepared at pH = 10.3. Molecular addition of parahydrogen was




Figure 4. PHIP of $1\text{-}^{13}\text{C-PEP-}d_2$ to HP $1\text{-}^{13}\text{C-PLAC-}d_2$. (a) Diagram of molecular hydrogenation of $1\text{-}^{13}\text{C-PEP-}d_2$, deuteration of 95% (SI, section 2.4), with parahydrogen to form HP $1\text{-}^{13}\text{C-PLAC-}d_2$. (b) ^{13}C spectrum of PHIP HP $1\text{-}^{13}\text{C-PLAC-}d_2$ (52 μmoles in ~3 mL and ~12 μmoles of unreacted $1\text{-}^{13}\text{C-PEP-}d_2$ corresponding to ~82% conversion) acquired at 5.75 mT *in situ* of the automated PHIP polarizer using 7 atm of parahydrogen and ~92 °C. (c) Signal reference ^{13}C spectrum of sodium $1\text{-}^{13}\text{C-acetate}$ (165 mmol in 52 mL of D_2O) with polarization $P = 4.70 \times 10^{-4}\%$ at acquisition. (d) T_1 decay of ^{13}C hyperpolarization of HP $1\text{-}^{13}\text{C-PLAC-}d_2$.

performed using ultrahigh purity H₂ with >90% para-state.³⁶ After polarization transfer to ¹³C, Figure 1, PHIP product 1-¹³C-PLAC-*d*₂ 8'' was detected *in situ* of the polarizer by single-scan ¹³C NMR spectroscopy, Figure 4b, with a similar RF probe body as previously reported.^{32,37} The enhancement factor and %*P* were calculated as before³² yielding *P* = 15.6 ± 3.0% corresponding to the ¹³C signal and polarization enhancement factor $\epsilon = (3.1 \pm 0.6) \times 10^7$ when compared to ¹³C thermal nuclear spin polarization of only 5.0×10^{-7} % at 5.75 mT and 289 K with *in situ* *T*₁ (5.75 mT) = 51 ± 2 s (SI), which would be impractical to detect at such a low field without a hyperpolarization procedure; it would take an estimated >1 billion years to record the same signal from this sample without sensitivity enhancement.

Percent conversion of 1-¹³C-PEP-*d*₂ to HP 1-¹³C-PLAC-*d*₂ determined by ¹³C NMR of solutions ejected from the PHIP polarizer immediately after *in situ* NMR detection (SI, Table S1) was ~82% (Figure 4b), and it was used to calculate the effective HP 1-¹³C-PLAC-*d*₂ concentration used for calculation of % ¹³C hyperpolarization *P* and ϵ above (SI). Importantly, *in situ* NMR detection required several extra seconds. As a result, the apparent % conversion (measured by high-resolution NMR spectroscopy using samples ejected from PHIP polarizer) was affected by this additional reaction time, leading to an overestimate of % conversion and an underestimate of % *P* and ϵ . Therefore, the reported values of ¹³C *P* = 15.6 ± 3.0% and $\epsilon = (3.1 \pm 0.6) \times 10^7$ represent the lower limit of the achieved ¹³C hyperpolarization. Consequently, the actual polarization level may be significantly higher, but the conversion yield is lower. Improving % conversion of PEP to PLAC and approaching ~100% conversion is feasible similarly to PHIP of succinate¹³ by further increasing the reaction temperature or parahydrogen partial pressure.

CONCLUSION

Deuterium labeling of PHIP precursor, Figures 1 and 2, significantly improved ¹³C PLAC hyperpolarization. PHIP conducted under identical conditions demonstrated the polarization increase by 14.3 ± 4.5-fold, as well as the *T*₁ increase from 36 ± 2 s to 58 ± 2 s at 47.5 mT. This effect of PHIP precursor deuteration was systematically quantified for the first time. Furthermore, acceleration of hydrogenation reaction by performing PHIP at elevated temperatures enabled preparation of highly concentrated and highly ¹³C polarized 1-¹³C-PLAC-*d*₂, e.g., >15.6% of 52 μmoles in 3 mL. This represents ~100-fold increase in the payload of produced ¹³C magnetization per unit volume, when compared to previously reported 1% polarization of 7 μmoles in 3 mL for 1-¹³C-PLAC. These improvements make ¹³C HP 1-¹³C-PLAC-*d*₂ suitable for *in vivo* applications. The synthetic methodology presented here allows for large-scale (tens of grams) production of 1-¹³C-PEP-*d*₂, which produces highly polarized 1-¹³C-PLAC-*d*₂ via PHIP. The produced ¹³C hyperpolarization enables >7 orders of sensitivity enhancements of ¹³C NMR signal and nuclear spin polarization. As an example, a small quantity of analyte (52 μmoles) was detected in a large volume (60 mL) reactor with high signal-to-noise ratio (SNR ~ 200) using direct ¹³C detection at 5.75 mT.

ASSOCIATED CONTENT

Supporting Information

Synthetic and PHIP buffers preparation protocols, ¹H, ¹³C NMR, and HR-MS spectra, and raw PHIP data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

*E-mail: roman.shchepin@vanderbilt.edu.

*E-mail: eduard.chekmenev@vanderbilt.edu.

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank NIGMS 1R21 GM107947-01 (K.W.W.), NIH ICMIC 5P50 CA128323-03 (K.W.W.), 5R00 CA134749-03, 3R00CA134749-02S1, and DoD CDMRP Breast Cancer Program Era of Hope Award W81XWH-12-1-0159/BC112431 for funding support.

REFERENCES

- (1) Ardenkjaer-Larsen, J. H.; Fridlund, B.; Gram, A.; Hansson, G.; Hansson, L.; Lerche, M. H.; Servin, R.; Thaning, M.; Golman, K. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, *100*, 10158–10163.
- (2) Goldman, M.; Johannesson, H. C. *R. Phys.* **2005**, *6*, 575–581.
- (3) Kurhanewicz, J.; Vigneron, D.; Brindle, K.; Chekmenev, E.; Comment, A.; Cunningham, C.; DeBerardinis, R.; Green, G.; Leach, M.; Rajan, S.; Rizi, R.; Ross, B.; Warren, W. S.; Malloy, C. *Neoplasia* **2011**, *13*, 81–97.
- (4) Hövener, J.-B.; Chekmenev, E.; Harris, K.; Perman, W.; Robertson, L.; Ross, B.; Bhattacharya, P. *Magn. Reson. Mater. Phys., Biol. Med.* **2009**, *22*, 111–121.
- (5) Hövener, J.-B.; Chekmenev, E.; Harris, K.; Perman, W.; Tran, T.; Ross, B.; Bhattacharya, P. *Magn. Reson. Mater. Phys., Biol. Med.* **2009**, *22*, 123–134.
- (6) Bowers, C. R.; Weitekamp, D. P. *Phys. Rev. Lett.* **1986**, *57*, 2645–2648.
- (7) Albers, M. J.; Bok, R.; Chen, A. P.; Cunningham, C. H.; Zierhut, M. L.; Zhang, V. Y.; Kohler, S. J.; Tropp, J.; Hurd, R. E.; Yen, Y.-F.; Nelson, S. J.; Vigneron, D. B.; Kurhanewicz, J. *Cancer Res.* **2008**, *68*, 8607–8615.
- (8) Eisenschmid, T. C.; Kirss, R. U.; Deutsch, P. P.; Hommeltoft, S. I.; Eisenberg, R.; Bargon, J.; Lawler, R. G.; Balch, A. L. *J. Am. Chem. Soc.* **1987**, *109*, 8089–8091.
- (9) Bowers, C. R.; Weitekamp, D. P. *J. Am. Chem. Soc.* **1987**, *109*, 5541–5542.
- (10) Reineri, F.; Viale, A.; Giovenzana, G.; Santelia, D.; Dastru, W.; Gobetto, R.; Aime, S. *J. Am. Chem. Soc.* **2008**, *130*, 15047–15053.
- (11) Goldman, M.; Johannesson, H.; Axelsson, O.; Karlsson, M. C. R. *Chimie* **2006**, *9*, 357–363.
- (12) Bhattacharya, P.; Chekmenev, E. Y.; Perman, W. H.; Harris, K. C.; Lin, A. P.; Norton, V. A.; Tan, C. T.; Ross, B. D.; Weitekamp, D. P. *J. Magn. Reson.* **2007**, *186*, 150–155.
- (13) Chekmenev, E. Y.; Hövener, J.; Norton, V. A.; Harris, K.; Batchelder, L. S.; Bhattacharya, P.; Ross, B. D.; Weitekamp, D. P. *J. Am. Chem. Soc.* **2008**, *130*, 4212–4213.
- (14) Zacharias, N. M.; Chan, H. R.; Sailasuta, N.; Ross, B. D.; Bhattacharya, P. *J. Am. Chem. Soc.* **2012**, *134*, 934–943.
- (15) Chekmenev, E. Y.; Chow, S. K.; Tofan, D.; Weitekamp, D. P.; Ross, B. D.; Bhattacharya, P. *J. Phys. Chem. B* **2008**, *112*, 6285–6287.

- (16) Bhattacharya, P.; Chekmenev, E. Y.; Reynolds, W. F.; Wagner, S.; Zacharias, N.; Chan, H. R.; Bünger, R.; Ross, B. D. *NMR Biomed.* **2011**, *24*, 1023–1028.
- (17) Reineri, F.; Viale, A.; Ellena, S.; Alberti, D.; Boi, T.; Giovenzana, G. B.; Gobetto, R.; Premkumar, S. S. D.; Aime, S. *J. Am. Chem. Soc.* **2012**, *134*, 11146–11152.
- (18) Shchepin, R. V.; Coffey, A. M.; Waddell, K. W.; Chekmenev, E. Y. *J. Am. Chem. Soc.* **2012**, *134*, 3957–3960.
- (19) Goldman, M.; Johannesson, H.; Axelsson, O.; Karlsson, M. *Magn. Reson. Imaging* **2005**, *23*, 153–157.
- (20) Trantzschel, T.; Bernarding, J.; Plaumann, M.; Lego, D.; Gutmann, T.; Ratajczyk, T.; Dillenberger, S.; Buntkowsky, G.; Bargon, J.; Bommerich, U. *Phys. Chem. Chem. Phys.* **2012**, *14*, 5601–5604.
- (21) Shchepin, R. V.; Pham, W.; Chekmenev, E. Y. *J. Labelled Comp. Radiopharm.* **2014**, under review.
- (22) Golman, K.; Petersson, J. S. *Acad. Radiol.* **2006**, *13*, 932–942.
- (23) Day, S. E.; Kettunen, M. I.; Gallagher, F. A.; Hu, D. E.; Lerche, M.; Wolber, J.; Golman, K.; Ardenkjaer-Larsen, J. H.; Brindle, K. M. *Nat. Med.* **2007**, *13*, 1382–1387.
- (24) Hurd, R. E.; Yen, Y.-F.; Mayer, D.; Chen, A.; Wilson, D.; Kohler, S.; Bok, R.; Vigneron, D.; Kurhanewicz, J.; Tropp, J.; Spielman, D.; Pfefferbaum, A. *Magn. Reson. Med.* **2010**, *63*, 1137–1143.
- (25) Lupo, J. M.; Chen, A. P.; Zierhut, M. L.; Bok, R. A.; Cunningham, C. H.; Kurhanewicz, J.; Vigneron, D. B.; Nelson, S. J. *Magn. Reson. Imaging* **2010**, *28*, 153–162.
- (26) Mayer, D.; Yen, Y. F.; Josan, S.; Park, J. M.; Pfefferbaum, A.; Hurd, R. E.; Spielman, D. M. *NMR Biomed.* **2012**, *25*, 1119–1124.
- (27) Chen, A. P.; Kurhanewicz, J.; Bok, R.; Xua, D.; Joun, D.; Zhang, V.; Nelson, S. J.; Hurd, R. E.; Vigneron, D. B. *Magn. Reson. Imaging* **2008**, *26*, 721–726.
- (28) Hirschbein, B. L.; Mazonod, F. P.; Whitesides, G. M. *J. Org. Chem.* **1982**, *47*, 3765–3766.
- (29) White, R. H. *Biochemistry* **1978**, *17*, 3833–3840.
- (30) Guzman, M. I.; Colussi, A. J.; Hoffmann, M. R. *J. Phys. Chem. A* **2006**, *110*, 931–935.
- (31) Dotson, G. D.; Dua, R. K.; Clemens, J. C.; Wooten, E. W.; Woodard, R. W. *J. Biol. Chem.* **1995**, *270*, 13698–13705.
- (32) Waddell, K. W.; Coffey, A. M.; Chekmenev, E. Y. *J. Am. Chem. Soc.* **2011**, *133*, 97–101.
- (33) Kadlecsek, S.; Vahdat, V.; Nakayama, T.; Ng, D.; Emami, K.; Rizi, R. *NMR Biomed.* **2011**, *24*, 933–942.
- (34) Cai, C.; Coffey, A. M.; Shchepin, R. V.; Chekmenev, E. Y.; Waddell, K. W. *J. Phys. Chem. B* **2013**, *117*, 1219–1224.
- (35) Golman, K.; in't Zandt, R.; Thaning, M. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 11270–11275.
- (36) Feng, B.; Coffey, A. M.; Colon, R. D.; Chekmenev, E. Y.; Waddell, K. W. *J. Magn. Reson.* **2012**, *214*, 258–262.
- (37) Coffey, A. M.; Shchepin, R. V.; Wilkens, K.; Waddell, K. W.; Chekmenev, E. Y. *J. Magn. Reson.* **2012**, *220*, 94–101.