



Published in final edited form as:

Pharm Res. 2014 July ; 31(7): 1644–1655. doi:10.1007/s11095-013-1267-1.

Interindividual Variability in the Cardiac Expression of Anthracycline Reductases in Donors with- and without- Down Syndrome

Adolfo Quiñones-Lombraña^{*,†}, Daniel Ferguson^{*,†}, Rachael Hageman Blair[‡], James L. Kalabus[†], Almedina Redzematovic[†], and Javier G. Blanco[†]

[†]Department of Pharmaceutical Sciences, School of Pharmacy and Pharmaceutical Sciences, The State University of New York at Buffalo, Buffalo, New York, United States of America

[‡]Department of Biostatistics, School of Public Health and Health Professions, The State University of New York at Buffalo, Buffalo, New York, United States of America

Abstract

Purpose—The intracardiac synthesis of anthracycline alcohol metabolites (e.g., daunorubicinol) contributes to the pathogenesis of anthracycline-related cardiotoxicity. Cancer patients with Down syndrome (DS) are at increased risk for anthracycline-related cardiotoxicity. We profiled the expression of anthracycline metabolizing enzymes in hearts from donors with- and without- DS.

Methods—Cardiac expression of CBR1, CBR3, AKR1A1, AKR1C3 and AKR7A2 was examined by quantitative real time PCR, quantitative immunoblotting, and enzyme activity assays using daunorubicin. The *CBR1* polymorphism rs9024 was investigated by allelic discrimination with fluorescent probes. The contribution of CBRs/AKR proteins to daunorubicin reductase activity was examined by multiple linear regression.

Results—*CBR1* was the most abundant transcript (average relative expression; DS: 81%, non-DS: 58%), and AKR7A2 was the most abundant protein (average relative expression; DS: 38%, non-DS: 35%). Positive associations between cardiac CBR1 protein levels and daunorubicin reductase activity were found for samples from donors with- and without- DS. Regression analysis suggests that sex, CBR1, AKR1A1, and AKR7A2 protein levels were significant contributors to cardiac daunorubicin reductase activity. *CBR1* rs9024 genotype status impacts on cardiac *CBR1* expression in non-DS hearts.

Conclusions—CBR1, AKR1A1, and AKR7A2 protein levels point to be important determinants for predicting the synthesis of cardiotoxic daunorubicinol in heart.

Keywords

Anthracyclines; anthracycline-related cardiotoxicity; Down syndrome; carbonyl reductases; aldo-keto reductases

Javier G. Blanco, Department of Pharmaceutical Sciences, University at Buffalo. The State University of New York, 470 Kapoor Hall, Buffalo, NY 14214 - 8033, Phone: 716 - 645 - 4820, jgblanco@buffalo.edu.

*Denotes equal contribution

INTRODUCTION

In humans, carbonyl reductases (CBRs) and aldo-keto reductases (AKRs) play predominant roles during the metabolism of more than 30 clinically relevant drugs including the antipsychotic agent haloperidol, the opioid receptor antagonist naltrexone, the antiemetic dolasetron, the anti-inflammatory ketoprofen, and the anticancer anthracyclines doxorubicin and daunorubicin (1–3). The use of anthracyclines for the chemotherapy of a variety of solid and hematological cancers is limited by the development of cardiotoxicity in some patients (4). For example, a study of children ($n = 115$) who survived leukemia revealed that 60% of those treated with anthracyclines developed preclinical abnormalities in cardiac structure and function (5). Of note, pediatric cancer patients with Down syndrome (DS, trisomy 21) constitute a population at particularly greater risk for anthracycline-related cardiotoxicity, and a safe dose of anthracyclines for cancer patients with DS remains to be defined (6, 7). For example, children with DS represent 15% of pediatric acute myeloid leukemia (AML) cases, and a report from the Children's Oncology group has documented clinically symptomatic cardiomyopathy in 17.5% of DS-AML patients treated with an anthracycline-containing regimen (7, 8).

The complex pathogenesis of anthracycline-related cardiotoxicity is mediated by a combination of oxidative stress and metabolic perturbations in myocardial tissue that are induced by the C-13 anthracycline alcohol metabolites, whose formation is catalyzed by specific CBRs and AKRs (4, 9–12). A growing cumulus of evidence indicates that two monomeric CBRs, namely CBR1 and CBR3, together with the members of the AKRs superfamily AKR1A1, AKR1C3 and AKR7A2 catalyze the synthesis of cardiotoxic C-13 anthracycline alcohol metabolites (e.g., daunorubicinol and doxorubicinol) (13–16). In addition specific *CBRs/AKRs* genetic variants may contribute to the unpredictable pharmacological profile of anthracyclines in cancer patients (17–19). For example, a recent study from the Children's Oncology group described the impact of functional single nucleotide polymorphisms in *CBR1* and *CBR3* on the risk of anthracycline-related cardiomyopathy in childhood cancer survivors (20). Thus, interindividual variability in the expression of CBRs and AKRs would impact the intracardiac formation of cardiotoxic C-13 anthracycline alcohol metabolites, and consequently the pharmacodynamics of anthracycline drugs. Furthermore, the *CBR1* and *CBR3* genes are located in the DS critical region of chromosome 21 (21q21-21q22.3). The altered expression of *CBRs* as a result of the gene dosage effect may contribute to the increased risk of anthracycline-related cardiotoxicity in cancer patients with DS (21).

In spite of the prominent contributions of CBRs and AKRs towards the pharmacodynamics of anthracycline drugs, reports documenting gene expression levels and protein abundance in cardiac tissue are limited to the analysis of individual samples or pooled tissue samples (13, 22, 23). Thus, the main goal of this study was to document the extent of interindividual variability in the expression of CBR1, CBR3, AKR1A1, AKR1C3 and AKR7A2 in a collection of heart samples from donors with- and without- DS. The expression of CBRs and AKRs was examined by quantitative real time PCR (qRT-PCR) with specific primers, quantitative immunoblotting with specific antibodies, and enzyme activity assays using the anthracycline substrate daunorubicin. We also examined the impact of a functional

polymorphism in *CBRI* (rs9024), known to impact *CBRI* expression and daunorubicinol synthesis in liver, on cardiac *CBRI* gene expression and enzymatic activity for the substrate daunorubicin (21, 24, 25).

MATERIAL AND METHODS

Human heart samples

The Institutional Review Board of the State University of New York at Buffalo approved this research. Heart samples from donors with- (n = 9) and without- DS (n = 30) were procured from The National Disease Research Interchange (NDRI, funded by the National Center for Research Resources), The Cooperative Human Tissue Network (CHTN, funded by the National Cancer Institute), and the National Institute of Child Health and Human Development (NICHD) Brain and Tissue Bank. The postmortem to tissue recovery interval was 10 h. Samples (2 – 20 g, myocardium, left ventricle only) were frozen immediately after recovery and stored in liquid nitrogen until further processing. The main demographics from donors with- and without- DS are summarized in Supplemental Table I. Down syndrome status (yes/no) and relevant diagnoses (Supplemental Table II) were obtained from anonymous medical histories. Heart samples were processed following standardized procedures to isolate DNA and RNA as described (24, 26).

Array CGH analysis

Down syndrome status was confirmed by array comparative genomic hybridization (aCGH). Briefly, genomic DNA (3.0 µg) from test samples and a euploid reference DNA sample were fluorescently labeled and hybridized to high resolution Agilent 244K aCGH arrays containing +236,000 coding and non-coding human probes. Changes in DNA copy number were determined by evaluating log₂ ratios across whole chromosomes. aCGH assays were performed at the Genomics core facility, Roswell Park Cancer Institute (Buffalo, NY).

Quantitative real-time PCR

Cardiac *CBRI*, *CBR3*, *AKR1A1*, *AKR1C3*, and *AKR7A2* mRNA expression was analyzed by qRT-PCR with gene specific primers (Table I) following the MIQE guidelines (27). Briefly, 5 ng of total RNA was reverse transcribed and amplified using the one-step QuantiTect SYBR Green RT-PCR kit (Qiagen, Venlo, The Netherlands). Target genes and *ACTB* (reference gene) were amplified in parallel with the following cycling parameters: 50°C for 30 min (reverse transcription), 95°C for 10 min (*Taq* DNA polymerase activation), 40 cycles of 95°C for 15 s (denaturation), 56°C for 30 s (annealing) and 72°C for 30 s (extension). PCR amplification products were analyzed by electrophoresis in 1% agarose gels stained with ethidium bromide. PCR amplification products were cloned into pCR 2.1- TOPO vectors (Invitrogen, Carlsbad, CA), and their identities were verified by DNA sequencing. Cloned PCR products were diluted to generate standards for calibration curves. In all cases, the regression coefficients of the standard curves were $r^2 \geq 0.95$. The amplification efficiencies of the target genes (e.g., *CBRI*, *AKR1A1*) and the reference gene (*ACTB*) were comparable and ranged between 96 – 110%. For each gene, experimental samples and standards for calibration curves were analyzed in triplicate. For each heart sample, the copy number of the target gene and *ACTB* were calculated using the average Ct values and direct

extrapolation from the calibration curves (Fig. 1). In all cases, cardiac mRNA levels were expressed as copy number ratios using the following expression:

$$\text{Copy number ratio} = \frac{\text{Target gene copy number} \times 10^6}{ACTB \text{ copy number}}$$

Quantitative immunoblotting

Heart tissue cytosols (~50 mg) were obtained using the Minute Cytoplasmic and Nuclear Extraction Kit (Invent Biotechnologies, Eden Prairie, MN) following the manufacturer instructions. Protein concentration was determined using a BCA assay (Pierce, Rockford, IL) with bovine serum albumin as a standard.

Cytosolic proteins were denatured with NuPAGE LDS sample buffer (Invitrogen) containing NuPAGE sample reducing agent (Invitrogen), and boiled at 70°C for 10 minutes prior to use. Briefly, cytosolic protein (12.5 µg for CBR1, AKR1A1, AKR1C3 and AKR7A2, and 40 µg for CBR3) and corresponding recombinant protein standards (Abcam, Cambridge, England) were separated by gel electrophoresis using NuPAGE Novex 4 – 12% Bis-Tris precast gels, and transferred onto nitrocellulose membranes using the iBlot Gel Transfer Device (Invitrogen). Membranes were blocked with 5% non-fat milk in 0.2% Tween 20-PBS for 1 hour at room temperature and then probed with specific polyclonal anti-human CBR1 (1:3000; Santa Cruz Biotechnology, Dallas, TX), CBR3 (1:500; Santa Cruz Biotechnology), AKR1A1 (1:3000; Abnova, Taipei City, Taiwan), AKR1C3 (1:3000; Abnova), and AKR7A2 (1:3000; Abnova) antibodies overnight at 4°C. Next, membranes were incubated with a secondary goat or rabbit anti-IgG antibodies conjugated with horseradish peroxidase (1:10000; Santa Cruz Biotechnology) for 1.5 hours at room temperature. To normalize for protein loading, membranes were stripped with Restore Western Blot Stripping Buffer (Thermo Scientific, Waltham, MA), and re-probed with anti-GAPDH antibody (1:1000; Santa Cruz Biotechnology). Immunoreactive bands were visualized with the ECL Plus Western blotting substrate (GE Healthcare, Little Chalfont, England) on a ChemiDoc XRS imager (Bio-Rad, Berkeley, CA). Densitometric analysis was performed using Image Lab software (Bio-Rad). Cardiac CBR1, CBR3, AKR1A1, AKR1C3, and AKR7A2 protein levels were estimated by direct extrapolation from the corresponding recombinant protein standard curves. Detection of recombinant CBR1, CBR3, AKR1A1, AKR1C3, and AKR7A2 proteins was linear (range: 0.01 – 0.1 µg; $r^2 > 0.99$). The limits of detection (LOD) and limits of quantification (LOQ) were calculated based on the standard deviation of the band intensity and the slope of the calibration curve (Supplemental Fig. 1). Results are as follows: $LOD_{CBR1}=0.34$ ng, $LOQ_{CBR1}=0.80$ ng; $LOD_{CBR3}=0.49$ ng, $LOQ_{CBR3}=1.49$ ng; $LOD_{AKR1A1}=0.58$ ng, $LOQ_{AKR1A1}=1.76$ ng; $LOD_{AKR1C3}=0.33$ ng, $LOQ_{AKR1C3}=0.98$ ng; $LOD_{AKR7A2}=0.44$ ng, $LOQ_{AKR7A2}=1.34$ ng.

Kinetic analysis

Anthracycline reductase activity was measured in heart cytosols using the substrate daunorubicin as described previously (21). Validation experiments with pooled heart cytosols showed that 400 µM daunorubicin [S] ensured conditions of V_{max} (zero-order

kinetics). Incubation mixtures (0.5 mL) contained potassium phosphate buffer (0.1 M; pH 7.4), cytosolic protein (400 µg) and daunorubicin (400 µM; Sigma-Aldrich, St. Louis, MO). Mixtures were equilibrated for 5 minutes at 37°C. Reactions were initiated by the addition of NADPH (200 µM; Sigma). Reaction mixtures were flash frozen with liquid nitrogen after 2 minutes, and stored at -80°C until daunorubicinol quantification. Daunorubicinol was quantified using a high performance liquid chromatography (UPLC) fluorescent detection method (28). The metabolite was extracted, and then quantified using a daunorubicinol standard (Toronto Research Chemicals, Toronto, ON) as previously described (29).

CBR1 Genotyping

The *CBR1* polymorphism (rs9024) was investigated by allelic discrimination with fluorescent probes and real-time PCR (Assays-by-Design; Applied Biosystems, Foster City, CA) as described previously (24, 30). Rs9024 genotype status in DS samples (i.e., trisomy 21) was determined with a validated allelic discrimination assay as described (21).

Data analysis

Statistical comparisons were performed using Excel 2007 (Microsoft Office; Microsoft, Redmond, WA) and GraphPad Prism version 4.03 (GraphPad Software Inc., La Jolla, CA). The Kolmogorov–Smirnov test was used to analyze the normality of data sets. The Mann-Whitney *U* test or Student's *t* test were used to compare group means. Pearson's coefficient of correlation (r_p) was used to analyze data sets with normal distributions. Comparison tests were considered significant at the $P < 0.05$.

Multiple linear regression models were fit using the R programming language (<http://www.r-project.org/>). Multiple linear regression analysis was restricted to non-DS hearts due to sample size limitations. For the regression modeling, the observations that have a missing value in at least one of the protein expression measurements were eliminated from the data set. CBR3 was not considered for the regression modeling due to the number of missing values. In the union of missing data for the remaining proteins: CBR1, AKR1A1, AKR1C3, and AKR7A2, there were 12 observations with missing values for at least one of the measured proteins. These observations were eliminated for model fitting, and the remaining subset of 18 observations (9 males and 9 females), was used for the regression. The full model was represented as:

$$\text{Daunorubicin Activity} \sim \beta_0 + \beta_1 \times \text{Sex} + \beta_2 \times \text{CBR1} + \beta_3 \times \text{AKR1A1} + \beta_4 \times \text{AKR1C3} + \beta_5 \times \text{AKR7A2} + \varepsilon,$$

where β_j for $j = 0, 1, \dots, 5$ represent the regression coefficients for the predictors in the model, β_0 is the intercept, and $\varepsilon \sim N(0,1)$, is an error term. CBR1, AKR1A1, AKR1C3, and AKR7A2 represent the normalized protein concentration (nmol/gram cytosolic protein) in heart. The term Sex was represented as a binary factor.

RESULTS

***CBR1*, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* mRNA expression in hearts from donors with- and without- DS**

In all cases, DS status from medical records (i.e., trisomy 21) was confirmed by aCGH. Cardiac *CBR1*, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* mRNA levels in donors with- and without- DS are shown in Fig. 1 (panels: a – e). Analysis of the average relative expression levels showed that *CBR1* mRNA is the most abundant transcript in heart samples from donors with- (81%) and without- DS (58%. Fig. 1f and 1g). On average, the remaining transcripts showed similar trends in their relative expression levels in samples from donors with- ($CBR3 \approx AKR1C3 > AKR1A1 > AKR7A2$. Fig. 1g) and without- DS ($CBR3 > AKR1C3 > AKR1A1 > AKR7A2$. Fig. 1f). The cardiac expression of *CBR1*, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* mRNA displayed considerable interindividual variability in samples from donors with- and without- DS (Table II). The cardiac expression of *CBRs* and *AKRs* mRNAs did not significantly differ between samples from donors with- and without- DS (Table II).

***CBR1*, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* protein expression in hearts from donors with- and without- DS**

Cardiac *CBR1*, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* protein expression levels in samples from donors with- and without- DS are shown in Fig. 2 (panels: a – e). Eight out of 9 heart samples from donors with- DS exhibited *CBR3* protein levels below the LOQ (Supplemental Fig. 2). *AKR7A2* showed the highest average expression levels in hearts from both groups, accounting for 33% of total *CBRs*-*AKRs* in donors with- DS and 30% of total *CBRs*-*AKRs* in donors without- DS. On average, the pattern of relative cardiac expression was similar in samples from donors with- ($AKR7A2 > AKR1A1 > AKR1C3 > CBR1 > CBR3$. Fig. 2g) and without- DS ($AKR7A2 > AKR1A1 \approx AKR1C3 > CBR1 > CBR3$. Fig. 2f). Table III shows cardiac *CBRs* and *AKRs* protein levels in samples from donors with- and without- DS. The average cardiac expression of *CBR1* was 57% higher in samples from donors with- DS than in samples from donors without- DS (Fig. 2 and Table III). The expression levels of *AKR1A1*, *AKR1C3* and *AKR7A2* did not differ between samples from donors with- and without- DS (Table III).

Maximal daunorubicin reductase activity in hearts from donors with- and without- DS

Fig. 3A shows maximal daunorubicin reductase activities in hearts from donors with- and without- DS. Variable levels of daunorubicin reductase activity were observed in cardiac samples from donors with- DS (range: 0.9 – 6.0 nmol/mg-min; 7-fold) and without- DS (range: 0.9 – 7.4 nmol/mg-min; 8-fold. Fig. 3a). Statistical comparisons of the means demonstrated that cardiac maximal daunorubicin reductase activities were similar between both groups (Activity_{non-DS} = 2.5 ± 1.5 vs. Activity_{DS} = 2.3 ± 1.9, $P = 0.28$. Fig. 3a). Linear regression analysis was used to test for associations between *CBRs*-*AKRs* protein levels and cardiac maximal daunorubicin reductase activities (Table IV). There existed a significant positive association between *CBR1* protein levels and daunorubicin reductase activity in samples from donors with- and without- DS (Table IV and Fig. 3b and 3c).

Our collection included three samples from pediatric donors with- DS (Supplemental Table III). Age-related changes in the cardiac expression of CBRs/AKRs during the developmental continuum encompassing birth and adolescence may impact the metabolism of anthracycline substrates in pediatric patients. Exploratory analyses revealed no differences in the cardiac expression of *CBRs/AKRs* (mRNA, protein and daunorubicin reductase activity) in samples from pediatric donors in comparison to samples from non-pediatric donors (Supplemental Table III).

Multiple linear regression analysis was used to examine the impact of cardiac CBRs/AKRs protein levels on daunorubicin reductase activities in samples from donors without- DS. The results (Supplemental Table IV) indicate Sex is the most significant factor ($P < 0.001$), followed by AKR7A2, CBR1, and AKR1A1 ($P < 0.05$). AKR1C3 was not significant and removed from the model in an effort to identify the most parsimonious model of daunorubicin reductase activity (Table V). As expected, the removal of this term did not change the overall model, and the terms that were significant in the full model remained significant in the reduced model (Table V).

***CBR1* rs9024 genotype-phenotype associations in heart samples from donors with- and without- DS**

Tables VI and VII show cardiac *CBR1* expression in samples from donors with- and without- DS stratified by *CBR1* rs9024 genotype status. No statistical comparisons are shown for donors with- DS due to sample size limitations. Visual inspection of the data shown in Fig. 4 suggests the absence of an evident association between rs9024 genotype status and cardiac *CBR1* mRNA and protein expression in samples from donors with- DS (Fig. 4a and 4b). Similarly, visual inspection does not suggest any association between *CBR1* rs9024 genotype status and maximal cardiac daunorubicin reductase activity in samples from donors with- DS (Fig. 4c). The analysis of donors without- DS showed that, on average, relative *CBR1* mRNA levels tend to be higher in samples with *CBR1* rs9024 homozygous G/G genotype compared to samples with homozygous A/A genotype; the difference between genotype groups did not reach statistical significance (Table VII and Fig. 5a). Further analysis showed a trend towards higher CBR1 protein levels (≈ 4 -fold) in samples from donors with *CBR1* rs9024 homozygous G/G genotype in comparison to samples with homozygous A/A genotype (Table VII and Fig. 5b). In line, heart cytosols from donors with homozygous *CBR1* rs9024 G/G genotypes displayed higher maximal rates of daunorubicinol synthesis (≈ 1.6 -fold) than cytosols from donors with homozygous A/A genotype (Table VII and Fig. 5c).

DISCUSSION

There is a paucity of reports describing the expression of prominent drug metabolizing enzymes in tissues from subjects with- DS. Within the context of multi-agent chemotherapy that includes anthracyclines for the treatment of pediatric leukemia, there is still considerable uncertainty about the optimal dose requirements for patients with- DS (31). We propose that variability in the cardiac expression of specific CBRs and AKRs may contribute to the unpredictable pharmacodynamics of anthracycline drugs. The goal of this

study was to document the expression of *CBR1*, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* in a collection of heart samples from donors with- and without- DS.

CBR1, *CBR3*, *AKR1A1*, *AKR1C3* and *AKR7A2* mRNAs are expressed at variable levels in heart samples from donors with- and without- DS (Fig. 1 and Table II). Of all the transcripts, *CBR3* mRNA exhibited the widest range of expression in samples from donors with- (34-fold) and without- DS ($>1.3 \times 10^5$ fold). The cardiac expression of *CBR1* and *CBR3* mRNAs are of particular interest due to the location of the *CBR1* and *CBR3* genes in chromosome 21, and the expected ≈ 1.5 -fold increase in gene expression due to “gene dosage effect” in tissue samples with trisomy 21. Studies on transcriptional profiles in the DS setting suggest that the gene-dosage effect for most of the chromosome 21 transcripts ($\approx 70\%$) is compensated by natural variation in gene expression (32). The compensation effect appears to be dependent upon the cellular context. For example, the expression of *CBR1* and *CBR3* mRNAs is increased in trisomic fibroblasts (≈ 1.5 -fold) but not in trisomic lymphoblastoid cell lines (33). In this study, the expression of *CBR3* mRNA was similar in samples from donors with- and without- DS. Thus, it appears that natural interindividual variability in *CBR3* mRNA expression compensates the expected gene dosage effect in trisomic heart tissues. In contrast, *CBR1* mRNA expression tended to be higher (1.7-fold) in samples from donors with- DS than in samples from donors without- DS (Fig. 1 and Table II). Trisomic heart samples also showed a concomitant ≈ 1.6 -fold increase in *CBR1* protein expression (Fig. 2 and Table III). These findings are in agreement with our previous observation documenting increased cardiac *CBR1* mRNA and *CBR1* protein expression in a subset of 4 samples from donors with- DS (21). It is assumed that the overexpression of “dosage-sensitive” genes from chromosome 21 is involved in specific DS phenotypes (e.g., cognitive impairments, facial dysmorphisms, and congenital heart disease (34). In this context, our data suggest that *CBR1* overexpression in the cardiac tissue from donors with- DS may contribute to phenotypic abnormalities in trisomic hearts.

At the protein level, *AKR7A2* was the most abundant anthracycline reductase in hearts from donors with- and without- DS, representing $\approx 30\%$ of the total reductase content (Fig. 2). In non-DS hearts, linear regression analysis of protein content versus maximal enzymatic activity for the substrate daunorubicin suggests that *AKR7A2* content accounts for $\approx 8\%$ of the total variance in cardiac daunorubicin reductase activity (Table IV). A similar analysis indicates that cardiac *CBR1* protein content accounts for $\approx 21\%$ of the variance in daunorubicin reductase activity in samples from donors without- DS. The contribution of *CBR1* expression to daunorubicin reductase activity in hearts from donors with- DS appears to be substantial. In this case, the regression analysis suggests that *CBR1* content accounts for $\approx 90\%$ of the variance in cardiac daunorubicin reductase activity in hearts from donors with- DS (Fig. 3 and Table IV). These findings suggest that *CBR1* protein expression is an important predictor of daunorubicin synthesis in human heart.

CBR1 rs9024 genotype status has been associated with the hepatic expression of *CBR1* and anthracycline reductase activity for the substrate doxorubicin. In this study, the impact of *CBR1* rs9024 genotype status on cardiac *CBR1* expression was apparent in the group of heart samples from donors without- DS (Fig. 4 and Table VII). Samples with homozygous G/G genotype exhibited higher daunorubicin reductase activity ($\approx 60\%$ increase) than

samples with homozygosis for the A allele. At first glance, the impact of *CBR1* rs9024 on cardiac daunorubicin reductase activity may appear relatively modest. However, certain therapeutic regimens such as combination chemotherapies for pediatric acute myeloid leukemia require the administration of several doses of daunorubicin ($n > 5$). In this “chronic” context, the effect size of *CBR1* rs9024 genotype status may become more relevant in terms of favoring the intracardiac build-up of cardiotoxic daunorubicinol. It appears that *CBR1* rs9024 genotype status does not impact the cardiac expression of *CBR1* in heart samples from donors with- DS. A recent study described regulation of *CBR1* expression by microRNAs (hsa-miR-574-5p and hsa-miR-921) dependent on the allele status of the *CBR1* rs9024 (25). New evidence suggests that the expression of specific microRNAs is dysregulated in lymphocytes from subjects with- DS (35). Thus, it is plausible that alterations in the cardiac expression of hsa-miR-574-5p and/or hsa-miR-921 in the DS setting would in turn disrupt the regulation of *CBR1* dependent upon rs9024 genotype status. Our laboratory is exploring this intriguing possibility by examining the expression of hsa-miR-574-5p and hsa-miR-921 in hearts and cell lines from donors with- and without- DS.

There are some limitations in this study, the main one being the relatively small number of cardiac samples from donors with- DS. Our procurement rates for samples from donors with- DS is low (≈ 1 sample every 9 – 12 months), even after working with national cooperative resources such as NDRI, CHTN, and the NICHD Brain and Tissue Bank. The scarcity of representative tissue samples from donors with- DS continues to impair the execution of large-scale quantitative studies to tackle fundamental topics related to the metabolism and disposition of commonly used drugs in this group of subjects (36, 37). Second, studies based on the use of tissue samples from cadaveric donors are prone to many issues including, but not limited to sample degradation, incomplete/null medical histories and/or demographics, use of concomitant medications/drugs/smoking status. Nevertheless, the collection of samples that is the object of this study may still represent an informative window to assess the extent of variability in the cardiac expression of anthracycline reductases in subjects with- and without- DS.

Weiss highlighted the need for performing comprehensive cardiac drug metabolism studies to characterize the system response at the organ level during pharmacotherapy with common drugs (e.g., anthracyclines) (38). In this study, multiple regression analysis suggests that sex together with *CBR1*, *AKR1A1*, and *AKR7A2* protein expression account for $\approx 70\%$ of the total variance in cardiac daunorubicin reductase activity. Thus, the expression of specific CBRs and AKRs appears to drive the cardiac synthesis of daunorubicinol. Future modeling efforts should be directed towards the identification of distinct cardiac CBR/AKR expression profiles that dictate low and/or high rates of synthesis of cardiotoxic C-13 anthracycline alcohol metabolites. The present study represents a necessary step for the creation of novel quantitative methods to predict the variable pharmacodynamics of anthracyclines in human heart by integrating *genetic* and *phenotypic* data (e.g., CBR/AKR protein abundance, rs9024, miRNA levels, DS status, etc) for the specific CBRs and AKRs involved in the metabolism of these clinically useful drugs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institute of General Medical Sciences [GM073646].

ABBREVIATIONS

aCGH	Array comparative genomic hybridization
ACTB	Actin B
AKR1A1	aldo-keto reductase family 1, member A1
AKR1C3	aldo-keto reductase family 1, member C3
AKR7A2	aldo-keto reductase family 7, member A2
AKRs	aldo-keto reductases
CBR1	carbonyl reductase 1
CBR3	carbonyl reductase 3
CBRs	carbonyl reductases
DS	Down syndrome
LOD	limit of detection
LOQ	limit of quantification

References

1. Penningand TM, Drury JE. Human aldo-keto reductases: Function, gene regulation, and single nucleotide polymorphisms. *Arch Biochem Biophys.* 2007; 464:241–250. [PubMed: 17537398]
2. Hoffmannand F, Maser E. Carbonyl reductases and pluripotent hydroxysteroid dehydrogenases of the short-chain dehydrogenase/reductase superfamily. *Drug Metab Rev.* 2007; 39:87–144. [PubMed: 17364882]
3. Rosemondand MJ, Walsh JS. Human carbonyl reduction pathways and a strategy for their study in vitro. *Drug Metab Rev.* 2004; 36:335–361. [PubMed: 15237858]
4. Menna P, Paz OG, Chello M, Covino E, Salvatorelli E, Minotti G. Anthracycline cardiotoxicity. *Expert Opin Drug Saf.* 2012; 11(Suppl 1):S21–36. [PubMed: 21635149]
5. Wouters KA, Kremer LCM, Miller TL, Herman EH, Lipshultz SE. Protecting against anthracycline-induced myocardial damage: a review of the most promising strategies. *British Journal of Haematology.* 2005; 131:561–578. [PubMed: 16351632]
6. Krischer JP, Epstein S, Cuthbertson DD, Goorin AM, Epstein ML, Lipshultz SE. Clinical cardiotoxicity following anthracycline treatment for childhood cancer: the Pediatric Oncology Group experience. *J Clin Oncol.* 1997; 15:1544–1552. [PubMed: 9193351]
7. O'Brien MM, Taub JW, Chang MN, Massey GV, Stine KC, Raimondi SC, Becton D, Ravindranath Y, Dahl GV. Cardiomyopathy in children with Down syndrome treated for acute myeloid leukemia: a report from the Children's Oncology Group Study POG 9421. *J Clin Oncol.* 2008; 26:414–420. [PubMed: 18202418]
8. Tauband JW, Ravindranath Y. What's up with down syndrome and leukemia-A lot! *Pediatr Blood Cancer.* 2011; 57:1–3. [PubMed: 21480471]

9. Menna P, Salvatorelli E, Minotti G. Cardiotoxicity of Antitumor Drugs. *Chem Res Toxicol.* 2008; 21:978–989. [PubMed: 18376852]
10. Mushlin PS, Cusack BJ, Boucek RJ Jr, Andrejuk T, Li X, Olson RD. Time-related increases in cardiac concentrations of doxorubicinol could interact with doxorubicin to depress myocardial contractile function. *Br J Pharmacol.* 1993; 110:975–982. [PubMed: 8298821]
11. Olson RD, Mushlin PS, Brenner DE, Fleischer S, Cusack BJ, Chang BK, Boucek RJ Jr. Doxorubicin cardiotoxicity may be caused by its metabolite, doxorubicinol. *Proc Natl Acad Sci U S A.* 1988; 85:3585–3589. [PubMed: 2897122]
12. Stewart DJ, Grewaal D, Green RM, Mikhael N, Goel R, Montpetit VA, Redmond MD. Concentrations of doxorubicin and its metabolites in human autopsy heart and other tissues. *Anticancer Res.* 1993; 13:1945–1952. [PubMed: 8297100]
13. Bains OS, Grigliatti TA, Reid RE, Riggs KW. Naturally occurring variants of human aldo-keto reductases with reduced in vitro metabolism of daunorubicin and doxorubicin. *J Pharmacol Exp Ther.* 2010; 335:533–545. [PubMed: 20837989]
14. Bains OS, Takahashi RH, Pfeifer TA, Grigliatti TA, Reid RE, Riggs KW. Two Allelic Variants of Aldo-Keto Reductase 1A1 Exhibit Reduced in Vitro Metabolism of Daunorubicin. *Drug Metab Dispos.* 2008; 36:904–910. [PubMed: 18276838]
15. Lakhman SS, Ghosh D, Blanco JG. Functional significance of a natural allelic variant of human carbonyl reductase 3 (CBR3). *Drug Metab Dispos.* 2005; 33:254–257. [PubMed: 15537833]
16. Miura T, Nishinaka T, Terada T. Different functions between human monomeric carbonyl reductase 3 and carbonyl reductase 1. *Mol Cell Biochem.* 2008; 315:113–121. [PubMed: 18493841]
17. Blanco JG, Leisenring WM, Gonzalez-Covarrubias VM, Kawashima TI, Davies SM, Relling MV, Robison LL, Sklar CA, Stovall M, Bhatia S. Genetic polymorphisms in the carbonyl reductase 3 gene CBR3 and the NAD(P)H:quinone oxidoreductase 1 gene NQO1 in patients who developed anthracycline-related congestive heart failure after childhood cancer. *Cancer.* 2008; 112:2789–2795. [PubMed: 18457324]
18. Lal S, Sandanaraj E, Wong ZW, Ang PC, Wong NS, Lee EJ, Chowbay B. CBR1 and CBR3 pharmacogenetics and their influence on doxorubicin disposition in Asian breast cancer patients. *Cancer Sci.* 2008; 99:2045–2054. [PubMed: 19016765]
19. Lal S, Mahajan A, Chen WN, Chowbay B. Pharmacogenetics of target genes across doxorubicin disposition pathway: a review. *Curr Drug Metab.* 2010; 11:115–128. [PubMed: 20302569]
20. Blanco JG, Sun CL, Landier W, Chen L, Esparza-Duran D, Leisenring W, Mays A, Friedman DL, Ginsberg JP, Hudson MM, Neglia JP, Oeffinger KC, Ritchey AK, Villaluna D, Relling MV, Bhatia S. Anthracycline-Related Cardiomyopathy After Childhood Cancer: Role of Polymorphisms in Carbonyl Reductase Genes--A Report From the Children's Oncology Group. *J Clin Oncol.* 2012; 30:1415–1421. [PubMed: 22124095]
21. Kalabus JL, Sanborn CC, Jamil RG, Cheng Q, Blanco JG. Expression of the anthracycline-metabolizing enzyme carbonyl reductase 1 in hearts from donors with Down syndrome. *Drug Metab Dispos.* 2010; 38:2096–2099. [PubMed: 20729274]
22. Kassner N, Huse K, Martin HJ, Godtel-Armbrust U, Metzger A, Meineke I, Brockmoller J, Klein K, Zanger UM, Maser E, Wojnowski L. Carbonyl reductase 1 is a predominant doxorubicin reductase in the human liver. *Drug Metab Dispos.* 2008; 36:2113–2120. [PubMed: 18635746]
23. O'Connor T, Ireland LS, Harrison DJ, Hayes JD. Major differences exist in the function and tissue-specific expression of human aflatoxin B1 aldehyde reductase and the principal human aldo-keto reductase AKR1 family members. *Biochem J.* 1999; 343(Pt 2):487–504. [PubMed: 10510318]
24. Gonzalez-Covarrubias V, Zhang J, Kalabus JL, Relling MV, Blanco JG. Pharmacogenetics of human carbonyl reductase 1 (CBR1) in livers from black and white donors. *Drug Metab Dispos.* 2009; 37:400–407. [PubMed: 19022938]
25. Kalabus JL, Cheng Q, Blanco JG. MicroRNAs Differentially Regulate Carbonyl Reductase 1 (CBR1) Gene Expression Dependent on the Allele Status of the Common Polymorphic Variant rs9024. *PLoS One.* 2012; 7:e48622. [PubMed: 23133646]

26. Sultan M, Piccini I, Balzereit D, Herwig R, Saran NG, Lehrach H, Reeves RH, Yaspo ML. Gene expression variation in Down's syndrome mice allows prioritization of candidate genes. *Genome Biol.* 2007; 8:R91. [PubMed: 17531092]
27. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem.* 2009; 55:611–622. [PubMed: 19246619]
28. de Jong J, Guerand WS, Schoofs PR, Bast A, van der Vijgh WJ. Simple and sensitive quantification of anthracyclines in mouse atrial tissue using high-performance liquid chromatography and fluorescence detection. *J Chromatogr.* 1991; 570:209–216. [PubMed: 1797830]
29. Fetterly GJ, Aras U, Lal D, Murphy M, Meholick PD, Wang ES. Development of a preclinical PK/PD model to assess antitumor response of a sequential aflibercept and doxorubicin-dosing strategy in acute myeloid leukemia. *The AAPS journal.* 2013; 15:662–673. [PubMed: 23550025]
30. Cheng Q, Yang W, Raimondi SC, Pui CH, Relling MV, Evans WE. Karyotypic abnormalities create discordance of germline genotype and cancer cell phenotypes. *Nat Genet.* 2005; 37:878–882. [PubMed: 16041371]
31. Seewald L, Taub JW, Maloney KW, McCabe ER. Acute leukemias in children with Down syndrome. *Mol Genet Metab.* 2012; 107:25–30. [PubMed: 22867885]
32. Ait Yahya-Graison E, Aubert J, Dauphinot L, Rivals I, Prieur M, Golfier G, Rossier J, Personnaz L, Creau N, Blehaut H, Robin S, Delabar JM, Potier MC. Classification of human chromosome 21 gene-expression variations in Down syndrome: impact on disease phenotypes. *American journal of human genetics.* 2007; 81:475–491. [PubMed: 17701894]
33. Prandini P, Deutsch S, Lyle R, Gagnebin M, Delucinge Vivier C, Delorenzi M, Gehrig C, Descombes P, Sherman S, Dagna Bricarelli F, Baldo C, Novelli A, Dallapiccola B, Antonarakis SE. Natural gene-expression variation in Down syndrome modulates the outcome of gene-dosage imbalance. *American journal of human genetics.* 2007; 81:252–263. [PubMed: 17668376]
34. Patterson D. Molecular genetic analysis of Down syndrome. *Human genetics.* 2009; 126:195–214. [PubMed: 19526251]
35. Xu Y, Li W, Liu X, Chen H, Tan K, Chen Y, Tu Z, Dai Y. Identification of dysregulated microRNAs in lymphocytes from children with Down syndrome. *Gene.* 2013
36. McCabeand LL, McCabe ER. Down syndrome: issues to consider in a national registry, research database and biobank. *Mol Genet Metab.* 2011; 104:10–12. [PubMed: 21501965]
37. Oster-Granite ML, Parisi MA, Abbeduto L, Berlin DS, Bodine C, Bynum D, Capone G, Collier E, Hall D, Kaeser L, Kaufmann P, Krischer J, Livingston M, McCabe LL, Pace J, Pfenninger K, Rasmussen SA, Reeves RH, Rubinstein Y, Sherman S, Terry SF, Whitten MS, Williams S, McCabe ER, Maddox YT. Down syndrome: national conference on patient registries, research databases, and biobanks. *Mol Genet Metab.* 2011; 104:13–22. [PubMed: 21835664]
38. Weiss M. Functional characterization of drug uptake and metabolism in the heart. *Expert Opin Drug Metab Toxicol.* 2011; 7:1295–1306. [PubMed: 21870896]




Fig. 1. Cardiac *CBR1*, *CBR3*, *AKR1A1*, *AKR1C3*, and *AKR7A2* mRNA expression in samples from donors with- ($n = 9$) and without- DS ($n = 30$) (Panels: a – e). Each symbol depicts the average of individual samples. Horizontal lines indicate group means. Insets show the corresponding calibration curves. Samples and standards for calibration curves were analyzed in triplicates. Relative abundance of cardiac *CBRs* and *AKRs* mRNAs in donors without- (f) and with- DS (g).




Fig. 2. Cardiac CBR1, CBR3, AKR1A1, AKR1C3, and AKR7A2 protein expression in samples from donors with- and without- DS. Each symbol depicts the average of individual samples. Horizontal lines indicate group means. Samples and standards for calibration curves were analyzed in duplicates for CBR3 and in triplicates for CBR1, AKR1A1, AKR1C3, AKR7A2. Samples exhibiting protein levels below the LOQs were excluded (Supplemental Fig. 1). Insets show representative immunoblots for recombinant standards (left) and

cytosolic CBRs-AKRs plus GADPH (right). Relative abundance of cardiac CBRs and AKRs proteins in donors without- (f) and with- DS (g).




Fig. 3. Daunorubicin reductase activity in donors with- (n=9) and without- DS (n=30) (a). Each symbol depicts the average of individual heart samples. Each sample was analyzed in duplicates. Group means are indicated by horizontal lines. Linear regression analysis of maximal daunorubicin reductase activity versus cardiac CBR1 protein expression in samples from donors without- and with- DS (panels b and c).




Fig. 4. Impact of *CBR1* rs9024 genotype status on: cardiac *CBR1* expression (a), cardiac *CBR1* protein expression (b), and daunorubicin reductase activity (c) in donors with- DS. Each symbol represents individual heart samples. Horizontal lines indicate group means.




Fig. 5. Impact of *CBR1* rs9024 genotype status on: cardiac *CBR1* expression (a), cardiac *CBR1* protein expression (b), and daunorubicin reductase activity (c) in donors without- DS. Each symbol represents individual heart samples. Horizontal lines indicate group means.

Table IList of PCR primers for the amplification of cardiac *CBRs* and *AKRs* mRNAs

Gene	Forward primer	Reverse primer	Size (bp)
<i>CBR1</i>	5'-TCAAGCTGAAGTGACGATGA-3'	5'-GGTGCACCTCCCTTCTTTGTA-3'	239
<i>CBR3</i>	5'-AACCTCATGGGAGAGTGGTG-3'	5'-TCCTCGATAAGACCGTGACC-3'	231
<i>AKR1A1</i>	5'-CCTGGTCAGGTAAAAGCAGC-3'	5'-CCCGCTCAAAGGCATAAGG-3'	286
<i>AKR1C3</i>	5'-CCTCCAGAGGTTCCGAGAAG-3'	5'-GTGGACCAAAGCTTTGAAGTG-3'	173
<i>AKR7A2</i>	5'-CCTCCAGAGGTTCCGAGAAG-3'	5'-GTGGACCAAAGCTTTGAAGTG-3'	168
<i>ACTB</i>	5'-GGACTTCGAGCAAGAGATGG-3'	5'-AGCACTGTGTTGGCGTACAG-3'	234

Table II

Cardiac *CBRI*, *CBR3*, *AKR1A1*, *AKR1C3*, and *AKR7A2* mRNA expression in samples from donors without- and with- DS

Transcript	Non-DS			DS			P value
	Copy number ratio		Range	Copy number ratio		Range	
	Mean \pm SD	Range		Mean \pm SD	Range		
<i>CBRI</i>	19283.8 \pm 30824.3	245.5–130786.4	33029.3 \pm 36187.0	3103.1–96061.0	0.23		
<i>CBR3</i>	6975.3 \pm 12271.4	0.4–51914.3	2992.8 \pm 3465.6	334.5–11487.5	0.46		
<i>AKR1A1</i>	2078.4 \pm 2396.1	66.0–10865.2	1073.6 \pm 955.5	144.2–3048.4	0.42		
<i>AKR1C3</i>	3986.8 \pm 5264.9	14.3–24131.8	2913.3 \pm 1631.6	227.0–4732.8	0.69		
<i>AKR7A2</i>	1026.1 \pm 955.7	42.3–3747.9	965.8 \pm 590.6	143.9–1883.0	0.98		

Table III

Cardiac CBR1, CBR3, AKR1A1, AKR1C3, and AKR7A2 protein expression in samples from donors without- and with- DS

Transcript	Non-DS		DS		P value
	nmol/g cytosolic protein		nmol/g cytosolic protein		
	Mean ± SD	Range	Mean ± SD	Range	
CBR1	11.9 ± 8.9	1.2 – 30.2	18.8 ± 6.7	14.3 – 31.1	0.04
CBR3	6.3 ± 5.5	1.0 – 15.8	2.3	N/A	N/A
AKR1A1	27.4 ± 17.3	0.0 – 74.1	28.3 ± 15.4	15.0 – 63.0	0.86
AKR1C3	26.0 ± 22.4	0.5 – 70.3	18.4 ± 23.2	0.3 – 61.4	0.32
AKR7A2	31.0 ± 21.0	5.4 – 68.1	33.8 ± 19.6	6.8 – 61.2	0.97

N/A., not applicable (*n* = 1)

Table IV

Linear regression analysis of cardiac daunorubicin reductase activities versus CBRs/AKRs protein levels in samples from donors without- and with- DS

Protein	Non-DS		DS	
	Pearson correlation coefficient	<i>P</i> value	Pearson correlation coefficient	<i>P</i> value
CBR1	0.46	0.04	0.96	0.002
CBR3	0.20	0.58	N/A	N/A
AKR1A1	0.00	0.99	-0.03	0.95
AKR1C3	-0.16	0.41	0.32	0.41
AKR7A2	0.28	0.19	-0.22	0.58

N/A., not applicable ($n = 1$).

Table V

Results of the reduced multiple regression model fit.

Term	β estimate	Std. Error	t-stat	P value	Sig. Codes
Intercept	-2.527	1.023	-2.471	0.028	***
Sex	2.560	0.571	4.481	0.001	****
CBR1	2.380	1.001	2.377	0.033	**
AKR1A1	1.739	0.592	2.939	0.012	**
AKR7A2	1.242	0.357	3.479	0.004	***

The significant codes are as follows:

**** for $P < 0.001$,

*** for $P < 0.01$,

** $P < 0.05$,

* $P < 0.1$. The r^2 is 0.73

Table VI

Cardiac *CBR1* mRNA expression, CBR1 protein expression and daunorubicin reductase activity in hearts from donors with- DS stratified by *CBR1* rs9024 genotype status

<i>CBR1</i> rs9024 genotype (n)	<i>CBR1</i> mRNA copy number ratio Mean \pm SD	<i>CBR1</i> protein (nmol/g cytosolic protein) Mean \pm SD	Daunorubicin reductase activity (nmol daunol/mg. min) Mean \pm SD
AA (2)	36423.3 \pm 38775.3	18.8 \pm 4.7	1.5 \pm 0.9
AG (6)	59954.5 \pm 9322.3	31.1	4.5 \pm 1.9
GG (21)	11138.5 \pm 47942.6	14.7 \pm 0.7	1.1 \pm 0.2

Table VII

Cardiac *CBR1* mRNA expression, CBR1 protein expression and daunorubicin reductase activity in hearts from donors without- DS stratified by *CBR1* rs9024 genotype status

<i>CBR1</i> rs9024 genotype (n)	<i>CBR1</i> mRNA copy number ratio Mean ± SD	<i>CBR1</i> protein (nmol/g cytosolic protein) Mean ± SD	Daunorubicin reductase activity (nmol daunol/mg. min) Mean ± SD
AA (2)	2719.1 ± 2206.0	3.1 ± 0.8	1.7 ± 0.6
AG (6)	29148.5 ± 50373.5	11.4 ± 6.4	2.3 ± 0.9
GG (21)	18726.8 ± 25996.4	13.2 ± 9.6	2.7 ± 1.7