

ORIGINAL INVESTIGATION

Children's Exposure to Secondhand and Thirdhand Smoke Carcinogens and Toxicants in Homes of Hookah Smokers

Nada O. F. Kassem DrPH¹, Reem M. Daffa MPH¹, Sandy Liles MPH¹, Sheila R. Jackson MPH¹, Noura O. Kassem MPH¹, Maram A. Younis MSc¹, Setoo Mehta MPH¹, Menglan Chen MS², Peyton Jacob III PhD³, Steve G. Carmella BA², Dale A. Chatfield PhD⁴, Neal L. Benowitz MD³, Georg E. Matt PhD^{1,5}, Stephen S. Hecht PhD², Melbourne F. Hovell PhD¹

¹Center for Behavioral Epidemiology and Community Health, Graduate School of Public Health, Division of Health Promotion, San Diego State University, San Diego, CA; ²Masonic Cancer Center, University of Minnesota, Minneapolis, MN; ³Departments of Medicine and Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, CA; ⁴Department of Chemistry, San Diego State University, San Diego, CA; ⁵Department of Psychology, San Diego State University, San Diego, CA

Corresponding Author: Nada O. F. Kassem, DrPH, Center for Behavioral Epidemiology and Community Health, Graduate School of Public Health, San Diego State University, 9245 Sky Park Court, Suite 230, San Diego, CA 92123, USA. Tel: 619-370-7488; Fax: 858-505-8614; E-mail: nadakassem@hotmail.com

Received October 21, 2013; accepted January 19, 2014

ABSTRACT

Introduction: We examined homes of hookah-only smokers and nonsmokers for levels of indoor air nicotine (a marker of secondhand smoke) and indoor surface nicotine (a marker of thirdhand smoke), child uptake of nicotine, the carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), and the toxicant acrolein by analyzing their corresponding metabolites cotinine, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) and NNAL-glucuronides (total NNAL) and 3-hydroxypropylmercapturic acid.

Methods: Data were collected at 3 home visits during a 7-day study period from a convenience sample of 24 households with a child 5 years or younger. Three child urine samples and 2 air and surface samples from the living room and the child bedroom were taken in homes of nonsmokers ($n = 5$) and hookah-only smokers ($n = 19$) comprised of daily hookah smokers ($n = 8$) and weekly/monthly hookah smokers ($n = 11$).

Results: Nicotine levels in indoor air and on surfaces in the child bedrooms in homes of daily hookah smokers were significantly higher than in homes of nonsmokers. Uptake of nicotine, NNK, and acrolein in children living in daily hookah smoker homes was significantly higher than in children living in nonsmoker homes. Uptake of nicotine and NNK in children living in weekly/monthly hookah smoker homes was significantly higher than in children living in nonsmoker homes.

Conclusions: Our data provide the first evidence for uptake of nicotine, the tobacco-specific lung carcinogen NNK, and the ciliotoxic and cardiotoxic agent acrolein in children living in homes of hookah smokers. Our findings suggest that daily and occasional hookah use in homes present a serious, emerging threat to children's long-term health.

INTRODUCTION

Secondhand smoke (SHS), a by-product of tobacco smoking, is an indoor toxic air contaminant, contains human carcinogens associated with illnesses in children (California Environmental Protection Agency, 2006; Centers for Disease Control and Prevention [CDC], 2010a; Environmental Protection Agency, 1992). There are no known safe levels of SHS exposure; SHS is a toxic mix of more than 7,000 chemicals that kills each year more than 600,000 nonsmokers globally (CDC, 2010a; World Health Organization [WHO], 2013). The WHO (1999) estimated

that 700 million or nearly half of the world's children may be exposed to SHS, particularly at home. In the United States, during 2007–2008, the National Health and Nutrition Examination Survey (NHANES) found that about half (53.6%) of children aged 3–11 years were exposed to SHS (CDC, 2010b). Children living in homes of smokers are additionally at risk of exposure to thirdhand smoke (THS). THS consists of residual or aged tobacco smoke particles and toxicants, including nicotine, which remains after tobacco has been smoked and adheres to and is emitted from household surfaces (Matt et al., 2011a). Hang et al. (2013) reported that exposure to THS is genotoxic in human cell

doi:10.1093/ntr/ntu016

Advance Access publication March 3, 2014

© The Author 2014. Published by Oxford University Press on behalf of the Society for Research on Nicotine and Tobacco. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

Hookahs and secondhand smoke

lines. Research on SHS and THS tends to focus on cigarettes; however, hookah smoking, another method of tobacco use, as a source of SHS and THS has not been studied.

A hookah smoker inhales tobacco smoke through a hose attached to a hookah stem topped with hookah tobacco covered with perforated aluminum foil and inserted into a glass base partially filled with water (Khater, Abd El-Aziz, Al-Sewaidan, & Chaouachi, 2008; Shihadeh, 2003). Burning charcoal is placed on top of the aluminum foil to heat the hookah tobacco, which produces the smoke that the user inhales. Tobacco used for hookah smoking is either unflavored or flavored. Flavored tobacco, known as “Moassel,” is a mixture of fruit-flavored tobacco, molasses, artificial flavoring substances, and humectants such as glycerol (Khater et al., 2008; Maziak, Ward, Afifi Soweid, & Eissenberg, 2004; Schubert, Luch, & Schulz, 2013).

Hookah tobacco smoking is on the rise globally (Cobb, Ward, Maziak, Shihadeh, & Eissenberg, 2010; Maziak, 2011). In the United States, in 2013, 26.6% of male and 23.2% of female college students nationally reported ever hookah use (American College Health Association, 2013). Among adults, in 2008, the California Tobacco Surveys showed that 11.2% of males and 2.8% of females ever used hookah (Al-Delaimy et al., 2010). Among middle and high school students, in 2011, the National Youth Tobacco Survey showed that 8.1% of males and 6.6% of females nationally ever used hookah (Amrock, Gordon, Zelikoff, & Weitzman, 2014).

Many hookah smokers are smoking hookah in home settings. In the United States, 43.4%–79.0% of hookah smoker university students surveyed reported smoking hookah at home or in their dormitory (Heinz et al., 2013; Lipkus, Eissenberg, Schwartz-Bloom, Prokhorov, & Levy, 2011). In Syria, nearly half (49.2%) of daily hookah smokers surveyed (mean age, 30.1 years) reported that their usual place of smoking hookah was mainly home (Maziak, Ward, & Eissenberg, 2004). This is potentially alarming particularly with respect to potential exposure of children living in these homes.

Hookah tobacco smoking has been associated with increased risk for periodontal disease, lung and oral cancers, coronary heart and pulmonary disease (Akl et al., 2010; Raad et al., 2011; Shaikh, Vijayaraghavan, Sulaiman, Kazi, & Shafi, 2008). Studies have identified carcinogens and toxicants such as polyaromatic hydrocarbons and volatile aldehydes in hookah tobacco smoke and biomarkers of these toxicants in the urine of hookah smokers (Al Rashidi, Shihadeh, & Saliba, 2008; Daher et al., 2010; Jacob et al., 2011, 2013).

Nicotine and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) are the most important tobacco-specific markers of SHS and SHS exposure (Benowitz & Jacob, 1994; Hecht, 2002). Nicotine is the main addictive constituent of tobacco products; its metabolite, cotinine, is the most widely used biological marker of recent SHS exposure (Benowitz & Jacob 1994). NNK is a potent tobacco-specific pulmonary carcinogen (Hecht, 2002). Its metabolites, 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) and NNAL-glucuronides (total NNAL) are consistently elevated in adult nonsmokers and children exposed to SHS (Bernert et al., 2010; Hecht, 2002). Another toxic compound, acrolein, is a strong ciliotoxic agent, also believed to contribute to lung carcinogenesis via DNA damage and inhibition of DNA repair, as well as to cardiovascular disease via its oxidant properties (Feng, Hu, Hu, & Tang, 2006; Luo et al., 2007). Acrolein is one of the most abundant, reactive toxicants in cigarette smoke (Feng et al.,

2006). A major metabolite of acrolein used for biomonitoring is 3-hydroxypropylmercapturic acid (3-HPMA), which can be measured in urine (Carmella et al., 2007).

We are not aware of any previous studies that have examined SHS, THS, and carcinogen and other toxicant uptake by children who live in homes of hookah-only smokers. These measurements are crucial for assessing the possible negative health consequences to children who live in these homes.

METHODS

Study Design

A cross-sectional nonequivalent group comparison study was employed. Research assistants (RAs) collected data from a convenience sample ($N = 24$) of households with hookah-only smokers ($n = 19$): (daily hookah smokers [$n = 8$], weekly/monthly hookah smokers [$n = 11$]), and nonsmokers ($n = 5$). Data were collected during three home visits during a 7-day study period between 2010 and 2011: tobacco use, demographics, a 7-Day Home Tobacco Use Diary, a 7-Day Child Observation Diary, home and household characteristics, two environmental samples (air and surface) per home, and three urine samples per child. Adult participants received \$150 as an incentive. The study was approved by the Institutional Review Board of San Diego State University (SDSU).

Inclusion Criteria

Homes were eligible if participants had lived in their current home for at least 6 months and their home was either a “smoker home” or a “nonsmoker home” as described previously (Matt et al., 2011b). “Smoker homes” were those in which residents have smoked hookah-only tobacco indoors or outdoors (on the patio or balcony) during at least 5 of the past 6 months, including the current most recent month, and have not smoked cigarettes or any other tobacco products anywhere inside the home or outdoors in the past 6 months. “Nonsmoker homes” were those in which no smokers lived and no visitors were allowed to smoke indoors or outdoors. Eligible households consisted of either a male or female, 18 years or older, hookah-only smoker or nonsmoker living with a healthy child 5 years old or younger. Hookah-only smokers were eligible if they smoked only hookah at least once a month and did not smoke any other tobacco product during the past 6 months. Nonsmokers and their children were eligible if they were not exposed to SHS from any tobacco product indoors or outdoors at their home for at least the past 6 months. “Hookah-only” smokers are referred to as “hookah smokers” throughout the remaining manuscript.

Recruitment Efforts and Screening

The RAs recruited participants from the community via intercept brief screening interviews at community colleges, universities, malls, Arab American churches and mosques, Middle Eastern restaurants and grocery stores, and community events in the cities of San Diego, La Mesa, and El Cajon, CA. The RAs contacted eligible households by phone to invite them to participate and coordinate the first home visit to obtain an informed consent and start the study (see [Supplementary Material](#) for details on parent reported measures, child SHS exposure measures, and home visits protocol).

Environmental Measures

Air Nicotine

Two air samples were collected with passive diffusion monitor badges to measure air nicotine, one in the living room and one in the child bedroom. A blank nonanalyzed badge was placed in a third room as a bogus pipeline (Matt et al., 2004). Using a home sketch floor plan drawn by the RAs identifying locations of windows, doors, smoking area, the RAs placed the monitors in each home during the first home visit for the duration of 7 days. Monitors were hung in the air about 2 m from the floor, out of reach of the children and away from doors, windows, and corners and hours in the room recorded. The badges consisted of a modified 37-mm diffusive cassette with a sodium bisulfate-treated Teflon-coated glass fiber filter. Badges were transported in ice coolers to SDSU laboratory for analyses. Nicotine was extracted into hexane and analyzed on a gas chromatograph with a nitrogen detector similar to published methods (Bernert, McGuffey, Morrison, & Pirkle, 2000). The limit of detection (LOD) was 0.02 $\mu\text{g}/\text{m}^3$.

Surface Nicotine

RAs collected two surface wipes, one from the living room for both indoor smokers and outdoor smokers, and one from the child bedroom area. Surface wipes were taken from window sills or door panels during the first home visit. To preserve the nicotine, wipes were wetted with 10 drops (1 ml) of freshly prepared 0.1% ascorbic acid using sterile pipettes, as described previously (Matt et al., 2004). Surface area wipe samples were collected with a cotton wipe (cosmetic 100% cotton) using vertical and horizontal strokes covering 100 cm^2 within a 10 \times 10 cm disposable frame. Wipes were placed in sanitized 20 ml amber glass bottles and transported in ice coolers to SDSU laboratory for analysis. Blanks were not used due to a small budget. Nicotine extracts from wipes samples were determined using liquid chromatography tandem mass spectrometry (LC-MS/MS) with electrospray ionization (see details in Matt et al., 2011b). The LOD was 5 ng/100 cm^2 (0.5 $\mu\text{g}/\text{m}^2$).

Biological Measures

Three child urine samples were collected by parents on Days 1, 3, and 6 to measure urinary cotinine, total NNAL, and 3-HPMA. Parents stored the urine samples in the freezer until pickup within 48 hr by RAs and transferred in a cooler to our research center laboratory. Urine samples were aliquoted, stored in a freezer (-20°C), then sent frozen in dry ice to three laboratories.

Laboratory urine analyses for cotinine were conducted by LC-MS/MS at SDSU with a LOD of 0.05 ng/ml as previously described (Bernert et al., 2000). Laboratory urine analyses for total NNAL were conducted by LC-MS/MS at the Clinical Pharmacology Laboratory, University of California San Francisco, with a LOD of 0.25 pg/ml as previously described (Jacob et al., 2008). Laboratory urine analyses for 3-HPMA were conducted by LC-atmospheric pressure chemical ionization-MS/MS-selected reaction monitoring at the Masonic Cancer Center, University of Minnesota, with an LOD of 2 pmol/ml as previously described (Carmella et al., 2007).

Statistical Analysis

Mann-Whitney *U* tests were used to identify differences in air nicotine, surface nicotine, and biomarkers of hookah tobacco

SHS exposure in children living in the three types of homes; Wilcoxon signed-rank tests to identify differences in air and surface nicotine between living rooms and child bedrooms within all homes; Spearman's rho (ρ) correlations to identify associations between air nicotine, surface nicotine, and biomarkers; and independent *t* tests to identify differences in demographics and hookah smoking habits by type of home or type of smoker. Geometric means (GMs) and 95% confidence intervals (CIs) were computed for number of hookah heads and hours smoked in 7 days, air nicotine, surface nicotine, and biomarkers. For nondetectable values, the mean of zero and the LOD was used. All statistical tests were two-tailed with an alpha level of .05. Data analyses were conducted using SPSS version 19.

RESULTS

Child, parent/guardian, and home characteristics are presented in Table 1. All 24 children were in excellent/good physical and mental health. About half were males with a median age of 3 years. Hookah smokers were mostly parents (94.7%, $n = 18$), males (73.7%, $n = 14$), and of Middle Eastern descent (94.7%, $n = 18$). Both hookah smokers and nonsmokers had been living in the same home for more than 6 months.

Hookah smoking habits are presented in Table 2. Hookah-only smokers were comprised of 8 daily and 11 weekly/monthly smokers. All hookah smokers owned a hookah at home and smoked only flavored hookah tobacco "Moassel" and no other tobacco products. Table 3 presents the GM and 95% CIs of air and surface nicotine levels in living rooms and child bedrooms and number of hookah heads and hours smoked in 7 days. A hookah head was defined as one hookah tobacco serving, which was equivalent to 10–20 g of hookah tobacco (Monzer, Sepetdjian, Saliba, & Shihadeh, 2008).

Air Nicotine Levels

Living Rooms

Nicotine was detected in the air of the living rooms of 88% (7 of 8 homes) of daily hookah smoker homes, 60% (6 of 10 homes; one monitor missing) of the weekly/monthly hookah smoker homes compared to 20% (1 of 5 homes) of the nonsmoker homes. In daily hookah smoker homes, GM air nicotine levels were 14.3 and 4.8 times higher, respectively, than those found in the living rooms of nonsmoker homes and weekly/monthly hookah smoker homes. In weekly/monthly hookah smoker homes, GM air nicotine levels were 3 times higher than those found in nonsmoker homes.

Child Bedrooms

Nicotine was detected in the air of the child bedrooms of 88% (7 of 8 homes) of daily hookah smoker homes, 55% (6 of 11 homes) of weekly/monthly smoker homes compared to none (0 of 5 homes) of the nonsmoker homes. In daily hookah smoker homes, the GM air nicotine levels were significantly 41 and 13.7 times higher, respectively, than those found in the child bedrooms of nonsmoker homes and weekly/monthly hookah smoker homes. In weekly/monthly hookah smoker homes, GM air nicotine levels were 3 times higher than those found in the child bedrooms of nonsmoker homes; however, the difference was not significant.

Hookahs and secondhand smoke

Table 1. Characteristics of Child Participants, Parents/Guardians, Households, and Homes ($N = 24$)^{a,b}

	Nonsmoker households ($n = 5$), n (%)	Hookah-only smoker households ($n = 19$), n (%)	p^c value	Frequency of hookah-only smoking		p value
				Daily hookah smoker ($n = 8$), n (%)	Weekly/monthly hookah smoker ($n = 11$), n (%)	
Child						
Age (years)						
Mean \pm <i>SD</i>	3.30 \pm 0.7	2.68 \pm 1.1	.262	2.38 \pm 1.1	2.91 \pm 1.2	.324
Median (range)	3.0 (2.5–4)	3.0 (1–4.5)		2.3 (1–4)	3.0 (1–4)	
Gender						
Male	2 (40.0)	9 (47.4)		3 (37.5)	6 (54.5)	
Female	3 (60.0)	10 (52.6)		5 (62.5)	5 (45.5)	
Physical health						
Excellent	4 (80.0)	10 (52.6)		3 (37.5)	7 (63.6)	
Good	1 (20.0)	9 (47.4)		5 (62.5)	4 (36.4)	
Mental health						
Excellent	4 (80.0)	11 (57.9)		4 (50.0)	7 (63.6)	
Good	1 (20.0)	8 (42.1)		4 (50.0)	4 (36.4)	
Parent or guardian						
Age (years)						
Mean \pm <i>SD</i>	36.4 \pm 1.5	39.2 \pm 8.2	.181	43.1 \pm 5.3	36.3 \pm 8.9	.069
Median	37.0	40.0		42	34	
Gender						
Male	0 (0.0)	14 (73.7)		4 (50.0)	10 (90.9)	
Female	5 (100)	5 (26.3)		4 (50.0)	1 (9.1)	
Ethnicity						
Middle Eastern	0 (0.0)	18 (94.7)		8 (100)	10 (90.9)	
Non-Middle Eastern White	5 (100)	1 (5.3)		0 (0.0)	1 (9.1)	
Relationship to child						
Mother	5 (100)	5 (26.3)		0 (0.0)	5 (45.5)	
Father	0 (0.0)	13 (68.4)		7 (87.5)	6 (54.5)	
Other	0 (0.0)	1 (5.3)		1 (12.5)	0 (0.0)	
Household and home						
Months residing in home						
Mean \pm <i>SD</i>	43.0 \pm 43.4	20.1 \pm 18.4	.308	15.4 \pm 10.8	23.5 \pm 22.3	.355
Median	18.0	13.0		12.5	13.0	
Number of hookah smokers						
One	0 (0.0)	10 (52.6)		4 (50.0)	6 (60.0)	
Two	0 (0.0)	9 (47.4)		4 (50.0)	4 (40.0)	
Type of home						
House	4 (80.0)	7 (36.8)		1 (12.5)	6 (54.5)	
Apartment	1 (20.0)	12 (63.2)		7 (87.5)	5 (45.5)	
Number of bedrooms						
Two	1 (20.0)	10 (52.6)		5 (62.5)	5 (45.5)	
Three or more	4 (80.0)	9 (47.4)		3 (37.5)	6 (54.5)	

^aHookah Smoking History Questionnaire, we used open-ended questions for continuous variables.

^bHome and Household Characteristics Form.

^cIndependent samples t test for significance at $p < .05$.

Correlations among air nicotine, surface nicotine, and biomarkers are presented in Table 4. Air nicotine levels in the living rooms were positively correlated with the total number of hookah heads smoked during the 7 days, and air nicotine levels in the child bedrooms were positively correlated with air nicotine levels in the living rooms.

Surface Nicotine Levels

Living Rooms

Nicotine was detected on surfaces in the living rooms of 100% (8 of 8 homes) of daily hookah smoker homes, 91% (10 of 11

homes) of weekly/monthly smoker homes compared to 40% (2 of 5 homes) of nonsmoker homes. In daily hookah smoker homes, GM surface nicotine levels in living rooms were significantly 61.1 times higher than those found in nonsmoker homes and 5.4 times higher than found in weekly/monthly hookah smoker homes. In week/monthly hookah smoker homes, GM surface nicotine levels were significantly 11.4 times higher than those found in the living rooms of nonsmoker homes.

Child Bedrooms

Nicotine was detected on surfaces in the child bedrooms of 100% (8 of 8 homes) of daily hookah smoker homes, 91% (10 of 11

Table 2. Hookah Smoking Habits of Parents/Guardians ($N = 24$)^a

	Nonsmoker ($n = 5$), n (%)	Hookah-only smoker ($n = 19$), n (%)	Frequency of hookah-only smoking		p value ^b
			Daily hookah smoker ($n = 8$), n (%)	Weekly/monthly hookah smoker ($n = 11$), n (%)	
Do you currently smoke hookah daily, weekly, monthly, or not at all?					
Daily	N/A	8 (42.1)	8 (100)	0 (0.0)	
Weekly		5 (26.3)	0 (0.0)	5 (45.5)	
Monthly		6 (31.6)	0 (0.0)	6 (54.5)	
How many hookah heads do you usually smoke on the day you smoke?					
Means \pm SD (hookah heads)	N/A	1.47 \pm 0.772	2.13 \pm 0.835	1 \pm 0.0	<.001
Median (range)		1 (1–3)	2 (1–3)	1 (1–1)	
How long do you smoke hookah on the day you smoke?					
Means \pm SD (minutes)	N/A	87 \pm 84.2	106.9 \pm 121.4	72.7 \pm 43.6	.397
Median (range)		60 (15–300)	45 (15–300)	60 (30–180)	
Which days of the week do you usually smoke hookah at home?					
Monday–Thursday	N/A	12 (66.7)	7 (87.5)	5 (50.0)	
Friday		12 (66.7)	7 (87.5)	5 (50.0)	
Saturday		16 (88.9)	8 (100.0)	8 (80.0)	
Sunday		14 (77.8)	8 (100.0)	6 (60.0)	
What times do you usually smoke hookah at home?					
Mornings (6 a.m.–12:59 p.m.)	N/A	4 (21.1)	4 (50.0)	0 (0.0)	
Afternoons (1 p.m.–5:59 p.m.)		7 (36.8)	6 (75.0)	1 (9.1)	
Evenings (6 p.m.–12:59 a.m.)		14 (73.7)	4 (50.0)	10 (90.9)	
Nights (1 a.m.–5:59 a.m.)		2 (10.5)	1 (12.5)	1 (9.1)	
Did you smoke hookah during the past 7 days?					
Yes	N/A	16 (84.2)	8 (100)	8 (72.7)	
No		3 (15.8)	0 (0.0)	3 (27.3)	
Do you currently own a hookah at your home?					
Yes	0 (0.0)	19 (100)	8 (100)	11 (100)	
No	5 (100)	0 (0.0)	0 (0.0)	0 (0.0)	
What type of hookah tobacco do you currently smoke?					
Smoke only flavored hookah tobacco (Moassel)	N/A	19 (100)	8 (100)	11 (100)	
Smoke only unflavored hookah tobacco (Ajami)		0 (0.0)	0 (0.0)	0 (0.0)	
Smoke both equally		0 (0.0)	0 (0.0)	0 (0.0)	
In the past 6 months, have any other tobacco products (cigarettes, cigar, pipe, etc.) been smoked inside your home?					
Yes	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	
No	5 (100)	19 (100)	8 (100)	11 (100)	
In the past 6 months, was hookah being smoked:					
Mostly inside your home	N/A	8 (42.1)	2 (25.0)	6 (54.5)	
Only outdoors at your home		11 (57.9)	6 (75.0)	5 (45.5)	
If smoked outside home on patio or balcony ^c					
How often patio or balcony doors were open?					
Almost always/sometimes	N/A	11 (100)	6 (100)	5 (100)	
Never		0 (0.0)	0 (0.0)	0 (0.0)	

^aHookah Smoking History Questionnaire, we used open-ended questions for continuous variables.

^bIndependent samples t test for significance at $p < .05$; statistically significant p values are noted in bold.

^cHookah Smoking Session Form.

homes) of weekly/monthly smoker homes compared to 40% (2 of 5 homes) of nonsmoker homes. In daily hookah smoker homes, GM surface nicotine levels in child bedrooms were significantly 47.5 times higher than those found in nonsmoker homes and 1.9 times higher than found in weekly/monthly hookah smoker

homes. In weekly/monthly hookah smoker homes, GM surface nicotine levels were significantly 24.5 times higher than those found in the child bedrooms of nonsmoker homes.

Surface nicotine levels in the living rooms were positively correlated, respectively, with total number of hours hookah

Table 3. Continued

	Hookah-only smoker homes by frequency of smoking							
	Nonsmoker homes (<i>n</i> = 5), GM (95% CI) ^a	Hookah-only smoker homes (<i>n</i> = 19), GM (95% CI)	<i>p</i> value ^b	Daily hookah smoker homes (<i>n</i> = 8), GM (95% CI)	<i>p</i> value ^c	Weekly/monthly hookah smoker homes (<i>n</i> = 11), GM (95% CI)	<i>p</i> value ^d	<i>p</i> value ^e
Surface wipes nicotine (µg/m ²) ⁿ								
Living room	1.37 (0.00–5.85)	31.98 (11.75–84.33)	.014	83.67 (12.50–529.94)	.013	15.62 (6.36–36.54)	.047	.083
Range	0.02–13.85	0.01–2,741.74		0.09–2,741.74		0.01–158.95		
% , Freq/ <i>n</i> above LOD	40%, 2/5	95%, 18/19		100%, 8/8		91%, 10/11		
Child's room	1.36 (0.00–6.61)	44.06 (16.87–112.63)	.012	64.60 (13.25–301.01)	.013	33.29 (9.48–111.18)	.036	.620
Range	0.01–20.82	0.01–7,180.90		8.10–7,180.90 ^o		0.01–1,033.76		
% , Freq/ <i>n</i> above LOD	40%, 2/5	95%, 18/19		100%, 8/8		91%, 10/11		
<i>p</i> ^p	.953	.809		.638		.307		

Note. LOD = limit of detection.

^aGM (95% CI) = geometric mean and 95% confidence interval.

^{b–e}*p* values were derived from Mann–Whitney *U* tests on raw data; two-tailed with an alpha level *p* < .05. *p*^b, *p*^c, *p*^d, *p*^e, respectively, ^bhookah smokers vs. nonsmokers, ^cdaily smokers vs. nonsmokers, ^dweekly/monthly smokers vs. nonsmokers, ^edaily smokers vs. weekly/monthly smokers.

^fParent reported number of hookah heads and number of hours hookah smoked in 7 days were calculated from the 7-Day Home Tobacco Use Diary.

^gNicotine air samples were collected with passive diffusion monitor badges during 7 days and were placed in the living room and child room.

^hFor example, 548 ng × 1 µg/1,000 ng/(0.252 m³) = 2.18 µg/m³.

ⁱPercentage (%) of samples with nicotine levels above the LOD: nicotine air LOD = 0.02 µg/m³ per 7 days, nicotine surface wipes LOD = 0.5 µg/m².

^jFreq/*n* = frequency of samples with levels above the LOD/sample size of samples per group.

^kFor example, 1 surface wipe/living room/home × 8 homes = 8 wipes.

^l(1 monitor missing). Air nicotine levels were quantified in nanograms (ng) and were reported in micrograms/cubic meter (µg/m³) as follows: 25 cm³/min (rate of monitor air sampling) × 1,440 min/day × 1 m³/106 cm³ × 7 day = 0.252 m³.

^mAir nicotine living room vs. child room.

ⁿSurface wipes were collected from the living room and child room, one wipe per room using 10 cm by 10 cm surface area for wiping (100 cm² = 0.01 m²). Surface nicotine levels were quantified in ng and were reported in micrograms/meter squared (µg/m²).

^oFor example, 71,808.9 ng × 1 µg/1,000 ng/(0.01 m²) = 7,180.9 µg/m². Statistically significant *p* values are noted in bold.

^pSurface nicotine living room vs. child bedroom using Wilcoxon log-rank tests on raw data.

Table 4. Spearman's Rho Correlations of Child Urinary Biomarkers of Hookah-Only Tobacco SHS Exposure With Air and Surface Nicotine

	Air nicotine living room	Air nicotine child bedroom	Surface nicotine living room	Surface nicotine child bedroom	Cotinine	Total NNAL	3-HPMA	Total ^a hookah smoked	Total ^a number hookah smoked	Number of bedrooms in homes	Total ^a hours child exposed to SHS	Child age
Air nicotine living room	1	.483 ^b	.322	.133	.254	.415	.043	.302	.512	-.119	NA	NA
Air nicotine child bedroom		.019^c	.134	.544	.242	.049	.847	.162	.013	.590	NA	NA
Surface nicotine living room		1	.416	.297	.374	.417	.343	.294	.377	-.322	NA	NA
Surface nicotine child bedroom			.043	.158	.072	.042	.101	.164	.069	.125	NA	NA
			1	.635	.358	.508	.403	.404	.469	-.122	NA	NA
Surface nicotine living room				.001	.086	.011	.051	.050	.021	.572	NA	NA
Surface nicotine child bedroom				1	.173	.397	.203	.343	.471	-.160	NA	NA
Cotinine					.419	.055	.340	.101	.020	.455	NA	NA
					1	.625	.157	.596	.598	-.352	.670	-.313
Total NNAL						.001	.465	.002	.002	.092	< .001	.137
						1	.217	.353	.582	-.300	.521	-.432
3-HPMA							.310	.090	.003	.154	.009	.035
							1	.140	.131	-.490	.314	-.359
								.513	.541	.015	.136	.085

Note. 3-HPMA = 3-hydroxypropylmercapturic acid; NA = not applicable; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; SHS = secondhand smoke. Biomarkers units: cotinine ng/mg creatinine, NNAL pg/mg creatinine, 3-HPMA pmol/mg creatinine.

^aIn 7 days.

^bSpearman's rho correlation coefficient (ρ).

^c*p* value; statistically significant *p* values are noted in bold.

smoked and total number of hookah heads smoked in 7 days. Surface nicotine levels in the child bedrooms were positively correlated with surface nicotine levels in living rooms and to total number of hookah heads smoked in 7 days.

Child Exposure to Hookah SHS

Urinary levels of cotinine, total NNAL, and 3-HPMA are presented in Table 5. We included uncorrected and creatinine-corrected maximum values of the biomarkers (GM of the highest level of three spot urine samples collected during 1 week), and the average values (GM of the average level of three spot urine samples collected during 1 week). Children living in daily hookah smoker homes were exposed to hookah SHS 1.6 times longer than children living in weekly/monthly hookah smoker homes.

Urine cotinine was detected in 75% (18 of 24 samples) of samples from children living in daily hookah smoker homes compared to 85% (28 of 33 samples) of samples from children living in weekly/monthly hookah smoker homes and 60% (9 of 15 samples) of samples from children living in nonsmoker homes. GM urine cotinine levels in children living in daily hookah smoker homes were significantly 6.5 times higher than those found in children living in nonsmoker homes and 1.8 times higher than in children living in weekly/monthly hookah smoker homes. GM urine cotinine levels in children living in week/monthly hookah smoker homes were significantly 3.7 times higher than those found in children living in nonsmoker homes.

Urine total NNAL levels were detected in 71% (17 of 24 samples) of samples from children living in daily hookah smoker homes compared to 67% (22 of 33 samples) of samples from children living in weekly/monthly hookah smoker homes and 7% (1 of 15 samples) of urine from children living in nonsmoker homes. GM urine total NNAL levels in children living in daily hookah smoker homes were significantly 37.3 higher than those found in children living in nonsmoker homes and 2.2 times higher than in children living in weekly/monthly hookah smoker homes. GM urine total NNAL levels in children living in week/monthly hookah smoker homes were significantly 17 times higher than those found in children living in nonsmoker homes.

Urine 3-HPMA was detected in all children. GM urine 3-HPMA levels in children living in daily hookah smoker homes were significantly 1.9 times higher than those found in children living in nonsmoker homes and 1.4 times higher than in children living in weekly/monthly hookah smoker homes. GM urine 3-HPMA levels in children living in week/monthly hookah smoker homes were 1.37 times higher than those found in children living in nonsmoker homes.

DISCUSSION

This is the first study to investigate nicotine levels in indoor air and on surfaces inside homes of hookah-only smokers, and carcinogen and other toxicant uptake in children living in these homes. Our results demonstrate significantly higher exposures to nicotine, NNK, and acrolein in children who live in such homes compared to children who live in nonsmoker homes. These results point to potential dangers of hookah smoking in homes with children.

We compared our data from homes of daily hookah smokers to data from studies on daily cigarette smoker households with children. Hookah smoking differs from cigarette smoking in that daily hookah smokers smoke less frequently per day, however, for longer time per smoking session (WHO, 2006). Our sample of daily hookah smokers reported that on the day they smoke, they usually smoked an average of two hookah heads for an average of 100 min (Table 2).

All hookah smokers in our study smoked “Moassel,” which contains about 30% tobacco (Khater et al., 2008). Analysis of nicotine content of 11 commercial brands of “Moassel” showed that the average nicotine content of “Moassel” was 3.35 mg/g tobacco (range, 1.8–6.3 mg/g) (Hadidi & Mohammed, 2004). Accordingly, the average nicotine content of one hookah head of 20 g Moassel is 67 mg/hookah head (range, 36–126 mg) compared to 10.2 mg/cigarette (Hadidi & Mohammed, 2004; Kozlowski et al., 1998). To date, the U.S. Food and Drug Administration (FDA) has not regulated hookah tobacco or its many flavors and additives. Although the Family Smoking Prevention and Tobacco Control Act, which gave the FDA the authority to regulate tobacco products to protect public health, became law in 2009, the FDA has been waiting for evidence of toxicological exposures to initiate regulatory practices (FDA, 2013). This study contributes to the evidence of hookah tobacco SHS toxicological exposures in children.

In contrast to beliefs that hookah tobacco is less harmful than cigarettes, the CDC (2011) reported that hookah smoking is not a safe alternative to smoking cigarettes (Akl et al., 2013). We detected relatively high levels of nicotine on surfaces in living rooms and child bedrooms of 18 out of 19 hookah smoker homes. Children living in these homes are at risk for exposure to tobacco THS throughout their homes. GM surface nicotine levels in daily hookah smoker homes were higher than levels found in indoor daily cigarette smoker homes (living rooms: GM, 83.67 vs. 51.3 $\mu\text{g}/\text{m}^2$; child bedrooms: GM, 64.6 vs. 41.9 $\mu\text{g}/\text{m}^2$, respectively) (Matt et al., 2011b).

Nicotine contamination levels on home surfaces are due to accumulation of THS over time (Quintana et al., 2013). A culturally based aspect of hookah smoking behavior may have influenced the relatively high levels of nicotine on surfaces in homes of hookah smokers. Our study sample of hookah smokers was comprised mainly of Arab Americans. Reported and observed behaviors of hookah smoker participants who smoke outdoors may have allowed hookah tobacco smoke to drift inside the home. Based on home sketches and the location of smoking, outdoor hookah smokers (11 of 19 smokers) reported smoking on the patio or the balcony near their living room door. They also reported that patio or balcony doors were either almost always or sometimes open when smoking (Table 2). Reasons for having the doors open, as identified by the RAs through direct observation, were to allow family members in and out of the home to bring food and drinks, to welcome incoming guests for socializing, or for children to play indoors and outdoors.

The majority of smokers reported that they usually smoked hookah during the afternoon or evening hours and over the weekend, times used for socializing (Table 2). This paper focused on frequency of hookah smoking; however, future investigations are needed to compare SHS and THS levels inside homes of indoor versus outdoor hookah smoker households.

Table 5. Urinary Levels of Cotinine, Total NNAL, and 3-HPMA in Children (≤5 Years) Living in Homes of Hookah-Only Smokers and Nonsmokers (N = 24)

	Hookah-only smoker homes by frequency of smoking			
	Nonsmoker homes (n = 5), GM (95% CI) ^a	Hookah-only smoker homes (n = 19), GM (95% CI)	Daily hookah smoker Homes (n = 8), GM (95% CI)	Weekly/monthly hookah smoker homes (n = 11), GM (95% CI)
Child exposure to hookah SHS				
Hours exposed in 7 days ^f	0 (0-0)	3.39 (2.01-5.39)	4.42 (1.92-9.06)	2.76 (1.35-5.04)
Median		2.17	4.21	1.92
Range (hours)		0.02-19	0.33-19	0.02-15.17
Cotinine				
Maximum ^g				
ng/ml	0.34 (0.07-0.68)	0.92 (0.51-1.44)	1.21 (0.38-2.52)	0.73 (0.35-1.22)
ng/mg creatinine	0.90 (0.23-1.92)	3.26 (1.82-5.43)	4.13 (1.52-9.48)	2.71 (1.24-5.15)
Average ^h				
ng/ml	0.16 (0.06-0.26)	0.65 (0.30-1.08)	0.88 (0.21-1.93)	0.49 (0.17-0.91)
ng/mg creatinine	0.44 (0.16-0.79)	2.08 (1.07-3.58)	2.85 (0.83-7.12)	1.62 (0.71-3.00)
Range in ng/mg	0.05-2.86	0.05-28.44	0.05-28.44	0.05-15.85
% Freq/n above LOD ^j	60%, 9/15 ⁱ	81%, 46/57	75%, 18/24	85%, 28/33
Total NNAL				
Maximum				
pg/ml ^k	0.20 (0.0-0.57)	3.40 (1.59-6.45)	4.70 (1.10-14.46)	2.64 (1.06-5.45)
pg/mg creatinine	0.56 (0.20-1.03)	8.53 (3.92-17.47)	12.42 (2.98-44.24)	6.43 (2.53-14.63)
Average				
pg/ml	0.11 (0.0-0.24)	2.55 (1.15-4.87)	3.70 (0.80-11.30)	1.89 (0.71-3.90)
pg/mg creatinine	0.28 (0.14-0.43)	6.68 (2.93-13.99)	10.43 (2.27-38.98)	4.75 (1.80-10.81)
Range in pg/mg	0.08-1.43	0.08-242.20	0.15-242.20	0.08-66.32
% Freq/n above LOD	7%, 1/15	68%, 39/57	71%, 17/24	67%, 22/33

(Continued)

Table 5. Continued

	Hookah-only smoker homes by frequency of smoking			
	Nonsmoker homes (n = 5), GM (95% CI) ^a	Hookah-only smoker homes (n = 19), GM (95% CI)	Daily hookah smoker Homes (n = 8), GM (95% CI)	Weekly/monthly hookah smoker homes (n = 11), GM (95% CI)
			p value ^b	p value ^c
3-HPMA				
Maximum pmol/ml ^l	991 (479–2,050)	1,085 (749–1,573)	.546	.999
pmol/mg creatinine	2,306 (1,423–3,737)	3,665 (2,847–4,718)	.110	.057
Average				
pmol/ml	616 (339–1,121)	714 (495–1,031)	.414	.770
pmol/mg creatinine	1,600 (1,112–2,303)	2,488 (1,890–3,277)	.095	.040
Range in pmol/mg	553–5,864	300–8,889	778–7,866	300–8,889
% Freq/n above LOD	100%, 15/15	100%, 57/57	100%, 24/24	100%, 33/33

Note. 3-HPMA = 3-hydroxypropylmercapturic acid; LOD = limit of detection; NNAL = 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol; SHS = secondhand smoke.

^aGM (95% CI) = geometric mean and 95% confidence interval.

^{b–e}p values were derived from Mann–Whitney U tests on raw data; two-tailed with an alpha level $p < .05$; statistically significant p values are noted in bold. ^b, ^c, ^d, ^ep values are noted in bold. ^b, ^c, ^d, ^ep values are noted in bold. ^b, ^c, ^d, ^ep values are noted in bold. ^b, ^c, ^d, ^ep values are noted in bold. ^b, ^c, ^d, ^ep values are noted in bold.

^fhookah smokers vs. nonsmokers, ^gdaily smokers vs. nonsmokers, ^hweekly/monthly smokers vs. nonsmokers, ⁱdaily smokers vs. weekly/monthly smokers.

^jNumber of hours child exposed to hookah SHS in 7 days was calculated from reported data from two diaries: 7-Day Home Tobacco Use Diary and 7-Day Child observation Diary.

^kMaximum = GM of the highest level of three spot urine samples collected during 1 week, uncorrected and corrected by creatinine.

^lAverage = GM of the average level of three spot urine samples collected during 1 week, uncorrected and corrected by creatinine.

^mPercentage (% rounded up) of urine samples above the LOD: cotinine LOD = 0.05 ng/ml, NNAL LOD = 0.25 pg/ml, and 3-HPMA LOD = 2 pmol/ml.

ⁿFreq/n = frequency of samples with levels above the LOD/sample size of samples per group, for example, three urine sample per child × 5 homes = 15 urine samples.

^oMolecular weight for NNAL is 209; to convert from pg/ml to pmol/ml divide by 209, for example, 0.25 pg/ml = 0.0012 pmol/ml.

^pMolecular weight for 3-HPMA is 221; to convert from pmol/ml to pg/ml multiply by 221, for example, 2 pmol/ml × 221 = 442 pg/ml.

Hookahs and secondhand smoke

Also, seven of eight daily hookah smokers lived in apartments. Of these, almost half (47.4%) reported having two hookah smokers per household. This is of concern in an apartment complex because hookah tobacco smoke could enter other apartments. Nonsmokers who reside in multiunit housing may share the same air space as those who smoke in adjacent units (King, Travers, Cummings, Mahoney, & Hyland, 2010). King et al. (2010) documented that SHS can transfer between living units within the same multiunit housing building. Therefore, emerging smoke-free policies in multiunit housing restricting cigarette smoking to protect residents from exposure to SHS should take into consideration hookah smoking.

Urinary levels of cotinine (GM, 0.88 ng/ml) and total NNAL (GM, 3.7 pg/ml) in children living in homes of daily hookah smokers were lower than those reported in previous studies in children exposed to cigarette SHS. In a study of children ages 10 years or younger living in homes of daily cigarette smokers ($N = 79$) in Minnesota, the GM urinary levels of total cotinine and total NNAL were 11.9 ng/ml and 0.08 pmol/ml (16.7 pg/ml), respectively (Thomas et al., 2011). Elementary school children with parental reported exposure to SHS ($N = 38$) in Minnesota had GM urinary levels of total cotinine and total NNAL of 12.6 ng/ml and 0.04 pmol/ml (8.4 pg/ml), respectively (Hecht et al., 2001). Children in Moldova 5–10 years old ($N = 7$) with reported exposure to SHS at home had GM urinary levels of total cotinine and total NNAL of 4.6 ng/ml and 0.061 pmol/ml (12.8 pg/ml) (Stepanov, Hecht, Duca, & Mardari, 2006). The majority of daily hookah smokers in our sample smoked outdoors ($n = 2$ indoor smokers, $n = 6$ outdoor smokers), a larger sample of indoor hookah smokers is warranted in future studies.

Consistent with previous studies, we found a significant positive correlation between urinary cotinine and total NNAL (Hecht et al., 2001; Stepanov et al., 2006; Thomas et al., 2011). We also found significant positive correlations between the child urinary cotinine and total NNAL levels and indicators of tobacco smoke exposure, namely, number of hookah heads smoked and number of hours the child was exposed to hookah tobacco SHS per week. These correlations were stronger than previously observed between child urinary cotinine and total NNAL, respectively, and number of cigarettes (mean \pm SD, 9.5 ± 5.3 cigarettes) smoked per day at home ($r = .37$, $p < .001$; $r = .30$, $p < .01$) (Thomas et al., 2011).

Urinary total NNAL levels were positively correlated with surface nicotine in the living rooms and child bedrooms. Thomas et al. (2014) detected the presence of NNK on surfaces in most homes occupied by cigarette smokers. Therefore, more research is needed to identify associations between THS and NNK uptake in nonsmokers living in smokers' homes.

Urinary total NNAL levels were negatively correlated with child age. Data from the NHANES (2007–2008) showed that children had significantly higher concentrations of total NNAL than did adults aged 20 years or older ($p < .001$) (Bernert et al., 2010). Younger children are more likely to spend more time at home than older children, and less likely to be able to remove themselves from environments in which smoking occurs, thereby being exposed at home for a longer time to tobacco SHS (Bernert et al., 2010).

All children had 3-HPMA in their urine, independent of exposure to hookah SHS. Acrolein occurs endogenously as a lipid peroxidation product and naturally in foods (International Agency for Research on Cancer [IARC], 1995; Pan & Chung,

2002). It is also formed during the combustion of fossil fuels, wood and tobacco and during the heating of cooking oils (IARC, 1995; Yu, Chiu, Au, Wong, & Tang, 2006). Nazaroff and Singer (2004) found that population intake of acrolein from residential SHS appears to be higher than from ambient sources. Our findings are consistent with studies that demonstrate higher acrolein uptake in smokers compared to nonsmokers (Carmella et al., 2007; Hecht, Yuan, & Hatsukami, 2010). Levels of urinary 3-HPMA in our sample of children exposed to daily hookah SHS were significantly higher, respectively, than levels found in children living in nonsmoker homes (GM, 2,966 vs. 1,600 pmol/mg; $p = .040$) and were 1.6 times higher than levels found in a previous study of 21 adult nonsmokers (mean, 1,900 pmol/mg) (Carmella et al., 2007).

We found higher levels of 3-HPMA in children living in smaller houses or apartments. Urinary 3-HPMA levels were negatively correlated with the number of bedrooms in homes. Using smoking machines, Daher et al. (2010) found that a single hookah use session emitted $1,135 \pm 97$ μ g acrolein in the sidestream smoke; however, two studies found inconsistent average yields of acrolein in the gas phase of mainstream hookah smoke generated using 10 g of flavored hookah tobacco, a low yield of 11.3 and a high yield of 892 μ g/smoking session (Al Rashidi et al., 2008; Schubert, Heinke, Bewersdorff, Luch, & Schulz, 2012).

Our findings inform targeted trials to reduce SHS exposure among children, especially those with illnesses, who live in homes of hookah smokers (Stotts et al., 2011, 2013; Tyc et al., 2012). Such trials are important in the context of the studies successfully completed in the last 20 years showing that SHS exposure can be reduced when families are coached to do so (Hovell et al., 1994, 2000, 2002, 2009, 2011, 2013; Klepeis et al., 2013; Meltzer, Hovell, Meltzer, Atkins, & de Peyster, 1993).

Limitations and Recommendations

This study is limited by a small sample size and the use of a convenience sample, thus limiting power and generalizability. Most measures had wide CIs that most likely explain some of the nonsignificant findings. Our hookah smoking participants were almost exclusively Middle Easterners and the nonsmokers were non-Middle Eastern Whites. The detection of low levels of nicotine in two of five homes of nonsmokers could be due to drifting tobacco smoke or that surface nicotine levels were not corrected with field blanks. A field blank is a cotton round that is handled and transported with actual samples but is not used to wipe target surfaces (Quintana et al., 2013). Quintana et al. (2013) recommended that nicotine levels detected on field blanks are subtracted from those found on actual surface wipes to control for contamination from extraneous nicotine sources other than those targeted by the measure.

Future efforts are needed with larger sample sizes among various populations, housing arrangements, climates, taking into consideration the additive effect of migrating smoke, indoor versus outdoor smoking, and the use of surface wipe blanks to provide a more refined assessment of toxicant and carcinogen exposure from hookah tobacco SHS.

CONCLUSIONS

Our data provide the first evidence that children living in homes of hookah smokers are at risk of exposure to nicotine and the

tobacco-specific carcinogen NNK. All children had detectable levels of 3-HPMA with the highest levels in children living in homes of daily hookah smokers. Exposure to acrolein in non-smokers warrants further research. Our findings suggest that daily and occasional hookah use in homes present a potentially serious threat to children's long-term health. Our results call for clinical trials to assist hookah smoking families to reduce exposure to children and/or quit smoking completely, call for regulatory actions to limit toxicants in hookah products, and call to action for the implementation of voluntary smoke-free home rules.

SUPPLEMENTARY MATERIAL

Supplementary material can be found online at <http://www.ntr.oxfordjournals.org>

FUNDING

This work was supported by the Flight Attendant Medical Research Institute (YCSA 52364 to NOFK) and American Cancer Society (116623-RSG-09-098-01-CNE to NOFK); Instrumentation and analytical chemistry at University of California San Francisco were supported by the Flight Attendant Medical Research Institute Bland Lane Center of Excellence on Secondhand Smoke (to Rita F. Redberg); the National Institutes of Health (S10 RR026437 to PJ and P30 DA012393 to Reese T. Jones); California Tobacco-Related Disease Research Grants Program (17RT-0162H, 19CA-0164, 13RT-0161H to GEM); U.S. National Cancer Institute (CA-81301 to SSH); the National Institutes of Health (R01 HL103684-01A1, R01 CA138192-01A1 to MFH).

DECLARATION OF INTERESTS

NLB has provided expert testimony in tobacco litigation related to nicotine addiction in the past 5 years.

ACKNOWLEDGMENTS

We thank our participants, the staff of the three laboratories, and the research staff of the Center for Behavioral Epidemiology and Community Health: Aimen Khalil, MD, Jodi Kudas, Dana Dorman, and Jan Rivera.

REFERENCES

- Akl, E. A., Gaddam, S., Gunukula, S. K., Honeine, R., Jaoude, P. A., & Irani, J. (2010). The effects of waterpipe tobacco smoking on health outcomes: A systematic review. *International Journal of Epidemiology*, *39*, 834–857. doi:10.1093/ije/dyq002
- Akl, E. A., Jawad, M., Lam, W. Y., Co, C. N., Obeid, R., & Irani, J. (2013). Motives, beliefs and attitudes towards waterpipe tobacco smoking: A systematic review. *Harm Reduction Journal*, *10*, 12. doi:10.1186/1477-7517-10-12
- Al-Delaimy, W. K., White, M. M., Mills, A. L., Pierce, J. P., Emory, K., Boman, M., ... Edland, S. (2010). Final Summary Report of: Two Decades of the California Tobacco Control Program: California Tobacco Survey, 1990–2008. La Jolla, CA: University of California, San Diego. Retrieved from http://libraries.ucsd.edu/ssds/pub/CTS/cpc00009/CDPH_CTS2008_summary_report_final.pdf
- Al Rashidi, M., Shihadeh, A., & Saliba, N. A. (2008). Volatile aldehydes in the mainstream smoke of the narghile waterpipe. *Food and Chemical Toxicology*, *46*, 3546–3549. doi:10.1016/j.fct.2008.09.007
- American College Health Association. (2013). American College Health Association-National College Health Assessment II: Reference Group Executive Summary Spring 2013. Hanover, MD. Retrieved from www.acha.org/docs/ACHA-NCHA-II_ReferenceGroup_ExecutiveSummary_Spring2013.pdf
- Amrock, S. M., Gordon, T., Zelikoff, J. T., & Weitzman, M. (2014). Hookah use among adolescents in the United States: Results of a National Survey. *Nicotine Tobacco Research*, *16*, 231–237. doi:10.1093/ntr/ntt160
- Benowitz, N. L., & Jacob, P., III. (1994). Metabolism of nicotine to cotinine studied by a dual stable isotope method. *Clinical Pharmacology and Therapeutics*, *56*, 483–493. doi:10.1038/clpt.1994.169
- Bernert, J. T., McGuffey, J. E., Morrison, M. A., & Pirkle, J. L. (2000). Comparison of serum and salivary cotinine measurements by a sensitive high performance liquid chromatography-tandem mass spectrometry method as an indicator of exposure to tobacco smoke among smokers and nonsmokers. *Journal of Analytical Toxicology*, *24*, 333–339. doi:10.1093/jat/24.5.333
- Bernert, J. T., Pirkle, J. L., Xia, Y., Jain, R. B., Ashley, D. L., & Sampson, E. J. (2010). Urine concentrations of a tobacco-specific nitrosamine carcinogen in the U.S. population from secondhand smoke exposure. *Cancer Epidemiology, Biomarkers and Prevention*, *19*, 2969–2977. doi:10.1158/1055-9965.EPI-10-0711
- California Environmental Protection Agency. (2006). Environmental tobacco smoke: A toxic air contaminant. A fact sheet. Retrieved from www.arb.ca.gov/toxics/ets/factsheetets.pdf
- Carmella, S., Chen, M., Zhang, Y., Zhang, S., Hatsukami, D., & Hecht, S. (2007). Quantitation of acrolein-derived (3-hydroxypropyl)mercapturic acid in human urine by liquid chromatography-atmospheric pressure chemical ionization tandem mass spectrometry: Effects of cigarette smoking. *Chemical Research in Toxicology*, *20*, 986–990. doi:10.1021/tx700075y
- Centers for Disease Control and Prevention. (2010a). How tobacco smoke causes disease: The biology and behavioral basis for smoking-attributable disease: A report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. Retrieved from www.cdc.gov/tobacco/data_statistics/sgr/2010/consumer_booklet/pdfs/consumer.pdf
- Centers for Disease Control and Prevention. (2010b). Vital signs: Nonsmokers' exposure to secondhand smoke—United States, 1999–2008. *Morbidity and Mortality Weekly Report*, *59*, 1141–1146. Retrieved from www.cdc.gov/tobacco/data_statistics/mmwr/byyear/2010/mm59e0907a2/intro.htm
- Centers for Disease Control and Prevention. (2011). Fact sheet: Hookahs - smoking & tobacco use. Retrieved from www.cdc.gov/tobacco/data_statistics/fact_sheets/tobacco_industry/hookahs/
- Cobb, C., Ward, K. D., Maziak, W., Shihadeh, A. L., & Eissenberg, T. (2010). Waterpipe tobacco smoking: An emerging health crisis in the United States. *American Journal of Health Behavior*, *34*, 275–285. doi:10.5993/AJHB.34.3.3
- Daher, N., Saleha, R., Jaroudia, I., Sheheitlia, H., Badra, T., Sepetdjanb, E., ... Shihadeha, A. (2010). Comparison of

- carcinogen, carbon monoxide, and ultrafine particle emissions from narghile waterpipe and cigarette smoking: Sidestream smoke measurements and assessment of secondhand smoke emission factors. *Atmospheric Environment*, *44*, 8–14. doi:10.1016/j.atmosenv.2009.10.004
- Environmental Protection Agency. (1992). Respiratory health effects of passive smoking: Lung cancer and other disorders (EPA Publication No. EPA/600/6-90/006F). U.S. Environmental Protection Agency, Office of Research and Development, Office of Health and Environmental Assessment, Washington, DC. Retrieved from www.epa.gov/smokefree/pubs/etsfs.html
- Feng, Z., Hu, W., Hu, Y., & Tang, M. (2006). Acrolein is a major cigarette-related lung cancer agent: Preferential binding at p53 mutational hotspots and inhibition of DNA repair. *Proceedings of the National Academy of Sciences of the United States of America*, *103*, 15725–15726. doi:10.1073/pnas.0607031103
- Hadidi, K. A., & Mohammed, F. I. (2004). Nicotine content in tobacco used in hubble-bubble smoking. *Saudi Medical Journal*, *25*, 912–917. PMID 15235699.
- Hang, B., Sarker, A. H., Havel, C., Saha, S., Hazra, T. K., Schick, S., ... Jacob, P. III. (2013). Thirdhand smoke causes DNA damage in human cells. *Mutagenesis*, *28*, 381–391. doi:10.1093/mutage/get013
- Hecht, S. S. (2002). Human urinary carcinogen metabolites: Biomarkers for investigating tobacco and cancer. *Carcinogenesis*, *23*, 907–922. doi:10.1093/carcin/23.6.907
- Hecht, S. S., Ye, M., Carmella, S. G., Fredrickson, A., Adgate, J. L., Greaves, I. A., ... Sexton, K. (2001). Metabolites of a tobacco-specific lung carcinogen in the urine of elementary school-aged children. *Cancer Epidemiology, Biomarkers and Prevention*, *10*, 1109–1116. PMID 11700257.
- Hecht, S. S., Yuan, J. Y., & Hatsukami, D. (2010). Applying tobacco carcinogen and toxicant biomarkers in product regulation and cancer prevention. *Chemical Research in Toxicology*, *23*, 1001–1008. doi:10.1021/tx100056m
- Heinz, A. J., Giedgowd, G. E., Crane, N. A., Veilleux, J. C., Conrad, M., Braun, A. R., ... Kassel, J. D. (2013). A comprehensive examination of hookah smoking in college students: Use patterns and contexts, social norms and attitudes, harm perception, psychological correlates and co-occurring substance use. *Addictive Behaviors*, *38*, 2751–2760. doi:10.1016/j.addbeh.2013.07.009
- Hovell, M. F., Adams, M. A., Hofstetter, C. R., Martinez-Donate, A. P., González-Pérez, G. J., Rovniak, L. S., & Boman-Davis, M. C. (2013). Complete home smoking bans and anti-tobacco contingencies: A natural experiment. *Nicotine Tobacco Research*, *16*, 186–196. doi:10.1093/ntr/ntt130
- Hovell, M. F., Meltzer, S. B., Wahlgren, D. R., Matt, G. E., Hofstetter, C. R., ... Pirkle J.L. (2002). Asthma management and environmental tobacco smoke exposure reduction in Latino children: A controlled trial. *Pediatrics*, *110*, 946–956. doi:10.1542/peds.110.5.946
- Hovell, M. F., Meltzer, S.B., Zakarian, J. M., Wahlgren, D. R., Emerson, J. A., Hofstetter, C. R., ... Atkins, C. J. (1994). Reduction of environmental tobacco smoke exposure among asthmatic children: A controlled trial. *Chest*, *106*, 440–446. doi:10.1378/chest.106.2.440. Erratum in: *Chest* (1995), *107*, 1480. doi:10.1378/chest.107.5.1480-b
- Hovell, M. F., Wahlgren, D. R., Liles, S., Jones, J. A., Hughes, S. C., Matt, G. E., ... Ding, D. (2011). Providing coaching and cotinine results to preteens to reduce their secondhand smoke exposure: A randomized trial. *Chest*, *140*, 681–689. doi:10.1378/chest.10-2609.
- Hovell, M. F., Zakarian, J. M., Matt, G. E., Hofstetter, C. R., Bernert, J. T., & Pirkle, J. (2000). Effect of counseling mothers on their children's exposure to environmental tobacco smoke: Randomized controlled trial. *British Medical Journal*, *321*, 337–342. doi:http://dx.doi.org/10.1136/bmj.321.7257.337
- Hovell, M. F., Zakarian, J. M., Matt, G. E., Liles, S., Jones, J. A., Hofstetter, C. R., ... Benowitz, N. L. (2009). Counseling to reduce children's secondhand smoke exposure and help parents quit smoking: A controlled trial. *Nicotine Tobacco Research*, *11*, 1383–1394. doi:10.1093/ntr/ntp148
- International Agency for Research on Cancer. (1995). Cleaning, some chlorinated solvents and other industrial chemicals. *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, *63*, 337–372. Retrieved from <http://monographs.iarc.fr/ENG/Monographs/vol63/>
- Jacob, P., III, Abu Raddaha, A. H., Dempsey, D., Havel, C., Peng, M., Yu, L., & Benowitz, N. L. (2011). Nicotine, carbon monoxide, and carcinogen exposure after a single use of a waterpipe. *Cancer Epidemiology, Biomarkers and Prevention*, *20*, 2345–2353. doi:10.1158/1055-9965.EPI-11-0545
- Jacob, P., III, Abu Raddaha, A. H., Dempsey, D., Havel, C., Peng, M., Yu, L., & Benowitz, N. L. (2013). Comparison of nicotine and carcinogen exposure with waterpipe and cigarette smoking. *Cancer Epidemiology, Biomarkers and Prevention*, *22*, 765–772. doi:10.1158/1055-9965.EPI-12-1422
- Jacob, P., III, Havel, C., Lee, D. H., Yu, L., Eisner, M., & Benowitz, N. (2008). Subpicogram per milliliter determination of the tobacco-specific carcinogen metabolite 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol in human urine using liquid chromatography-tandem mass spectrometry. *Analytical Chemistry*, *80*, 8115–8121. doi:10.1021/ac8009005
- Khater, A. E., Abd El-Aziz, N. S., Al-Sewaidan, H. A., & Chauachi, K. (2008). Radiological hazards of Narghile (hookah, shisha, goza) smoking: Activity concentrations and dose assessment. *Journal of Environmental Radioactivity*, *99*, 1808–1814. doi:10.1016/j.jenvrad.2008.07.005
- King, B. A., Travers, M. J., Cummings, K. M., Mahoney, M. C., & Hyland, A. J. (2010). Secondhand smoke transfer in multiunit housing. *Nicotine and Tobacco Research*, *12*, 1133–1141. doi:10.1093/ntr/ntq162
- Klepeis, N. E., Hughes, S. C., Edwards, R. D., Allen, T., Johnson, M., Chowdhury, Z., ... Hovell, M. F. (2013). Promoting smoke-free homes: A novel behavioral intervention using real-time audio-visual feedback on airborne particle levels. *PLoS ONE*, *8*, e73251. doi:10.1371/journal.pone.0073251
- Kozlowski, L. T., Mehta, N. Y., Sweeney, C. T., Schwartz, S. S., Vogler, G. P., Jarvis, M. J., & West, R. J. (1998). Filter ventilation and nicotine content of tobacco in cigarettes from Canada, the United Kingdom, and the United States. *Tobacco Control*, *7*, 369–375. doi:10.1136/tc.7.4.369
- Lipkus, I. M., Eissenberg, T., Schwartz-Bloom, R. D., Prokhorov, A. V., & Levy, J. (2011). Affecting perceptions of harm and addiction among college waterpipe tobacco smokers. *Nicotine Tobacco Research*, *13*, 599–610. doi:10.1093/ntr/ntp049
- Luo, J., Hill, B. G., Gu, Y., Cai, J., Srivastava, S., Bhatnagar, A., & Prabhu, S. D. (2007). Mechanisms of acrolein-induced myocardial dysfunction: Implications for environmental and endogenous aldehyde exposure. *American Journal of Physiology, Heart and Circulatory Physiology*, *293*, H3673–H3684. doi:10.1152/ajpheart.00284.2007
- Matt, G. E., Quintana, P. J., Destailhats, H., Gundel, L. A., Sleiman, M., Singer, B. C., ... Hovell, M. F. (2011a). Thirdhand tobacco smoke: Emerging evidence and arguments for a multidisciplinary research agenda. *Environmental*

- Health Perspectives*, 119, 1218–1226. doi:10.1289/ehp.1103500
- Matt, G. E., Quintana, P. J., Hovell, M. F., Bernert, J. T., Song, S., Novianti, N., ... Larson, S. (2004). Households contaminated by environmental tobacco smoke: Sources of infant exposures. *Tobacco Control*, 13, 29–37. doi:10.1136/tc.2003.003889
- Matt, G. E., Quintana, P. J., Zakarian, J. M., Fortmann, A. L., Chatfield, D. A., Hoh, E., ... Hovell, M. F. (2011b). When smokers move out and non-smokers move in: Residential thirdhand smoke pollution and exposure. *Tobacco Control*, 20, e1. doi:10.1136/tc.2010.037382
- Maziak, W. (2011). The global epidemic of waterpipe smoking. *Addictive Behaviors*, 36, 1–5. doi:10.1016/j.addbeh.2010.08.030
- Maziak, W., Eissenberg, T. E., & Ward, K. D. (2004). Factors related to level of narghile use: The first insights on tobacco dependence in narghile users. *Drug and Alcohol Dependence*, 76, 101–106. doi:10.1016/j.drugalcdep.2004.04.007
- Maziak, W., Ward, K. D., Afifi Soweid, R. A., & Eissenberg, T. (2004). Tobacco smoking using a waterpipe: A re-emerging strain in a global epidemic. *Tobacco Control*, 13, 327–333. doi:10.1136/tc.2004.008169
- Meltzer, S. B., Hovell, M. F., Meltzer, E. O., Atkins, C. J., & de Peyster, A. (1993). Reduction of secondary smoke exposure in asthmatic children: Parent counseling. *The Journal of Asthma*, 30, 391–400. doi:10.3109/02770909309056743
- Monzer, B., Sepetdjian, E., Saliba, N., & Shihadeh, A. (2008). Charcoal emissions a source of CO and carcinogenic PAH in mainstream narghile waterpipe smoke. *Food and Chemical Toxicology*, 46, 2991–2995. doi:10.1016/j.fct.2008.05.031
- Nazaroff, W. W., & Singer, B. C. (2004). Inhalation of hazardous air pollutants from environmental tobacco smoke in US residences. *Journal of Exposure Analysis and Environmental Epidemiology*, 14(Suppl. 1), S71–S77. doi:10.1038/sj.jea.7500361
- Pan, J., & Chung, F. L. (2002). Formation of cyclic deoxyguanosine adducts from omega-3 and omega-6 polyunsaturated fatty acids under oxidative conditions. *Chemical Research in Toxicology*, 15, 367–372. doi:10.1021/tx010136q
- Quintana, P. J., Matt, G. E., Chatfield, D., Zakarian, J. M., Fortmann, A. L., & Hoh, E. (2013). Wipe sampling for nicotine as a marker of thirdhand tobacco smoke contamination on surfaces in homes, cars, and hotels. *Nicotine Tobacco Research*, 15, 1555–1563. doi:10.1093/ntr/ntt014
- Raad, D., Gaddam, S., Schunemann, H. J., Irani, J., Abou Jaoude, P., Honeine, R., & Akl, E. A. (2011). Effects of waterpipe tobacco smoking on lung function: A systematic review and meta-analysis. *Chest*, 139, 764–774. doi:10.1378/chest.10-0991
- Schubert, J., Heinke, V., Bewersdorff, J., Luch, A., & Schulz, T. (2012). Waterpipe smoking: The role of humectants in the release of toxic carbonyls. *Archives of Toxicology*, 86, 1309–1316. doi:10.1007/s00204-012-0884-5
- Schubert, J., Luch, A., & Schulz, T.G. (2013). Waterpipe smoking: Analysis of the aroma profile of flavored waterpipe tobaccos. *Talanta*, 115, 665–674. doi:10.1016/j.talanta.2013.06.022
- Shaikh, R. B., Vijayaraghavan, N., Sulaiman, A. S., Kazi, S., & Shafi, M. S. (2008). The acute effects of waterpipe smoking on the cardiovascular and respiratory systems. *Journal of Preventive Medicine and Hygiene*, 49, 101–107. Retrieved from <http://ksu.edu.sa/sites/KSUArabic/Research/ncys/Documents/r44.pdf> PMID 19278135.
- Shihadeh, A. (2003). Investigation of mainstream smoke aerosol of the argileh waterpipe. *Environmental Toxicology and Chemistry*, 41, 143–152. doi:10.1016/S0278-6915(02)00220-X
- Stotts, A. L., Evans, P. W., Green, C. E., Northrup, T. F., Dodrill, C. L., Fox, J. M., ... Hovell, M. F. (2011). Secondhand smoke risk in infants discharged from a NICU: Potential for significant health disparities? *Nicotine Tobacco Research*, 13, 1015–1022. doi:10.1093/ntr/ntt116
- Stotts, A. L., Green, C., Northrup, T. F., Dodrill, C. L., Evans, P., Tyson, J., ... Hovell, M. F. (2013). Feasibility and efficacy of an intervention to reduce secondhand smoke exposure among infants discharged from a neonatal intensive care unit. *Journal of Perinatology*, 33, 811–816. doi:10.1038/jp.2013.43
- Stepanov, I., Hecht, S. S., Duca, G., & Mardari, I. (2006). Uptake of the tobacco-specific lung carcinogen 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone by Moldovan children. *Cancer Epidemiology, Biomarkers and Prevention*, 15, 7–11. doi:10.1158/1055-9965.EPI-05-0293
- Thomas, J. L., Guo, H., Carmella, S. G., Balbo, S., Han, S., Davis, A., ... Hecht, S. S. (2011). Metabolites of a tobacco-specific lung carcinogen in children exposed to secondhand or thirdhand tobacco smoke in their homes. *Cancer Epidemiology, Biomarkers and Prevention*, 20, 1213–1221. doi:10.1158/1055-9965.EPI-10-1027
- Thomas, J. L., Hecht, S. S., Luo, X., Ming, X., Ahluwalia, J. S., & Carmella, S. G. (2014). Thirdhand tobacco smoke: A tobacco-specific lung carcinogen on surfaces in smokers' homes. *Nicotine and Tobacco Research*, 16, 26–32. doi:10.1093/ntr/ntt110
- Tyc, V., Huang, Q., Nicholson, J., Schultz, B., Hovell, M. F., Lensing, S., ... Zhang, H. (2012). Intervention to reduce secondhand smoke exposure among children with cancer: A controlled trial. *Psychooncology*, 22, 1104–1111. doi:10.1002/pon.3117
- U. S. Food and Drug Administration (FDA). (2013). Overview of the family smoking prevention and tobacco control act. Consumer fact sheet. Retrieved from www.fda.gov/downloads/TobaccoProducts/GuidanceComplianceRegulatoryInformation/UCM336940.pdf
- World Health Organization. (1999). International consultation on environmental tobacco smoke and child health: Consultation report. Retrieved from www.who.int/tobacco/research/en/ets_report.pdf
- World Health Organization. (2006). Tobacco use in Shisha: Studies on waterpipe smoking in Egypt/WHO Regional Office for the Eastern Mediterranean, Egyptian Smoking Prevention Research Institute. Retrieved from www.emro.who.int/tfi/PDF/shisha.pdf
- World Health Organization. (2013). Tobacco. Fact sheet No. 339. Retrieved from www.who.int/mediacentre/factsheets/fs339/en/index.html
- Yu, I. T., Chiu, Y. L., Au, J. S., Wong, T. W., & Tang, J. L. (2006). Dose-response relationship between cooking fumes exposures and lung cancer among Chinese nonsmoking women. *Cancer Research*, 66, 4961–4967. doi:10.1158/0008-5472.CAN-05-2932