



Published in final edited form as:

Harv Rev Psychiatry. 2014 ; 22(3): 149–161. doi:10.1097/HRP.000000000000014.

Childhood Maltreatment, Emotional Dysregulation, and Psychiatric Comorbidities

Yael Dvir, MD, Julian D. Ford, PhD, Michael Hill, BS, and Jean A. Frazier, MD

Division of Child and Adolescent Psychiatry, Department of Psychiatry, University of Massachusetts Medical School (Drs. Dvir and Frazier, and Mr. Hill); University of Connecticut School of Medicine (Dr. Ford)

Abstract

Affect dysregulation, defined as the impaired ability to regulate and/or tolerate negative emotional states, and has been associated with interpersonal trauma and post-traumatic stress. Affect regulation difficulties also play a role in many other psychiatric conditions, including anxiety disorders and mood disorders, specifically major depression in youth and bipolar disorder throughout the life span.

Exposure to traumatic events and interpersonal trauma in childhood is associated with a wide range of psychosocial, developmental, and medical impairments in children, adolescents and adults, with emotional dysregulation being a core feature that may help to account for this heightened risk. In order to understand how the developmental effects of childhood maltreatment contribute to emotional dysregulation and psychiatric sequelae, we review emotional regulation and its developmental neurobiology, and examine the research evidence of associations between childhood traumatization, emotional dysregulation, and psychiatric co-morbidities in children, adolescents and adults.

Keywords

childhood maltreatment; comorbidity; emotional regulation/dysregulation; interpersonal trauma; post-traumatic stress; posttraumatic stress disorder

Multiple interdependent processes are involved in emotions and their regulation, spanning the biological, psychological, and interpersonal domains.¹ A fundamental interplay exists between emotional and cognitive operations, which include selecting and modifying situations that have emotional significance, deploying attention, integrating information, making judgments and decisions, and selecting behavioral responses. While some emotions are generated automatically, others are elicited and fully apprehended only after considerable “meaning analysis.” In the same way, emotional regulation may be automatic or, instead, controlled with effortful cognitive processing.^{2,3} Emotion involves the coordination of physiological reactions, memory, cognitive appraisals, and behavior. Emotional regulation can be defined as having control not only over how and when, but also

the intensity and positive/negative valence with which, emotions are felt, experienced, and expressed. Emotions and their regulation occur continuously over time and may include changes in all three domains (behavior, experience, and physiology).³

Humans have the capacity to increase, maintain, or decrease negative and positive emotions in both conscious and unconscious ways that can facilitate or impede the attainment of behavioral goals.³⁻⁵ We can modify our responses to emotions by planning ahead to select or modify emotionally laden situations or by focusing on the response once an emotion has been experienced. Attentional deployment can determine what aspect of a situation is being focused on, and may result in distraction (which can decrease awareness and intensity of negative emotions) or in rumination (which may increase their salience, persistence, and intensity). Changing a thought that is associated with a situation can determine the attached meaning. Although this technique is often used to decrease emotional intensity, it can also magnify a response or alter the emotion altogether. Both emotional states and goal-directed behavior can be modulated by decreasing negative, or increasing positive, expressive behavior, or by addressing physiological responses—through the use of, for example, cognitive-behavioral therapy or medications that decrease negative affective states such as anxiety and depression.³ Other features of emotional regulation are emotional awareness, the capacity to recognize and distinguish emotions experienced by oneself and others, and social cognition, the ability to process social emotions, including empathy, moral reasoning, theory of mind, and emotional comprehension. Both emotional awareness and social cognition appear to be key elements in the ability to regulate emotions, and are impaired in a number of psychiatric conditions.⁶

[1] NEUROBIOLOGY OF EMOTION AND ITS REGULATION

The generation of emotion occurs as an interaction of bottom-up (brain stem and limbic system to higher cortical regions) and top-down (prefrontal cortex to midbrain and brain stem areas via the amygdala) series of actions.⁵⁻¹⁰ Neuroimaging data suggest that responses to emotional tasks are widely distributed throughout the brain, that higher cortical areas are not limited to emotional regulation, and that limbic regions are not restricted to emotional activation. Emotional regulation involves a widely distributed functional network with bidirectional associations among many emotion-relevant brain regions.^{5,11,12} In a meta-analysis of 162 neuroimaging studies of emotion, Kober and colleagues¹¹ identified regions within the medial, orbital, and inferior lateral frontal cortices as brain regions that were activated consistently during administration of emotional tasks. The amygdala, ventral striatum, thalamus, hypothalamus, and periaqueductal gray (brain regions that have been found to be important in emotion in animals) were identified as areas with multiple activations. The maturation of neural and neuroendocrine arousal systems associated with emotion throughout childhood and adolescence can explain the decrease in emotional lability and increase in self-control that occurs over this early period of development. These processes include maturation of parasympathetic regulation in early childhood and the developmental changes in the hypothalamic-pituitary-adrenocortical axis. Maturation of these systems is shaped by early experiences and caregiver responsiveness. Developmental influences promoting enhanced emotional regulation for children as they grow older include

the acquisition of language as a means to understand and communicate emotions, and maturation of other cognitive functions, including the attentional system.⁷

These brain systems are shaped by early experiences and reflect developmental history. It is possible that early life adversity changes the threshold of limbic reactivity or changes perceptual and cognitive appraisals related to threat.^{7,13} For example, children growing up in early adversity are more likely to be emotionally reactive to stress and also less capable of emotional regulation.¹⁴ Brain-imaging studies in those who experienced childhood maltreatment point to frontolimbic circuits as the most affected brain regions.¹⁵ Consistent with this view, two different forms of emotional dysregulation with distinct patterns of cortical and subcortical activation and symptom profiles in posttraumatic stress disorder (PTSD) have been identified.^{16,17} Undermodulated emotion (e.g., anxiety, hyperarousal, dysphoria) is related to the classic pattern of heightened amygdala activation and reduced prefrontal inhibitory activation. Overmodulated emotion (e.g., dissociation, emotional numbing), by contrast, is related to extensive midline prefrontal inhibition of limbic activity.¹⁷ Support for a dissociative subtype of PTSD consistent with overmodulation has been reported in a recent World Health Organization World Mental Health Survey¹⁸ and in recent clinical and neurobiological research.¹⁹

Recent neuroimaging research has identified a different set of patterns of brain activity than the fear/stress circuits—which may be involved in emotional regulation and dysregulation in PTSD and other psychiatric disorders. This *default mode network* involves activation of the medial temporal and prefrontal cortices and also limbic areas (e.g., hippocampus) that are integrated in the posterior cingulate.^{20,21} The default mode pattern of brain activation occurs when attention is focused away from the external environment and goal-directed behavior and toward self-referential thought.²¹ Reduced levels of default mode activation have been found in individuals with borderline personality disorder who are experiencing pain²² and in persons with PTSD when at rest,²⁰ and are also a predictor of risk of developing PTSD in acutely traumatized individuals.²³ Deficits in default mode network connectivity have been hypothesized to be associated with sequelae of exposure to childhood adversity, including emotional dysregulation and deficits in attentional shifting, mindfulness, and self-referential encoding.^{20,22} As such, default mode processing potentially represents a brain substrate of emotional dysregulation that is distinct from the classic fear-conditioning pathway in anxiety disorders.²⁴

[2] EMOTIONAL REGULATION AND ASSOCIATED PSYCHIATRIC MORBIDITY

Problems with emotional regulation play a role in the development, maintenance, and treatment of many psychiatric conditions. Maladaptive coping with challenging emotions is common in depression, bipolar disorder, borderline personality disorder, substance-use disorders, eating disorders, and somatoform disorders, among others.²⁵ In children, some forms of externalizing problems have been linked with emotional dysregulation.² In addition, reactive aggression, which is driven by negative emotional states, is associated with higher cortisol reactivity²⁶ and decreased emotion regulation. Maltreated children are at higher risk for both reactive aggression and deficits in emotional regulation.^{27,28} More

generally, executive dysfunction and emotional control (including effortful, cognitively mediated control) appear to be inversely related,² and internalizing problems such as anxiety, depression, withdrawal, and somatic complaints are conceptually related to other forms of emotional dysregulation that involve difficulty in controlling attention and cognition. For example, rumination and attention bias toward negative stimuli is associated with internalizing disorders.²

From a developmental perspective, an important question is whether emotional dysregulation is a risk factor for, or a consequence of, psychopathology. McLaughlin and colleagues²⁹ investigated this question by prospectively assessing depression, anxiety, aggressive behavior, disordered eating, and emotional regulation (measures of emotional understanding and expression, and rumination in reaction to distress) in 1065 adolescents. They found that emotional dysregulation contributed to multiple and varied psychiatric and psychosocial impairments in adolescence (anxiety, aggression, and disordered eating) but that emotional dysregulation was not primarily a consequence of psychopathology; such pathology did not predict changes in emotional regulation over time.

Childhood dysregulation may have detrimental effects across the lifespan by increasing the risk of emotional dysregulation in adulthood. For example, a 14-year prospective, follow-up study of 2076 Dutch children found that those with severe challenges regarding emotional regulation (attention problems, aggressive behavior, and anxious-depression as rated on the Child Behavior Checklist) were at increased risk for problems with regulating affect, behavior, and cognition in early adulthood.³⁰ Retrospective studies have shown that problems with emotional regulation in childhood secondary to exposure to complex trauma are associated with dysregulation in multiple domains of informational processing (physiological, sensory, emotional, and cognitive) and self- and relational dysregulation throughout adulthood.¹³

[1] CHILDHOOD TRAUMATIZATION AND PSYCHIATRIC COMORBIDITIES IN CHILDREN AND ADULTS

The consequences of exposure to interpersonal trauma vary from individual to individual and also, over time, for the same individual. Many traumatized children do not develop PTSD or any other disorder. Nevertheless, individuals who experience interpersonal trauma in childhood are at increased risk for numerous psychiatric disorders, including attachment disorders,¹³ PTSD, depression, and anxiety disorders,³¹ oppositional or conduct disorders,^{32,33} eating disorders,³⁴ substance abuse,^{35,36} a dissociative variant of PTSD,¹⁷ and personality disorders.¹ Traumatized children also are at risk for self-harm, sexualized behavior, anger, poor impulse control, and attention difficulties.¹ Revictimization is common in maltreated children and adolescents, and is associated with increased risk for PTSD and other comorbidities, such as depressive and substance use disorders.³⁷

Various terms have been used to describe children who have experienced early, recurrent, and severe interpersonal trauma—including *developmental trauma disorder*,³⁸ *complex trauma*,³⁹ and *polyvictimization*.⁴⁰ Those children often present with extreme dysregulation in the physical, affective, behavioral, cognitive, and interpersonal domains. Currently, the

category of developmental trauma disorder is being evaluated for clinical utility, and a national field trial is under way. In addition, the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5) has incorporated a number of changes aimed at increasing the developmental sensitivity of PTSD diagnoses in children and adolescents. In preschool-aged children, these changes include a broader range of acute emotional reactions (diminished interest may manifest as restricted play, and detachment may manifest as social withdrawal) and a reduced threshold for internalized experiences that young children may not have the language to describe. In school-aged children and adolescents, trauma may now include loss (including placement in foster care) and the injury or death of a parent. In addition, for all ages, PTSD symptoms have been broadened to include dysregulation in the bodily, affective (including a range of emotions beyond anxiety), cognitive, behavioral, relational, and self/identity domains.^{41,42}

Table 1 summarizes some pertinent studies. Findings of two cross-sectional studies suggest an association between childhood maltreatment/other forms of childhood trauma and subsequent psychiatric morbidity. Psychiatric assessment of children removed from their parents' care due to severe maltreatment revealed that more than one in three (35%) met criteria for PTSD and those children were also at risk for attention-deficit/hyperactivity disorder (ADHD), other anxiety disorders, brief psychotic disorder or psychotic disorder not otherwise specified, and suicidal ideation.⁴³ Assessment of trauma history and PTSD symptoms in youth presenting with oppositional-defiant disorder (ODD), ADHD, or adjustment disorder suggested that those with externalizing disorders (both ODD and ADHD) were more likely to have experienced maltreatment and that they had more PTSD symptoms, even after controlling for overlapping ODD and PTSD criterion D symptoms (hyperarousal/hypervigilance).⁴⁴

Two large prospective, longitudinal studies also suggested associations between maltreatment in early childhood, and the development of PTSD and other psychiatric and behavioral disorders later in adolescence. In a community sample of youth followed from kindergarten through grade 11,⁵ adolescents identified at baseline as having experienced physical abuse had significant social and psychiatric difficulties at follow-up.⁴⁵ Even when risk factors associated with maltreatment (socioeconomic status, single-parent family, family stress, maternal social support, and the child's temperament and prenatal/postnatal health) were taken into consideration, maltreated adolescents were significantly more likely to be absent from school, less likely to expect higher education, and experienced more social difficulties and social withdrawal. They also suffered from more anxiety, depression, PTSD, dissociation, and thought problems. The Great Smoky Mountains Study, a longitudinal study of psychopathology and use of medical services in childhood, assessed children from age 9 through 16 for traumatic events, posttraumatic stress (PTS) symptoms and DSM-IV disorders. Although traumatic events were common and had been experienced by over two-thirds of the children by age 16, only a minority of these children (13.4%) had developed PTS symptoms, and less than 0.5% of the children had developed PTSD. PTS symptoms were most likely to develop in those who experienced violent or sexual traumas, and were predicted for those with prior and multiple traumatic exposures, preexisting anxiety disorders, and family adversity. PTS symptoms were rare and brief in those who

experienced a first traumatic event and were not predicted by parental psychopathology. It is interesting to note that trauma exposure was strongly associated with anxiety and depression, and nearly doubled the rates of other psychiatric disorders (all diagnostic groups excluding substance abuse). This association was most pronounced in children who experienced PTS symptoms.⁴⁶

The adverse impact of trauma exposure on risk of psychiatric morbidity may begin early in childhood. Almost a quarter (23.4%) of prospectively followed toddlers had already experienced a potentially traumatic event; those exposed were more likely to exhibit internalizing and externalizing behaviors and had more severe symptoms; and approximately 20% of those exposed had PTSD.⁴⁷ A cross-sectional examination of these associations in preschool children suggested that after controlling for socioeconomic factors, exposure to domestic violence was associated with internalizing (depression, separation anxiety, PTS) and externalizing (conduct problems) disorders. Parental mood and anxiety symptoms partially or fully accounted for these associations, suggesting a potential intergenerational effect of dysregulation in children's posttraumatic psychopathology.⁴⁸ Another prospective study following children from age of three to school-aged found that exposure to domestic and neighborhood violence at or before the age of three was correlated with PTS, internalizing and externalizing symptom severity, and social difficulties at school age, independent of economic disadvantage and exposure to violence within the past year.⁴⁹ Various other studies have shown that adults who experienced early life trauma continue to be at risk for PTSD, other anxiety⁵⁵ and affective disorders,⁵⁰ addictions,⁵⁶⁻⁵⁸ psychotic illnesses,^{34,59} personality disorders,⁵² dissociative identity disorder,⁶⁰ and suicidal behavior⁶¹ as well as for revictimization and multiple medical problems, including diabetes, heart disease, immune disorders, and chronic obstructive pulmonary disease.^{1,62-64}

Adults self-reporting childhood adversity, especially emotional abuse, were found to have increased risk of lifetime and current depressive disorders in adulthood, even decades after occurrence of adversity.⁵⁰ Adults with severe mood disorders who report adverse childhood experiences were found to have increased high-risk behaviors, substance abuse, revictimization, and PTSD, and more severe psychiatric illness that presented at a younger age.⁵¹

Although all psychiatric disorders are likely to involve dysregulation, perhaps the clearest example of psychopathology involving emotional dysregulation is bipolar disorder. Over the last decade, childhood bipolar disorder has been a controversial topic debated in the research and clinical communities; it is now largely agreed, however, that some youth do meet full DSM criteria for bipolar disorder and that this phenotype does exist. While it is possible that this debate has affected the rate of bipolar disorder diagnosed in persons with trauma exposure, the available evidence suggests that childhood trauma histories are frequent and that they affect the clinical presentation of bipolar disorder. Childhood exposure to interpersonal trauma is reported in about half of adult patients with bipolar disorder, across several studies. Adults with bipolar disorder and a history of childhood trauma have more frequent prepubertal onset of affective symptoms, significantly younger age at bipolar illness onset, and more severe symptoms.⁶⁵⁻⁶⁹ In a recent publication, Connor and Doerfler⁷⁰ identified higher rates of comorbid PTSD in a small single-site clinical sample of

youth with bipolar disorder, diagnosed using standardized assessments and DSM-IV-TR criteria (n = 27), compared to youth with disruptive behavioral disorders and negative mood (n = 96). Emerging evidence suggests that both polyvictimization and the presence of comorbid bipolar disorder in adolescents lead to PTSD presentations that are less responsive to treatment and less likely to remit.⁷¹ To date, however, no study has systematically evaluated the association between childhood exposure to interpersonal trauma and early-onset bipolar disorder.

Emotional dysregulation is also prominent in personality disorders. The association between exposure to interpersonal trauma and personality disorders is one that has received, and continues to receive, substantial attention. Individuals with severe, disabling personality disorders (borderline, schizotypal, and paranoid) self-report more types of interpersonal trauma and childhood physical abuse than those with other personality disorders.^{52,53} More specifically, adults with borderline personality disorder and paranoid personality disorder have the highest rates of exposure to trauma (especially childhood sexual trauma), high rates of PTSD, and younger age at first exposure to trauma.^{52,53} There is also an association between childhood trauma by primary care giver and emotional dysregulation in adults with borderline personality disorder.⁷²⁻⁷⁴

Further, two forms of emotional dysregulation were identified in an adult psychiatric inpatient sample, consistent with the distinction described by Lanius and colleagues¹⁷ of emotional *undermodulation* and *overmodulation*—although the sample also suggested a more nuanced view of the role of dissociation in emotional dysregulation. Patients diagnosed with borderline personality disorder were primarily characterized by undermodulation of emotion, positive psychoform dissociation symptoms (such as flashbacks), and complex PTSD symptoms. By contrast, those diagnosed with somatoform disorders tended to present with overmodulated emotional responding and negative psychoform dissociation symptoms (such as amnesia).⁷²⁻⁷⁴ In addition, specific difficulties with different affect capacities were found to be associated with borderline personality disorder, somatoform disorders, and other psychiatric disorders, suggesting that patterns of emotional dysregulation can be identified in relation to specific types of psychopathology.⁵⁴

Studies have consistently found a correlation between the severity of personality disorder (including but not limited to borderline personality disorder) and the severity and developmental impact of trauma exposure, as evident by younger age at first exposure to maltreatment, more assaultive and personal trauma, and more types of traumatic events.^{52,53}

To summarize, maltreated children are at increased risk for internalizing disorders (including PTS symptoms and PTSD), externalizing disorders, and psychosocial impairment at initial presentation and also when longitudinally followed or retrospectively assessed in adolescence and adulthood. Not all trauma exposure results in PTS or PTSD, but it does appear to increase risk of a variety of other disorders. In adults, the potential sequelae of exposure to adversity in childhood involve a heightened risk and severity of affective disorders (including but not limited to bipolar disorder) and also of other forms of severe psychopathology, including personality disorders.

[1] CHILDHOOD MALTREATMENT AND EMOTIONAL DYSREGULATION

Emotional regulation can be viewed as a developmental task that is highly influenced by the ability to develop appropriate secure attachments. Emotional regulation appears to develop in the context of responsive caregiving and peer involvement in early life.¹ Not only do caregivers provide for their children's basic survival needs, but interactions with caregivers are necessary for the development of bodily self-regulation.¹ In humans, childhood maltreatment, especially repeated trauma, disrupts the acquisition of appropriate emotional regulation and interpersonal skills.^{64,75} This disruption of skill acquisition may occur as a result of psychological experiences but is also a sign of the neurobiological effects of maltreatment.^{75,76} These effects include molecular alterations to stress hormone response systems, which, in turn, affects myelination, neuronal morphology, neurogenesis, and synaptogenesis in different brain regions, resulting in functional changes in left hemisphere development, decreased right/left hemisphere integration, increased limbic electrical irritability, and diminished functional activity in the cerebellar vermis.^{70,76}

A number of studies have empirically examined the associations between emotional dysregulation and a history of traumatic exposure in youth, and have highlighted the importance of early caregiver responsiveness and how deficits in responsiveness may compromise the development of interpersonal communication and interpretation of social cues. Key study findings are further described in Table 2, and summarized below. In two studies, Shipman and colleagues^{77,80} attempted to identify which processes in emotional development differ in maltreated and non-maltreated children. They found that girls who experienced sexual abuse were less able to understand and regulate emotions. They also expected to be receiving less emotional support and have more interpersonal conflict in response to expression of negative emotional states—in particular, when expressing sadness to parents or anger to peers.⁷⁷ When compared to healthy controls, children (boys and girls) who experienced neglect had lesser ability to understand negative emotions such as anger and sadness, and fewer adaptive emotional-regulation skills. Furthermore, these children expected their mothers to respond negatively to expression of anger and sadness, and possibly as a result, they attempted to hold back their display of such emotions.⁸⁰ Both studies suggest that maltreated children, whether they experience sexual trauma or neglect, show deficits or delays in understanding and regulating emotions, and that they anticipate negative reactions reaction to display of sadness and anger. This point is an important one, as such skills can be taught as part of clinical interventions aimed at treating victimized children.^{87,88}

The development of secure attachment with caregivers early in childhood has been theorized to be essential to the development of emotional regulation.⁸⁹ Disruption of the formation of secure internal representations (described by Bowlby as “working models”)⁹⁰ by interpersonal trauma such as maltreatment may therefore substantially compromise the acquisition of emotional-regulation capacities in childhood. Shields and colleagues⁷⁸ examined whether maltreated children had difficulties in forming secure caregiver representations and regulating emotion, and whether such difficulties were associated with problems in social adjustment. They found that children who experienced maltreatment had disorganized, vague, and negatively toned internal representations of caregivers and had

problems with emotional dysregulation, aggression, and decreased social competence (peer rejection).

Pollak and colleagues⁹¹ examined the development of emotion recognition in maltreated children. Their studies identify a number of differences in mechanisms linking early-life exposure to potentially traumatic experiences and emerging emotions. Neglected preschool children had more difficulty discriminating emotional expressions and identifying discrete emotions, whereas physically abused children displayed a response bias for angry facial expressions and showed the most variance across emotions. Physically abused 8–11 year olds had difficulties disengaging attention from angry facial cues.⁹² Despite these deficits in emotion recognition, however, physically abused children were more accurately able to recognize early facial expression of anger, when few physiological cues were available, suggesting a readiness to detect subtle behavioral signs of anger.⁹³

A few studies found that a history of maltreatment affects children's development of emotional regulation and social adjustment. Maughan and colleagues⁷⁹ found that in response to a simulation of angry adult interactions, 80% of maltreated children, compared to 37% of the non-maltreated controls, were observed to have problems with emotional regulation. These problems were associated with maternal reports of child behavior problems and of anxious and depressed symptoms. Kim and colleagues⁸⁵ described a bidirectional relationship between emotional dysregulation and social dysfunction in maltreated youth. Emotional dysregulation was associated with a history of neglect, physical, and sexual abuse (separately and in combination) and with earlier age at onset of abuse. Emotional-regulation deficits were also linked to externalizing behaviors, which, in turn, contributed to peer rejection, and vice versa. Not surprisingly, in the same study, emotional regulation ability was positively correlated with peer acceptance and lower rates of internalizing behaviors.

Rogosch and colleagues⁸¹ conceptualized emotional and behavioral dysregulation—as defined by affective negativity, irritability, lability, suicidal/self-harm behavior, and impulsivity and extreme conflict and struggle in interpersonal relationships with peers and adults—as potential precursors to borderline personality disorder in children, and found consistent evidence of a relationship between history of maltreatment and all of the indices of dysregulation.

Studies examining the associations between trauma history and emotional dysregulation have also been conducted in young adult populations, specifically with college students. These studies describe a consistent correlation between trauma exposure and emotional dysregulation, as well as a correlation between PTS symptom severity and emotional dysregulation.^{64,83,84} Tull and colleagues⁸³ described a correlation between PTS/PTSD and several measures of emotional dysregulation, including difficulty accepting and recognizing emotions, difficulties utilizing strategies to manage and regulate negative emotion when upset, and impulsivity. Burns and colleagues⁶⁴ found that emotional abuse was the most powerful predictor of emotional dysregulation, perhaps because it interferes with the acquisition of developmentally appropriate emotional-regulation skills. In Ford and colleagues' sample,⁹⁴ more than 50% reported at least one symptom of dysregulation, which

was, in turn, associated with risk of PTSD and other anxiety or affective disorders. Dysregulation also was uniquely associated with the severity of interpersonal trauma history, particularly with past abuse or multiple interpersonal traumas. By contrast, non-interpersonal trauma was associated with risk of PTSD and dissociative symptoms, but not with dysregulation.⁹⁴ These findings suggest that interventions aimed at improving emotional-regulation strategies may be an important aspect of treatment (and prevention) for young women who have experienced potentially traumatic adversity in childhood.^{64,87,88}

Emotional dysregulation can also be understood as the underlying mechanism for risky sexual behavior and sexual revictimization among young adult victims of child sexual and physical abuse. Messman-Moore⁸⁴ demonstrated that a history of childhood physical and sexual abuse was associated with both risky sexual behavior and increased risk for being raped as an adolescent or adult; more than a quarter of these women had been sexually revictimized, with emotional dysregulation appearing to be a mediating factor. A higher lifetime number of sexual partners (including strangers) and other risky sexual behaviors were predicted by measures of emotional dysregulation. It also appears that the severity of physical and sexual revictimizations is predicted by childhood sexual abuse.⁹⁵ Recent evidence suggests that patterns of exposure to sexual abuse and dysregulation may be intergenerational. Mothers with a history of sexual abuse in their own childhoods were found to be at risk for impairment in their internal attachment representations and in their attachment behavior with their daughters,⁹⁶ and their daughters showed impairment in their capacities for emotional regulation⁹⁷ and were also at risk for exposure to sexual abuse themselves.⁹⁸

Emotional dysregulation and other impaired self-capacities in adults have been shown to be associated with past exposure in childhood to emotional abuse and a lack of emotional support by parental figures.⁹⁹ A large study examined the contribution of emotional dysregulation, childhood trauma, and negative affect to a range of clinical psychopathology, and found that childhood trauma severity, negative affect, and emotional dysregulation all were correlated. Childhood trauma severity also was associated with PTS, drug abuse, depression, and suicidality.⁸⁶ In a study of women who were abused in childhood, PTS severity was associated with functional impairment.⁸² After controlling for PTSD severity, however, emotional dysregulation and interpersonal problems both were associated with functional impairment, and together had an adverse influence on psychosocial functioning equal to that of PTSD symptoms.⁸² Thus, emotional regulation, interpersonal problems, and PTSD may be three distinct, but interrelated, adverse sequelae of childhood trauma exposure.

[1] CONCLUSIONS

These studies suggest a complex and bidirectional relationship between childhood trauma exposure and emotional dysregulation. Trauma exposure in childhood is associated with reduced ability to understand and regulate emotions (potentially mediated by relational/attachment difficulties with caregivers and peers) and with heightened levels of internalizing and externalizing psychopathology and impaired social functioning beginning in childhood and continuing into adulthood. Adults with childhood trauma histories are at risk for

emotional-regulation problems, PTS, other internalizing-disorder symptoms, functional impairment, and revictimization.

Emotions are generated and regulated in response to situations or thoughts, based on past experience, and can be automatic or controlled. Emotions have a neurobiological substrate and are generated and regulated through bottom-up and top-down processes, involving a complex interplay between cortical and limbic brain regions. Cognitive control and executive function play important roles in these processes, which develop in childhood. Emotional dysregulation has been linked to many forms of psychopathology, including externalizing and internalizing problems in children, and is associated with increased risk across diagnoses. Hence, emotional dysregulation appears to be an important target for therapeutic interventions and focus for children and adults alike.

Substantial evidence connects the experience of childhood exposure to interpersonal traumas, such as maltreatment or family violence, with emotional dysregulation and with multiple related psychiatric sequelae and comorbidities (and not just PTSD). Because childhood exposure to interpersonal trauma is associated with emotional dysregulation across a variety of psychiatric disorders, further clinical and scientific study is needed to determine how best to understand and treat childhood trauma-related emotional dysregulation in a variety of psychiatric disorders, including but not limited to PTSD.

Given that research into emotion generation, recognition, and regulation is expanding to cover wider areas of neuroscience and psychiatric conditions, standardized assessments of emotional dysregulation, both for evaluation/treatment and research purposes, are critically needed. Specifically, clinical and translation research is needed to assess trauma-related emotional dysregulation and to determine what treatments are effective for different psychiatric disorders. Preventive measures and targeted therapeutic interventions, both psychotherapeutic and pharmacologic, may enhance our abilities to address emotional dysregulation, a problem that impairs the lives of so many who have a history of interpersonal trauma.

Acknowledgments

Dr. Frazier receives research support from Glaxo Smith Kline, Seaside Therapeutics, Pfizer, Inc., and Roche Pharmaceuticals (Dr. Frazier).

REFERENCES

1. Ford JD. Treatment implications of altered affect regulation and information processing following child maltreatment. *Psychiatr Ann.* 2005; 35:410–419.
2. Eisenberg N, Spinrad TL, Eggum ND. Emotion-related self-regulation and its relation to children's maladjustment. *Annu Rev Clin Psychol.* 2010; 6:495–525. [PubMed: 20192797]
3. Gross JJ. Emotion regulation: affective, cognitive, and social consequences. *Psychophysiology.* 2002; 39:281–291. [PubMed: 12212647]
4. Kim SH, Hamann S. Neural correlates of positive and negative emotion regulation. *J Cogn Neurosci.* 2007; 19:776–798. [PubMed: 17488204]
5. Ochsner KN, Bunge SA, Gross JJ, Gabrieli JD. Rethinking feelings: an fMRI study of the cognitive regulation of emotion. *J Cogn Neurosci.* 2002; 14:1215–1229. [PubMed: 12495527]

6. Lanius RA, Bluhm RL, Frewen PA. How understanding the neurobiology of complex post-traumatic stress disorder can inform clinical practice: a social cognitive and affective neuroscience approach. *Acta Psychiatr Scand.* 2011; 124:331–348. [PubMed: 21854369]
7. Thompson RA. Emotion and emotion regulation: two sides of the developing coin. *Emot Rev.* 2011; 3:53–61.
8. Pavuluri MN, Herbener ES, Sweeney JA. Affect regulation: a systems neuroscience perspective. *Neuropsychiatr Dis Treat.* 2005; 1:9–15. [PubMed: 18568120]
9. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception I: the neural basis of normal emotion perception. *Biol Psychiatry.* 2003; 54:504–514. [PubMed: 12946879]
10. Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception II: implications for major psychiatric disorders. *Biol Psychiatry.* 2003; 54:515–528. [PubMed: 12946880]
11. Kober H, Barrett LF, Joseph J, Bliss-Moreau E, Lindquist K, Wager TD. Functional grouping and cortical-subcortical interactions in emotion: a meta-analysis of neuroimaging studies. *Neuroimage.* 2008; 42:998–1031. [PubMed: 18579414]
12. Ochsner KN, Ray RR, Hughes B, et al. Bottom-up and top-down processes in emotion generation: common and distinct neural mechanisms. *Psychol Sci.* 2009; 20:1322–1331. [PubMed: 19883494]
13. Andersen SL, Lyss PJ, Dumont NL, Teicher MH. Enduring neurochemical effects of early maternal separation on limbic structures. *Ann N Y Acad Sci.* 1999; 877:756–759. [PubMed: 10415699]
14. D'Andrea W, Ford JD, Stolbach B, Spinazzola J, van der Kolk BA. Understanding interpersonal trauma in children: why we need a developmentally appropriate trauma diagnosis. *Am J Orthopsychiatry.* 2012; 82:187–200. [PubMed: 22506521]
15. Hart H, Rubia K. Neuroimaging of child abuse: a critical review. *Front Hum Neurosci.* 2012; 6:1–24. [PubMed: 22279433]
16. Steuwe C, Lanius RA, Frewen PA. Evidence for a dissociative subtype of PTSD by latent profile and confirmatory factor analyses in a civilian sample. *Depress Anxiety.* 2012; 29:689–700. [PubMed: 22653764]
17. Lanius RA, Vermetten E, Loewenstein RJ, et al. Emotion modulation in PTSD: clinical and neurobiological evidence for a dissociative subtype. *Am J Psychiatry.* 2010; 167:640–647. [PubMed: 20360318]
18. Stein DJ, Koenen KC, Friedman MJ, Hill E, et al. Dissociation in posttraumatic stress disorder: evidence from the world mental health surveys. *Biol Psychiatry.* 2013; 73:302–312. [PubMed: 23059051]
19. Lanius RA, Brand B, Vermetten E, Frewen PA, Spiegel D. The dissociative subtype of posttraumatic stress disorder: rationale, clinical and neurobiological evidence, and implications. *Depress Anxiety.* 2012; 29:701–708. [PubMed: 22431063]
20. Bluhm RL, Williamson PC, Osuch EA, et al. Alterations in default network connectivity in posttraumatic stress disorder related to early-life trauma. *J Psychiatry Neurosci.* 2009; 34:187–194. [PubMed: 19448848]
21. Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci.* 2008; 1124:1–38. [PubMed: 18400922]
22. Kluetsch RC, Schmahl C, Niedtfeld I, et al. Alterations in default mode network connectivity during pain processing in borderline personality disorder. *Arch Gen Psychiatry.* 2012; 69:993–1002. [PubMed: 22637967]
23. Lanius RA, Bluhm RL, Coupland NJ, et al. Default mode network connectivity as a predictor of post-traumatic stress disorder symptom severity in acutely traumatized subjects. *Acta Psychiatr Scand.* 2010; 121:33–40. [PubMed: 19426163]
24. Lanius RA, Frewen PA, Vermetten E, Yehuda R. Fear conditioning and early life vulnerabilities: two distinct pathways of emotion dysregulation and brain dysfunction in PTSD. *Eur J Psychotraumatol.* Epub 2010 Dec 10.
25. Berking M. Emotion regulation and mental health recent findings, current challenges, and future directions. *Curr Opin Psychiatry.* 2012; 25:128–134. [PubMed: 22262030]

26. Lopez-Duran NL, Olson SL, Hajal NJ, Felt BT, Vazquez DM. Hypothalamic pituitary adrenal axis functioning in reactive and proactive aggression in children. *J Abnorm Child Psychol*. 2009; 37:169–182. [PubMed: 18696227]
27. Ford JD, Fraleigh LA, Albert DB, Connor DF. Child abuse and autonomic nervous system hyporesponsivity among psychiatrically impaired children. *Child Abuse Negl*. 2010; 34:507–515. [PubMed: 20605634]
28. Shields A, Cicchetti D. Reactive aggression among maltreated children: the contributions of attention and emotion. *J Clin Child Psychol*. 1998; 27:381–395. [PubMed: 9866075]
29. McLaughlin KA, Hatzenbuehler ML, Mennin DS, Nolen-Hoeksema S. Emotion dysregulation and adolescent psychopathology: a prospective study. *Behav Res Ther*. 2011; 49:544–554. [PubMed: 21718967]
30. Althoff RR, Verhulst FC, Rettew DC, Hudziak JJ, van der Ende J. Adult outcomes of childhood dysregulation: a 14-year follow-up study. *J Am Acad Child Adolesc Psychiatry*. 2010; 49:1105–1116. [PubMed: 20970698]
31. Ford JD, Elhai JD, Ruggiero KJ, Frueh BC. Refining posttraumatic stress disorder diagnosis: Evaluation of symptom criteria with the national survey of adolescents. *J Clin Psychiatry*. 2009; 70:748–755. [PubMed: 19389336]
32. Ford JD, Connor DF, Hawke J. Complex trauma among psychiatrically impaired children: A cross-sectional, chart-review study. *J Clin Psychiatry*. 2009; 70:1155–1163. [PubMed: 19573498]
33. Kerig PK, Ward RM, Vanderzee KL, Arnzen Moeddel M. Posttraumatic stress as a mediator of the relationship between trauma and mental health problems among juvenile delinquents. *J Youth Adolesc*. 2009; 38:1214–1225. [PubMed: 19669901]
34. Brady KT, Killeen TK, Brewerton T, Lucerini S. Comorbidity of psychiatric disorders and posttraumatic stress disorder. *J Clin Psychiatry*. 2000; 61:22–32. [PubMed: 10795606]
35. Dube SR, Miller JW, Brown DW, et al. Adverse childhood experiences and the association with ever using alcohol and initiating alcohol use during adolescence. *J Adolesc Health*. 2006; 38:3089–3096.
36. Strine TW, Dube SR, Edwards VJ, et al. Associations between adverse childhood experiences, psychological distress, and adult alcohol problems. *Am J Health Behav*. 2012; 36:408–423. [PubMed: 22370441]
37. Macdonald A, Danielson CK, Resnick HS, Saunders BE, Kilpatrick DG. PTSD and comorbid disorders in a representative sample of adolescents: the risk associated with multiple exposures to potentially traumatic events. *Child Abuse Negl*. 2010; 34:773–783. [PubMed: 20869115]
38. Ford JD, Grasso D, Greene C, Levine J, Spinazzola J, van der Kolk B. Clinical significance of a proposed developmental trauma disorder diagnosis: results of an international survey of clinicians. *J Clin Psychiatry*. 2013; 74:841–849. [PubMed: 24021504]
39. Cook A, Spinazzola J, Ford JD, et al. Complex trauma in children and adolescents. *Psychiatr Ann*. 2005; 35:390–398.
40. Finkelhor D, Ormrod RK, Turner HA. Poly-victimization: a neglected component in child victimization. *Child Abuse Negl*. 2007; 31:7–26. [PubMed: 17224181]
41. Scheeringa MS, Zeanah CH, Cohen JAM. PTSD in children and adolescents: towards an empirically based algorithm. *Depress Anxiety*. 2011; 28:770–782. [PubMed: 20734362]
42. Friedman MJ, Resick PA, Bryant RA, Brewin CR. Considering PTSD for DSM-5. *Depress Anxiety*. 2011; 28:750–769. [PubMed: 21910184]
43. Famularo R, Fenton T, Kinscherff R, Augustyn M. Psychiatric comorbidity in childhood post traumatic stress disorder. *Child Abuse Negl*. 1996; 20:953–961. [PubMed: 8902292]
44. Ford JD, Racusin R, Ellis CG, et al. Child maltreatment, other trauma exposure, and posttraumatic symptomatology among children with oppositional defiant and attention deficit hyperactivity disorders. *Child Maltreat*. 2000; 5:205–217. [PubMed: 11232267]
45. Lansford JE, Dodge KA, Pettit GS, Bates JE, Crozier J, Kaplow J. A 12-year prospective study of the long-term effects of early child physical maltreatment on psychological, behavioral, and academic problems in adolescence. *Arch Pediatr Adolesc Med*. 2002; 156:824–830. [PubMed: 12144375]

46. Copeland WE, Keeler G, Angold A, Costello EJ. Traumatic events and posttraumatic stress in childhood. *Arch Gen Psychiatry*. 2007; 64:577–584. [PubMed: 17485609]
47. Mongillo EA, Briggs-Gowan M, Ford JD, Carter AS. Impact of traumatic life events in a community sample of toddlers. *J Abnorm Child Psychol*. 2009; 37:455–468. [PubMed: 19034643]
48. Briggs-Gowan MJ, Carter AS, Clark R, Augustyn M, McCarthy KJ, Ford JD. Exposure to potentially traumatic events in early childhood: differential links to emergent psychopathology. *J Child Psychol Psychiatry*. 2010; 51:1132–1140. [PubMed: 20840502]
49. Briggs-Gowan MJ, Carter AS, Ford JD. Parsing the effects violence exposure in early childhood: modeling developmental pathways. *J Pediatr Psychol*. 2012; 37:11–22. [PubMed: 21903730]
50. Chapman DP, Whitfield CL, Felitti VJ, Dube SR, Edwards VJ, Anda RF. Adverse childhood experiences and the risk of depressive disorders in adulthood. *J Affect Disord*. 2004; 82:217–225. [PubMed: 15488250]
51. Lu W, Mueser KT, Rosenberg SD, Jankowski MK. Correlates of adverse childhood experiences among adults with severe mood disorders. *Psychiatr Serv*. 2008; 59:1018–1026. [PubMed: 18757595]
52. Yen S, Shea MT, Battle CL, et al. Traumatic exposure and posttraumatic stress disorder in borderline, schizotypal, avoidant, and obsessive-compulsive personality disorders: findings from the collaborative longitudinal personality disorders study. *J Nerv Ment Dis*. 2002; 190:510–518. [PubMed: 12193835]
53. Golier JA, Yehuda R, Bierer LM, et al. The relationship of borderline personality disorder to posttraumatic stress disorder and traumatic events. *Am J Psychiatry*. 2003; 160:2018–2024. [PubMed: 14594750]
54. van Dijke A, van der Hart O, van Son M, Buhning M, van der Heijden P, Ford JD. Cognitive and affective dimensions of difficulties in emotional functioning in somatoform disorders and borderline personality disorder. *Psychopathology*. 2012; 46:153–162. [PubMed: 23006331]
55. Cogle JR, Timpano KR, Sachs-Ericsson N, Keough ME, Riccardi CJ. Examining the unique relationships between anxiety disorders and childhood physical and sexual abuse in the national comorbidity survey-replication. *Psychiatry Res*. 2010; 177:150–155. [PubMed: 20381878]
56. Anda RF, Whitfield CL, Felitti VJ, et al. Adverse childhood experiences, alcoholic parents, and later risk of alcoholism and depression. *Psychiatr Serv*. 2002; 53:1001–1009. [PubMed: 12161676]
57. Dube SR, Anda RF, Felitti VJ, Edwards VJ, Croft JB. Adverse childhood experiences and personal alcohol abuse as an adult. *Addict Behav*. 2002; 27:713–725. [PubMed: 12201379]
58. Dube SR, Felitti VJ, Dong M, Chapman DP, Giles WH, Anda RF. Childhood abuse, neglect, and household dysfunction and the risk of illicit drug use: the adverse childhood experiences study. *Pediatrics*. 2003; 111:564–572. [PubMed: 12612237]
59. Whitfield CL, Dube SR, Felitti VJ, Anda RF. Adverse childhood experiences and hallucinations. *Child Abuse Negl*. 2005; 29:797–810. [PubMed: 16051353]
60. Teicher M, Andersen S, Polcari A, Anderson C, Navalta C. Developmental neurobiology of childhood stress and trauma. *Psychiatr Clin North Am*. 2002; 25:397–426. [PubMed: 12136507]
61. Dube SR, Anda RF, Felitti VJ, Chapman DP, Williamson DF, Giles WH. Childhood abuse, household dysfunction, and the risk of attempted suicide throughout the life span: findings from the adverse childhood experiences study. *JAMA*. 2001; 286:3089–3096. [PubMed: 11754674]
62. Anda RF, Brown DW, Dube SR, Bremner JD, Felitti VJ, Giles WH. Adverse childhood experiences and chronic obstructive pulmonary disease in adults. *Am J Prev Med*. 2008; 34:396–403. [PubMed: 18407006]
63. Anda RF, Felitti VJ, Bremner JD, et al. The enduring effects of abuse and related adverse experiences in childhood. *Eur Arch Psychiatry Clin Neurosci*. 2006; 256:174–186. [PubMed: 16311898]
64. Burns EE, Jackson JL, Harding HG. Child maltreatment, emotion regulation, and posttraumatic stress: the impact of emotional abuse. *J Aggress Maltreat Trauma*. 2010; 19:801–819.
65. Brown GR, McBride L, Bauer MS, Williford WO. Cooperative Studies Program 430 Study Team. Impact of childhood abuse on the course of bipolar disorder: a replication study in U.S. veterans. *J Affect Disord*. 2005; 89:57–67. [PubMed: 16213029]

66. Goldberg JF, Garno JL. Development of posttraumatic stress disorder in adult bipolar patients with histories of severe childhood abuse. *J Psychiatr Res.* 2005; 39:595–601. [PubMed: 16019032]
67. Goldberg JF, Garno JL. Age at onset of bipolar disorder and risk for comorbid borderline personality disorder. *Bipolar Disord.* 2009; 11:205–208. [PubMed: 19267703]
68. Garno JL, Goldberg JF, Ramirez PM, Ritzler BA. Impact of childhood abuse on the clinical course of bipolar disorder. *Br J Psychiatry.* 2005; 186:121–125. [PubMed: 15684234]
69. Etain B, Henry C, Bellivier F, Mathieu F, Leboyer M. Beyond genetics: childhood affective trauma in bipolar disorder. *Bipolar Disord.* 2008; 10:867–876. [PubMed: 19594502]
70. Connor DF, Doerfler LA. Characteristics of children with juvenile bipolar disorder or disruptive behavior disorders and negative mood: can they be distinguished in the clinical setting? *Ann Clin Psychiatry.* 2012; 24:261–270. [PubMed: 23145382]
71. Ford JA. Trauma exposure and posttraumatic stress disorder (PTSD) in the lives of adolescents. *J Am Acad Child Adolesc Psychiatry.* 2013; 52:780–783. [PubMed: 23880488]
72. van Dijke A, Ford JD, van der Hart O, van Son M, van der Heijden P, Buhning M. Affect dysregulation in borderline personality disorder and somatoform disorder: differentiating under- and over-regulation. *J Pers Disord.* 2010; 24:296–311. [PubMed: 20545496]
73. Van Dijke A, Ford JD, Van der Hart O, Van Son M, van der Heijden P, Böhning M. Complex posttraumatic stress disorder in patients with borderline personality disorder and somatoform disorders. *Psychol Trauma.* 2011; 4:162–168.
74. van Dijke A, van der Hart O, Ford JD, van Son M, van der Heijden P, Buhning M. Affect dysregulation and dissociation in borderline personality disorder and somatoform disorder: differentiating inhibitory and excitatory experiencing states. *J Trauma Dissociation.* 2010; 11:424–443. [PubMed: 20938867]
75. Cicchetti D, Rogosch FA. Adaptive coping under conditions of extreme stress: multilevel influences on the determinants of resilience in maltreated children. *New Dir Child Adolesc Dev.* 2009; 124:47–59. [PubMed: 19536787]
76. De Bellis MD. Developmental traumatology: the psychobiological development of maltreated children and its implications for research, treatment, and policy. *Dev Psychopathol.* 2001; 13:539–564. [PubMed: 11523847]
77. Shipman K, Zeman J, Penza S, Champion K. Emotion management skills in sexually maltreated and nonmaltreated girls: a developmental psychopathology perspective. *Dev Psychopathol.* 2000; 12:47–62. [PubMed: 10774595]
78. Shields A, Ryan RM, Cicchetti D. Narrative representations of caregivers and emotion dysregulation as predictors of maltreated children's rejection by peers. *Dev Psychol.* 2001; 37:321–337. [PubMed: 11370909]
79. Maughan A, Cicchetti D. Impact of child maltreatment and interadult violence on children's emotion regulation abilities and socioemotional adjustment. *Child Dev.* 2002; 73:1525–1542. [PubMed: 12361317]
80. Shipman K, Edwards A, Brown A, Swisher L, Jennings E. Managing emotion in a maltreating context: a pilot study examining child neglect. *Child Abuse Negl.* 2005; 29:1015–1029. [PubMed: 16159666]
81. Rogosch FA, Cicchetti D. Child maltreatment, attention networks, and potential precursors to borderline personality disorder. *Dev Psychopathol.* 2005; 17:1071–1089. [PubMed: 16613431]
82. Cloitre M. Beyond PTSD: emotion regulation and interpersonal problems as predictors of functional impairment in survivors of childhood abuse. *Behav Ther.* 2005; 36:119–124.
83. Tull MT, Barrett HM, McMillan ES, Roemer L. A preliminary investigation of the relationship between emotion regulation difficulties and posttraumatic stress symptoms. *Behav Ther.* 2007; 38:303–313. [PubMed: 17697854]
84. Messman-Moore TL, Walsh KL, DiLillo D. Emotion dysregulation and risky sexual behavior in revictimization. *Child Abuse Negl.* 2010; 34:967–976. [PubMed: 21030084]
85. Kim J, Cicchetti D. Longitudinal pathways linking child maltreatment, emotion regulation, peer relations, and psychopathology. *J Child Psychol Psychiatry.* 2010; 51:706–716. [PubMed: 20050965]

86. Bradley B, DeFife JA, Guarnaccia C, et al. Emotion dysregulation and negative affect: association with psychiatric symptoms. *J Clin Psychiatry*. 2011; 72:685–691. [PubMed: 21658350]
87. Ford JD, Steinberg KL, Hawke J, Levine J, Zhang W. Randomized trial comparison of emotion regulation and relational psychotherapies for PTSD with girls involved in delinquency. *Clin Child Adolesc Psychol*. 2012; 41:27–37.
88. Ford JD, Hawke J. Trauma affect regulation psychoeducation group and milieu intervention outcomes in juvenile detention facilities. *J Aggress Maltreat Trauma*. 2012; 21:365–384.
89. Bowlby, J. *A secure base: parent-child attachment and healthy human development*. New York: Basic; 1988.
90. Bretherton, I.; Munholland, K. Internal working models in attachment relationships: elaborating a central construct in attachment theory. In: Cassidy, J.; Shaver, PR., editors. *Handbook of attachment: theory, research, and clinical applications*. New York: Guilford; 2008. p. 102-127.
91. Pollak SD, Cicchetti D, Hornung K, Reed A. Recognizing emotion in faces: developmental effects of child abuse and neglect. *Dev Psychol*. 2000; 36:679–688. [PubMed: 10976606]
92. Pollak SD, Tolley-Schell SA. Selective attention to facial emotion in physically abused children. *J Abnorm Psychol*. 2003; 112:323–338. [PubMed: 12943012]
93. Pollak SD, Messner M, Kistler DJ, Cohn JF. Development of perceptual expertise in emotion recognition. *Cognition*. 2009; 110:242–247. [PubMed: 19059585]
94. Ford JD, Stockton P, Kaltman S, Green BL. Disorders of extreme stress (DESNOS) symptoms are associated with type and severity of interpersonal trauma exposure in a sample of healthy young women. *J Interpers Violence*. 2006; 21:1399–1416. [PubMed: 17057159]
95. Barnes JE, Noll JG, Putnam FW, Trickett PK. Sexual and physical revictimization among victims of severe childhood sexual abuse. *Child Abuse Negl*. 2009; 33:412–420. [PubMed: 19596434]
96. Kim K, Trickett PK, Putnam FW. Childhood experiences of sexual abuse and later parenting practices among non-offending mothers of sexually abused and comparison girls. *Child Abuse Negl*. 2010; 34:610–622. [PubMed: 20541259]
97. Noll JG, Trickett PK, Harris WW, Putnam FW. The cumulative burden borne by offspring whose mothers were sexually abused as children: descriptive results from a multigenerational study. *J Interpers Violence*. 2009; 24:424–449. [PubMed: 18451100]
98. Kim K, Trickett PK, Putnam FW. Attachment representations and anxiety: differential relationships among mothers of sexually abused and comparison girls. *J Interpers Violence*. 2011; 26:498–521. [PubMed: 20442447]
99. Briere J, Rickards S. Self-awareness, affect regulation, and relatedness: differential sequels of childhood versus adult victimization experiences. *J Nerv Ment Dis*. 2007; 195:497–503. [PubMed: 17568298]

Table 1

Studies Describing Exposure to Traumatic Stressors or Maltreatment, with Psychiatric Comorbidities in Children and Adults

| Study | Subjects | Diagnostic instruments | Findings |
|--|--|---|--|
| Trauma exposure in childhood | | | |
| Famularo et al. (1996) ⁴³ | n = 117 Age: 6–12 years Maltreated children presenting to juvenile/family court | Diagnostic Interview for Children and Adolescents, Revised Version | 35% PTSD Those with PTSD had increased risk for ADHD, anxiety disorders, psychotic symptom, suicidal ideation |
| Ford et al. (2000) ⁴⁴ | n = 165 Age: 6–17 years ADHD, oppositional defiant disorder, adjustment disorder | CBCL PTSD Checklist Traumatic Events Screening Inventory | Children with oppositional defiant disorder suffered more physical abuse & sexual abuse, & had increased PTSD symptom severity Rates of physical & sexual abuse and symptom severity as measured by CBCL: oppositional defiant disorder + ADHD > oppositional defiant disorder > ADHD > adjustment disorder |
| Lansford et al. (2002) ⁴⁵ | n = 585 youth 79% participated from kindergarten through 11th grade | Parent interview at kindergarten School records at 11th grade CBCL at 11th grade Adolescent Behavior Questionnaire | n = 69 (11.8%) had suffered physical abuse prior to kindergarten; had increased school absence, social difficulties, social withdrawal, anxiety, depression, PTSD, dissociation, thought problems; had decreased expectations of a higher education |
| Copeland et al. (2007) ⁴⁶ | n = 1420 Age: 9, 11, & 13 years at intake Followed annually until age 16 | CBCL Child and Adolescent Psychiatric Assessment | By age 16, 37% had exposure to multiple traumatic events; of those 13.4% had posttraumatic stress symptoms; 0.5% had PTSD Trauma had increased anxiety & depression, & had doubled other psychiatric disorders (all diagnostic groups except for substance abuse) |
| Trauma exposure in early childhood | | | |
| Mongillo et al. (2006) ⁴⁷ | n = 917 Age: 18–36 months | The Infant-Toddler Social & Emotional Assessment CBCL | From 6–36 months, 23.4% experienced at least one adverse/potentially traumatic event Trauma had led to increased symptom severity, internalizing, & externalizing; 20% PTSD |
| Briggs-Gowan et al. (2010) ⁴⁸ | n = 213 Age: 2–4 years | Preschool Age Psychiatric Assessment Child Life Events Scale | Exposure to domestic violence had resulted in increased internalizing (depression, separation anxiety, posttraumatic stress) & externalizing (conduct problems) |
| Briggs-Gowan et al. (2012) ⁴⁹ | n = 437 Age: 1 year Followed annually until age 3 | Child Life Events Scale The Infant-Toddler Social & Emotional Assessment CBCL Adaptive Social Behavior Ratings Scale | Exposure to domestic violence & neighborhood violence before the age of 3 years had resulted in increased posttraumatic stress symptoms & externalizing, & more social difficulties |
| Adverse childhood experiences | | | |
| Chapman et al. (2004) ⁵⁰ | n = 2946 adults | Adverse Childhood Experience Study Questionnaires Screening instrument for depressive disorders developed for the Medical Outcomes Study Psychiatric Screening Questionnaires for Primary Care Patients | Increased adverse childhood experiences increased the risk of lifetime & current depressive disorders |
| Lu et al. (2008) ⁵¹ | n = 254 adults | SCID Sexual Abuse Exposure Questionnaire PTSD Checklist Dartmouth Assessment of Lifestyle Instrument Revised Conflict Tactics Scales | Increased adverse childhood experiences associated with high-risk behaviors, substance abuse, revictimization, PTSD, & increased psychiatric illness severity at a younger age |
| Adult sequelae of childhood trauma exposure | | | |

| Study | Subjects | Diagnostic instruments | Findings |
|---------------------------------------|---|---|---|
| Yen et al. (2002) ⁵² | Adults Collaborative Longitudinal Personality Disorder Study (subgroup) | Diagnostic Interview for DSM-IV Personality Disorders SCID SCID Trauma Addendum | Adults with Borderline personality disorder had more trauma, childhood sexual abuse/trauma, more PTSD; first exposure to trauma was at a younger age |
| Golier et al. (2003) ⁵³ | n = 180 outpatients with personality disorder | Trauma History Questionnaire SCID, PTSD Module | Outpatient adults with Borderline personality disorder had more childhood/adulthood physical abuse & current PTSD Trauma had stronger association with paranoid personality disorder |
| Van Dijke et al. (2012) ⁵⁴ | n = 472 psychiatric inpatients | Composite International Diagnostic Interview Borderline Personality Disorder Severity Index Bermond-Vorst Alexithymia Questionnaire | Childhood trauma inflicted by a primary caregiver increased emotional dysregulation in inpatients with borderline personality disorder |

ADHD, attention-deficit/hyperactivity disorder; CBCL, Child Behavior Check List; PTSD, posttraumatic stress disorder; SCID, Structured Clinical Interview for DSM.

Table 2

Studies Describing Childhood Maltreatment and Emotional Dysregulation

| Study | Subjects | Measures of emotional dysregulation | Findings |
|---|---|---|---|
| Shields et al. (1998) ²⁸ | Maltreated children: n = 141 Healthy controls: n = 87 Age: 6–12 years | CBCL Teacher Report Form Emotion Regulation Checklist | Maltreated children had increased aggression & decreased attention, emotional regulation, & socially appropriate expression of emotions Physically abused children showed increased reactive aggression |
| Shipman et al. (2000) ⁷⁷ | Sexual abuse: n = 21 girls Healthy controls | Emotional understanding interview Children's Emotion Management Scales (Anger & Sadness) Emotion Regulation Checklist Emotion Management Interview | Sexually abused girls had decreased emotion regulation & understanding, & expected to be receiving less emotional support; had increased interpersonal conflict in response to expression of negative emotional states (sadness to parents, anger to peers) |
| Shields et al. (2001) ⁷⁸ | Maltreated children: n = 76 Healthy controls: n = 45 Age: 8–12 years | Rochester Parenting Stories | Maltreated children had increased emotion dysregulation & aggression, & decreased social competence |
| Maughan et al. (2002) ⁷⁹ | Maltreated children: n = 88 Healthy controls: n = 51 Age: 4–6 years | Person-oriented emotional-regulation patterns in response to simulated inter-adult anger CBCL | Maltreated children had increased emotion dysregulation |
| Shipman et al. (2005) ⁸⁰ | Maltreated children: n = 24 Healthy controls: n = 24 Age: 6–12 years | Questionnaires + interview-assessed emotional understanding & regulation | Maltreated children had decreased understanding of negative emotions (anger & sadness) & decreased emotional-regulation skills |
| Rogosch et al. (2005) ⁸¹ | Maltreated children: n = 185 Healthy controls: n = 175 School-aged | Maltreatment Classification System Attention Network Test for children Perceptions of Peers and Self Relationship Stance Questionnaire Children's Depression Inventory Teacher's report form Emotion Regulation Checklist Student-Teacher Relationship Scale Peer sociometric ratings | Maltreated children had higher mean & levels on borderline personality disorder precursors |
| Cloitre et al. (2005) ⁸² | n = 164 women | Childhood Maltreatment Interview Schedule Sexual Assault and Additional Interpersonal Violence Schedule Social Adjustment Scale Self Report Modified PTSD Symptoms Scale General Expectancy for Negative Mood Regulation Scale Inventory of Interpersonal Problems | Increased PTSD symptoms predicted greater functional impairment Decreased emotional regulation & interpersonal problems together: equal to PTSD symptoms in impact on psychosocial functioning |
| Tull et al. (2007) ⁸³ | n = 108 college students with history of trauma | Trauma Events Questionnaire PTSD Checklist Difficulties in Emotion Regulation Scale Positive and Negative Affect Schedule | Students with posttraumatic stress symptoms had increased emotion dysregulation Students with PTSD had much greater emotion dysregulation |
| Burns et al. (2010) ⁶⁴ | n = 912 female college students | Childhood Trauma Questionnaire Difficulties in Emotion Regulation Scale Trauma Symptom Inventory | Increased reports of sexual, physical, or emotional abuse were linked to increased emotion dysregulation & posttraumatic stress symptoms |
| Messman-Moore et al. (2010) ⁸⁴ | n = 752 female college students | Computer Assisted Maltreatment Inventory Difficulties in Emotion Regulation Scale Cognitive Appraisal of Risky Events–Revised Sexual Experiences Survey | Childhood physical & sexual abuse increased the risk for adolescent/adult rape Emotional dysregulation was a mediating factor for revictimization Sexually risky behavior was predicted by emotion dysregulation |
| Kim et al. (2010) ⁸⁵ | Maltreated children: n = 215 | Maltreatment Classification System Emotion Regulation Checklist Peer nominations | Increased emotion dysregulation associated with neglect; physical & sexual |

| Study | Subjects | Measures of emotional dysregulation | Findings |
|-------------------------------------|--|---|--|
| | Healthy controls: n = 206 Age: 6–12 years | Teacher's report form | abuse; multiple types of maltreatment experiences; younger age at onset Externalizing behavior were associated with increased emotion dysregulation & peer rejection (bi-directional) |
| Bradley et al. (2011) ⁸⁶ | n = 530 adults | Childhood Trauma Questionnaire Positive and Negative Affect Schedule Emotion Dysregulation Scale Clinician-administered PTSD Scale | Increased childhood trauma was associated with increased negative affect, posttraumatic stress symptoms, drug abuse, depression, suicidality & emotional dysregulation |

CBCL, Child Behavior Check List; PTSD, posttraumatic stress disorder.