



Comparison of Methodologies to Quantify Phytate Phosphorus in Diets Containing Phytase and Excreta from Broilers

L. de P. Naves*, P. B. Rodrigues, A. G. Bertechini, A. D. Corrêa¹, D. H. de Oliveira, E. C. de Oliveira, W. F. Duarte², and M. R. R. da Cunha³

Animal Science Department, Federal University of Lavras (UFLA), Lavras, Minas Gerais, 37200-000, Brazil

ABSTRACT: The use of a suitable methodology to quantify the phytate phosphorus (P_{phy}) content in both the feed and the excreta from broilers is required to enable accurate calculation of the catalytic efficiency of the phytase supplemented in the feed. This study was conducted to compare 2 analytical methodologies (colorimetry and also high-performance liquid chromatography with a refractive index detector) in order to calculate the phytase efficiency by utilizing the results from the methodology that was shown to be the most appropriate. One hundred and twenty broilers were distributed in a (4+1)×2 factorial arrangement, corresponding to 4 diets that were equally deficient in P supplemented with increasing levels of phytase (0, 750, 1,500, and 2,250 units of phytase activity – FTU – per kg of feed) plus 1 positive control diet without phytase, supplied to male and female birds. The result indicated that the colorimetric methodology with an extraction ratio of 1:20 (mass of sample in g:volume of the solvent extractor in mL) was shown to be the most adequate. There was no interaction between the phytase level and the sex of the broilers ($p>0.05$). Males consumed 12% more P_{phy} than did females ($p<0.01$), but the sex of the broilers did not affect ($p>0.05$) the excretion and retention coefficient of P_{phy} . The increase in the phytase level of the diet reduced (linear, $p<0.01$) the P_{phy} excretion. The greatest P_{phy} retention was estimated at 87.85% when the diet contained 1,950 FTU/kg ($p<0.01$), indicating that it is possible to reduce the inorganic P in the formulation at an amount equivalent to 87.85% of the P_{phy} content present in the feed, which, in this research, corresponds to a decrease in 2.86 g of P/kg of the feed. (**Key Words:** Broiler, Myo-inositol Phosphate, Phosphorus, Phytase, Phytate, Phytic Acid)

INTRODUCTION

Although the economic and environmental benefits obtained through the use of phytase in broiler diets are reported in the literature (Nagata et al., 2011; Santos et al., 2011a,b; Gomide et al., 2012), its catalytic efficiency to make the phytate phosphorus (P_{phy}) of the feed available is not yet well established, and, according to Angel et al.

(2002), this should be done while considering the P_{phy} concentration in both the feed and the excreta. Among the analytical methods used for the phytate determination, those that utilize colorimetric and refraction principles stand out (Wu et al., 2009). However, these methodologies have not yet been compared with each other for samples of feed and excreta. This comparison is important because the use of an inappropriate analytical methodology might lead to misinterpretations of the phytase efficiency.

The colorimetric method is based on a reaction between the ferric ion (Fe^{3+}) and the sulfosalicylic acid (wade reagent [WR]) with the formation of a complex that can be spectrophotometrically monitored at 500 nm. In the presence of phytate, Fe^{3+} binds preferentially to P_{phy} , reducing the color intensity of WR. However, as the Fe^{3+} does not discriminate between the P_{phy} or the inorganic P (P_{inorg}) present in the sample, it is necessary that, before the colorimetric dosing, the sample go through an anion

* Corresponding Author: Luciana de Paula Naves. Tel: +55-35-3829-1691, Fax: +55-35-3829-1231, E-mail: luciana.naves@hotmail.com

¹ Chemistry Department, UFLA, Lavras, Minas Gerais, 37200-000, Brazil.

² Biology Department, UFLA, Lavras, Minas Gerais, 37200-000, Brazil.

³ Chromatography Laboratory, Ezequiel Dias Foundation, Belo Horizonte, Minas Gerais, 30510-010, Brazil.

Submitted Aug. 29, 2013; Revised Dec. 10, 2013; Accepted Jan. 24, 2014

exchange resin to separate P_{phy} from P_{inorg} (Latta and Eskin, 1980). After the correlation with a standard curve of phytic acid (IP_6), the result obtained is multiplied by 0.282 to express the content of P_{phy} in the sample because this constant corresponds to the molar ratio of P in the IP_6 molecule (Angel et al., 2002). However, it is necessary to assess whether this methodology overestimates the P_{phy} in samples containing phosphate esters with fewer than six phosphate groups, which is expected in excreta from broilers fed diets with phytase because according to Greiner et al. (2002), the products of the enzyme degradation of IP_6 can include different phosphorylation degrees of the myo-inositol (IP_5 to IP_1). In contrast, the high-performance liquid chromatography with a refractive index detector (HPLC-IR) discriminated the fractions from IP_3 to IP_6 in rice bran treated with phytase (Cúneo et al., 2000), arousing the interest in evaluating it in experiments with broilers.

Thus, the objectives of this study were to select the most adequate methodology for the determination of the P_{phy} content in both feed and excreta from broilers, to calculate the supplementation level of phytase in which there is the highest retention coefficient of P_{phy} and to estimate the P concentration value that can be reduced in the feed formulation when this level of phytase is used.

MATERIALS AND METHODS

Experimental materials and procedures

One metabolism experiment was conducted in Lavras, state of Minas Gerais, Brazil (21°13'52" S; 44°58'16"W), with broiler chickens in the period from 15 to 28 days of age to compare two analytical methodologies employed to quantify the content of P_{phy} in both the feeds and excreta from birds and to determine the catalytic efficiency of the phytase supplemented in the diet by utilizing the results from the methodology that was shown to be the most appropriate.

All the experimental procedures adopted were approved by the Committee of Ethics in the Use of Animals (protocol code. 004/11). The experiment was conducted in a completely randomized design, distributed in a (4+1)×2 factorial arrangement, corresponding to 4 diets equally deficient in available P (P_{av}) supplemented with increasing levels of phytase (0, 750, 1,500, and 2,250 units of phytase activity – FTU – per kg of feed) plus 1 positive control diet without phytase, formulated according to Rostagno et al. (2005). The 6-phytase utilized in this study (EC 3.1.3.26 synthesized by *Aspergillus oryzae*) was a powder commercial product marketed by DSM Company, with determined activity of 9,980 FTU/g of product, according to the protocol of Engelen et al. (1994). All 5 diets were supplied to male and female broilers for a total of 10 treatments with 3 replications of 4 birds per pen. The use of

male and female birds within a cage (mixed housing) could increase the coefficient of variation of the experiment; thus, the sex was considered a factor within the factorial design in order to minimize the variation between repetitions of the same treatment, promoting an experiment with a smaller mean square error.

One hundred and twenty broiler chicks (60 birds of each sex) of the Cobb-500 lineage were acquired at one day of age, sexed, and raised in a conventional shed for broilers up to 14 days of age and received a diet based on corn and soybean meal that was formulated to meet their nutritional requirements (Rostagno et al., 2005). On the 15th day of age, the broilers were weighed individually, separated by weight ranges, and transferred to the cages (experimental unit; with dimensions 50×50×50 cm) in a metabolism room. The experimental units were arranged to the average initial weight of the broilers, which was similar to one another (0.465 g±0.002). The room offered illumination and thermostatically controlled temperature. Each cage was provided with a pressure-type drinker, an individual gutter-type feeder with border, and a tray to collect the excreta. Experimental diets (Table 1) and water were offered *ad libitum* in the period from 15 to 28 days of age.

The experimental period comprised 13 days, ten days of which were destined for adaptation to the facilities and experimental diets, followed by three days (Rodrigues et al., 2005) of total collection of excreta, per experimental unit. Feed and leftovers were weighed and recorded on the 25th and 28th days of the age of broilers for subsequent calculations of feed intake during that period.

Sample collection and chemical analysis

The excreta were collected daily in the morning, grouped according to the experimental unit in identified plastic bags, and stored in a freezer up to the last day of collection when the excreta were weighed, homogenized, and pre dried in a forced-ventilation oven at 55°C for 72 h. The pre-dried excreta were ground, and the total dry matter content was obtained by considering the loss of moisture during the pre-drying stage and also at 105°C until constant weight (method 934.01 of the AOAC, 2005).

The P_{phy} content of the excreta was determined by colorimetry according to Frühbeck et al. (1995) with modifications and also by HPLC-IR according to Nappi et al. (2006), with adaptations. Both methodologies were utilized to analyze the extracts obtained in two extraction ratios of mass of sample:volume of extractor solvent (1:20 and 3:20, in g/mL). Parallel to this, three samplings of corn, soybean meal, and the positive and negative control diets were collected, ground, and analyzed in a manner that was similar to the one employed for the excreta. Thus, 1 or 3 g of sample was suspended in 20 mL of 0.66 M HCl followed by mechanical agitation at room temperature during two

Table 1. Composition and nutritional contents of the experimental diets

Ingredient (g/kg of natural matter)	PC ¹	NC ²	NC supplemented with phytase ³
			750/1,500/2,250
Maize	596.20	604.86	604.86
Soybean meal, 48% CP	335.29	333.71	333.71
Soybean oil	31.61	28.68	28.68
Salt	4.82	4.82	4.82
L-lysine HCl 78%	1.67	1.70	1.70
DL-methionine 99%	2.29	2.28	2.28
L-threonine 98.5%	0.35	0.35	0.35
Mineral premix ⁴	0.50	0.50	0.50
Vitamin premix ⁵	0.40	0.40	0.40
Choline chloride 60%	0.45	0.45	0.45
Calcitic limestone	8.74	16.10	16.10
Dicalcium phosphate	17.19	5.65	5.65
Inert (kaolin)	0.500	0.500	0.43/0.35/0.28
Phytase ⁶	0.000	0.000	0.07/0.15/0.22
Calculated nutritional composition (g/kg of natural matter)			
Metabolizable energy (kcal/kg)	3,050	3,050	3,050
Crude protein	201.2	201.2	201.2
Digestible lysine	11.10	11.10	11.10
Digestible methionine+cysteine	7.94	7.94	7.94
Sodium	2.10	2.10	2.10
Calcium	8.55	8.55	8.55
Total phosphorus	6.39	4.26	4.26
Available phosphorus ⁷	4.26	2.13	2.13
Phytate phosphorus ⁸	2.13	2.13	2.13
Phytase activity ^{3,6}	0	0	750/1,500/2,250

¹ PC, positive control diet formulated to meet the nutritional requirements of broilers, in the period from 15 to 28 days of age, according to Rostagno et al. (2005).

² NC, negative control diet deficient in available phosphorus.

³ Enzyme supplementation expressed in FTU/kg of diet, in which 1 FTU corresponds to one unit of phytase activity.

⁴ Content kg⁻¹ of diet: 55 mg Zn; 48 mg Fe; 10 mg Cu; 78 mg Mn; 0.70 mg I; 0.18 mg Se.

⁵ Content kg⁻¹ of diet: 4.8 mg vit. B₂; 8,000 UI vit. A; 2,000 UI vit. D₃; 1.2 mg vit. B₁; 2.4 mg vit B₆; 10.8 µg vit. B₁₂; 0.024 mg biotin; 0.64 mg folic acid; 11.6 mg pantothenic acid; 1.92 mg vit. K₃; 16.2 UI vit. E.

⁶ Ronozyme NP-(CT). 6-phytase (EC 3.1.3.26) synthesized by microorganism *Aspergillus oryzae* (DSM Company, 17594). Enzyme with declared activity of 10,000 FTU/g of product and determined activity of 9,980 FTU/g of product, according to the protocol of Engelen et al. (1994).

⁷ Considering that one-third of the total phosphorus present in the corn and soybean meal correspond to available phosphorus and that the phosphorus supplied by the dicalcium phosphate is 100% available (Rostagno et al., 2005).

⁸ Considering that two-thirds of the total phosphorus present in the corn and soybean meal correspond to phytate phosphorus (Rostagno et al., 2005).

hours. Afterward, the extract was centrifuged at 14,881×g for 30 min at 15°C, and the supernatant was collected and filtered through Qualy 14 µm pore filter paper. At the 1:20 extraction, the filtered supernatant was diluted in function of the concentration of phytate present in the sample and of the standard curve of IP₆ (colorimetric method). The dilutions (v:v in mL) utilized for the corn, soybean meal, and diets were, respectively, 3:25, 1:25, and 2:25. However, for the excreta, the dilution varied from 1:25 to 2:25 in function of the treatment because when a higher level of phytase is supplemented in the diet, there is a lower content of P_{phy} in the excreta requiring, consequently, a lower dilution level. After the dilution, the pH was adjusted with

1.0 M NaOH to 6.0. In the case of the 3:20 extraction, the filtered solution collected after centrifugation had its pH adjusted to 6.0 without going through the dilution step conducted at the 1:20 extraction.

The sample purification to separate the P_{phy} from P_{inorg} was conducted in glass columns (0.7×15 cm) packed with 0.5 g of anion exchange resin (AG 1-X4, 100-200 dry mesh, Bio-Rad) and washed with 15 mL of 0.6 M NaCl and 30 mL of ultrapure water, respectively. After this resin activation process, 10 mL of a sample with pH 6.0 were applied to the column, and the eluate was discarded. Subsequently, 15 mL of 0.6 M NaCl were utilized for the elution of the polyphosphates retained in the resin, and the

pH of the eluate was collected and adjusted to 3.0 with 0.1 M HCl because this pH facilitates the link between phosphate and WR (Latta and Eskin, 1980).

In the sequence of the colorimetric method, an aliquot of the eluate that had been collected (3 mL for the 1:20 extraction or 100 to 600 μ L for the 3:20 extraction; in these latter cases, ultrapure water was pipetted to maintain a final volume of 3 mL) was mixed with 1 mL of WR (consisting of 0.03% of ferric chloride hexahydrate and 0.3% of sulfosalicylic acid). The tubes were agitated in a vortex, and absorbance was read at 500 nm using ultrapure water to zero the spectrophotometer. The standard curve of IP₆ was made by pipetting increasing volumes of the stock solution (200 mg/L) of IP₆ (P0109, batch 057k0049, Sigma-Aldrich), so as to obtain concentrations varying from 0 to 150 μ g; each tube received 1 mL WR, and the final volume was equivalent to 4 mL. The conversion of the concentration of phytate into P_{phy} was done considering the molar ratio of the P in the molecule of IP₆ of 28.2% (Angel et al., 2002).

The remainder of the eluate that had been collected (15 mL eluted from the anion exchange resin minus the aliquot utilized for the colorimetric determination) was evaporated at 40°C till that, a solid residue was obtained, which was subsequently re-suspended in 2 mL of the mobile phase, centrifuged twice sequentially (15,000 \times g for 10 min at 4°C), filtered in a GV hydrophilic membrane (0.22 μ m pore and 25 mm diameter, Durapore), and stored at 4°C until quantification of the myo-inositol phosphates by HPLC-IR.

The mobile phase consisted of 900 mL of 50 mM formic acid, 900 mL of methanol, 45 mL of 5 mM EDTA, and 54 mL of tetrabutylammonium hydroxide 20%. The pH of the mobile phase was adjusted to 4.3 with sulfuric acid and before being injected in the chromatograph, the mobile phase was vacuum filtered and sonicated (degassed) for 50 min. The analyses were performed in a Shimadzu chromatograph that was equipped with a refractive index detector (RID10Ai) and a C₁₈ CAPCELL PAK column (5 μ m \times 4.6 mm \times 150 mm) with a mobile phase at a flow of 0.5 mL/min. The temperature of the oven was maintained at 45°C, and the injection volume was 20 μ L.

For the identification of phosphorylated fractions, the retention time of the standard compounds myo-inositol hexaphosphate or phytic acid (IP₆, code P0109, batch 057k0049); myo-inositol-1,3,4,5,6-pentaphosphate (IP₅, code I9261, batch 114K12831); myo-inositol-1,3,4,6-tetraphosphate (IP₄, code I9386, batch 60K17321); and myo-inositol-1,4,5-triphosphate (IP₃, code. I8761, batch 21H84341), obtained under the same conditions of analysis, were compared with the retention times of the peaks obtained in the samples. The IP₅ was utilized as an internal standard for quantification of the phosphorylated fractions (IP₃ to IP₆) present in the samples using the following

equation (Sandberg and Ahderinne, 1986):

$$\text{Concentration of IP}_Y \text{ in the sample (in mg/mL)} \\ = \frac{0.32 \times \text{IP}_Y \text{ peak area in the sample}}{\text{Peak area of the internal standard}}$$

In this equation, IP_Y corresponds to the different fractions of myo-inositol phosphates (IP₃, IP₄, IP₅, or IP₆). The conversion of the concentration of IP₆ in P was calculated, considering the molar ratio of P in the IP₆ molecule, which is 28.2%. For fractions IP₅, IP₄, and IP₃, the molar ratios of 26.7, 24.8, and 22.1%, respectively, were utilized.

The protein content in the extracts (mg of protein per ml of extract) was dosed according to method of Bradford (1976) by utilizing bovine serum albumin as a standard. This analysis was conducted in samples extracts centrifuged and no centrifuged to investigate the hypothesis of that phytate can form complexes with proteins of manner that these phytate-protein complexes can interfere in the phytate quantification.

Balance and coefficient of hydrolysis (or retention) of phytate phosphorus

These parameters were calculated on a dry matter basis, utilizing the P_{phy} contents determined by the two methodologies evaluated. The P_{phy} intake (g/bird) was calculated by multiplying the feed intake (g/bird) by the P_{phy} content (%) in the feed. To calculate the absolute excretion of P_{phy} (g/bird), the amount of excreta (g/bird) was multiplied by the P_{phy} content (%) in the excreta. The hydrolysis coefficient of P_{phy} was calculated by equation:

$$\text{Coefficient of hydrolysis of P}_{\text{phy}} \\ = \frac{(\text{P}_{\text{phy}} \text{ intake} - \text{excretion of P}_{\text{phy}}) \times 100}{\text{P}_{\text{phy}} \text{ intake}}$$

Statistical analysis

Each cage was considered one experimental unit, and an independent statistical analysis was conducted for each analytical methodology evaluated. The data of the intake, excretion, and coefficient of hydrolysis of P_{phy} were subjected to analysis of variance (p<0.05) by utilizing command General Linear Model of software SAS (2004), and when significant, polynomial regression models were utilized (p<0.05) to evaluate the effect of increasing supplementation of phytase on these parameters. The significance of the effect of the sex of birds was evaluated by the F test itself (p<0.05). In addition, the positive control diet was compared with the other experimental diets by the Dunnett test (p<0.05).

RESULTS AND DISCUSSION

Concentrations of phytate and phytate phosphorus determined in ingredients, diets and excreta

The initial proposal of this research was to compare the two analytical methodologies (colorimetry and HPLC-IR) only at the extraction ratio of 1:20 (mass of sample in g:volume of extractor solvent in mL), as Brito (2008) had already managed to quantify phytate in diets and excreta from broilers by colorimetry at this extraction ratio. However, with the 1:20 extraction, it was not possible to detect all the peaks with regard to the phosphorylated fractions of myo-inositol by the HPLC-IR method (especially for the samples of excreta from broilers fed the diets containing the highest levels of phytase) and, besides, the peaks identified in the chromatogram presented an intensity that was inferior to the limit of quantification of the analyte in the analysis conditions. Therefore, the extract was concentrated so as to enable the analytes of interest to be quantified via HPLC. During this procedure of concentration of the extract, it was observed that the highest possible ratio was of 3:20 because at higher ratios, the sample would become pasty, which is undesirable, as this could compromise the process of solubilization of the analytes.

For the colorimetric method, the extraction at the ratio of 3:20 resulted in the determination of lower concentrations of phytate and, consequently, lower contents of P_{phy} (Tables 2 and 3), probably due to the interference from the proteins present in this extract. According to Frühbeck et al. (1995), the pH of the diluted aliquot should be adjusted to 6.0 because this value is above of the isoelectric point of most of the vegetable proteins; which means that in theory, these proteins would present negative net charge and there would be electrostatic repulsion with the phytate, thus preventing the formation of complexes between inositol phosphate (IP) and protein. However, it appears that the formation of these phytate-protein complexes occurred, even adjusting the pH to 6.0.

To investigate the formation of these phytate-protein complexes, samples extracted at the ratio of 3:20 were centrifuged, and a great amount of precipitate (pellet) was visually verified, with only 52.07 ± 2.32 μg protein per ml of extract determined in the supernatant (the non-centrifuged extracts resulting from extraction ratios of 1:20 and 3:20 had presented 109.41 ± 1.30 and 800.73 ± 1.71 μg protein/mL, respectively). Moreover, no trace of phytate was detected in the centrifuged supernatant, reinforcing the hypothesis that phytate was complexed with the proteins from the sample, and precipitated after centrifugation due to the high molecular weight of these phytate-protein complexes.

Furthermore, with the 1:20 extraction, the sample would

naturally go through the anion exchange resin by the action of gravity; however, with the 3:20 extraction, the sample appeared blurred, and for their passage through the resin, a pressure pump had to be used. Therefore, at the extraction ratio of 3:20, the complexation with proteins results in an underestimation of the phytate content because a portion of the IP-protein complexes probably does not go through resin, and those which might be eluted with the saline solution, cannot be properly quantified due to their inadequate reaction with the WR. In addition to the proteins, phytate can also complex with minerals (Kornegay, 2001), which might interfere in the dosage of phytate; however, as the protein content in the diet is much higher than the mineral content, it is likely that the effect of protein is more pronounced.

When the extract at the rate of 1:20 was dosed by the colorimetric method (Table 2), the contents of P_{phy} determined for corn and soybean meal in this study were approximately 30% and 44% higher than the mean values presented in the last edition of the Brazilian Tables for Poultry and Swine: 0.217 and 0.437 g P_{phy} per 100 g DM of corn and soybean meal, respectively (Rostagno et al., 2011); however, the methodology utilized for the determination of these contents was not published. In addition to the methodology, the differences observed can also be due to environmental, genetic, and/or cultivation variations, which can alter the content of P_{phy} in the seeds and grains. For example, Wadt et al. (2010) collected 94 samples of corn originated from the South, Southeast, and Midwest regions of the Brazil in the summer 2009 harvest, and they verified differences not only within the same region but also between cultivation regions, with P_{phy} contents determined ranging from 1.0 to 2.6 g/kg of corn, which represents a variation of up to 160% in the P_{phy} contents determined in the samples. Therefore, in this study, the colorimetric methodology at the extraction ratio of 1:20 was shown to be the most adequate method for quantifying the P_{phy} .

The retention times (in minutes) for standards IP_3 , IP_4 , IP_5 , and IP_6 were 6.793, 10.386, 15.113, and 18.791, respectively. The P_{phy} contents determined in corn, soybean meal, and diets by the HPLC-IR methodology at the extraction ratio of 3:20 were inferior to those determined by the colorimetric method at the extraction ratio of 1:20 (Table 2), which also should be due to the complexation of phytate with other molecules present in the sample, especially proteins. However, this was not observed for the excreta, which can be justified by the profile of esters of myo-inositol phosphate present in the sample (Table 3).

In general, the higher the degree of phosphorylation of myo-inositol, the higher its power of complexation because the greater the number of phosphate groups in the molecule, the greater the number of negative charges resulting from

Table 2. Contents of phytate and phytate phosphorus in ingredients and diets, determined according to two analytical methodologies and extraction ratios

Content (g/100 g of dry matter)	Corn	Soybean meal	NC diet ¹	PC diet ²
HPLC-IR ^{3,4} 3:20 (m:v)				
Total myo-inositol phosphates ⁵	0.031	0.549	0.173	0.179
Phytate phosphorus ⁶	0.009	0.154	0.049	0.050
P of IP ₃	0.000	0.000	0.000	0.000
P of IP ₄	0.000	0.000	0.000	0.000
P of IP ₅	0.000	0.018	0.008	0.008
P of IP ₆	0.009	0.136	0.041	0.042
Colorimetry 3:20 (m:v)				
Phytate	0.107	0.375	0.112	0.106
Phytate phosphorus ⁷	0.030	0.106	0.032	0.030
Colorimetry 1:20 (m:v)				
Phytate	1.000	2.230	1.341	1.348
Phytate phosphorus ⁷	0.282	0.629	0.378	0.380

m:v, mass of sample in g:volume of extractor solvent in mL.

IP_x, myo-inositol phosphates, wherein x corresponds to the degree of phosphorylation of the molecule.

¹ Negative control (NC) diet formulated with 2.13 g of available phosphorus/kg of diet.

² Positive control (PC) diet formulated with 4.26 g of available phosphorus/kg of diet.

³ HPLC-IR, high-performance liquid chromatography with refractive index detector.

⁴ At the extraction ratio of 1:20 (m:v), the concentration of myo-inositol phosphates in the aliquot injected in the chromatograph was below of the limit of detection and/or quantification.

⁵ Corresponding to the sum of fractions IP₃, IP₄, IP₅, and IP₆.

⁶ Phytate phosphorus (%) = (% of IP₃×0.221)+(%) of IP₄×0.248)+(%) of IP₅×0.267)+(%) of IP₆×0.282).

⁷ Phytate phosphorus (%) = determined phytate content (%)×0.282.

the deprotonation of these phosphate groups (Kornegay, 2001). In this sense, Yu et al. (2012) recently evaluated the effect of the degree of phosphorylation of myo-inositol on the potential of the molecule to complex with the soy protein and bovine β-casein and concluded that, for both proteins, IP₆ has a high chelating power which is strongly reduced by the loss of one phosphate group (IP₅), becoming practically null in the other phosphorylated fractions (IP₄ to IP₁). Therefore, due to the fractions with fewer than five phosphate groups in the excreta (Table 3), the interference of the proteins in these samples was lower than in the ingredients and diets.

In the corn, only fraction IP₆ was detected; whereas in soybean meal and the diets, the myo-inositol phosphates were distributed between IP₅ and IP₆, but with predominance of IP₆ (Table 2). Consequently, probably there were high rates of complexing of these fractions with other molecules or ions, and once complexed, the physico-chemical properties of these complexes are different from isolated fractions, thus causing alterations in the retention times, impeding these complexes from being identified in the chromatogram, leading to underestimation of the results. This interference by the formation of complexes was lower in the excreta due to the predominance of fractions with low complexing capacity such as IP₄ and IP₃, which are products from the enzymatic action of phytase on the phytate.

Balance and coefficient of hydrolysis (or retention) of phytate phosphorus

Although it is possible to make comparisons between the methodologies to quantify P_{phy} based on the contents determined, these data are inappropriate to evaluate the phytase efficiency to make P_{phy} available for broilers. Therefore, it is necessary to take into account not only the P_{phy} content determined in the diets and excreta, but also the feed intake and the amount of excreta produced in the collection period. As previously discussed, the methodologies of colorimetry and HPLC-IR at the extraction ratio of 3:20 presented problems in the determination of the P_{phy} content, which is reflected in inadequate balance and retention of P_{phy} (Table 4).

In this regard, the colorimetric methodology with the extraction ratio of 1:20 was shown to be the most adequate, and, therefore, presented the most reliable results on the balance and retention of P_{phy} by the broilers (Table 4). There was no interaction (p>0.05) between the levels of phytase and sex for intake, excretion, or the coefficient of hydrolysis of P_{phy}. In addition, the birds sex affected (p<0.01) only the intake of P_{phy}, so that the males consumed 12% more P_{phy} than did the females. Therefore, it is possible to consider that the mean values obtained to calculate the regression equations for the excretion and the retention coefficient were originated from six replicates (three for each sex). The increase in the phytase level in the diet did not affect (p>0.05) intake, but linearly reduced

Table 3. Contents of phytate and phytate phosphorus (P_{phy}) in excreta from male and female broilers, determined according to two analytical methodologies and extraction ratios

Content (g/100 g of dry matter)	Calculated nutritional composition (g/kg of natural matter)				PC diet ²
	0	750	1,500	2,250	
HPLC-IR ^{3,4} 3:20 (m:v)					
Total myo-inositol phosphates ⁵					
Male	3.840	3.994	2.750	2.719	3.784
Female	4.527	2.866	2.212	2.505	4.480
Phytate phosphorus ⁶					
Male	0.856	0.888	0.614	0.608	0.844
Female	1.010	0.640	0.494	0.560	1.000
P of IP ₃					
Male	0.799	0.849	0.567	0.559	0.782
Female	0.938	0.590	0.458	0.512	0.921
P of IP ₄					
Male	0.036	0.024	0.029	0.030	0.043
Female	0.054	0.033	0.024	0.030	0.059
P of IP ₅					
Male	0.014	0.013	0.014	0.016	0.015
Female	0.016	0.013	0.011	0.015	0.018
P of IP ₆					
Male	0.007	0.002	0.004	0.003	0.004
Female	0.002	0.004	0.001	0.003	0.002
Colorimetry 3:20 (m:v)					
Phytate					
Male	0.244	0.168	0.066	0.090	0.316
Female	0.118	0.076	0.070	0.083	0.344
Phytate phosphorus ⁷					
Male	0.069	0.047	0.019	0.025	0.089
Female	0.033	0.021	0.020	0.023	0.097
Colorimetry 1:20 (m:v)					
Phytate					
Male	1.519	1.456	0.458	0.655	2.425
Female	1.566	0.757	0.809	0.753	3.004
Phytate phosphorus ⁷					
Male	0.428	0.411	0.129	0.185	0.684
Female	0.442	0.213	0.228	0.212	0.847

m:v, mass of sample in g:volume of extractor solvent in mL.

IP_x, myo-inositol phosphates, wherein x corresponds to the degree of phosphorylation of the molecule.

¹ Negative control (NC) diet, without phytase, formulated with 2.13 g of available phosphorus/kg of diet.

² Positive control (PC) diet, without phytase, formulated with 4.26 g of available phosphorus/kg of diet.

³ HPLC-IR, high-performance liquid chromatography with refractive index detector.

⁴ At the extraction ratio of 1:20 (m:v), the concentration of myo-inositol phosphates in the aliquot injected in the chromatograph was below of the limit of detection and/or quantification.

⁵ Corresponding to the sum of fractions IP₃, IP₄, IP₅, and IP₆.

⁶ Phytate phosphorus (%) = (% of IP₃×0.221)+(%) of IP₄×0.248)+(%) of IP₅×0.267)+(%) of IP₆×0.282).

⁷ Phytate phosphorus (%) = determined phytate content (%)×0.282.

($p < 0.01$) the P_{phy} excretion up to 57.75%, which represents good results, as it demonstrates the possibility of producing broilers with less P deposition in the environment using phytase in the diet.

In the diets without supplemented phytase, there was good use of the P_{phy} present in the diet, which was similar to the observations by Han et al. (2009), who attributed this

result to the action of the endogenous phytase that can be synthesized by cells of the intestinal epithelium of broilers (Applegate et al., 2003). Besides, the decrease of P_{av} from 4.26 to 2.13 g/kg of the diet increased the coefficient of hydrolysis of the P_{phy} from 43.37% to 69.67%, which was similar to the results reported by Onyango et al. (2006), who evaluated the effect of two levels of P in the diet

Table 4. Balance (g/bird) and retention (%) of phytate phosphorus (in dry matter) of male and female broilers fed diets deficient in available phosphorus supplemented with increasing levels of phytase, utilizing two analytical methodologies and extraction ratios for the determination of the content of phytate phosphorus in the diets and excreta

	NC diet ¹ supplemented with phytase (FTU/kg)				PC diet ²	RMSE	Significance (p-value)		
	0	750	1,500	2,250			Diet	Sex	Diet×sex
HPLC-IR ^{3,4} 3:20 (m:v)									
Intake ⁵									
Male ^a	0.180*	0.191	0.190	0.180*	0.193	0.009	0.0098	<0.0001	0.9705
Female ^b	0.159*	0.172	0.168	0.157*	0.176				
Excretion									
Male ⁶	0.806	0.868 ^a	0.609 ^{a,*}	0.550*	0.878	0.104	<0.0001	0.0101	0.0481
Female ⁷	0.785	0.568 ^{b,*}	0.425 ^{b,*}	0.440*	0.952				
Retention ⁸	-370.8	-292.5*	-186.3*	196.4*	-396.7	64.646	<0.0001	0.4757	0.0706
Colorimetry 3:20 (m:v)									
Intake ⁹									
Male ^a	0.110	0.117	0.116	0.110	0.116	0.005	0.0270	<0.0001	0.9651
Female ^b	0.098	0.105	0.103	0.096	0.106				
Excretion									
Male ¹⁰	0.045 ^{a,*}	0.046 ^{a,*}	0.019*	0.023*	0.093	0.005	<0.0001	<0.0001	0.0003
Female	0.025 ^{b,*}	0.019 ^{b,*}	0.017*	0.018*	0.092				
Retention									
Male ¹¹	59.05 ^{b,*}	60.38 ^{b,*}	84.13*	79.05*	19.83	4.694	<0.0001	0.0018	0.0002
Female	74.46 ^{a,*}	81.68 ^{a,*}	83.59*	80.59*	12.88				
Colorimetry 1:20 (m:v)									
Intake									
Male ^a	1.394	1.456	1.467	1.398	1.468	0.066	0.0599	<0.0001	0.9640
Female ^b	1.235	1.315	1.299	1.221	1.341				
Excretion ¹²	0.400*	0.293*	0.157*	0.169*	0.794	0.076	<0.0001	0.1570	0.0513
Retention ¹³	69.67*	79.17*	88.45*	86.88*	43.37	5.732	<0.0001	0.9872	0.0646

m:v, mass of sample in g:volume of extractor solvent in ml. FTU, units of phytase activity. RMSE, root mean square error.

¹ Negative control (NC) diet, without phytase, formulated with 2.13 g of available phosphorus/kg of diet.

² Positive control (PC) diet, without phytase, formulated with 4.26 g of available phosphorus/kg of diet.

³ HPLC-IR, high-performance liquid chromatography with refractive index detector.

⁴ At the extraction ratio of 1:20 (m:v), the concentration of myo-inositol phosphates in the aliquot injected in the chromatograph was below of the limit of detection and/or quantification.

⁵ Phytate phosphorus intake = $(-1.00701 \times 10^{-8})x^2 + 0.0000218x + 0.170$ ($R^2 = 0.99$).

⁶ Phytate phosphorus excretion by male broilers = $-0.000137x + 0.863$ ($R^2 = 0.75$).

⁷ Phytate phosphorus excretion by female broilers = $-0.000157x + 0.731$ ($R^2 = 0.83$).

⁸ Phytate phosphorus retention = $0.0840x - 355.889$ ($R^2 = 0.87$).

⁹ Phytate phosphorus intake = $(6.17 \times 10^{-9})x^2 + (1.33 \times 10^{-5})x + 0.104$ ($R^2 = 0.99$).

¹⁰ Phytate phosphorus excretion by male broilers = $-0.0000125x + 0.047$ ($R^2 = 0.68$).

¹¹ Phytate phosphorus retention by male broilers = $0.0112x + 58.089$ ($R^2 = 0.71$).

¹² Phytate phosphorus excretion = $-0.000110x + 0.379$ ($R^2 = 0.87$).

¹³ Phytate phosphorus retention = $-0.00000492x^2 + 0.0120x + 69.14$ ($R^2 = 0.97$).

* Differs from the positive control diet by the Dunnett test ($p < 0.05$).

^{a,b} Means in a column with different superscripts letters differ by the F test ($p < 0.05$).

(without phytase supplementation) on the activity of phytase present in the intestinal mucosa (duodenum) of broilers, in the period from 8 to 20 days of age, and observed that the activity of this enzyme rose from 29.46 to 43.25 nmols of $_{inorg}P$ /mg of protein per min when the concentration of total P in the diet was reduced from 7.0 to 3.6 g/kg of diet. These results suggest that there is an influence of dietary P levels on the activity and efficacy of intestinal phytase, although the mechanisms are not yet fully understood (Bedford and Partridge, 2010). One

possible biochemical explication for these results is based in the mechanism of enzymatic inhibition in the presence of the product from catalysis of manner that the increase in the P_{inorg} concentration in the diet causes a down-regulation in the activity of the intestinal phytase (Onyango et al., 2006).

The increase in the inclusion of phytase in the diet increased ($p < 0.01$) the retention of P_{phy} , corroborating with the research of Han et al. (2009), who also verified increasing improvement in the P_{phy} retention in function of increase in the phytase level in the diet. The greatest

retention coefficient was estimated at 87.85% when the diet contained 1,950 FTU/kg of diet (Coefficient of retention of the $P_{phy} = -0.00000492x^2 + 0.01919x + 69.14$; $R^2 = 0.97$). Thus, as the mean content of P_{phy} determined in the diet was 3.25 g P_{phy} /kg of the diet in natural matter (considering the mean determined value of 3.78 g P_{phy} /kg of diet in DM and the content of moisture of $13.88\% \pm 0.45$), it is possible to reduce the P content in the diet in 87.85% (or 2.86 g of P per kg) by adding 1,950 FTU/kg of the diet.

IMPLICATIONS

The extraction ratio and methodology of dosage utilized to determine the P_{phy} content in both the diets and excreta from broilers interfere directly with the results of P_{phy} retention. The combination of the extraction ratio of 1:20 (mass of sample in g:volume of extractor solvent in mL) and colorimetric methodology is the one that yielded the best results. The broiler sex did not affect the P_{phy} utilization, in the period from 15 to 28 days of age. Supplementation of 1,950 FTU/kg corresponds to a reduction of 2.86 g in P/kg of the diet.

ACKNOWLEDGMENTS

The work was supported by the CNPq (National Council for Scientific and Technological Development) and by the FAPEMIG (Foundation to Support Research of the State of Minas Gerais), Brazil.

REFERENCES

- Angel, R., N. M. Tamim, T. J. Applegate, A. S. Dhandu, and L. E. Ellestad. 2002. Phytic acid chemistry: Influence on phytin-phosphorus availability and phytase efficacy. *J. Appl. Poult. Res.* 11:471-480.
- AOAC. 2005. Official Methods of Analysis. 18th edn. Association of Official Analytical Chemists, Gaithersburg, USA.
- Applegate, T. J., R. Angel, and H. L. Classen. 2003. Effect of dietary calcium, 25-hydroxycholecalciferol, or bird strain on small intestinal phytase activity in broiler chickens. *Poult. Sci.* 82:1140-1148.
- Bedford, M. R. and G. G. Partridge. 2010. *Enzymes and Farm Animal Nutrition*. 2nd edn. CABI, Cambridge, England.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72:248-254.
- Brito, J. A. 2008. Vitamin D₃ (cholecalciferol) and 25-hydroxycholecalciferol (25-OHD₃) in Broiler Diets. Ph.D. Thesis. Federal University of Lavras, Lavras, Minas Gerais, Brazil.
- Cúneo, F., J. Amaya-Farfan, and F. Carraro. 2000. Phytate distribution in stabilized rice bran treated with exogenous phytase. *Ciênc. Tecnol. Aliment.* 20:94-98.
- Engelen, A. J., F. C. Van der Heeft, P. H. G. Randsdorp, and E. L. C. Smit. 1994. Simple and rapid determination of phytase activity. *J. AOAC Int.* 77:760-764.
- Frühbeck, G., R. Alonso, F. Marzo, and S. Santidrián. 1995. A modified method for the indirect quantitative analysis of phytate in foodstuffs. *Anal. Biochem.* 225:206-212.
- Gomide, E. M., P. B. Rodrigues, L. de P. Naves, V. M. P. Bernardino, L. M. dos Santos, and A. A. P. Garcia. 2012. Diets with reduced levels of nutrients supplemented with phytase and amino acids for broilers. *Ciênc. agrotec.* 36:100-107.
- Greiner, R., A. Farouk, M. L. Alminger, and N. G. Carlsson. 2002. The pathway of dephosphorylation of myo-inositol hexakisphosphate by phytate-degrading enzymes of different *Bacillus* spp. *Can. J. Microbiol.* 48:986-994.
- Han, J. C., X. D. Yang, H. X. Qu, M. Xu, T. Zhang, W. L. Li, J. H. Yao, Y. R. Liu, B. J. Shi, J. F. Zhou, and X. Y. Feng. 2009. Evaluation of equivalency values of microbial phytase to inorganic phosphorus in 22- to 42-day-old broilers. *J. Appl. Poult. Res.* 18:707-715.
- Kornegay, E. T. 2001. Digestion of phosphorus and other nutrients: the role of phytases and factors influencing their activity. In: *Enzymes in Farm Animal Nutrition* (Ed. M. R. Bedford and G. G. Partridge). Finnfed International, Marlborough, Wiltshire, UK. p. 237-271.
- Latta, M. and M. Eskin. 1980. A simple and rapid colorimetric method for phytate determination. *J. Agric. Food Chem.* 28:1313-1315.
- Nagata, A. K., P. B. Rodrigues, R. R. Alvarenga, M. G. Zangerônimo, K. F. Rodrigues, and G. F. R. Lima. 2011. Energy and protein levels in diets containing phytase for broilers from 22 to 42 days of age: performance and nutrient excretion. *R. Bras. Zootec.* 40:1718-1724.
- Nappi, G. U., M. R. Ribeiro-Cunha, J. V. Coelho, and L. Jokl. 2006. Validation methods to determine phytic and oxalic acids in "multimisturas". *Ciênc. Tecnol. Aliment.* 26:811-820.
- Onyango, E. M., E. K. Asem, and O. Adeola. 2006. Dietary cholecalciferol and phosphorus influence intestinal mucosa phytase activity in broilers chicks. *Br. Poult. Sci.* 47:632-639.
- Rodrigues, P. B., R. S. Martinez, R. T. F. Freitas, A. G. Bertechini, and E. T. Fialho. 2005. Effect of collection time and methodologies on the digestibility and energy value of poultry diets. *R. Bras. Zootec.* 34:882-889.
- Rostagno, H. S., L. F. T. Albino, J. L. Donzele, P. C. Gomes, R. F. de Oliveira, D. C. Lopes, A. S. Ferreira, and S. L. T. Barreto. 2005. *Brazilian Table for Poultry and Pigs: Food Composition and Nutritional Requirements*. 2ed. Federal University of Viçosa, Minas Gerais, Brazil.
- Rostagno, H. S., L. F. T. Albino, J. L. Donzele, P. C. Gomes, R. F. de Oliveira, D. C. Lopes, A. S. Ferreira, S. L. T. Barreto, and R. F. Euclides. 2011. *Brazilian Table for Poultry and Pigs: Food Composition and Nutritional Requirements*. 3ed. Federal University of Viçosa, Minas Gerais, Brazil.
- Sandberg, A. S. and R. Ahderinne. 1986. HPLC method for determination of inositol tri-, tetra-, penta and hexaphosphates in foods and intestinal contents. *J. Food Sci.* 51:547-550.
- Santos, L. M. dos, P. B. Rodrigues, R. R. Alvarenga, L. de P. Naves, R. Hespanhol, G. F. R. Lima, M. C. C. Lara, and L. R. Silva. 2011a. Calcium and available phosphorus levels in diets supplemented with phytase for broilers in the growing and finishing phases. *R. Bras. Zootec.* 40:2486-2495.

- Santos, L. M. dos, P. B. Rodrigues, R. T. F. Freitas, A. G. Bertechini, E. T. Fialho, E. M. Gomide, and L. de P. Naves. 2011b. Calcium and available phosphorus levels in diets with phytase for broilers in the pre-starter and starter phases. *R. Bras. Zootec.* 40:2476-2485.
- SAS Institute Inc. 2004. Statistical Analysis System User's Guide: Version 9.1. SAS Institute Inc., Cary, NC, USA.
- Yu, S., A. Cowieson, C. Gilbert, P. Plumstead, and S. Dalsgaard. 2012. Interactions of phytate and myo-inositol phosphate esters (IP₁₋₅) including IP₅ isomers with dietary protein and iron and inhibition of pepsin. *J. Anim. Sci.* 90:1824-1832.
- Wadt, G., T. T. Santos, G. A. Gomes, A. Cowieson, and M. R. Bedford. 2010. Total and phytate phosphorus in corn samples collected from different regions of the Brazil. In: IV Latin American Congress on Animal Nutrition, São Pedro, São Paulo, Brazil, pp. 7-8.
- Wu, P., J-C. Tian, C. E. C. Walker, and F-C. Wang. 2009. Determination of phytic acid in cereals – A brief review. *Int. J. Food Sci. Technol.* 44:1671-1676.