

Effects of Different Sizes of Glass Beads on the Release of Sporocysts from *Eimeria tenella* Oocysts

Myung-Jo You*

Laboratory of Veterinary Parasitology, College of Veterinary Medicine and Bio-safety Research Center, Chonbuk National University, Jeonju 561-756, Korea

Abstract: The oocyst wall is severed by means of mechanical injury or chemical agents. This study reports the percentage of in vitro sporocyst release following mechanical shaking in the presence of varying sizes of glass beads. Glass beads measured 0.5, 1, and 3 mm in diameter and were shaken with the oocysts for different times ranging from 5 sec to 5 min. Approximately 80% of sporocysts were released with 5 min of shaking in the presence of 3 mm glass beads, as well as 30 sec with 0.5 mm beads and 1 mm glass beads. The release of sporocysts of *E. tenella* was most efficient using 1 mm glass beads and treatment times of 30 sec to 1 min. Therefore, the use of 1 mm glass beads with 30 sec to 1 min of agitation is recommended in order to maximize sporocyst release and recovery and to improve the yield of viable sporozoites for use in biochemical, tissue culture, and immunological applications of coccidia.

Key words: *Eimeria tenella*, glass bead, oocyst, sporocyst

Avian *Eimeria* spp. are recognized as the parasitic protozoans with the greatest economic impact on poultry production worldwide [1]. Protozoan parasites of the genus *Eimeria* multiply in the intestinal tract and cause tissue damage and increase susceptibility to other diseases [2]. Sporulation of *Eimeria* spp. oocysts is affected by moisture, temperature, and oxygen levels [3]. Oocysts are enclosed in a thick outer shell and consist of a single cell that undergoes the process of sporulation to enter the infective stage after approximately 48 hr. Four sporocysts, each containing 2 sporozoites, are generally discernible within the oocysts. The oocyst wall has 2 distinguishable layers: an inner and outer layers [4]. *In vivo*, the oocyst wall may be opened through mechanical injury, such as being crushed in the gizzard of a bird. A high percentage of sporocysts are quickly released from crushed oocysts within the gizzard [5], and the sporozoites then enter the intestine. Sporozoites of *Eimeria tenella* invade chick cecal folds ~4 hr after infection, followed by sporozoite migration into the crypt epithelium. The first asexual replication occurs 48 hr later [6]. At least 2 generations

of asexual reproduction occur before the sexual phase, where motile microgametes seek out and unite with macrogametes. The zygote matures into oocysts, which are then released from the intestinal mucosa and expelled in the feces. The life cycle of *E. tenella* is typical of all *Eimeria*.

To break the wall of *Eimeria* spp. oocysts for PCR analysis, glass beads, a mortar and a pestle, or chemical agents may be used [7]. *Eimeria* antigens are obtained from suspensions of oocysts in PBS by disruption with glass beads followed by repeated freezing and thawing [8]. The excystation of sporozoites from *Eimeria* oocysts is an important step for in vitro studies of these parasites. The aim of the present study was to assess the degree of released sporocysts of *E. tenella* oocysts using different sizes of glass beads and treatment durations.

Oocysts were field-isolated samples (Namwon Farm, Namwon, Korea). The correct species assignment and purity of samples were confirmed by singular ITS-PCR. Genes of *E. tenella* were collected using a Nucleic Acid Purification Kit (Exprep™ Plasmid DNA, GeneAll, Seoul, Korea). Genes were sequenced at the Jeonju Biomaterials Institute and submitted to GenBank (accession no. FJ447468). Since 2009, our laboratory has maintained generations of *E. tenella* on chickens. Feces were collected from 5 to 7 days post inoculation. Oocysts were suspended at $\sim 3.0 \times 10^5$ per milliliter in 2% (w/v) potassium dichromate in a conical flask. Oocysts were sporulated under

•Received 15 October 2013, revised 10 March 2014, accepted 4 April 2014.

*Corresponding author (tick@chonbuk.ac.kr)

© 2014, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

air supply within 5-8 days [9] and were checked for complete sporulation. A 200- μ l sample of 3.0×10^5 oocysts per milliliter was added to 800 μ l of saturated salt solution. The oocysts suspension was then mixed at a ratio of 1:1 with glass beads (0.5, 1, and 3 mm in diameter) [10] and vortexed for 5 sec, 30 sec, 1 min, or 5 min. For vortexing, Vortex-2 Genie (G 560, Scientific Industries, Springfield, Massachusetts, USA) was used. During vortexing, the conical tube was kept angular using a maximum speed (3,200 rpm). Same vortex speed was fixed in all groups. After oocysts were vortexed and added to 4 ml of saturated salt solution, the remaining oocysts were counted by the standard MacMaster method [11]. The data were subjected to ANOVA with SAS (Ver. 8x). Duncan's multiple range tests was used to determine significant differences among the various durations or bead sizes.

Treatment of the oocytes with 1 mm glass bead (Fig. 1A, B) for 5 sec caused approximately 20% sporocysts release (Fig. 1C), while treatment for 1 min caused the release of more than 90% of sporocysts (Fig. 1D). Release rates of sporocysts using 0.5 mm glass beads ($56.8 \pm 7.3\%$) were higher than those with other bead sizes after 5 sec of treatment ($P < 0.01$). Release rates of sporocysts with 0.5 mm and 1 mm glass beads were not significantly different after 30 sec of treatment, but release rates using 0.5 mm and 1 mm glass beads were higher than 3 mm after 1 min and 5 min of treatment ($P < 0.01$). Change of treatment times was highest at 5 min ($P < 0.01$). For all bead sizes, the release rates of sporocysts increased as the agitation time increased (Fig. 2).




Fig. 1. Effect of glass beading on the excystation of *Eimeria tenella* oocysts. (A) Sporulated oocysts. $\times 1,000$. (B) No treatment of oocysts. $\times 400$. (C) Oocysts were vortexed with 1 mm glass bead for 5 sec. $\times 400$. (D) Oocysts were vortexed with 1 mm glass bead for 1 min. $\times 400$.

During in vitro experiments, oocysts may be made to release 80% of viable sporocysts [16]. There are several methods that can be used to rupture the cell wall of *Eimeria* species; the most common includes sonication, microwave, heating, and pestle and mortar. The greatest advantage of using glass beads for wall rupture is that they are an efficient way to inexpensively process large quantities of oocysts. The concept of mechanical crushing with glass beads came from the in vivo mechanism by which oocysts are crushed in the gizzard [12]. Our study for oocyst disruption of *Eimeria* did not require enzymes. Landers [13] demonstrated that excystation of *E. nieschulzi* could not be achieved using pepsin or trypsin. It was also found that, for release of *E. septata* sporocysts, hydrochloric acid, pepsin, and trypsin were required [14]. Pre-treatment of *E. nieschulzi* oocysts with pepsin destabilizes the oocyst wall structure, ensuring that mechanical disruption by vortexing with glass beads is more efficient [15]. Chai et al. [16] described that the wall structure of *E. tenella* oocysts was ruptured using a teflon-coated tissue homogenizer. The high mechanical resistance of *Eimeria* spp. oocysts required a greater number of glass beads during vortexing. Dulski and Turner [17] described a protocol similar to ours for disrupting *E. tenella* oocysts using 3 mm glass beads. They found that roughly 60% of oocysts were broken by orbital shaking at 200 rpm. Sporocysts have also been released from oocysts by agitating oocyst suspensions with 4 mm glass beads [18]. *E. tenella* oocysts placed in an equal volume of 0.5 mm glass beads and exposed to the maximum speed of agitation resulted in invisible sporocysts [19]. If too many glass beads are used, sporocysts may be damaged; in order to ensure a high yield of undamaged sporocysts, the agita-




Fig. 2. Sporocyst release rate of *Eimeria tenella* oocysts at different glass beads and treatment times. Oocysts were vortexed with 0.5, 1, and 3 mm glass beads for 5 sec, 30 sec, 1 min, and 5 min. Oocysts release of sporocysts is depicted as the mean release of 4 times \pm SD. *Significance ($P < 0.01$).

tion procedure must be stopped after a very specific duration.

After the oocysts wall is crushed, the sporocysts and glass beads are collected separately. The use of smaller glass beads may result in greater difficulty of recovering sporocysts. Our results suggest that sporocyst release from oocysts was efficient after vortexing with 0.5 mm glass beads for 5 to 30 sec, with 1 mm glass beads for 30 sec to 1 min and with 3 mm glass beads for 1 to 5 min. However, the use of 0.5 mm glass beads presented greater difficulties in recovering the sporocysts compared to larger glass beads. Therefore, the present study recommended the use of 1 mm glass beads for 30 sec to 1 min in order to maximize sporocyst release and recovery. Oocyst walls of *Eimeria* spp. have various characteristics. Some *Eimeria* species are more susceptible to chemical agents while others are more susceptible to mechanical injury. Because each *Eimeria* species is unique, with each being lysed upon application of different degrees of either mechanical or chemical stresses, different methods should be tested to improve the yield of viable sporozoites for use in biochemical, tissue culture, and immunological applications of coccidia.

ACKNOWLEDGMENT

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (MEST) (No. KRF-2008-313-E00610).

CONFLICT OF INTEREST

The author has no conflict of interest related to this work.

REFERENCES

- Ruff MD. Important parasites in poultry production systems. *Vet Parasitol* 1999; 84: 337-347.
- Morris GM, Gasser RB. Biotechnological advances in the diagnosis of avian coccidiosis and the analysis of genetic variation in *Eimeria*. *Biotechnol Adv* 2006; 24: 590-603.
- Allen PC, Fetterer RH. Recent advances in biology and immunobiology of *Eimeria* species and in diagnosis and control of infection with these coccidian parasites of poultry. *Clin Microbiol Rev* 2002; 15: 58-65.
- Ferguson DJ, Belli SI, Smith NC, Wallach MG. The development of the macrogamete and oocyst wall in *Eimeria maxima*: immunolight and electron microscopy. *Int J Parasitol* 2003; 33: 1329-1340.
- Doran DJ, Farr MM. Excystation of the poultry coccidium, *Eimeria acervulina*. *J Protozool* 1962; 9: 154-161.
- Daszak P. Zoite migration during infection: parasite adaptation to host defences. *Parasitol Today* 1999; 15: 67-72.
- Kaya G, Dale C, Maudlin I, Morgan K. A novel procedure for total nucleic acid extraction from small numbers of *Eimeria* species oocysts. *Turkiye Parazitoloj Derg* 2007; 31: 180-183.
- Rose ME, Hesketh P, Rennie M. Coccidiosis: rapid depletion of circulating lymphocytes after challenge of immune chickens with parasite antigens. *Infect Immun* 1984; 45: 166-171.
- Waldenstedt L, Elwinger K, Lunden A, Thebo P, and Uggla A. Sporulation of *Eimeria maxima* oocysts in litter with different moisture contents. *Poultry Sci* 2001; 80: 1412-1415.
- Kurth M, Entzeroth R. Improved excystation protocol for *Eimeria nieschulzi* (Apicomplexa, Coccidia). *Parasitol Res* 2008; 102: 819-822.
- Dunn A, Keymer A. Factors affecting the reliability of the McMaster technique. *J Helminthol* 1986; 60: 260-262.
- Jordan FTW, Pattison M. *Poultry Diseases*, 6th ed. London, UK. W.B. Saunders Company Ltd. 1996, p 975.
- Landers EJ. Studies on excystation of coccidial oocysts. *J Parasitol* 1960; 46: 195-200.
- Kowalik S, Zahner H. *Eimeria separata*: method for the excystation of sporozoites. *Parasitol Res* 1999; 85: 496-499.
- Krucken J, Hosse RJ, Mouafo AN, Entzeroth R, Bierbaum S, Marinovski P, Hain K, Greif G, Wunderlich F. Excystation of *Eimeria tenella* sporozoites impaired by antibody recognizing gametocyte/oocyst antigens GAM22 and GAM56. *Eukaryotic Cell* 2008; 7: 202-211.
- Chai JY, Lee SH, Kim UH, Yun CK. Development of *Eimeria tenella* in MDCK cell culture with a note on enhancing effect of preincubation with chicken spleen cells. *Korean J Parasitol* 1989; 27: 87-100.
- Dulski P, Turner M. The purification of sporocysts and sporozoites from *Eimeria tenella* oocysts using percoll density gradients. *Avian Dis* 1988; 32: 235-239.
- Wagenbach GE. Purification of *Eimeria tenella* sporozoites with glass bead columns. *J Parasitol* 1969; 55: 833-838.
- Tomley F. Techniques for isolation and characterization of apical organelles from *Eimeria tenella* sporozoites. *Methods* 1997; 13: 171-176.

