

Perspectives in Pharmacology

Animal Models That Best Reproduce the Clinical Manifestations of Human Intoxication with Organophosphorus Compounds

Edna F. R. Pereira, Yasco Aracava, Louis J. DeTolla, Jr., E. Jeffrey Beecham, G. William Basinger, Jr., Edgar J. Wakayama, and Edson X. Albuquerque

Division of Translational Toxicology, Department of Epidemiology and Public Health (E.F.R.P., Y.A., E.X.A.), and Program of Comparative Medicine and Departments of Pathology, Medicine, and Epidemiology and Public Health (L.J.D.), University of Maryland School of Medicine, Baltimore, Maryland; Countervail Corporation, Charlotte, North Carolina (E.J.B., G.W.B.); and Biomedical Advanced Research and Development Authority and Office of the Assistant Secretary for Preparedness and Response, Department of Health and Human Services, Washington, DC (E.J.W.)

Received March 25, 2014; accepted June 5, 2014

ABSTRACT

The translational capacity of data generated in preclinical toxicological studies is contingent upon several factors, including the appropriateness of the animal model. The primary objectives of this article are: 1) to analyze the natural history of acute and delayed signs and symptoms that develop following an acute exposure of humans to organophosphorus (OP) compounds, with an emphasis on nerve agents; 2) to identify animal models of the clinical manifestations of human exposure to OPs; and 3) to review the mechanisms that contribute to the immediate and delayed OP neurotoxicity. As discussed in this study, clinical manifestations of an acute exposure of humans to OP compounds can be faithfully reproduced in rodents and nonhuman primates. These manifestations include an acute cholinergic crisis in addition to signs of neurotoxicity that develop

long after the OP exposure, particularly chronic neurologic deficits consisting of anxiety-related behavior and cognitive deficits, structural brain damage, and increased slow electroencephalographic frequencies. Because guinea pigs and non-human primates, like humans, have low levels of circulating carboxylesterases—the enzymes that metabolize and inactivate OP compounds—they stand out as appropriate animal models for studies of OP intoxication. These are critical points for the development of safe and effective therapeutic interventions against OP poisoning because approval of such therapies by the Food and Drug Administration is likely to rely on the Animal Efficacy Rule, which allows exclusive use of animal data as evidence of the effectiveness of a drug against pathologic conditions that cannot be ethically or feasibly tested in humans.

The discovery of the usefulness of organophosphorus (OP) compounds as pesticides in the late 1930s unfortunately led to the identification of some of the most toxic compounds synthesized by humankind, including tabun, soman, sarin,

cyclosarin, VR, and VX, which would become collectively known as nerve agents (Tucker, 2006). These nerve agents have been stockpiled and used as weapons of mass destruction in chemical warfare and in terrorist attacks against civilians. The death toll and poor health outcomes of the alleged use of sarin against civilians in Damascus, Syria, as recently as August 2013, and the catastrophic results of the use of tabun or sarin during the Second Sino-Japanese War (1937–1945), the 1980s Iraq-Iran conflict, and the 1990s terrorist attacks in Japan are well documented (Romano and King, 2001; Coupland and Leins, 2005; Tucker, 2006; Dolgin, 2013; Patrick et al., 2013; http://www.un.org/disarmament/content/slideshow/Secretary_General_Report_of_CW_Investigation.pdf).

The fatalities and poor health conditions resulting from acute and/or chronic exposures of humans to OP pesticides have also become a major public health concern. There are estimates that more than 3 million cases of acute OP pesticide

This work was supported in part by funds from the National Institutes of Health National Institute of Neurological Disorders and Stroke CounterACT Program [Grant U01-NS059344]; and by federal funds from the Biomedical Advanced Research and Development Authority, Office of the Assistant Secretary for Preparedness and Response, Office of the Secretary, Department of Health and Human Services, under contract with Countervail Corp. [Contract HHSO100201100030C].

The opinions and assertions contained herein are the private views of the coauthors and do not represent the official position or policies of the Biomedical Advanced Research and Development Authority, Office of the Assistant Secretary for Preparedness and Response, Office of the Secretary, Department of Health and Human Services, the CounterACT program, the National Institute of Neurological Disorders and Stroke, the National Institutes of Health, or the US Government.

dx.doi.org/10.1124/jpet.114.214932

ABBREVIATIONS: ACh, acetylcholine; AChE, acetylcholinesterase; ADC, apparent diffusion coefficient; AMN, atropine methylnitrate; CNS, central nervous system; EEG, electroencephalogram; FDA, Food and Drug Administration; LC₅₀, median lethal concentration; MRI, magnetic resonance imaging; OP, organophosphorus; PTSD, post-traumatic stress disorder; T₂, spin-spin relaxation time; VEP, visual-evoked potential.

poisoning occur per year, with more than 1 million cases being attributed to occupational exposure (Karalliedde and Senanayake, 1989). Just as alarming are the conservative estimates that pesticide self-poisoning worldwide accounts for more than 250,000 deaths per year, which corresponds to about one-third of the world's suicide cases (Gunnell et al., 2007). Finally, epidemiologic studies have provided compelling evidence that acute or continued exposure to subacute levels of OP pesticides is associated with increased risks of debilitating neurologic disorders in both adults and developing children (Levin and Rodnitzky, 1976; Savage et al., 1988; Dahlgren et al., 2004; Rosas and Eskenazi, 2008; Engel et al., 2011).

Acute signs and symptoms of OP poisoning result primarily, although not exclusively, from their common action as irreversible inhibitors of acetylcholinesterase (AChE), the enzyme that hydrolyzes the neurotransmitter acetylcholine (ACh) (Albuquerque et al., 1985). Persistent stimulation of muscarinic receptors by accumulated ACh leads to a muscarinic syndrome characterized by miosis, profuse secretions, bradycardia, bronchoconstriction, hypotension, and diarrhea. Overstimulation of nicotinic receptors triggers tachycardia and skeletal muscle fasciculation, whereas their subsequent desensitization contributes to muscle weakness. The broad range of central nervous system (CNS)-related acute effects includes anxiety, restlessness, confusion, ataxia, tremors, seizures, and central cardiorespiratory paralysis (reviewed in Hurst et al., 2012; see also Yokoyama et al., 1998). Therefore, treatment of acute OP intoxication relies heavily on the use of an oxime to reactivate OP-inhibited AChE, atropine to block the overactivation of muscarinic receptors, and benzodiazepines to reduce the incidence and intensity of OP-induced convulsions and the resulting neuropathology. However, the inadequate outcomes of these treatments have been extensively scrutinized (Buckley et al., 2004). For example, AChE inhibited by some OPs, including the nerve agent soman, ages quickly and cannot be reactivated by clinically available oximes (Kassa, 2002). Some clinical trials have also revealed no clinical benefit of the use of oximes against poisoning with different pesticides, despite the fact that blood AChE was reactivated (Buckley et al., 2011). In addition, as discussed later in this article, recurrent seizures develop even after OP-induced acute seizures are controlled with benzodiazepines (Sekijima et al., 1997). Thus, there is an urgent need to develop new therapeutic strategies to treat and/or prevent the immediate and delayed toxicity of OP compounds.

The public health relevance of OP toxicity to humans worldwide, the need for a comprehensive understanding of the AChE-unrelated mechanisms that contribute to the pathologic conditions triggered by exposure of males and females at different ages to OPs, and the urgency to develop efficacious medical countermeasures to treat and/or prevent these conditions underscore the need to identify optimal animal models in which the clinical manifestations of human OP toxicity can be replicated. In the present article, we provide a comprehensive analysis of the natural history of acute and delayed signs and symptoms that develop following exposure of humans to OP compounds, with an emphasis on nerve agents. This analysis, then, is followed by a discussion of how faithfully the clinical manifestations of human exposure to OP compounds can be reproduced in different animal models of OP intoxication and a brief review of mechanisms that contribute to the immediate and delayed OP neurotoxicity. This information placed in perspective lays the groundwork for future

research aimed at developing safe and effective therapeutic interventions against the acute and delayed toxic effects induced by OP pesticides and nerve agents.

Clinical Manifestations of Human Exposure to OP Compounds: Acute and Long-Term Clinical Signs and Symptoms

Studies of humans who had been experimentally exposed to nerve agents under controlled conditions provided some of the first evidence of the direct effects of nerve agents on neurologic functions in humans. For example, Bowers et al. (1964) analyzed the clinical signs presented by 96 young male volunteers on active duty in the Army or the Air Force following their percutaneous exposure to a low dose of an OP compound likely to be VX. Although the volunteers did not develop overt signs of acute toxicity, they presented with a syndrome that was broadly referred to as a "state of altered awareness" and was characterized by difficulty in sustaining attention and slowing of intellectual and motor processes, in addition to subjective feelings of agitation, anxiety, and confusion (Bowers et al., 1964). In general, these clinical manifestations were associated with the inhibition of AChE in red blood cells, and symptomatic recovery was associated with the recovery of AChE activity (Grob and Harvey, 1953; Bowers et al., 1964). Long-term follow-up of these individuals is not available. However, case reports of humans occupationally or intentionally exposed to nerve agents, follow-up studies of the victims of the 1995 terrorist attack with sarin in the Tokyo subway, and case reports and follow-up studies of humans accidentally or intentionally exposed to OP pesticides, summarized below, all support the contention that persistent delayed neurologic deficits develop following an initial exposure to these chemicals.

Metcalf and Holmes (1969) reported that workers who had a history of exposure to OP compounds presented years later with attention deficits, memory impairment, and difficulty in maintaining alertness that were accompanied by increased slow electroencephalographic activity in the θ range. A subsequent study compared the electroencephalograms (EEGs) recorded from industrial workers years after their confirmed accidental exposure to sarin with those recorded from control subjects (Duffy et al., 1979). In this study, visual inspection of the EEGs revealed that the exposure to sarin was associated with an increase of the slow (δ and θ) frequencies and a reduction of the α frequency. Spectral analysis of the EEGs also revealed a significant increase of the high-frequency activity in the β range (12–30 Hz) years after the exposure to sarin (Duffy et al., 1979).

Follow-up studies of Japanese victims of the sarin attack in Matsumoto also reported that, long after the sarin exposure, victims of the attack presented with significant EEG alterations, some of which were consistent with recurring electrical seizures (Sekijima et al., 1997). One of the victims was a male subject who at the time of the attack presented severe disturbance of consciousness, developed convulsions, and required assisted ventilation. At a 1-year follow-up examination, sharp wave complexes, which are consistent with epileptiform activity (Westmoreland, 1998), were present in the EEG of this victim. Two other victims were women who at the time of the attack presented no convulsions, needed no artificial ventilation, and

had only mild disturbance of consciousness. During the 1-year follow-up examination, bursts of δ activity, which often correlate with focal brain lesions (Harmony et al., 1995), were present in their EEGs. Abnormal bursts of δ activity could still be detected in the EEG of one of the two female victims during a 2-year follow-up examination (Sekijima et al., 1997). During the 2-year follow-up examination, 14-Hz spikes were also detected in the EEG of a fourth victim, a female who at the time of the attack presented no convulsions and needed no respiratory support (Sekijima et al., 1997). Although the authors suggest that these 14-Hz spikes represented epileptic electroencephalographic changes, caution is warranted given that this patient was 16 years old at the time of the examination and 14-Hz spikes frequently appear in the EEG during adolescence (Klass and Westmoreland, 1985). These case reports clearly indicate that patients who do not develop acute convulsions in response to an acute exposure to nerve agents can, years later, present with abnormal slow δ waves that are suggestive of focal brain lesions. They also support the notion that patients who survive a severe case of nerve agent intoxication can years later present with epileptic EEG discharges.

It has been suggested that hypoxia resulting from motor convulsions and respiratory distress during the acute phase of OP intoxication contributes to the neurologic deficits that develop and persist long after the acute phase of intoxication subsides (Nozaki et al., 1995; Hatta et al., 1996). For example, Nozaki et al. (1995) reported the case of a patient who presented with amnesia 15 days and 6 months after recovering from convulsions induced by an exposure to VX. Hatta et al. (1996) reported the case of another patient who presented with retrograde amnesia and personality changes after recovering from clonic-tonic generalized convulsions and severe episodes of dyspnea induced by sarin during the terrorist attack in Tokyo. However, case reports and follow-up studies of people who did not experience overt CNS signs of intoxication following an OP exposure and yet developed persistent delayed neurologic deficits, as described below, support the contention that neurologic impairments can develop following an acute exposure to levels of nerve agents or OP pesticides that are not sufficient to induce convulsions and/or hypoxia.

The 1995 terrorist attack with sarin in the Tokyo subway is the largest documented exposure of a civilian population to a nerve agent. Approximately 95% of the 111 victims who were admitted to hospitals and diagnosed as moderately or severely intoxicated received the conventional therapeutic interventions for OP poisoning, including atropine to block the overactivation of muscarinic receptors and pralidoxime to reactivate sarin-inhibited AChE; diazepam was also used as needed to treat victims who developed convulsions (Okumura et al., 1996). Even though the treatments effectively countered the acute signs of toxicity, victims of the sarin attack who did not present with episodes of convulsion at the time of the attack presented years later with significant memory decline (Hood, 2001; Nishiwaki et al., 2001).

Although it is well recognized that post-traumatic stress disorder (PTSD) associated with the experience of a warfare attack can confound the evaluation of long-term health effects induced by the chemical used in the attack, some studies have provided evidence that PTSD alone cannot explain changes in neurophysiologic functions and disruption of the structural integrity of specific brain regions observed in victims who had been exposed to sarin in the Tokyo subway attack (Murata

et al., 1997; Yamasue et al., 2007). For example, 6–8 months after the sarin attack, event-related and visual-evoked potentials (P300 and VEP, respectively) were analyzed in 18 victims of the attack and 18 control subjects (Murata et al., 1997). P300 is an event-related potential associated with decision making, and VEP is an evoked potential that reflects the conduction time from the retina to the visual cortex. In victims of the attack, the latencies of both P300 and VEP were significantly longer than those measured in control subjects. The finding that the longer latencies of those potentials did not correlate with the victims' high PTSD scores strongly indicated that sarin, rather than PTSD, was the cause of the neurophysiologic alterations measured in those patients (Murata et al., 1997). Likewise, Nishiwaki et al. (2001) reported that first responders who were dispatched to the site of the attack in Tokyo presented 3 years later with memory deficits that were evident in the backward digit span test and were independent of PTSD symptoms. Finally, Yamasue et al. (2007) reported that 5 years after the attack victims presented with significant decrease in gray matter volume in the right insular cortex, the right temporal cortex, and the left hippocampus. In these victims, the volume of the left subinsular white matter was positively correlated with decreased serum cholinesterase levels measured after the incident, but not with the occurrence or severity of PTSD.

A case report published in 2010 corroborated the notion that exposure of humans to low levels of sarin can trigger delayed neurocognitive deficits (Loh et al., 2010). This is the report of a US Army sergeant who 8 months after experiencing mild signs of acute toxicity following an exposure to sarin in Iraq presented with attention deficits and impaired motor coordination in the absence of any signs or symptoms of emotional distress or mood disorders.

Persisting neurologic symptoms of memory loss, decreased concentration, irritability, and personality changes have also been observed in all members of a family who were accidentally exposed to the OP pesticide diazinon (Dahlgren et al., 2004). A pesticide company mistakenly sprayed the interior of the family's house with diazinon, and, soon after the exposure, all members of the family presented acute signs and symptoms of OP intoxication, including headaches, nausea, skin irritation, runny nose, and vomiting. They did not present with convulsion. Three months to 3 years later, all family members presented with cognitive deficits and mood disorders. In these cases, the persistent neurologic dysfunctions following the exposure to the OP pesticide clearly developed in the absence of acute convulsions and hypoxia. Population studies have provided additional evidence that signs of emotional distress and memory deficits develop in humans long after their recovery from acute poisoning by cholinesterase inhibitors or from a hypercholinergic syndrome following acute occupational exposure to OP pesticides (Levin and Rodnitzky, 1976; Savage et al., 1988; Rosenstock et al., 1991; Steenland et al., 1994; Wesseling et al., 2002; Roldán-Tapia et al., 2005).

In summary, a cholinergic crisis defines the acute phase of OP intoxication in humans, even though the prevalence of specific signs and symptoms is OP specific (e.g., Nozaki et al., 1995). In addition, although it is true that subtle differences in the prevalence of delayed signs and symptoms of OP neurotoxicity also exist among individuals exposed to different OP pesticides and nerve agents, it appears that mood disorders, specifically anxiety and depression, and cognitive

deficits, particularly attention deficits and memory impairment, are common persistent neurologic conditions seen in individuals long after their exposure to these chemicals (Metcalf and Holmes, 1969; Levin and Rodnitzky, 1976; Savage et al., 1988; Rosenstock et al., 1991; Steenland et al., 1994; Nozaki et al., 1995; Hatta et al., 1996; Hood, 2001; Nishiwaki et al., 2001; Wesseling et al., 2002; Dahlgren et al., 2004; Roldán-Tapia et al., 2005). In many cases, these delayed neurologic deficits have been observed in the absence of acute convulsions and/or hypoxia induced by the OP compounds. In some cases, these deficits have also been dissociated from the trauma associated with the exposure event and have been correlated with the following: 1) disruption of the structural integrity of different brain structures, including the hippocampus and the cingulate cortex, as seen in magnetic resonance imaging (MRI) studies of victims of the Tokyo subway terrorist attack with sarin (Yamasue et al., 2007); 2) decreased regional cerebral blood flow, particularly in the occipital lobes, as seen in a single-photon emission computed tomography study of subjects long after they had experienced an acute intoxication with OP pesticides (Mittal et al., 2011); and 3) increased slow electroencephalographic activity, as reported in individuals occupationally exposed to sarin (Metcalf and Holmes, 1969). Therefore, at a minimum, translationally relevant animal models should present acute and delayed signs and symptoms of OP toxicity that recapitulate those described in this work and should also predict the effectiveness and safety of medical countermeasures against OP toxicity in humans.

Animal Models of OP Intoxication

Signs of acute toxicity induced by OP pesticides and nerve agents in numerous animal species ranging from small rodents such as mice, rats, and guinea pigs to large mammals, including nonhuman primates, closely resemble those presented by humans exposed to these compounds. Following exposure of different animal species to high doses of nerve agents, a cholinergic crisis develops and is characterized by chewing and gnawing, profuse secretions, diarrhea, muscle fasciculation, restlessness, tremors, motor convulsions, and respiratory distress that can lead to death (Deshpande et al., 1986; Kawabuchi et al., 1989; Albuquerque et al., 2006; Maxwell et al., 2006; Despaigne et al., 2007). The rank order of lethal potencies (LD_{50}) of the nerve agents is invariably the same among the different species, with the LD_{50} increasing from VX to soman to sarin (Bajgar, 1992; Koplovitz and Stewart, 1994; Lenz et al., 2005; Maxwell et al., 2006; Fawcett et al., 2009). In addition, the LD_{50} of different nerve agents correlate well with their IC_{50} to inhibit *in vitro* AChE activity in brain extracts from rats and guinea pigs (Sivam et al., 1984; Fawcett et al., 2009), supporting the contention that, as in humans, AChE inhibition is an important determinant of the acute toxicity of nerve agents in animal models of acute OP intoxication.

Studies from various laboratories further confirmed that, similar to humans (Nozaki et al., 1995), different animal species present signs of acute toxicity that are nerve agent-dependent (Shih and McDonough, 1999; Shih et al., 2003; Fawcett et al., 2009; O'Donnell et al., 2011). For example, guinea pigs challenged with $0.8 \times LD_{50}$ VX, soman, or sarin present facial twitches, chewing, slight hyperlocomotion, and head tremors (Fawcett et al., 2009). With increasing doses of soman or sarin,

guinea pigs present progressively with forelimb clonus, increased secretions, muscle fasciculation, rearing, strong grinding, gnashing or clenching of the teeth (bruxism), unremitting tonic-clonic convulsions, and severe respiratory distress. In contrast, even at a dose as high as $1.2 \times LD_{50}$, VX does not induce recurrent or unremitting motor convulsions in guinea pigs. Motor convulsions induced by $1.2 \times LD_{50}$ VX are brief and show spontaneous termination (Fawcett et al., 2009). In earlier studies, Shih and McDonough (1999) and Shih et al. (2003) also reported that electrical seizures are usually detected in rats and guinea pigs following a subcutaneous injection of $\geq 2 \times LD_{50}$ soman, sarin, tabun, cyclosarin, VX, or VR, with the incidence of these seizures being higher with soman, sarin, tabun, cyclosarin, and VR than with VX. They further reported the following: 1) onset of electrical seizures is slowest with VX than with the other nerve agents, and 2) doses of anticonvulsants needed to control seizures induced by VX or sarin are significantly lower than those needed to control seizures induced by the other nerve agents.

A recent study conducted with guinea pigs treated with atropine methylnitrate (AMN) prior to their challenge with $1.0 \times LD_{50}$ sarin or VX further revealed that only among sarin-exposed animals was lethality proportional to the percentage of animals that developed electrical seizures (O'Donnell et al., 2011). In that study, approximately 70% of all guinea pigs injected with $1.0 \times LD_{50}$ sarin developed electrical seizures. Nearly 50% of sarin-challenged animals that developed electrical seizures did not survive the first 24-hour post-challenge, whereas all animals that did not develop seizures survived the sarin challenge (O'Donnell et al., 2011). In contrast, approximately 40% of the AMN-treated animals developed seizures when challenged with $1.0 \times LD_{50}$ VX, but all guinea pigs survived the VX challenge regardless of whether they developed seizures (O'Donnell et al., 2011). Because unremitting convulsions induced by soman and sarin contribute to their lethal potency, timely treatment of soman- or sarin-challenged animals with benzodiazepines significantly improves their survival and increases the effectiveness of the standard therapy consisting of the muscarinic antagonist atropine and such AChE reactivators as pralidoxime in reducing mortality induced by these nerve agents (Lallement et al., 1997, 1998, 1999; McDonough et al., 1999, 2000). This is in agreement with the well reported increased risk of death associated with status epilepticus in humans (Ferlisi and Hocker, 2013; Jette and Trevathan, 2014).

A single exposure of different animal species to high levels of soman, sarin, or VX causes profound neuropathology, with significant neuronal loss in various brain regions, including the hippocampus, amygdala, piriform cortex, and thalamus (Lemerrier et al., 1983; McLeod et al., 1984; Petras, 1994; Kadar et al., 1995; Shih et al., 2003). The finding that the severity of the neuropathology induced by the nerve agents correlates well with the duration and severity of convulsions led to the suggestion that early management of nerve agent-induced convulsions would be sufficient to reduce the neuropathology and the accompanying neurologic deficits (Martin et al., 1985; Hayward et al., 1990; Lallement et al., 1993, 1994, 1997, 1998; McDonough and Shih, 1997; McDonough et al., 2000; Raveh et al., 2002; Shih et al., 2003). However, spontaneous recurrent motor convulsions and electrical seizures in addition to neurodegeneration have been observed in rats long after acute seizures following an exposure to nerve

agents were terminated by benzodiazepines (de Araujo Furtado et al., 2010). Spontaneous recurrent motor convulsions and/or electrical seizures have also been reported in soman-exposed rats and nonhuman primates long after the acute toxicity of the nerve agent subsided (Romano et al., 2001; Despain et al., 2007). Therefore, treatment with anticonvulsants, although essential to control the acute tonic-clonic convulsions and improve survivability among animals exposed to high levels of nerve agents, is not sufficient to counter the development of delayed neurologic deficits. In this respect, it is noteworthy that neurodegeneration and the resulting neurologic dysfunctions observed in soman-intoxicated rodents can be significantly reduced by therapeutic interventions that, although unable to control the seizures, effectively decrease glutamate excitotoxicity (Filliat et al., 1999).

Neurologic dysfunctions and structural brain damage that resemble those presented by humans exposed to these agents have been successfully reproduced in rats, guinea pigs, and nonhuman primates exposed to these agents. And, of major significance, in many studies these neurologic deficits have been detected at doses of nerve agents that are not sufficient to induce overt signs of acute toxicity in the animals. This is a crucial point given that, as described in the previous section, follow-up clinical studies revealed neurologic deficits in human subjects who were occupationally or intentionally exposed to levels of nerve agents or OP pesticides that were not sufficient to trigger overt signs of acute toxicity.

Anxiety-related behavior, a clinical manifestation of OP exposure in humans, develops and lasts long after a single exposure of rats and guinea pigs to sarin or soman. Several studies revealed that rats and guinea pigs presented with anxiety-related behavior when tested in the open field and in the elevated T-maze days to months after an acute exposure to a low dose of soman (Sirkka et al., 1990; Baille et al., 2001; Mamczarz et al., 2010). In addition, Bajgar et al. (2004) reported that, 1 month after the inhalational exposure to $0.7 \times \text{LCt}_{50}$ sarin (i.e., 70% of the median lethal concentration), female guinea pigs presented increased locomotor activity and exacerbated stereotypic behaviors that are suggestive of anxiety (Choi et al., 2013).

Cognitive deficits have also been observed in animals exposed to nerve agents. One study reported that, following a single inhalation exposure to a level of sarin sufficiently low to trigger only mild or no signs of cholinergic hyperstimulation, rats showed significant impairment in spatial discrimination in the Y-maze (Kassa et al., 2002). Other studies reported that rats and mice that developed severe signs of acute toxicity, including convulsions, following a single subcutaneous exposure to $1.0\text{--}1.2 \times \text{LD}_{50}$ soman, showed immediate and delayed cognitive impairments in the Morris water maze (Filliat et al., 1999; Raveh et al., 2002). Of particular interest is a report that in asymptomatic, soman-challenged mice cognitive deficits could be detected at 3 months, but not 1 month following the challenge (Filliat et al., 2007). Guinea pigs, too, develop cognitive deficits long after their exposure to soman (Mamczarz et al., 2011; Pereira et al., 2012). Three months after surviving the challenge with $1.0 \times \text{LD}_{50}$ soman, prepubertal female guinea pigs present spatial learning impairment in the Morris water maze (Mamczarz et al., 2011) and adult male guinea pigs present both spatial learning and memory retention deficits (Pereira et al., 2012).

An *in vivo* MRI study of female guinea pigs that survived the challenge with $1.0 \times \text{LD}_{50}$ soman demonstrated significant and long-lasting damage to the structural integrity of brain regions known to play a central role in cognitive processing and in control of emotional behaviors (Gullapalli et al., 2010). Segmentation of the brain images obtained from all soman-exposed guinea pigs before and 7 days after the injection of the nerve agent revealed a significant increase in ventricular cerebrospinal fluid that was suggestive of brain atrophy. In animals that developed severe signs of intoxication in response to $1.0 \times \text{LD}_{50}$ soman, there was a significant, long-lasting increase of spin-spin relaxation time (T₂) values in the piriform cortex, amygdala, and thalamus in addition to a transient increase of T₂ values in the hippocampus. A significant amount of cell lysis may have accounted for the regional increase in T₂ values, given that loss of neurogranin mRNA signal (a neuronal marker) and large numbers of Fluoro-Jade-B-positive neurons were observed in the brains of the animals that were severely intoxicated with soman.

Using diffusion-weighted MRI to assess brain damage in mice exposed to a convulsive dose of soman ($172 \mu\text{g}/\text{kg}$), Testylier et al. (1999) demonstrated that 3 hours following the soman exposure there were changes in apparent diffusion coefficient (ADC) that were also compatible with edema in different brain regions. In agreement with the findings obtained from soman-exposed guinea pigs, the MRI and histopathological studies carried out in mice exposed to soman led to the conclusion that the cerebral edema induced by soman is primarily intracellular and cytotoxic.

Bhagat et al. (2001; 2005) used T₂-weighted imaging and diffusion-weighted imaging to evaluate the evolution of brain pathology following exposure of adult male Sprague-Dawley rats to a convulsive dose of soman ($180\text{--}200 \mu\text{g}/\text{kg}$; $1.8\text{--}2.0 \times \text{LD}_{50}$). The authors observed a significant 23% decrease in ADC at 12 hours after soman exposure in the hippocampus and thalamus. In these rats, ADC returned to normal values in 7 days. They also observed at 12 hours after soman exposure a significant decrease in T₂ values, which returned to near normal values at 24 hours. Although these studies lasted no longer than 7 days, the acute ADC changes seen in the hippocampus and thalamus of rats that survived the exposure to soman were compatible with those seen in the follow-up studies of the victims of the Tokyo terrorist attack with sarin and were suggestive of disruption of the integrity of white matter.

Studies from numerous laboratories have also shown that the relative contribution of δ waves to the EEG power spectrum increases significantly following an exposure of rats, mice, guinea pigs, and nonhuman primates to convulsing doses of soman (McDonough et al., 1998; Carpentier et al., 2001). This increase in the slow δ EEG frequency resembles that described in the previous section for humans exposed to sarin and other OP compounds and appears to be a predicting factor of the neuropathology that eventually develops in the brain of these animals (McDonough et al., 1998; Carpentier et al., 2001). Also, as reported for workers who had been occupationally exposed to sarin (Duffy et al., 1979), an increase in β activity was detected in the EEG of nonhuman primates 1 year after their exposure to sarin (Burchfiel et al., 1976; Burchfiel and Duffy, 1982).

In summary, it is clear that signs of acute OP intoxication in rats, mice, guinea pigs, and nonhuman primates closely resemble the clinical manifestations of acute OP intoxication in humans. Evidence also exists that delayed neurologic deficits,

particularly anxiety-related behavior and cognitive deficits, structural brain damage, and functional EEG alterations (specifically an increase of the slow EEG frequencies), seen in humans long after a single exposure to nerve agents can be reproduced in rats, mice, guinea pigs, and nonhuman primates exposed to these agents. The question, then, arises as to whether any one animal model is better than the others.

Undoubtedly, nonhuman primates, particularly macaques, hold a number of advantages over small mammals as animal models in biomedical research. The genetic, anatomic, physiologic, behavioral, and metabolic similarities between humans and nonhuman primates make the nonhuman primates the most appropriate animal model of a variety of human conditions, including OP intoxication (Lallement et al., 1998; Lenz et al., 2005; Anderson, 2008; Sun et al., 2008). It is particularly noteworthy that, because the anatomy and physiology of the nonhuman primate brain are very close to those of humans, neurobehavioral tasks normally used to assess cognitive functions in humans (e.g., delayed-matching-to-sample task) have been successfully adapted for nonhuman primates, and thereby increased the translational relevance of nonhuman primates in research (Buccafusco, 2008). Specifically as it relates to the field of OP toxicity, the sensitivity of nonhuman primates to different OPs closely resembles that of humans. For example, the intravenous LD₅₀ of sarin in Rhesus monkeys and humans are 15 and 14 $\mu\text{g}/\text{kg}$, respectively (Woodard et al., 1994; <http://www.iom.edu/~media/Files/Report%20Files/2007/Long-Term-Health-Effects-of-Participation-in-Project-SHAD-Shipboard-Hazard-and-Defense/SARINNERVEAGENT.pdf>). However, taking into consideration the principles of humane experimental research, less sentient species are needed, particularly, for proof-of-concept studies. In this respect, guinea pigs have stood out as an appropriate small animal model of OP intoxication. Specifically, the sensitivity of guinea pigs to OP compounds closely resembles that of humans. For example, the intravenous LD₅₀ of sarin in guinea pigs (24 $\mu\text{g}/\text{kg}$) is only 1.7-fold higher than that reported for humans (Spruit et al., 2000; <http://www.iom.edu/~media/Files/Report%20Files/2007/Long-Term-Health-Effects-of-Participation-in-Project-SHAD-Shipboard-Hazard-and-Defense/SARINNERVEAGENT.pdf>). Furthermore, many anatomic features of the human brain are closely related to those in the guinea pig brain, making the guinea pigs a robust small, nonprimate model in which to study multidimensional neurobehaviors of direct relevance to humans (see Lee et al., 2014, and references therein).

Most studies of the acute and delayed toxicity induced by an exposure to different levels of OP compounds have been carried out in rats and mice. The relatively small size, short generation time, and accelerated life span of rats and mice help to keep space and time required to perform a study manageable. In addition, most behavioral tests have been developed and validated in rats and mice; antibodies against thousands of proteins are commercially available for both species; and manipulation of mouse and rat DNA has proven to be feasible, allowing scientists to explore how specific genes function in health and disease (Boverhof et al., 2011; see also Lee et al., 2014, and references therein). However, the sensitivity of rats and mice to the acute toxicity of OP compounds is reportedly lower than that of humans. For example, the intravenous LD₅₀ of sarin in rats and mice (45–63 and 83 $\mu\text{g}/\text{kg}$, respectively) are approximately 3- to

6-fold higher than the intravenous LD₅₀ of sarin in humans (14 $\mu\text{g}/\text{kg}$; <http://www.iom.edu/~media/Files/Report%20Files/2007/Long-Term-Health-Effects-of-Participation-in-Project-SHAD-Shipboard-Hazard-and-Defense/SARINNERVEAGENT.pdf>; John et al., 2009). The species-specific sensitivity to the toxic effects of OP compounds has been correlated with the activity of circulating carboxylesterases—the liver enzymes that catalyze the metabolism and inactivation of OP compounds. In fact, Maxwell et al. (1987) reported that inhibition of carboxylesterases by cresylbenzodioxaphosphorin oxide eliminates the species-related differences in the LD₅₀ of soman.

The effectiveness of medical therapies against OP intoxication in humans is also species-specific and can be more reliably predicted in guinea pigs and nonhuman primates than in mice and rats. For example, pralidoxime, a reactivator of OP-bound AChE, confers significant protection against soman intoxication in rats and mice, but not in guinea pigs and nonhuman primates (Inns and Leadbeater, 1983). Likewise, some clinical trials have reported the poor effectiveness of pralidoxime to treat acute poisoning of humans with OP compounds (Buckley et al., 2011). Thus, to standardize testing of antidotes (particularly pretreatments) against OP intoxication, initial studies are generally carried out in rodents, including guinea pigs, and results are subsequently confirmed in nonhuman primates (Inns and Leadbeater, 1983; Maxwell et al., 1987).

An ever growing number of epidemiologic studies have provided evidence that in utero exposure to OP pesticides is associated with increased prevalence of attention deficit hyperactive disorder, pervasive developmental disorder, and reduced intelligence quotient among children (reviewed in Rosas and Eskenazi, 2008). A recent MRI study also revealed significant structural abnormalities in the brains of children prenatally exposed to OP pesticide chlorpyrifos (Rauh et al., 2012). Thus, the availability of small animal models of OP-induced developmental toxicity is needed to prioritize and standardize safe and efficient medical therapies to treat even the most sensitive sector of the population in the event of a terrorist attack involving deployment of nerve agents or OP pesticides.

Although prenatal and/or early neonatal exposure of rats and mice to OP pesticides has been shown to compromise their brain development and cause neurocognitive deficits later in life that resemble those seen in humans exposed in utero to these compounds (Levin et al., 2001, 2002; Turgeman et al., 2011), striking differences exist between the CNS development of mice or rats and that of humans, making it difficult to extrapolate sensitive gestational periods from rats and mice to humans. In short-gestation species, such as the rat and mouse, the majority of CNS growth spurt occurs postnatally, generally within the first 3 weeks after birth. In long-gestation species, including guinea pigs, nonhuman primates, and humans, this growth spurt starts mid-gestation in the uterus, peaks during the last third part of pregnancy, and ends shortly after birth (Dobbing and Sands, 1970, 1973; Byrnes et al., 2001). Thus, the guinea pig emerges as a more appropriate rodent model system than rats and mice to study the deleterious effects of developmental exposure to OP pesticides and nerve agents.

Molecular Mechanisms That Contribute to the Acute and Delayed Toxicity of OP Compounds

It is generally accepted that irreversible inhibition of AChE is the primary mechanism underlying the acute toxicity of OP

pesticides and nerve agents. However, the fact that the prevalence of the various signs of acute toxicity varies among different OP compounds in humans and in animal models strongly suggests that direct interactions of these compounds with other molecular targets contribute to their acute toxicity. The most direct evidence that AChE is only one of the proteins targeted by the OP compounds came from studies of mice with a null mutation in the gene encoding AChE; injection of the OP compound diisopropylfluorophosphate or VX in these mice triggers a cholinergic syndrome identical to that seen in wild-type mice (Xie et al., 2000; Duysen et al., 2001).

Among the non-AChE molecular targets for OP compounds are the nicotinic and muscarinic receptors (Albuquerque et al., 1985; Silveira et al., 1990; Jett et al., 1991; Bomser and Casida, 2001). Direct interactions of OP compounds with nicotinic receptors at the neuromuscular junction lead to receptor desensitization that can contribute to the muscle weakness and paralysis seen in the acute phase of OP poisoning (reviewed in Albuquerque et al., 1985). In addition, direct interactions of the nerve agents sarin and soman with presynaptic muscarinic receptors in the hippocampus reduce GABAergic transmission onto CA1 pyramidal neurons (Chebabo et al., 1999; Santos et al., 2003), an action that can certainly contribute to the occurrence of seizures in subjects intoxicated with the nerve agents. AChE-independent reduction of neuronal M2 muscarinic receptor function by OP pesticides also potentiates bronchoconstriction induced by stimulation of the vagus in guinea pigs (Lein and Fryer, 2005), an action that can be an important determinant of the respiratory distress induced by these compounds.

The initial irreversible inhibition of AChE by OP compounds can trigger mechanisms that eventually lead to the neurotoxicity that develops either in the absence of an overt acute phase of intoxication or long after the acute phase of OP poisoning has resolved. Accumulated ACh resulting from OP-induced AChE inhibition has been linked to activation of a number of signal transduction mechanisms (Bodjarian et al., 1992; Savolainen and Hirvonen, 1992) that can contribute to time-dependent up- or downregulation of expression of genes that code for proteins involved in sustaining cell viability. In fact, toxicogenomic studies have reported that expression of genes linked to neuroprotective and neurodegenerative pathways is altered time dependently following an injection of rats with either a subconvulsive or a convulsive dose of sarin or soman (Damodaran et al., 2006; RamaRao et al., 2011; Spradling et al., 2011). In this respect, it is noteworthy that the frequency of sister chromatid exchanges was found to be significantly higher in human lymphocytes from victims of the 1995 sarin attack in Japan than in control subjects at two time points: 2–3 months and 3 years after the attack (Li et al., 2004). The increased frequency of sister chromatid exchanges in sarin-exposed humans who did not present with overt signs of acute toxicity is consistent with the notion that the nerve agent per se, in the absence of convulsions and/or hypoxia, can have a significant impact on gene expression.

Direct interactions of OPs with molecular targets other than AChE have also been proposed to play a key role in the development of delayed OP neurotoxicity. For example, OP compounds can decrease neuronal viability by disrupting the axonal transport of nutrients from the cell body to the axon terminals of neurons (Prendergast et al., 2007; Grigoryan et al., 2008). Cellular mechanisms underlying the OP-induced

axonal transport impairment may include direct covalent binding of OP compounds to such structural proteins as tubulin, kinesin, and dynein (reviewed in Terry, 2012; Androustopoulos et al., 2013). In fact, the ability of chlorpyrifos-oxon to inhibit the polymerization of tubulin has been associated with neurodegeneration in the hippocampus (Prendergast et al., 2007). Additional mechanisms that have been proposed to contribute to the delayed toxicity of OP compounds include exacerbated oxidative stress, imbalanced intracellular Ca^{2+} homeostasis, increased signaling mediated by inflammatory mediators, such as interleukins and cytokines, changes in cellular signaling mediated by neurotrophin receptors and protein kinases, and mitochondrial disruption (reviewed in Banks and Lein, 2012; Terry, 2012; Androustopoulos et al., 2013). Elucidating the cause-consequence relationship between the various cellular and molecular mechanisms known to be affected by specific OP compounds and the clinical manifestations induced by an acute, subacute, and/or chronic exposure to these compounds is a critical step toward the design and development of effective therapeutic interventions to treat OP toxicity.

The Food and Drug Administration (FDA) uses the Animal Efficacy Rule to approve medical treatments to reduce and/or prevent serious and/or life-threatening conditions caused by exposure to a permanently disabling toxic agent, such as OP pesticides and nerve agents, in which human efficacy trials are not feasible or ethical. Based on this rule, the FDA can rely exclusively on animal studies to provide substantial evidence of drug effectiveness if the following take place: 1) the mechanisms of action of the toxicant are reasonably well understood; 2) the effects of the toxicants and the effectiveness of the treatments can be demonstrated in animal species expected to react with a response predictive for humans; 3) the animal study endpoints are clearly related to the desired benefits in humans; and 4) data on the pharmacokinetics and pharmacodynamics of the treatments in animals and humans can allow selection of an effective dose for humans. In fact, it was on the basis of the Animal Efficacy Rule that in 2003 the FDA approved pyridostigmine bromide as a pretreatment to prevent the toxicity of soman in military personnel under the threat of exposure to this agent (Aebersold, 2012).

Recognizing the limitations of available therapies against OP poisoning, we need to seek more effective medical countermeasures to treat and/or prevent the acute and delayed neurotoxicity of nerve agents and OP pesticides. Therefore, identification of animal models of the immediate and delayed clinical manifestations of an acute exposure of humans to OP compounds and their mechanisms of action is crucial for studies aimed at the discovery of effective medical countermeasures to treat and/or prevent those conditions.

Acknowledgments

The authors thank Mabel A. Zelle and Robert J. Wesley for assistance in editing the manuscript.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Pereira, Aracava, DeTolla, Beecham, Basinger, Wakayama, Albuquerque.

References

- Aebersold P (2012) FDA experience with medical countermeasures under the animal rule. *Adv Prev Med* **2012**:507571.
- Albuquerque EX, Deshpande SS, Kawabuchi M, Aracava Y, Idriss M, Rickett DL, and Boyne AF (1985) Multiple actions of anticholinesterase agents on cholinergic synapses: molecular basis for prophylaxis and treatment of organophosphate poisoning. *Fundam Appl Toxicol* **5**:S182–S203.

- Albuquerque EX, Pereira EFR, Aracava Y, Fawcett WP, Oliveira M, Randall WR, Hamilton TA, Kan RK, Romano JA, Jr, and Adler M (2006) Effective countermeasure against poisoning by organophosphorus insecticides and nerve agents. *Proc Natl Acad Sci USA* **103**:13220–13225.
- Anderson DM (2008) The non-human primate as a model for biomedical research, in *Sourcebook of Models for Biomedical Research* (Conn PM, ed) pp 251–258, Human Press, Totowa, NJ.
- Androustopoulos VP, Hernandez AF, Liesivuori J, and Tsatsakis AM (2013) A mechanistic overview of health associated effects of low levels of organochlorine and organophosphorus pesticides. *Toxicology* **307**:89–94.
- Baille V, Dorandeu F, Carpentier P, Bizot JC, Filliat P, Four E, Denis J, and Lallement G (2001) Acute exposure to a low or mild dose of soman: biochemical, behavioral and histopathological effects. *Pharmacol Biochem Behav* **69**: 561–569.
- Bajgar J (1992) Biological monitoring of exposure to nerve agents. *Br J Ind Med* **49**: 648–653.
- Bajgar J, Sevelová L, Krejčová G, Fusek J, Vachek J, Kassa J, Herink J, de Jong LP, and Benschop HP (2004) Biochemical and behavioral effects of soman vapors in low concentrations. *Inhal Toxicol* **16**:497–507.
- Banks CN and Lein PJ (2012) A review of experimental evidence linking neurotoxic organophosphorus compounds and inflammation. *Neurotoxicology* **33**:575–584.
- Bhagat YA, Obenaus A, Hamilton MG, and Kendall EJ (2001) Magnetic resonance imaging predicts neuropathology from soman-mediated seizures in the rodent. *Neuroreport* **12**:1481–1487.
- Bhagat YA, Obenaus A, Hamilton MG, Mikler J, and Kendall EJ (2005) Neuroprotection from soman-induced seizures in the rodent: evaluation with diffusion- and T2-weighted magnetic resonance imaging. *Neurotoxicology* **26**:1001–1013.
- Bodjarian N, Lallement G, Carpentier P, Baubichon D, and Blanchet G (1992) Soman-induced phosphoinositide hydrolysis in rat hippocampal slices: biochemical characterization. *Neurotoxicology* **13**:715–721.
- Bomser JA and Casida JE (2001) Diethylphosphorylation of rat cardiac M2 muscarinic receptor by chlorpyrifos oxon in vitro. *Toxicol Lett* **119**:21–26.
- Boverhof DR, Chamberlain MP, Elcombe CR, Gonzalez FJ, Heflich RH, Hernández LG, Jacobs AC, Jacobson-Kram D, Luijten M, and Maggi A, et al. (2011) Transgenic animal models in toxicology: historical perspectives and future outlook. *Toxicol Sci* **121**:207–233.
- Bowers MB, Jr, Goodman E, and Sim VM (1964) Some behavioral changes in man following anticholinesterase administration. *J Nerv Ment Dis* **138**:383–389.
- Buccafusco JJ (2008) Estimation of working memory in macaques for studying drugs for the treatment of cognitive disorders. *J Alzheimers Dis* **15**:709–720.
- Buckley NA, Eddleston M, Li Y, Bevan M, and Robertson J (2011) Oximes for acute organophosphate pesticide poisoning. *Cochrane Database Syst Rev* **2**:CD005085.
- Buckley NA, Karalliedde L, Dawson A, Senanayake N, and Eddleston M (2004) Where is the evidence for treatments used in pesticide poisoning? Is clinical toxicology fiddling while the developing world burns? *J Toxicol Clin Toxicol* **42**: 113–116.
- Burchfiel JL and Duffy FH (1982) Organophosphate neurotoxicity: chronic effects of sarin on the electroencephalogram of monkey and man. *Neurobehav Toxicol Teratol* **4**:767–778.
- Burchfiel JL, Duffy FH, and Van Sim M (1976) Persistent effects of sarin and diethylrin on the primate electroencephalogram. *Toxicol Appl Pharmacol* **35**:365–379.
- Byrnes ML, Reynolds JN, and Brien JF (2001) Effect of prenatal ethanol exposure during the brain growth spurt of the guinea pig. *Neurotoxicol Teratol* **23**:355–364.
- Carpentier P, Foquin A, Dorandeu F, and Lallement G (2001) Delta activity as an early indicator for soman-induced brain damage: a review. *Neurotoxicology* **22**: 299–315.
- Chebabo SR, Santos MD, and Albuquerque EX (1999) The organophosphate sarin, at low concentrations, inhibits the evoked release of GABA in rat hippocampal slices. *Neurotoxicology* **20**:871–882.
- Choi YJ, Kim JY, Yoo SB, Lee JH, and Jahng JW (2013) Repeated oral administration of capsaicin increases anxiety-like behaviours with prolonged stress-response in rats. *J Biosci* **38**:561–571.
- Coupland R and Leins KR (2005) Science and prohibited weapons. *Science* **308**:1841.
- Dahlgren JG, Takhar HS, Ruffalo CA, and Zwass M (2004) Health effects of diazinon on a family. *J Toxicol Clin Toxicol* **42**:579–591.
- Damodaran TV, Patel AG, Greenfield ST, Dressman HK, Lin SM, and Abou-Donia MB (2006) Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure. *Biochem Pharmacol* **71**:497–520.
- de Araujo Furtado M, Lumley LA, Robison C, Tong LC, Lichtenstein S, and Yourick DL (2010) Spontaneous recurrent seizures after status epilepticus induced by soman in Sprague-Dawley rats. *Epilepsia* **51**:1503–1510.
- Deshpande SS, Viana GB, Kauffman FC, Rickett DL, and Albuquerque EX (1986) Effectiveness of physostigmine as a pretreatment drug for protection of rats from organophosphate poisoning. *Fundam Appl Toxicol* **6**:566–577.
- Despain KE, McDonough JH, McMonagle JD, McGinley MJ, and Evans J (2007) The toxicity of soman in the African Green monkey (*Chlorocebus aethiops*). *Toxicol Mech Methods* **17**:255–264.
- Dobbing J and Sands J (1970) Growth and development of the brain and spinal cord of the guinea pig. *Brain Res* **17**:115–123.
- Dobbing J and Sands J (1973) Quantitative growth and development of human brain. *Arch Dis Child* **48**:757–767.
- Dolgin E (2013) Syrian gas attack reinforces need for better anti-sarin drugs. *Nat Med* **19**:1194–1195.
- Duffy FH, Burchfiel JL, Bartels PH, Gaon M, and Sim VM (1979) Long-term effects of an organophosphate upon the human electroencephalogram. *Toxicol Appl Pharmacol* **47**:161–176.
- Duysen EG, Li B, Xie W, Schopfer LM, Anderson RS, Broomfield CA, and Lockridge O (2001) Evidence for nonacetylcholinesterase targets of organophosphorus nerve agent: supersensitivity of acetylcholinesterase knockout mouse to VX lethality. *J Pharmacol Exp Ther* **299**:528–535.
- Engel SM, Wetmur J, Chen J, Zhu C, Barr DB, Canfield RL, and Wolff MS (2011) Prenatal exposure to organophosphates, paraoxonase 1, and cognitive development in childhood. *Environ Health Perspect* **119**:1182–1188.
- Fawcett WP, Aracava Y, Adler M, Pereira EFR, and Albuquerque EX (2009) Acute toxicity of organophosphorus compounds in guinea pigs is sex- and age-dependent and cannot be solely accounted for by acetylcholinesterase inhibition. *J Pharmacol Exp Ther* **328**:516–524.
- Ferlisi M and Hocker S (2013) What can we learn from status epilepticus registries? *Epilepsia* **54** (Suppl 6):72–73.
- Filliat P, Baubichon D, Burckhart MF, Pernot-Marino I, Foquin A, Masqueliez C, Perrichon C, Carpentier P, and Lallement G (1999) Memory impairment after soman intoxication in rat: correlation with central neuropathology: improvement with anticholinergic and antilutamatergic therapeutics. *Neurotoxicology* **20**:535–549.
- Filliat P, Coubard S, Pierard C, Liscia P, Beracochea D, Four E, Baubichon D, Masqueliez C, Lallement G, and Collombet JM (2007) Long-term behavioral consequences of soman poisoning in mice. *Neurotoxicology* **28**:508–519.
- Grigoryan H, Schopfer LM, Thompson CM, Terry AV, Masson P, and Lockridge O (2008) Mass spectrometry identifies covalent binding of soman, sarin, chlorpyrifos oxon, diisopropyl fluorophosphate, and FP-biotin to tyrosines on tubulin: a potential mechanism of long term toxicity by organophosphorus agents. *Chem Biol Interact* **175**:180–186.
- Grob D and Harvey AM (1953) The effects and treatment of nerve gas poisoning. *Am J Med* **14**:52–63.
- Gullapalli RP, Aracava Y, Zhuo J, Helal Neto E, Wang J, Makris G, Merchenthaler I, Pereira EFR, and Albuquerque EX (2010) Magnetic resonance imaging reveals that galantamine prevents structural brain damage induced by an acute exposure of guinea pigs to soman. *Neurotoxicology* **31**:67–76.
- Gunnell D, Eddleston M, Phillips MR, and Konradsen F (2007) The global distribution of fatal pesticide self-poisoning: systematic review. *BMC Public Health* **7**:357.
- Harmony T, Fernández-Bouzas A, Marosi E, Fernández T, Valdés P, Bosch J, Riera J, Bernal J, Rodríguez M, and Reyes A (1995) Frequency source analysis in patients with brain lesions. *Brain Topogr* **8**:109–117.
- Hatta K, Miura Y, Asukai N, and Hamabe Y (1996) Amnesia from sarin poisoning. *Lancet* **347**:1343.
- Hayward LJ, Wall HG, Jaax NK, Wade JV, Marlow DD, and Nold JB (1990) Decreased brain pathology in organophosphate-exposed rhesus monkeys following benzodiazepine therapy. *J Neurol Sci* **98**:99–106.
- Hood E (2001) The Tokyo attacks in retrospect: sarin leads to memory loss. *Environ Health Perspect* **109**:A542.
- Hurst CG, Newmark J, and Romano JA, Jr (2012) Chemical terrorism: Introduction, in *Harrison's Principles of Internal Medicine*, 18th ed (Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, and Loscalzo J eds) pp 1779–1788, McGraw-Hill, New York.
- Inns RH and Leadbeater L (1983) The efficacy of bispyridinium derivatives in the treatment of organophosphonate poisoning in the guinea-pig. *J Pharm Pharmacol* **35**:427–433.
- Jett DA, Abdallah EAM, El-Fakahany EE, Eldefrawi ME, and Eldefrawi AT (1991) High-affinity activation by paraoxon of a muscarinic receptor subtype in rat brain striatum. *Pestic Biochem Physiol* **39**:149–157.
- Jette N and Trevathan E (2014) Saving lives by treating epilepsy in developing countries. *Neurology* **82**:552–553.
- John H, Balszweit F, Kehe K, Worek F, and Thiermann H (2009) Toxicokinetics of chemical warfare agents: Nerve agents and vesicants, in *Handbook of Toxicology of Chemical Warfare Agents*, 1st ed (Gupta RC, ed) pp 755–790, Elsevier, London.
- Kadar T, Shapira S, Cohen G, Sahar R, Alkalay D, and Raveh L (1995) Sarin-induced neuropathology in rats. *Hum Exp Toxicol* **14**:252–259.
- Karalliedde L and Senanayake N (1989) Organophosphorus insecticide poisoning. *Br J Anaesth* **63**:736–750.
- Kassa J (2002) Review of oximes in the antidotal treatment of poisoning by organophosphorus nerve agents. *J Toxicol Clin Toxicol* **40**:803–816.
- Kassa J, Krejčová G, and Vachek J (2002) The impairment of spatial memory following low-level sarin inhalation exposure and antidotal treatment in rats. *Acta Med* **45**:149–153.
- Kawabuchi M, Boyne AF, Deshpande SS, and Albuquerque EX (1989) The reversible carbamate, (-)-physostigmine, reduces the size of synaptic end plate lesions induced by sarin, an irreversible organophosphate. *Toxicol Appl Pharmacol* **97**:98–106.
- Klass DW and Westmoreland BF (1985) Nonepileptogenic epileptiform electroencephalographic activity. *Ann Neurol* **18**:627–635.
- Koplovitz I and Stewart JR (1994) A comparison of the efficacy of HI6 and 2-PAM against soman, tabun, sarin, and VX in the rabbit. *Toxicol Lett* **70**:269–279.
- Lallement G, Baubichon D, Clarençon D, Galonnier M, Peoc'h M, and Carpentier P (1999) Review of the value of gacyclidine (GK-11) as adjuvant medication to conventional treatments of organophosphate poisoning: primate experiments mimicking various scenarios of military or terrorist attack by soman. *Neurotoxicology* **20**:675–684.
- Lallement G, Clarençon D, Brochier G, Baubichon D, Galonnier M, Blanchet G, and Mestries JC (1997) Efficacy of atropine/pralidoxime/diazepam or atropine/HI-6/prodiazepam in primates intoxicated by soman. *Pharmacol Biochem Behav* **56**:325–332.
- Lallement G, Clarençon D, Masqueliez C, Baubichon D, Galonnier M, Burckhart MF, Peoc'h M, and Mestries JC (1998) Nerve agent poisoning in primates: antilethal, anti-epileptic and neuroprotective effects of GK-11. *Arch Toxicol* **72**:84–92.
- Lallement G, Delamanche IS, Pernot-Marino I, Baubichon D, Denoyer M, Carpentier P, and Blanchet G (1993) Neuroprotective activity of glutamate receptor antagonists against soman-induced hippocampal damage: quantification with an omega 3 site ligand. *Brain Res* **618**:227–237.
- Lallement G, Pernot-Marino I, Baubichon D, Burckhart MF, Carpentier P, and Blanchet G (1994) Modulation of soman-induced neuropathology with an anticonvulsant regimen. *Neuroreport* **5**:2265–2268.
- Lee KN, Pellom ST, Oliver E, and Chirwa S (2014) Characterization of the guinea pig animal model and subsequent comparison of the behavioral effects of selective dopaminergic drugs and methamphetamine. *Synapse* **68**:221–233.

- Lein PJ and Fryer AD (2005) Organophosphorus insecticides induce airway hyper-reactivity by decreasing neuronal M2 muscarinic receptor function independent of acetylcholinesterase inhibition. *Toxicol Sci* **83**:166–176.
- Lemercier G, Carpentier P, Sentenac-Roumanou H, and Morelis P (1983) Histological and histochemical changes in the central nervous system of the rat poisoned by an irreversible anticholinesterase organophosphorus compound. *Acta Neuropathol* **61**:123–129.
- Lenz DE, Maxwell DM, Koplovitz I, Clark CR, Capacio BR, Cerasoli DM, Federko JM, Luo C, Saxena A, and Doctor BP, et al. (2005) Protection against soman or VX poisoning by human butyrylcholinesterase in guinea pigs and cynomolgus monkeys. *Chem Biol Interact* **157**:205–210.
- Levin HS and Rodnitzky RL (1976) Behavioral effects of organophosphate in man. *Clin Toxicol* **9**:391–403.
- Levin ED, Addy N, Nakajima A, Christopher NC, Seidler FJ, and Slotkin TA (2001) Persistent behavioral consequences of neonatal chlorpyrifos exposure in rats. *Brain Res Dev Brain Res* **130**:83–89.
- Levin ED, Addy N, Baruah A, Elias A, Christopher NC, Seidler FJ, and Slotkin TA (2002) Prenatal chlorpyrifos exposure in rats causes persistent behavioral alterations. *Neurotoxicol Teratol* **24**:733–741.
- Li Q, Hirata Y, Kawada T, and Minami M (2004) Elevated frequency of sister chromatid exchanges of lymphocytes in sarin-exposed victims of the Tokyo sarin disaster 3 years after the event. *Toxicology* **201**:209–217.
- Loh Y, Swanberg MM, Ingram MV, and Newmark J (2010) Case report: long-term cognitive sequelae of sarin exposure. *Neurotoxicology* **31**:244–246.
- Mamczarz J, Kulkarni GS, Pereira EFR, and Albuquerque EX (2011) Galantamine counteracts development of learning impairment in guinea pigs exposed to the organophosphorus poison soman: clinical significance. *Neurotoxicology* **32**:785–798.
- Mamczarz J, Pereira EFR, Aracava Y, Adler M, and Albuquerque EX (2010) An acute exposure to a sub-lethal dose of soman triggers anxiety-related behavior in guinea pigs: interactions with acute restraint. *Neurotoxicology* **31**:77–84.
- Martin LJ, Doebler JA, Shih TM, and Anthony A (1985) Protective effect of diazepam pretreatment on soman-induced brain lesion formation. *Brain Res* **325**:287–289.
- Maxwell DM, Brecht KM, and O'Neill BL (1987) The effect of carboxylesterase inhibition on interspecies differences in soman toxicity. *Toxicol Lett* **39**:35–42.
- Maxwell DM, Brecht KM, Koplovitz I, and Sweeney RE (2006) Acetylcholinesterase inhibition: does it explain the toxicity of organophosphorus compounds? *Arch Toxicol* **80**:756–760.
- McDonough JH, Jr and Shih TM (1997) Neuropharmacological mechanisms of nerve agent-induced seizure and neuropathology. *Neurosci Biobehav Rev* **21**:559–579.
- McDonough JH, Jr, Clark TR, Slone TW, Jr, Zoeffel D, Brown K, Kim S, and Smith CD (1998) Neural lesions in the rat and their relationship to EEG delta activity following seizures induced by the nerve agent soman. *Neurotoxicology* **19**:381–391.
- McDonough JH, Jr, McMonagle J, Copeland T, Zoeffel D, and Shih TM (1999) Comparative evaluation of benzodiazepines for control of soman-induced seizures. *Arch Toxicol* **73**:473–478.
- McDonough JH, Jr, Zoeffel LD, McMonagle J, Copeland TL, Smith CD, and Shih TM (2000) Anticonvulsant treatment of nerve agent seizures: anticholinergics versus diazepam in soman-intoxicated guinea pigs. *Epilepsia Res* **38**:1–14.
- McLeod CG, Jr, Singer AW, and Harrington DG (1984) Acute neuropathology in soman poisoned rats. *Neurotoxicology* **5**:53–57.
- Metcalfe DR and Holmes JH (1969) VII. Toxicology and physiology. EEG, psychological, and neurological alterations in humans with organophosphorus exposure. *Ann N Y Acad Sci* **160**:357–365.
- Mittal T, Gupta N, Kohli A, Bhalla A, Singh B, and Singh S (2011) Correlation of defects in regional cerebral blood flow determined by 99mTc SPECT with residual neurocognitive testing abnormalities during and 3 months post exposure in acutely poisoned patients with organophosphates. *Clin Toxicol* **49**:464–470.
- Murata K, Araki S, Yokoyama K, Okumura T, Ishimatsu S, Takasu N, and White RF (1997) Asymptomatic sequelae to acute sarin poisoning in the central and autonomic nervous system 6 months after the Tokyo subway attack. *J Neurol* **244**:601–606.
- Nishiwaki Y, Maekawa K, Ogawa Y, Asukai N, Minami M, and Omae K; Sarin Health Effects Study Group (2001) Effects of sarin on the nervous system in rescue team staff members and police officers 3 years after the Tokyo subway sarin attack. *Environ Health Perspect* **109**:1169–1173.
- Nozaki H, Aikawa N, Fujishima S, Suzuki M, Shinozawa Y, Hori S, and Nogawa S (1995) A case of VX poisoning and the difference from sarin. *Lancet* **346**:698–699.
- O'Donnell JC, McDonough JH, and Shih TM (2011) In vivo microdialysis and electroencephalographic activity in freely moving guinea pigs exposed to organophosphorus nerve agents sarin and VX: analysis of acetylcholine and glutamate. *Arch Toxicol* **85**:1607–1616.
- Okumura T, Takasu N, Ishimatsu S, Miyamoto S, Mitsuhashi A, Kumada K, Tanaka K, and Hinohara S (1996) Report on 640 victims of the Tokyo subway sarin attack. *Ann Emerg Med* **28**:129–135.
- Patrick K, Stanbrook M, and Flegel K (2013) Lest we forget: why the use of chemical weapons must not go unchallenged. *CMAJ* **185**:1299.
- Pereira EFR, Pescille JD, Gavruschenko L, Gamman R, Mamczarz J, and Albuquerque EX (2012) Cognitive impairment in adult male guinea pigs long after an acute exposure to the organophosphorus poison soman: Program No. 840.07/A67, in *2012 Neuroscience Meeting Planner*; 2012 Oct 13–17; New Orleans, LA. Society for Neuroscience Online.
- Petrus JM (1994) Neurology and neuropathology of Soman-induced brain injury: an overview. *J Exp Anal Behav* **61**:319–329.
- Prendergast MA, Self RL, Smith KJ, Ghayoumi L, Mullins MM, Butler TR, Buccafusco JJ, Gearhart DA, and Terry AV, Jr (2007) Microtubule-associated targets in chlorpyrifos oxon hippocampal neurotoxicity. *Neuroscience* **146**:330–339.
- RamaRao G, Bhattacharya BK, Kumar S, and Waghmare CK (2011) Gene expression and phosphoprotein profile of certain key neuronal signaling proteins following soman intoxication. *Toxicology* **290**:195–202.
- Rauh VA, Perera FP, Horton MK, Whyatt RM, Bansal R, Hao X, Liu J, Barr DB, Slotkin TA, and Peterson BS (2012) Brain anomalies in children exposed prenatally to a common organophosphate pesticide. *Proc Natl Acad Sci USA* **109**:7871–7876.
- Raveh L, Weissman BA, Cohen G, Alkalay D, Rabinovitz I, Sonog H, and Brandeis R (2002) Caramiphen and scopolamine prevent soman-induced brain damage and cognitive dysfunction. *Neurotoxicology* **23**:7–17.
- Roldán-Tapia L, Parrón T, and Sánchez-Santed F (2005) Neuropsychological effects of long-term exposure to organophosphate pesticides. *Neurotoxicol Teratol* **27**:259–266.
- Romano JA, Jr and King JM (2001) Psychological casualties resulting from chemical and biological weapons. *Mil Med* **166**:21–22.
- Romano, Jr JA, McDonough JH, Sheridan R, and Sidell FR (2001) Health effects of low-level exposure to nerve agents, in *Chemical Warfare Agents: Toxicity at Low Levels* (Somani SM and Romano Jr JA eds) pp 1–24, CRC Press, Boca Raton, FL.
- Rosas LG and Eskenazi B (2008) Pesticides and child neurodevelopment. *Curr Opin Pediatr* **20**:191–197.
- Rosenstock L, Keifer M, Daniell WE, McConnell R, and Claypoole K; The Pesticide Health Effects Study Group (1991) Chronic central nervous system effects of acute organophosphate pesticide intoxication. *Lancet* **338**:223–227.
- Santos MD, Pereira EFR, Aracava Y, Castro NG, Fawcett WP, Randall WR, and Albuquerque EX (2003) Low concentrations of pyridostigmine prevent soman-induced inhibition of GABAergic transmission in the central nervous system: involvement of muscarinic receptors. *J Pharmacol Exp Ther* **304**:254–265.
- Savage EP, Keefe TJ, Mounce LM, Heaton RK, Lewis JA, and Burcar PJ (1988) Chronic neurological sequelae of acute organophosphate pesticide poisoning. *Arch Environ Health* **43**:38–45.
- Savolainen KM and Hirvonen MR (1992) Second messengers in cholinergic-induced convulsions and neuronal injury. *Toxicol Lett* **64**:437–445.
- Sekijima Y, Morita H, and Yanagisawa N (1997) Follow-up of sarin poisoning in Matsumoto. *Ann Intern Med* **127**:1042.
- Shih TM, Duniho SM, and McDonough JH (2003) Control of nerve agent-induced seizures is critical for neuroprotection and survival. *Toxicol Appl Pharmacol* **188**:69–80.
- Shih TM and McDonough JH, Jr (1999) Organophosphorus nerve agents-induced seizures and efficacy of atropine sulfate as anticonvulsant treatment. *Pharmacol Biochem Behav* **64**:147–153.
- Silveira CL, Eldefrawi AT, and Eldefrawi ME (1990) Putative M2 muscarinic receptors of rat heart have high affinity for organophosphorus anticholinesterases. *Toxicol Appl Pharmacol* **103**:474–481.
- Sirkka U, Nieminen SA, and Ylitalo P (1990) Neurobehavioral toxicity with low doses of sarin and soman. *Methods Find Exp Clin Pharmacol* **12**:245–250.
- Sivam SP, Hoskins B, and Ho IK (1984) An assessment of comparative acute toxicity of diisopropyl-fluorophosphate, tabun, sarin, and soman in relation to cholinergic and GABAergic enzyme activities in rats. *Fundam Appl Toxicol* **4**:531–538.
- Spradling KD, Lumley LA, Robison CL, Meyerhoff JL, and Dillman JF, 3rd (2011) Transcriptional responses of the nerve agent-sensitive brain regions amygdala, hippocampus, piriform cortex, septum, and thalamus following exposure to the organophosphonate anticholinesterase sarin. *J Neuroinflammation* **8**:84.
- Spruit HE, Langenberg JP, Trap HC, van der Wiel HJ, Helmich RB, van Helden HP, and Benschop HP (2000) Intravenous and inhalation toxicokinetics of sarin stereoisomers in atropinized guinea pigs. *Toxicol Appl Pharmacol* **169**:249–254.
- Steenland K, Jenkins B, Ames RG, O'Malley M, Chrislip D, and Russo J (1994) Chronic neurological sequelae to organophosphate pesticide poisoning. *Am J Public Health* **84**:731–736.
- Sun W, Doctor BP, Lenz DE, and Saxena A (2008) Long-term effects of human butyrylcholinesterase pretreatment followed by acute soman challenge in cynomolgus monkeys. *Chem Biol Interact* **175**:428–430.
- Terry AV, Jr (2012) Functional consequences of repeated organophosphate exposure: potential non-cholinergic mechanisms. *Pharmacol Ther* **134**:355–365.
- Testylier G, Tonduli L, and Lallemand G (1999) Implication of gamma band in soman-induced seizures. *Acta Biotheor* **47**:191–197.
- Tucker JB (2006) IG Farber, in *War of Nerves: Chemical Warfare from World War I to Al-Qaeda* (Tucker JB ed) pp 24–41, Pantheon Books, New York.
- Turgeman G, Pinkas A, Slotkin TA, Tfilin M, Langford R, and Yanai J (2011) Reversal of chlorpyrifos neurobehavioral teratogenicity in mice by allographic transplantation of adult subventricular zone-derived neural stem cells. *J Neurosci Res* **89**:1185–1193.
- Wesseling C, Keifer M, Ahlbom A, McConnell R, Moon JD, Rosenstock L, and Hogstedt C (2002) Long-term neurobehavioral effects of mild poisonings with organophosphate and n-methyl carbamate pesticides among banana workers. *Int J Occup Environ Health* **8**:27–34.
- Westmoreland BF (1998) The EEG findings in extratemporal seizures. *Epilepsia* **39** (Suppl 4):S1–S8.
- Woodard CL, Calamaio CA, Kaminskis A, Anderson DR, Harris LW, and Martin DG (1994) Erythrocyte and plasma cholinesterase activity in male and female rhesus monkeys before and after exposure to sarin. *Fundam Appl Toxicol* **23**:342–347.
- Xie W, Stribley JA, Chatonnet A, Wilder PJ, Rizzino A, McComb RD, Taylor P, Hinrichs SH, and Lockridge O (2000) Postnatal developmental delay and supersensitivity to organophosphate in gene-targeted mice lacking acetylcholinesterase. *J Pharmacol Exp Ther* **293**:896–902.
- Yamasue H, Abe O, Kasai K, Suga M, Iwanami A, Yamada H, Tochigi M, Ohtani T, Rogers MA, and Sasaki T, et al. (2007) Human brain structural change related to acute single exposure to sarin. *Ann Neurol* **61**:37–46.
- Yokoyama K, Araki S, Murata K, Nishikitani M, Okumura T, Ishimatsu S, and Takasu N (1998) A preliminary study on delayed vestibulo-cerebellar effects of Tokyo subway sarin poisoning in relation to gender difference: frequency analysis of postural sway. *J Occup Environ Med* **40**:17–21.

Address correspondence to: Dr. Edson X. Albuquerque, Division of Translational Toxicology, Department of Epidemiology and Public Health, University of Maryland School of Medicine, 10 South Pine Street, Suite 902, Baltimore, MD 21201. E-mail: ealbuquerque@som.umaryland.edu