



Published in final edited form as:

Oncogene. 2013 August 29; 32(35): 4130–4138. doi:10.1038/onc.2012.425.

Tumor suppressive *miR-124* targets androgen receptor and inhibits proliferation of prostate cancer cells

Xu-Bao Shi¹, Lingru Xue¹, Ai-Hong Ma², Clifford G. Tepper², Regina Gandour-Edwards³, Hsing-Jien Kung², and Ralph W. deVere White^{1,†}

¹Department of Urology, University of California, Davis, School of Medicine, Sacramento, CA 95817, USA

²Department of Biochemistry and Molecular Medicine, University of California, Davis, School of Medicine, Sacramento, CA 95817, USA

³Department of Pathology, University of California, Davis, School of Medicine, Sacramento, CA 95817, USA

Abstract

Although prostate cancer (CaP) is the most frequently diagnosed malignant tumor in American men, the mechanisms underlying the development and progression of CaP remain largely unknown. Recent studies have shown that downregulation of *miR-124* occurs in several types of human cancer, suggesting a tumor suppressive function of *miR-124*. Until now, however, it has been unclear whether *miR-124* is associated with CaP. In the present study, we completed a series of experiments to understand the functional role of *miR-124* in CaP. We detected the expression level of *miR-124* in clinical CaP tissues, evaluated the influence of *miR-124* on the growth of CaP cells, and investigated the mechanism underlying the dysregulation of *miR-124*. We found that *i*) *miR-124* directly targets the androgen receptor (AR) and subsequently induces a upregulation of p53; *ii*) *miR-124* is significantly down-regulated in malignant prostatic cells compared to that in benign cells and DNA methylation causes the reduced expression of *miR-124*; and *iii*) *miR-124* can inhibit the growth of CaP cells *in vitro* and *in vivo*. Data from this study revealed that loss of *miR-124* expression is a common event in CaP, which may contribute to pathogenesis of CaP. Our studies also suggest that *miR-124* is a potential tumor suppressive gene in CaP, and restoration of *miR-124* expression may represent a novel strategy for CaP therapy.

Keywords

prostate cancer; androgen; androgen receptor; miRNA; tumor suppressor

[†]Address correspondence to: Department of Urology, University of California, Davis, School of Medicine 4860 Y Street, Suite 3500, Sacramento, California, 95817. rwdeverewhite@ucdavis.edu.

Conflict of interest: The authors declare no conflict of interest.

Introduction

Prostate cancer (CaP) is the most frequently diagnosed malignant tumor and was the second leading cause of cancer-related death in American men in 2011 (Siegel et al 2012). Metastatic CaP can be treated effectively with androgen ablation. However, one of the most troubling aspects of this disease is that after hormone treatment, the tumor inevitably progresses from an androgen-dependent (AD) to an incurable castration-resistant (CR) form (Sadar 2011). The mechanism underlying the progression has been poorly understood. In the past decades, therefore, an important task in CaP research is to elucidate the molecular alteration occurring in the development of CR tumors. Considerable insight into the progression of CaP has been recently achieved, including the discovery of aberrantly expressed microRNAs (miRNAs).

The human genome may encode over 1000 miRNAs which negatively regulate approximately 60% of human genes (Friedman et al 2009, Lewis et al 2005). Thus, miRNAs are involved in almost all important cellular processes, not only in physiological conditions but also in diseases including cancer (Volinia et al 2006). Indeed, a number of cancer-related miRNAs have been identified recently. Some miRNAs have also been reported to be aberrantly expressed in human CaP (Shi et al 2008). These miRNAs, which act as tumor suppressor genes or oncogenes, contribute to the pathogenesis of CaP by directly targeting some proliferation-related genes or apoptotic molecules (Shenouda and Alahari 2009). In spite of these exciting findings, the role of miRNA in the development and progression of CaP has been largely unexplored. Thus, identifying CaP-associated miRNAs and investigating their roles in CaP will help us understand the mechanisms related to the development and progression of this disease.

MiR-124 is a highly conserved miRNA whose *in vivo* function is poorly defined. This small non-coding RNA was first reported to be highly expressed in neuronal cells (Makeyev et al 2007). Recent studies revealed that *miR-124* was significantly downregulated in several types of human cancers (Ando et al 2009, Furuta et al, Lujambio et al 2007). Moreover, it regulates some proliferation-related genes such as cyclin-dependent kinase 6 (CDK6) (Agirre et al 2009), forkhead box A2 (Foxa2) (Baroukh et al 2007) and solute carrier family 16, member 1 (SLC16A1) (Agirre et al 2009). Thus, *miR-124* is classified as tumor suppressor miRNA (Agirre, 2009 #11). Until now, however, the role of *miR-124* in CaP has been totally unknown, although it was reported previously to be undetectable in 22Rv1 CaP cells (Mitchell et al 2008).

In the present study, we aimed to explore the role of *miR-124* in CaP. We found that *miR-124* directly targets the androgen receptor (AR). We also detected the expression level of *miR-124* in clinical CaP tissues and found that a majority of these CaP tissues express low levels of *miR-124*. In addition, we observed that synthetic *miR-124* mimics inhibit the proliferation of CaP cells. Since the AR plays a crucial role in the pathogenesis of CaP, our studies established a link between dysregulated *miR-124*, overexpressed AR and CaP cell proliferation. Our results imply that *miR-124* is a potential tumor suppressor gene and downregulation of *miR-124* contributes to the development and progression of CaP.

Results

MiR-124 directly targets the AR and inhibits proliferation of CaP cells

Overexpression of the AR is well known to play an important role in pathogenesis of CaP. We are very interested in determining whether certain miRNA(s) contribute to upregulation of the AR. From two algorithms, TargetScan (Release 5.2) and miRanda (August 2010 Release) that predict miRNA-binding sites in the first 436 base of the AR 3'UTR (NM_000044.2), we identified three broadly conserved miRNA-binding sites for six miRNAs (*miR-130/miR-301*, *miR-124/miR-506* and *miR-30/miR-384*). In order to assess the ability of these miRNAs to regulate AR expression, AR-positive C4-2B CaP cells cultured in androgen-deprived medium were separately treated with four chemically-modified miRNA mimics (*miR-124*, *miR-130*, *miR-384* or *miR-506*, purchased from Ambion). As shown in Fig. 1A, compared to miRNA negative control (miR-NC), treatment with *miR-124* mimic resulted in a reduction of AR protein by ~70%, while the other three miRNA mimics induced 20–40% decrease in AR expression, suggesting a potent downregulation of the AR by *miR-124*. To determine the influence of these miRNAs on proliferation of CaP cell, AR-positive CaP cell lines (LNCaP, C4-2B and 22Rv1) were transiently transfected with each of these miRNA mimics. Consistent with its effect on AR level, transfection of *miR-124* mimic inhibited proliferation to a greater extent than the other miRNA mimics tested (SI Fig. 1). We thus focused on *miR-124* in this study.

To determine whether *miR-124*-mediated downregulation of the AR affects the AR activity, both AR-positive LNCaP and C4-2B were treated with *miR-124* mimic. Western blot analysis demonstrated that *miR-124*-induced down-regulation of the AR and was concomitant with a reduced PSA level (Fig. 1B). Since AR regulates *miR-125b* (Shi et al 2007), we used C4-2B and cds2 cells which express increased *miR-125b* (Shi et al 2007) to examine the effect of *miR-124* mimic on endogenous *miR-125b*. We found that *miR-124* induced downregulation of *miR-125b* by 30% in C4-2B and 54% in cds2 (Fig. 1C). To verify that the putative *miR-124* binding site in the 3' UTR of AR mRNA is responsible for regulation by *miR-124*, the 3' UTR was cloned into a luciferase reporter vector and then cotransfected with *miR-124* mimic into C4-2B cells. Luciferase activity was measured two days after transfection. As shown in Fig. 1D, transfection of *miR-124* mimic resulted in 40% reduction of the enzyme activity, indicating a direct interaction between *miR-124* and AR mRNA. Taken together, these data indicate that *miR-124* targets the AR and affects the AR activity.

MiR-124 is significantly downregulated in prostate cancer cells

We determined the expression level of *miR-124* in CaP cells. The abundance of *miR-124* was first examined in seven prostate cell lines (2 benign and 5 malignant) using quantitative PCR (qPCR). We found a reduced expression of *miR-124* in the malignant lines compared to the benign lines (Fig. 2A). We also performed Northern blot analysis of these cell lines and similar results were obtained (SI Fig. 2A). Next, we tested whether *miR-124* is down-regulated in clinical prostate samples. We reviewed the miRNA expression profiling of our initial clinical tissue specimens and found that the abundance of *miR-124* is markedly reduced in four CaP samples relative to that in one benign prostatic hyperplasia (BPH) tissue

(SI Fig. 2B). We then determined whether downregulation of *miR-124* is common in clinical CaP tissues. The levels of *miR-124* were detected in 79 clinical prostatic tissues including 19 BPHs, 44 primary CaPs, 6 lymph node metastases, and 10 CR tumors. Although *miR-124* abundances have overlap in a small number of benign and primary tumor tissues, the average *miR-124* level was 2.7-fold less in primary CaPs than in BPHs (39 in CaPs vs 146 in BPHs, $p < 0.01$) (Fig. 2B). Interestingly, 10 CR tumors and 6 metastases expressed extremely low levels of *miR-124*. Compared to the primary CaP samples, the averaged *miR-124* level decreases by 77% in these advanced tumors (39 in CaPs vs 9 in CR & metastatic tumors, $p < 0.05$) (Fig. 2B), suggesting a potential involvement of *miR-124* in CaP progression. To confirm the downregulation of *miR-124*, 18 CaP samples and matched benign adjacent tissues were analyzed using qPCR. In comparison with matched benign tissues, 15 of 18 CaP samples (83%) expressed significantly reduced *miR-124* (Fig. 2C). In addition, we performed Northern blot analysis in five matched prostate tissues in which sufficient quantities of RNA were available. As expected, the three CaPs (8, 11 and 16) having low *miR-124* levels detected by qPCR exhibit lower signal intensity in Northern blots (SI Fig. 2C). Taken together, these data provide strong evidence that *miR-124* is significantly reduced in CaP cell lines and in a majority of clinical CaP samples, and also suggest that decreased expression of *miR-124* is common in human CaP tissues.

***MiR-124* and AR are negatively correlated in human CaP tissues**

Having demonstrated that *miR-124* targets the AR and is significantly downregulated in CaP cells, we therefore asked whether clinical CaP samples having low level of *miR-124* overexpressed the AR. To address the issue, eight additional matched pairs of CaP and BPH were examined for expression of AR protein using immunohistochemical (IHC) analysis. In these samples, *miR-124* level detected by qPCR was significantly decreased in CaP samples compared to that in BPH. Conversely, IHC results demonstrated that AR immunostaining was more intense in 7 of 8 CaP samples than that in BPH matches. These results reveal an inverse relationship between *miR-124* and AR expression levels in prostatic tissues. Fig. 3 shows the representative results from four matches. In CaP samples AR protein staining is much higher than that in matched BPH tissues (Fig. 3A), while the expression of *miR-124* is significantly decreased in CaP tissue compared to that in BPH tissues (Fig. 3B). Therefore, our data suggest that down-regulation of *miR-124* results in increased expression of the AR in clinical CaP tissues.

***MiR-124* is methylated in CaP cells**

The question remained: why is *miR-124* downregulated in CaP cells? We first confirmed that androgens do not regulate *miR-124*, based on our observation that *miR-124* levels were similar in androgen R1881-treated and untreated CaP cell lines (data not shown). Previous study has shown that DNA methylation regulates the expression of miRNAs (Sinkkonen et al 2008). Moreover, computer analysis demonstrated that the *miR-124-1* at 8p23.1 and the *miR-124-3* at 20q13.33 are embedded within CpG islands and the *miR-124-2* at 8q12.3 is 760 bp downstream of a CpG island (Agirre et al 2009). In order to determine that DNA methylation downregulates *miR-124* in CaP cells, we performed a demethylation experiment in three androgen-independent cell lines (22Rv1, C4-2B and cds2). Cells were treated with 50 μ M of 5'-Aza-2'-Deoxycytidine (Aza) for 4 days and *miR-124* was detected by qPCR. It

was found that Aza treatment induced 2- to 5-fold increase in the *miR-124* level compared to that in the untreated cells (Fig. 4A). Since *miR-124* targets the AR as shown in Figure 1, we tested whether Aza downregulates expression of the AR. It was found that treatment with Aza reduced the AR level by ~40% in LNCaP and ~60% in C4-2B cells (Fig. 4B). We then assessed the methylation status of two benign prostate cell lines (pRNS-1-1 and RWPE-1) and five CaP cell lines using methylation-specific PCR (MSP). The 5' DNA fragments of three *miR-124* genes were strongly amplified by MSP primers in all cell lines except RWPE-1 whose *miR-124-2* fragment was not amplified by the MSP primers (Fig. 4C). The fragments of *miR-124-1* and *miR-124-2* were also amplified weakly by unmethylated DNA primers in some of these cell lines (Fig. 4C). MSP data suggest different extents of DNA methylation at the 5' regions of *miR-124* genes. In order to validate DNA methylation, the 5' DNA fragments of three *miR-124* genes in six cell lines were amplified using COBRA (combined bisulfite restriction analysis) primers, cloned and sequenced. It was found that 40-80% of CpG sites were methylated in *miR-124-1*, 20-57% in *miR-124-2* and 15-31% in *miR-124-3*, in five CaP cell lines, and the CpG methylation are 18%, 13% and 8%, respectively, in benign RWPE-1 cells (Fig. 4D).

We next determined the methylation status in 9 BPH tissues and 14 CaP tissues using COBRA. The 5' DNA fragments of three *miR-124* genes that contain 2, 5 and 6 CGCG sites, respectively, were amplified with COBRA primers and digested with the restriction enzyme *Bst*U1 which only cleaves methylated CGCG sites. It was found that *Bst*U1 was able to cut the *miR-124-1* PCR products in 13 of 14 (93%) CaP samples, *miR-124-2* in 11 of 14 (79%) samples and *miR-124-3* in 12 of 14 (86%) samples, while the *Bst*U1-digested BPH DNA fragments of *miR-124* genes are 11% (1/9), 22% (2/9) and 44% (4/9), respectively (Fig. 4E), indicating that significant *miR-124* methylation occurs in clinical CaP specimens. Taken together, our data suggest that DNA methylation occurring at the 5' regions of *miR-124* genes causes, at least in part, the reduced expression of *miR-124* in CaP cells.

***MiR-124* induces apoptosis**

Previous studies demonstrated that the AR upregulates *miR-125b*, while *miR-125b* represses expression of p53 (Le et al 2009) and facilitates proliferation of CaP cells (Shi et al, Shi et al 2007). Since *miR-124* targets the AR and downregulates *miR-125b*, we asked whether increased *miR-124* level increases apoptotic cell death. To address the issue, LNCaP and C4-2B cells were transfected with *miR-124* mimic and then cultured for 4 days. Apoptotic death cells were quantitatively detected by Annexin V binding assay. We observed that *miR-124* mimic induced 10.4% of LNCaP cells and 9.7% of C4-2B cells to undergo apoptotic cell death (Fig. 5A). The comparative figure for the miR-NC cells was 0.7% and 1.1%, respectively ($p < 0.01$). A representative Annexin V assay is shown in SI Fig. 3 in which *miR-124* mimic-induced apoptosis (upper-right and lower-right squares in quadrant gates) were 11.4% of LNCaP cells and 8.4% of C4-2B cells. To provide biochemical evidence for the occurrence of apoptosis, we determined whether *miR-124* mimic increases the expression of p53 and caspase 3. As expected, treatment of C4-2B cells with *miR-124* mimic induced a ~9.0-fold enhanced expression of p53 (Fig. 5B) and ~2.0-fold increase in caspase 3 (Fig. 5C). These data indicate that *miR-124*, at least in part, induces apoptosis by activating the p53 signaling pathway.

MiR-124 inhibits xenograft growth of CaP cells

Having demonstrated that *miR-124* inhibits proliferation of AR-positive CaP cells and induces apoptosis, we tested whether this miRNA exerts an inhibitory effect on xenograft growth of CaP cells *in vivo*. We employed 22Rv1 CaP cells to address this question due to their expression of low (Figure 2A) or undetectable (Mitchell et al 2008) level of *miR-124*, as well as their ability to efficiently form tumors in nude mice. Overexpression of *miR-124* in 22Rv1 was established using lentiviral vector system that expresses 28-fold elevation of *miR-124* relative to controls (data not shown). Xenograft tumors were generated by subcutaneously inoculating intact male nude mice with 22Rv1-lenti-*miR-124* cells or 22Rv1-lenti-vector cells. Tumors in *miR-124* group and control groups became palpable 18 days after inoculation, suggesting that *miR-124* did not affect the onset of 22Rv1 tumor. However, overexpression of *miR-124* significantly inhibited the growth of 22Rv1 tumors after three weeks when compared to controls (Fig. 6A). At the end of the experiments, qPCR analysis of *miR-124* level was performed in 3 lenti-*miR-124* tumors. It was found that *miR-124* levels were 23-fold increase in the *miR-124* tumors compared to the control tumors (Fig. 6B), similar to that in cultured 22Rv1-lenti-*miR-124* cells, revealing a stable expression of *miR-124 in vivo*. In addition, one lenti-vector tumor and two lenti-*miR-124* tumors were examined by Western blot analysis for their AR expression. As shown in Fig. 6C, lenti-*miR-124* tumors express moderate reduction of the full-length AR. Interestingly, overexpression of *miR-124* in 22Rv1 cells induced a downregulation of the truncated AR that may contribute to the aggressive features of 22Rv1 cells (Tepper et al 2002). Collectively, our results suggest that *miR-124* inhibits growth of 22Rv1 xenograft tumors by repressing AR expression.

Discussion

Although *miR-124* has been reported to be involved in several other cancer types (Ando et al 2009, Furuta et al, Lujambio et al 2007), its role in CaP is largely unknown. To date, only one study has reported that *miR-124* was undetectable in 22Rv1 CaP cells (Mitchell et al 2008). The present study explored the role of *miR-124* in CaP. We found that *miR-124* directly targets the AR, suggesting a potential tumor suppressor gene in CaP. In addition, our expression analysis revealed that *miR-124* was significantly downregulated in a majority of clinical CaP specimens tested, which suggests that dysregulation of *miR-124* may be a common event in patients with CaP and loss of *miR-124* may contribute to the progression of CaP.

In this study, we address the potential mechanisms underlying *miR-124* down-regulation in CaP cells. Since methylation of tumor suppressor genes (TSGs) is very common in both early and advanced stages of CaP (Graff et al 1995, Kang et al 2004, Lou et al 1999, Woodson et al 2004), *miR-124* acting as a potential TSG in CaP may be susceptible to epigenetic modification. Indeed, we observed DNA methylation of three *miR-124* genes in CaP cell lines and clinical CaP samples, an explanation for the reduction of *miR-124* levels. This is supported by the fact that demethylation treatment with Aza increased the abundance of *miR-124* in CaP cells. These data provide strong evidence that *miR-124* is downregulated in CaP in part as a result of DNA methylation. Our results are consistent with those reported

in other cancer types in which *miR-124* loci are aberrantly methylated (Agirre et al 2009, Furuta et al, Lujambio et al 2007). However, methylation of *miR-124* loci was not detected in a recent report (Rauhala et al 2010). In that study, Rauhala *et al.* treated CaP cells with 1 μM of Aza for two days followed by 300 nM of trichostatin A (TSA, a histone deacetylase inhibitor) for 24 hours. They identified 38 methylated miRNAs without *miR-124*. Mazar *et al* reported that Aza upregulates *miR-34b* expression in a dose-dependent manner and 1 μM of Aza did not induce elevation of *miR-34b* in melanoma cells (Mazar et al 2011). Since up to 100 μM of Aza failed to induce cytotoxicity in CaP cell lines (Walton et al 2008), we used 50 μM of Aza to treat CaP cells for four days, which induced significant elevation of *miR-124*. It is thus likely that our ability to detect the increased level of *miR-124* is due to the higher concentration of Aze used. Besides the epigenetic regulation, the expression level of *miR-124* can be affected by genetic factors, such as mutation and heterozygosity (LOH). Indeed, we detected a G \rightarrow C point mutation (or SNP) in the 5' region of *miR-124-2* (SI Fig. 4). Mutation (or SNP) in 5' region can reduce expression of mature miRNA by altered miRNA processing (Davis and Hata 2009, Jazdzewski et al 2008). In addition, a previous study revealed that tumor suppressive miRNAs are frequently located in LOH regions (Calin et al 2004). We know that *miR-124-1* is located at 8p23.1 and *miR-124-2* at 8q12.3. LOH in these two regions were reported to occur in approximately 50% of CaPs (Chang et al 2007, Matsuyama et al 2003, Perincheri et al 1999). Thus, LOH may affect the levels of *miR-124-1* & *miR-124-2*. In order to elucidate the mechanism underlying downregulation of *miR-124*, further study is needed to detect the frequencies of DNA mutation (or SNP) and LOH in three *miR-124* loci.

An interesting and clinically relevant discovery is that *miR-124* directly targets the AR. The AR contributes to the initiation and progression of CaP. A study has demonstrated that a moderately altered AR expression is sufficient to convert AD CaP cells to a CR form (Chen et al 2004). The AR has been reported to be upregulated in clinical CaPs, particularly those that are CR. Why AR is upregulated in CaP remains poorly understood. AR gene amplification appears to account for the AR upregulation in \sim 30% CR CaPs (Koivisto et al 1997); however, it is unknown how the AR is upregulated in the other CR tumors. Since the 3'UTR of the AR mRNA contains a conserved *miR-124* binding site, we performed several experiments to validate the regulation of the AR by *miR-124*. We also observed an inverse correlation between *miR-124* abundance and the AR expression level in clinical CaP samples tested. Therefore, our study demonstrates that dysregulation of *miR-124* may elevate the expression of the AR, which is a new mechanistic explanation for overexpression of the AR in CaP. In this present study, identification of *miR-124* as AR-targeting miRNA was based on the TargetScan and miRanda prediction programs. We noted that *miR-124*, as well as other three predicted miRNAs as shown in Figure 1A, were not selected as AR-targeting miRNAs in a recent study (Ostling et al 2011). In that study, the authors transfected five AR-positive CaP cell lines with 20 nM of human miRNA library and selected 71 miRNA candidates that affect the level of AR in all five cell lines. However, miRNA repression of target molecular expression is dose-dependent (Yamakuchi et al 2008). The dose of *miR-124* mimic used in this study is 100 nM. In addition, miRNA functions in a cell type- and context-dependent manner (Olive et al 2010), and some AR-targeting miRNAs may not equally affect the AR level in all five CaP cell lines. Therefore,

different treatment conditions and different selection criteria may account for the differential results.

Although we focus on the AR in this study, we realize that *miR-124* targets hundreds of human genes, and other potential targets may also be relevant for the tumor suppressive function of *miR-124* in CaP. We have identified a putative *miR-124* binding site in the 3'UTR of high mobility group AT-hook 1 gene (*HMGAI*). In a pilot study, transfection of C4-2B cells with *miR-124* mimic resulted in reduction of HMGA1 expression by 60%; furthermore, two clinical CaP specimens having low level of *miR-124* did overexpress cytoplasmic HMGA1 (see SI Fig. 5), suggesting a link between low *miR-124* and overexpression of HMGA1 in CaP. Previous studies revealed that HMGA1 was highly expressed in advanced CaPs (Takaha et al 2002, Tamimi et al 1993), and cytoplasmic HMGA1 directly interacts with p53, leading to p53 inactivation (Frasca et al 2006, Pierantoni et al 2006). Therefore, functional characterization of this *miR-124* target will provide new insight into elucidating the molecular alteration occurring in advanced CaP. Additionally, cyclin-dependent kinase 6 (CDK6) and forkhead box A2 (Foxa2) have been identified as promising targets of *miR-124* (Agirre et al 2009, Baroukh et al 2007). Both CDK6 and Foxa2 were previously reported to be elevated in CaP cells and correlate with tumor grade (Chen et al 1996, Mirosevich et al 2006, Qi et al). Moreover, these two proteins interact with the AR and enhance the AR activity (Lim et al 2005, Yu et al 2005). Further studies are required to understand whether upregulation of these two molecules in CaP cells are the result of decreased expression of *miR-124*.

Consistent with a previous report in nerve cells (Cao et al 2007), restoration of *miR-124* level by transfecting CaP cells with synthetic *miR-124* induces increased apoptosis. We also found that treatment of CaP cells with *miR-124* upregulated the expression of p53, which is due at least partially to *miR-124*-mediated inhibition of AR/*miR-125b* signaling (Shi et al 2007). Thus, upregulation of p53 signaling may play a key role in *miR-124*-induced apoptosis. Since *miR-124* induces apoptosis, we evaluated its effect on proliferation of CaP cells. Restoration of *miR-124* level significantly inhibited the growth of CaP cells. More interestingly, when lentiviral-transduced 22Rv1 CaP cells that stably express *miR-124* were injected subcutaneously into intact male nude mice, xenograft tumor growth also was significantly inhibited. Therefore, further testing of *miR-124* in pre-clinical models of CaP will help define its ultimate therapeutic potential for treatment of CaP.

In conclusion, this study has shown that loss of *miR-124* expression may be a common event in CaP. Since *miR-124* negatively regulates the AR, downregulation of *miR-124*, as found in CaP, increases expression of the receptor. Therefore, deregulation of *miR-124* as well as other AR-targeting miRNA candidates (Ostling et al 2011) contributes to high levels of AR expression in clinical CaP. Additionally, our data suggest that downregulation of *miR-124* may be involved in the pathogenesis of CaP. Our evidence supporting the ability of *miR-124* to induce apoptosis in CaP cells encourages the hope that restoring *miR-124* function in CaP cells, either by itself or in conjunction with other therapies, will offer improved survival, which will be through delayed development or treatment of CR prostate cancer.

Materials and Methods

Cell lines

Two immortalized benign prostatic epithelial cell lines (RWPE-1 and pRNS-1-1) and five CaP cell lines (LNCaP, C4-2B, cds2, 22Rv1 and LAPC-4) were employed in this study. RWPE-1 (provided by Dr. Mukta. Webber, Michigan State University, East Lansing, MI) and pRNS-1-1 (provided by Dr. Johng Rhim, University of the Health Sciences, Bethesda, MD) were maintained in keratinocyte serum-free medium supplemented with 50 mg/ml bovine pituitary extract and 5 ng/ml epidermal growth factor. CaP cell lines were maintained in RPMI 1640 medium containing 10% fetal bovine serum. Both C4-2B and cds2 cell lines are derivatives of LNCaP (Lin et al 2001, Shi et al 2004).

Clinical samples

Collection and use of CaP patient specimens were approved by the Institutional Review Board (UCD IRB#: 200312072-6). Primary CaP samples from radical prostatectomy, metastatic pelvic lymph nodes, CR tumor specimens and BPH tissues were obtained fresh from surgical excision by the Department of Urology, University of California, Davis. These CR tumors were described in our previous publication (de Vere White et al 1997): six from stage D patients and four from patients receiving combined androgen blockade therapy prior to radical prostatectomy. To ensure that portions of CaP tissues were enriched for tumor cells, a surgeon and a pathologist together sectioned the excised tissue, and a fresh piece was taken from the palpated cancer and rapidly frozen to -80°C until use. A permanent section was then cut from the immediately adjacent area for Hematoxylin and Eosin (H&E) staining. Based upon histological review of H&E section, it was confirmed that these CaP samples used in this study were tumor cell-enriched.

Reporter plasmid construction and luciferase assay

To construct reporter plasmids, a 0.45-kb DNA fragment of the *AR* 3'UTR containing the putative *miR-124* binding site was prepared by PCR. The *AR* 3'UTR fragment lacking the *miR-124* binding site was used as negative control. DNA fragments were cloned into the pMIR-REPORT Luciferase vector (Ambion) downstream of the reporter gene. The sequences and cloning direction of these PCR products were validated by DNA sequencing. For luciferase assay, cells (4×10^4 per well) were seeded into 24-well plates and cultured for 24 hours. The cells were then co-transfected with reporter plasmids and 100 nM chemically-modified *miR-124* mimic or miRNA negative control (miR-NC). The pRL-SV40 Renilla luciferase plasmid (Promega) was used as an internal control. Two days later, cells were harvested and lysed with passive lysis buffer (Promega). Luciferase activity was measured using a dual luciferase reporter assay (Promega). Luciferase activity was normalized by Renilla luciferase activity.

Western blot analysis

Cells were grown to 70–80% confluence and lysed in lysis buffer consisting of 25 mmol/L Tris-HCl (pH7.5), 150 mmol/L NaCl, 1 mmol/L EDTA, 1% Triton X-100, 0.1% SDS and 1× proteinase inhibitor cocktail, and equal concentrations of denatured protein samples were

loaded on a 10% SDS–polyacrylamide gel. After electrophoresis, proteins were transferred to an Immobilon PVDF membrane, and immunoblot analysis was conducted by the standard method.

qPCR

For quantitative expression analysis of miRNA, total RNA was isolated from fresh-frozen prostatic tissues, cultured cells or xenograft tumors using TRIzol reagent (Life Technologies, Inc.). The miRNA level was measured using a TaqMan miRNA assay kit (Applied Biosystems, Foster City, CA) following the manufacturer's instruction and PCR amplification was carried out in a 7900HT Fast Real-time PCR system (Applied Biosystems). Real-time PCR was performed in triplicate for each sample. The abundance of miRNA was measured using C_t (threshold cycle) following the approach described previously (Schmittgen and Livak 2008). C_t was calculated by subtracting the C_t of *U6* RNA from the C_t of the miRNA tested. C_t was calculated by subtracting the C_t of a reference sample from the C_t of each sample. For clinical tissues, the reference sample is one CaP specimen tested that expresses the lowest level of miRNA. Fold change was generated using the equation 2^{-C_t} .

Methylation assay

Genomic DNA was isolated from prostate cell lines using standard procedures, and from CaP cell-enriched tumor tissues following our previous method (Shi et al 1996). Bisulfite modification of DNA (1.0 μ g) was carried out by using an EZ DNA methylation-direct kit (Zymo Research, Irvine, CA). For the prostate cell lines, the methylation of 5' CpG island regions was detected by MSP using the primers specific for either methylated or unmethylated DNA. The methylated CpG dinucleotides were validated using bisulfite DNA sequencing. Amplified CpG islands were separately cloned into pCR 2.1 vector using TOPO TA Cloning Kit (Invitrogen) and at least three clones per PCR product were picked up for DNA sequencing. In clinical prostate tissues, the methylation status was assessed by COBRA assay. COBRA primer-amplified PCR products were purified using Amicon 30 filter (Millipore), digested by BstUI and separated on a 3% agarose gel (1% agarose plus 2% NuSieve GTG low-melting temperature agaros). The primers used for MSP and COBRA were designed based on “The Li Lab” program (<http://www.urogene.org/methprimer/index1.html>). The primer sequences and amplified size are listed in SI Table 1.

Annexin V binding assay

MiR-124-induced apoptosis was analyzed using a FACS Annexin V assay kit (Trevigen, Inc., Gaithersburg, MD) following the protocol provided by the manufacturer. Briefly, cells were transfected with *miR-124* mimic and incubated for 4 days. The cells were harvested, washed once with PBS, re-suspended in 100 μ l of 1 \times Annexin V binding buffer containing 1 μ l of Annexin V-FITC conjugate and 10 μ l of propidium iodide solution and then incubated for 30 min at room temperature in the dark. Subsequently, 400 μ l 1 \times Annexin V binding buffer was added and samples analyzed on a FACScan flow cytometer. Data analysis was performed using FACScan software (Becton Dickinson).

Animal experiment

A lentiviral *miR-124* expression vector that expresses a ~500-base human pre-*miR-124-1* and the empty lentiviral vector were purchased from System Biosciences (SBI, Mountain View, CA). Pseudovirus production and cell transduction were performed following the manufacturer's protocol. The resulting 22Rv1-*miR-124* and 22Rv1-vector cells were selected through fluorescence-activated cell sorting (FACS) and were maintained in medium described above. Overexpression of *miR-124* in infected 22Rv1 cells was confirmed by qPCR. Animal studies were performed in male athymic nu/nu mice (4- to 6-week-old; Harlan Laboratories). Mice were injected subcutaneously with suspensions of 2×10^6 22Rv1-*miR-124* cells or 22Rv1-vector cells in a mixture (1:1 vol/vol) of culture medium and Matrigel (Becton Dickinson). Tumor growth was monitored and tumor dimensions were measured twice per week. Tumor volumes were calculated according to the formula: $\frac{1}{2}$ (length \times width \times height). Animal studies were performed according to protocols approved by the Animal Care and Use Committee at University of California, Davis.

Statistical analysis

Statistical analysis was performed using GraphPad Prism software 5.0. Student's t-test was used to analyze the difference in *miR-124* levels between different groups. Statistical significance was set up to $p < 0.05$ in each test.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by NIH 1R01CA92068 (R. deVere White), Department of Defense PCRP PC080488 (R. deVere White) and The LANIE Foundation. We thank Dr. Mukta. Webber for providing RWPE-1 cell line and Dr. John Rhim for pRNS-1-1 cell line. We also gratefully thank Dr. Melanie C. Bradnam for her editorial assistance and Stephanie Soares for critically reading the manuscript.

Supported by NCI grant (CA92069) and DOD Grant (PC080488).


References

- Agirre X, Vilas-Zornoza A, Jimenez-Velasco A, Martin-Subero JI, Cordeu L, Garate L, et al. Epigenetic silencing of the tumor suppressor microRNA Hsa-miR-124a regulates CDK6 expression and confers a poor prognosis in acute lymphoblastic leukemia. *Cancer Res.* 2009; 69:4443–4453. [PubMed: 19435910]
- Ando T, Yoshida T, Enomoto S, Asada K, Tatematsu M, Ichinose M, et al. DNA methylation of microRNA genes in gastric mucosae of gastric cancer patients: its possible involvement in the formation of epigenetic field defect. *Int J Cancer.* 2009; 124:2367–2374. [PubMed: 19165869]
- Baroukh N, Ravier MA, Loder MK, Hill EV, Bounacer A, Scharfmann R, et al. MicroRNA-124a regulates Foxa2 expression and intracellular signaling in pancreatic beta-cell lines. *J Biol Chem.* 2007; 282:19575–19588. [PubMed: 17462994]
- Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, Yendamuri S, et al. Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci U S A.* 2004; 101:2999–3004. [PubMed: 14973191]
- Cao X, Pfaff SL, Gage FH. A functional study of miR-124 in the developing neural tube. *Genes Dev.* 2007; 21:531–536. [PubMed: 17344415]

- Chang BL, Liu W, Sun J, Dimitrov L, Li T, Turner AR, et al. Integration of somatic deletion analysis of prostate cancers and germline linkage analysis of prostate cancer families reveals two small consensus regions for prostate cancer genes at 8p. *Cancer Res.* 2007; 67:4098–4103. [PubMed: 17483320]
- Chen CD, Welsbie DS, Tran C, Baek SH, Chen R, Vessella R, et al. Molecular determinants of resistance to antiandrogen therapy. *Nat Med.* 2004; 10:33–39. [PubMed: 14702632]
- Chen Y, Robles AI, Martinez LA, Liu F, Gimenez-Conti IB, Conti CJ. Expression of G1 cyclins, cyclin-dependent kinases, and cyclin-dependent kinase inhibitors in androgen-induced prostate proliferation in castrated rats. *Cell Growth Differ.* 1996; 7:1571–1578. [PubMed: 8930407]
- Davis BN, Hata A. Regulation of MicroRNA Biogenesis: A miRiad of mechanisms. *Cell Commun Signal.* 2009; 7:18. [PubMed: 19664273]
- de Vere White R, Meyers F, Chi SG, Chamberlain S, Siders D, Lee F, et al. Human androgen receptor expression in prostate cancer following androgen ablation. *European urology.* 1997; 31:1–6. [PubMed: 9032526]
- Frasca F, Rustighi A, Malaguarnera R, Altamura S, Vigneri P, Del Sal G, et al. HMGA1 inhibits the function of p53 family members in thyroid cancer cells. *Cancer Res.* 2006; 66:2980–2989. [PubMed: 16540646]
- Friedman RC, Farh KK, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 2009; 19:92–105. [PubMed: 18955434]
- Furuta M, Kozaki KI, Tanaka S, Arai S, Imoto I, Inazawa J. miR-124 and miR-203 are epigenetically silenced tumor-suppressive microRNAs in hepatocellular carcinoma. *Carcinogenesis.* 31:766–776. [PubMed: 19843643]
- Graff JR, Herman JG, Lapidus RG, Chopra H, Xu R, Jarrard DF, et al. E-cadherin expression is silenced by DNA hypermethylation in human breast and prostate carcinomas. *Cancer Res.* 1995; 55:5195–5199. [PubMed: 7585573]
- Jazdzewski K, Murray EL, Franssila K, Jarzab B, Schoenberg DR, de la Chapelle A. Common SNP in pre-miR-146a decreases mature miR expression and predisposes to papillary thyroid carcinoma. *Proc Natl Acad Sci U S A.* 2008; 105:7269–7274. [PubMed: 18474871]
- Kang GH, Lee S, Lee HJ, Hwang KS. Aberrant CpG island hypermethylation of multiple genes in prostate cancer and prostatic intraepithelial neoplasia. *J Pathol.* 2004; 202:233–240. [PubMed: 14743506]
- Koivisto P, Kononen J, Palmberg C, Tammela T, Hyytinen E, Isola J, et al. Androgen receptor gene amplification: a possible molecular mechanism for androgen deprivation therapy failure in prostate cancer. *Cancer Res.* 1997; 57:314–319. [PubMed: 9000575]
- Le MT, Teh C, Shyh-Chang N, Xie H, Zhou B, Korzh V, et al. MicroRNA-125b is a novel negative regulator of p53. *Genes Dev.* 2009; 23:862–876. [PubMed: 19293287]
- Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell.* 2005; 120:15–20. [PubMed: 15652477]
- Lim JT, Mansukhani M, Weinstein IB. Cyclin-dependent kinase 6 associates with the androgen receptor and enhances its transcriptional activity in prostate cancer cells. *Proc Natl Acad Sci U S A.* 2005; 102:5156–5161. [PubMed: 15790678]
- Lin DL, Tarnowski CP, Zhang J, Dai J, Rohn E, Patel AH, et al. Bone metastatic LNCaP-derivative C4-2B prostate cancer cell line mineralizes in vitro. *Prostate.* 2001; 47:212–221. [PubMed: 11351351]
- Lou W, Krill D, Dhir R, Becich MJ, Dong JT, Frierson HF Jr, et al. Methylation of the CD44 metastasis suppressor gene in human prostate cancer. *Cancer Res.* 1999; 59:2329–2331. [PubMed: 10344738]
- Lujambio A, Roperio S, Ballestar E, Fraga MF, Cerrato C, Setien F, et al. Genetic unmasking of an epigenetically silenced microRNA in human cancer cells. *Cancer Res.* 2007; 67:1424–1429. [PubMed: 17308079]
- Makeyev EV, Zhang J, Carrasco MA, Maniatis T. The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Mol Cell.* 2007; 27:435–448. [PubMed: 17679093]

- Matsuyama H, Pan Y, Yoshihiro S, Kudren D, Naito K, Bergerheim US, et al. Clinical significance of chromosome 8p, 10q, and 16q deletions in prostate cancer. *Prostate*. 2003; 54:103–111. [PubMed: 12497583]
- Mazar J, Khaitan D, DeBlasio D, Zhong C, Govindarajan SS, Kopanathi S, et al. Epigenetic regulation of microRNA genes and the role of miR-34b in cell invasion and motility in human melanoma. *PloS one*. 2011; 6:e24922. [PubMed: 21949788]
- Mirosevich J, Gao N, Gupta A, Shappell SB, Jove R, Matusik RJ. Expression and role of Foxa proteins in prostate cancer. *Prostate*. 2006; 66:1013–1028. [PubMed: 16001449]
- Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A*. 2008; 105:10513–10518. [PubMed: 18663219]
- Olive V, Jiang I, He L. mir-17-92, a cluster of miRNAs in the midst of the cancer network. *The international journal of biochemistry & cell biology*. 2010; 42:1348–1354. [PubMed: 20227518]
- Ostling P, Leivonen SK, Aakula A, Kohonen P, Makela R, Hagman Z, et al. Systematic analysis of microRNAs targeting the androgen receptor in prostate cancer cells. *Cancer Res*. 2011; 71:1956–1967. [PubMed: 21343391]
- Perinchery G, Bukurov N, Nakajima K, Chang J, Hooda M, Oh BR, et al. Loss of two new loci on chromosome 8 (8p23 and 8q12-13) in human prostate cancer. *Int J Oncol*. 1999; 14:495–500. [PubMed: 10024682]
- Pierantoni GM, Rinaldo C, Esposito F, Mottolese M, Soddu S, Fusco A. High Mobility Group A1 (HMGA1) proteins interact with p53 and inhibit its apoptotic activity. *Cell Death Differ*. 2006; 13:1554–1563. [PubMed: 16341121]
- Qi J, Nakayama K, Cardiff RD, Borowsky AD, Kaul K, Williams R, et al. Siah2-dependent concerted activity of HIF and FoxA2 regulates formation of neuroendocrine phenotype and neuroendocrine prostate tumors. *Cancer Cell*. 18:23–38. [PubMed: 20609350]
- Rauhala HE, Jalava SE, Isotalo J, Bracken H, Lehmusvaara S, Tammela TL, et al. miR-193b is an epigenetically regulated putative tumor suppressor in prostate cancer. *Int J Cancer*. 2010; 127:1363–1372. [PubMed: 20073067]
- Sadar MD. Small molecule inhibitors targeting the “achilles' heel” of androgen receptor activity. *Cancer Res*. 2011; 71:1208–1213. [PubMed: 21285252]
- Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008; 3:1101–1108. [PubMed: 18546601]
- Shenouda SK, Alahari SK. MicroRNA function in cancer: oncogene or a tumor suppressor? *Cancer Metastasis Rev*. 2009; 28:369–378. [PubMed: 20012925]
- Shi XB, Xue L, Ma AH, Tepper CG, Kung HJ, White RW. miR-125b promotes growth of prostate cancer xenograft tumor through targeting pro-apoptotic genes. *Prostate*. 71:538–549. [PubMed: 20886540]
- Shi XB, Bodner SM, deVere White RW, Gumerlock PH. Identification of p53 mutations in archival prostate tumors. Sensitivity of an optimized single-strand conformational polymorphism (SSCP) assay. *Diagn Mol Pathol*. 1996; 5:271–278. [PubMed: 8955619]
- Shi XB, Ma AH, Tepper CG, Xia L, Gregg JP, Gandour-Edwards R, et al. Molecular alterations associated with LNCaP cell progression to androgen independence. *Prostate*. 2004; 60:257–271. [PubMed: 15176055]
- Shi XB, Xue L, Yang J, Ma AH, Zhao J, Xu M, et al. An androgen-regulated miRNA suppresses Bak1 expression and induces androgen-independent growth of prostate cancer cells. *Proc Natl Acad Sci U S A*. 2007; 104:19983–19988. [PubMed: 18056640]
- Shi XB, Tepper CG, White RW. MicroRNAs and prostate cancer. *J Cell Mol Med*. 2008; 12:1456–1465. [PubMed: 18624768]
- Siegel R, Ward E, Brawley O, Jemal A. Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin*. 2012; 61:212–236. [PubMed: 21685461]
- Sinkkonen L, Hugenschmidt T, Berninger P, Gaidatzis D, Mohn F, Artus-Revel CG, et al. MicroRNAs control de novo DNA methylation through regulation of transcriptional repressors in mouse embryonic stem cells. *Nat Struct Mol Biol*. 2008; 15:259–267. [PubMed: 18311153]

- Takaha N, Hawkins AL, Griffin CA, Isaacs WB, Coffey DS. High mobility group protein l(Y): a candidate architectural protein for chromosomal rearrangements in prostate cancer cells. *Cancer Res.* 2002; 62:647–651. [PubMed: 11830513]
- Tamimi Y, van der Poel HG, Denyn MM, Umbas R, Karthaus HF, Debruyne FM, et al. Increased expression of high mobility group protein l(Y) in high grade prostatic cancer determined by in situ hybridization. *Cancer Res.* 1993; 53:5512–5516. [PubMed: 8221692]
- Tepper CG, Boucher DL, Ryan PE, Ma AH, Xia L, Lee LF, et al. Characterization of a novel androgen receptor mutation in a relapsed CWR22 prostate cancer xenograft and cell line. *Cancer Res.* 2002; 62:6606–6614. [PubMed: 12438256]
- Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A.* 2006; 103:2257–2261. [PubMed: 16461460]
- Walton TJ, Li G, Seth R, McArdle SE, Bishop MC, Rees RC. DNA demethylation and histone deacetylation inhibition co-operate to re-express estrogen receptor beta and induce apoptosis in prostate cancer cell-lines. *Prostate.* 2008; 68:210–222. [PubMed: 18092350]
- Woodson K, Hanson J, Tangrea J. A survey of gene-specific methylation in human prostate cancer among black and white men. *Cancer Lett.* 2004; 205:181–188. [PubMed: 15036650]
- Yamakuchi M, Ferlito M, Lowenstein CJ. miR-34a repression of SIRT1 regulates apoptosis. *Proc Natl Acad Sci U S A.* 2008; 105:13421–13426. [PubMed: 18755897]
- Yu X, Gupta A, Wang Y, Suzuki K, Mirosevich J, Orgebin-Crist MC, et al. Foxa1 and Foxa2 interact with the androgen receptor to regulate prostate and epididymal genes differentially. *Ann N Y Acad Sci.* 2005; 1061:77–93. [PubMed: 16467259]

**Figure 1.**

MiR-124 negatively regulates androgen receptor (AR). **A**) Western blot analysis of the AR expression in C4-2B CaP cells that were treated with 100 nM of synthetic *miR-124*, *miR-130*, *miR-384* or *miR-506*. The numbers under the gels are the fold changes of AR protein relative to untreated C4-2B cells (untreat). Fold changes were calculated by scanning the AR bands and normalizing for β -actin bands. The upper portion is a schema of the first 436 bases of the AR 3' UTR (based on the RefSeq NM_000044.2). The digital numbers indicate the predicted three miRNA-binding sites targeted by six potential AR-targeting miRNAs. **B**) Western blot analysis of the expression of AR and PSA in LNCaP and C4-2B cells treated with 100 nM of *miR-124* mimic. Both mock and miRNA-NC were used as controls. **C**) Quantitative PCR (qPCR) assay of *miR-125b* levels in C4-2B and *cds2* cells treated with 100 nM of *miR-124* mimic or miR-NC. The values shown as mean \pm SE (n=3) are from three independent experiments performed in triplicate. **D**) Luciferase analysis in C4-2B cells. The assay was repeated three times with each assay being performed in four wells, and similar results were obtained each time. The representative results are shown as mean \pm SD (n = 4). The percentage represents enzyme activity in 100 nM *miR-124* mimic-transfected cells relative to that in 100 nM miR-NC-transfected cells. RLU, relative luciferase unit. BS3'UTR, AR 3'UTR fragment lacking the *miR-124* binding site.




Figure 2. Quantitative PCR (qPCR) assays of *miR-124* expression levels in CaP cells. **A)** *MiR-124* abundance in two benign prostate cell lines (pRNS-1-1 and RWPE-1) and five CaP cell lines (22Rv1, LNCaP, LAPC-4, cds2 and C4-2B). **B)** *MiR-124* expression levels in 19 benign prostatic hyperplasia (BPH) tissues, 44 primary CaPs, 6 lymph node metastases and 10 castration-resistant tumors. **C)** *MiR-124* levels in 18 matched benign and malignant prostate samples. The levels of *miR-124* of five pairs of prostate tissues (5, 8, 9, 11 & 16) were assayed using Northern blot analysis and similar results were obtained. The Northern blot results are shown in SI Figure 2B. “*”, no statistical difference ($p > 0.05$) between BPH and CaP tissues. In **A**, **B** and **C**, qPCR assays were repeated three times with each assay being performed in triplicate, and similar results were obtained each time. The values are shown as mean \pm SE ($n=3$).




Figure 3.

The expression levels of androgen receptor (AR) and *miR-124* in four matched prostate tissues (pt-1 to pt-4). **A**) Immunohistochemical analysis of the AR protein in four human CaP samples (*right*) and their benign prostate tissue (*left*). **B**) qPCR detection of the abundance of *miR-124* in both BPH and CaP tissues. The values are shown as mean \pm SE from three independent experiments.




Figure 4. Methylation of *miR-124* in CaP cells. **A)** Expression levels of *miR-124* in 22Rv1, C4-2B and cds2 CaP cell lines before and after treatment with 50 μ M of Aza. Results are expressed as fold change of *miR-124* relative to the untreated control. The assay was repeated three times with each assay being performed in three wells, and similar results were obtained each time. The representative results are shown as mean \pm SD ($n = 3$). **B)** Western blot analysis of the AR expression in LNCaP and C4-2B cells treated with 50 μ M of Aza. **C)** Methylation-specific PCR (MSP) assay of the 5' CpG islands of *miR-124-1*, *miR-124-2* and *miR-124-3* in seven prostatic cell lines: two benign lines (pRNS-1-1 and RWPE-1) and 5 malignant lines (22Rv1, LNCaP, LAPC-4, cds2 and C4-2B). **D)** Bisulfite sequencing analysis of *miR-124*

CpG island methylation in RWPE-1 cell line and five CaP cell lines. The top schematic view represents individual amplified 5' DNA fragments of three *miR-124* genes, which are located at -1895 to -1655 upstream of *pre-miR-124-1*, -1456 to -1188 of *pre-miR-124-2* and -1465 to -1246 of *pre-miR-124-3*. The vertical bars denote individual CpG dinucleotides. Methylation profiles of *miR-124* CpG island fragments in six cell lines tested were demonstrated in the bottom of the schema. CpGs are represented by open circles if unmethylated and by black circles if methylated. Each row exhibits methylated CpGs from at least three clones. The numbers on the right of each row are the percentage of methylated CpG dinucleotides. **E)** COBRA analysis of methylation of *miR-124-1*, *miR-124-2* and *miR-124-3* in 9 BPH tissues and 14 CaP samples. Purified PCR product was digested with *Bst*U1 that cuts methylated CGCG. The black arrows indicate the BPH samples having detectable methylation, and the white arrows indicate the CaP samples without methylation.





Figure 5. *MiR-124* induces apoptosis and upregulates p53. **A)** Annexin V assay of apoptosis. LNCaP and C4-2B cells were treated with 100 nM *miR-124* mimic or miR-NC for 4 days and stained with Annexin V and propidium iodide. Both early and late apoptotic cells are combined. The values are shown as mean \pm SE from three independent experiments. **B)** Western blotting analysis of p53 expression in 100 nM *miR-124* mimic-transfected C4-2B cells. Doxorubicin (Dox)-treated cells were used as positive control. **C)** Western blotting analysis of caspase 3 (Cas-3) in 100 nM *miR-124* mimic-transfected C4-2B cells. In both **B** and **C**, the numbers under the gels are the fold changes of p53 and Cas-3 in *miR-124* mimic-treated C4-2B cells relative to miR-NC-treated cells.

**Figure 6.**

Inhibition of xenograft tumor growth by *miR-124*. **A**). Five nude mice per group were injected subcutaneously with 2×10^6 22Rv1-*miR-124* cells, 22Rv1-vector cells, or untreated 22Rv1 cells. The growth of xenograft tumors were measured twice per week. Each time point represents mean \pm SD of five independent values (mm³). **B**) qPCR assay of *miR-124* level in xenograft tumors. The assay was repeated twice and similar results were obtained each time. The representative results are shown as mean \pm SD (n = 3). **C**) Western blot assay of the AR expression in two 22Rv1-*miR-124* tumors that express the full-length AR and a truncated AR (Tepper et al 2002).