



Correction

Correction: Cesarean Section and Rate of Subsequent Stillbirth, Miscarriage, and Ectopic Pregnancy: A Danish Register-Based Cohort Study

The *PLOS Medicine* Staff

There is an error in Table 2. The headings under ‘Previous Delivery’ are labeled incorrectly; ‘Vaginal’ and ‘Cesarean’ should be in the reverse order, as shown below.

Table 2. Causes of stillbirth by mode of delivery (1982–1996).

Type of Stillbirth	Previous Delivery	
	Cesarean	Vaginal
Explained stillbirth (n = 675)		
Antenatal complications	79 (18.8)	342 (81.2)
Complications of delivery	35 (24.5)	108 (75.5)
Congenital malformations	10 (11.2)	79 (88.8)
Maternal illness	5 (26.3)	14 (73.7)
Injury to mother	0 (0.00)	3 (100)
Unexplained stillbirth (n = 186)	31 (16.7)	155 (83.3)
Total	160 (18.6)	701 (81.4)

Data are *n* (percent). Type of stillbirth was classified as outlined by King-Hele et al. [38].
doi:10.1371/journal.pmed.1001670.t002

Reference

- O'Neill SM, Agerbo E, Kenny LC, Henriksen TB, Kearney PM, et al. (2014) Cesarean Section and Rate of Subsequent Stillbirth, Miscarriage, and Ectopic Pregnancy: A Danish Register-Based Cohort Study. *PLoS Med* 11(7): e1001670. doi:10.1371/journal.pmed.1001670

Citation: The *PLOS Medicine* Staff (2014) Correction: Cesarean Section and Rate of Subsequent Stillbirth, Miscarriage, and Ectopic Pregnancy: A Danish Register-Based Cohort Study. *PLoS Med* 11(7): e1001707. doi:10.1371/journal.pmed.1001707

Published: July 28, 2014

Copyright: © 2014 The *PLOS Medicine* Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.