

## Review article

<http://dx.doi.org/10.6065/apem.2014.19.2.57>  
Ann Pediatr Endocrinol Metab 2014;19:57-68

# Congenital hyperinsulinism: current status and future perspectives

Tohru Yorifuji, MD, PhD

Department of Pediatric  
Endocrinology and Metabolism,  
Children's Medical Center, Osaka  
City General Hospital, Osaka, Japan

The diagnosis and treatment of congenital hyperinsulinism (CHI) have made a remarkable progress over the past 20 years and, currently, it is relatively rare to see patients who are left with severe psychomotor delay. The improvement was made possible by the recent developments in the understanding of the molecular and pathological basis of CHI. Known etiologies include inactivating mutations of the  $K_{ATP}$  channel genes (*ABCC8* and *KCNJ11*) and *HNF4A*, *HNF1A*, *HADH*, and *UCP2* or activating mutations of *GLUD1*, *GCK*, and *SLC16A1*. The understanding of the focal form of  $K_{ATP}$  channel CHI and its detection by  $^{18}F$ -fluoro-L-DOPA positron emission tomography have revolutionized the management of CHI, and many patients can be cured without postoperative diabetes mellitus. The incidence of the focal form appears to be higher in Asian countries; therefore, the establishment of treatment systems is even more important in this population. In addition to diazoxide or long-term subcutaneous infusion of octreotide or glucagon, long-acting octreotide or lanreotide have also been used successfully until spontaneous remission. Because of these medications, near-total pancreatectomy is less often performed even for the diazoxide-unresponsive diffuse form of CHI. Other promising medications include pasireotide, small-molecule correctors such as sulfonylurea or carbamazepine, GLP1 receptor antagonists, or mammalian target of rapamycin inhibitors. Unsolved questions in this field include the identification of the remaining genes responsible for CHI, the mechanisms leading to transient CHI, and the mechanisms responsible for the spontaneous remission of CHI. This article reviews recent developments and hypothesis regarding these questions.

**Keywords:** Hyperinsulinism, Congenital, Hypoglycemia

## Introduction

Congenital hyperinsulinism (CHI) is the most common cause of persistent hypoglycemia in infancy, and severe hypoglycemia in infancy can cause permanent brain damage<sup>1-4</sup>; therefore, optimal management is extremely important. In the past, our armamentarium against severe CHI was very limited. With only diazoxide and near-total pancreatectomy as available options, many patients were left with psychomotor delay or epilepsy. Even worse, a number of patients developed postoperative insulin-dependent diabetes mellitus<sup>3,5,6</sup>.

Over the past 20 years, however, remarkable progress has been made in the diagnosis and management of CHI, which has directly translated into improved neurological outcomes for patients<sup>3,7</sup>. This improvement in the understanding of the pathogenesis of CHI and the development of diagnostic modalities have helped in deciding the optimal management strategy for each patient<sup>8-10</sup>. However, the situation is still far from ideal. Several unsolved questions and unmet needs remain.

In this review, I first discuss the diagnostic criteria and the practical treatment goals for CHI, which are the prerequisites for all subsequent management. Then, after a brief introduction of

Received: 4 April, 2014  
Accepted: 14 April, 2014

### Address for correspondence:

Tohru Yorifuji, MD, PhD  
Department of Pediatric  
Endocrinology and Metabolism,  
Children's Medical Center, Osaka  
City General Hospital, 2-13-22  
Miyakojima-Hondori, Miyakojima,  
Osaka 534-0021, Japan  
Tel: +81-6-6929-1221  
Fax: +81-6-6929-1090  
E-mail: t-yorifuji@hospital.city.osaka.jp

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ISSN: 2287-1012(Print)  
ISSN: 2287-1292(Online)

the mechanism of glucose-induced insulin secretion, I review the current status of the understanding and management of CHI. Finally, I list some of the unsolved questions in this field and introduce key findings that may guide us in the future.

## Diagnostic criteria and treatment goals for CHI

### 1. Diagnosis of CHI (Table 1)

To diagnose hyperinsulinemic hypoglycemia (HI), physicians rely both on clinical clues to identify hyperinsulinism and on laboratory tests to prove hyperinsulinemia.

#### 1) Clinical clues to suspect HI

The presence of HI may be suspected even when the patient is still in an emergency room by asking three questions: When did hypoglycemia develop after the last meal? Does the patient respond to glucagon injection? What is the amount of glucose infusion needed to keep the patient euglycemic?

#### (1) When did hypoglycemia develop after the last meal?

Euglycemia is maintained by a balance between hepatic glucose output and peripheral uptake induced by insulin. Hepatic glucose output is determined by three factors: food absorption, glycogenolysis, and gluconeogenesis. Fig. 1 shows the duration of glucose production by each of these mechanisms. In disorders of glycogenolysis, the patient typically becomes hypoglycemic after 4–5 hours of fasting. Similarly, in disorders of gluconeogenesis, the patient typically develops




Fig. 1. Glucose source during fasting.

Table 1. Diagnosis of hyperinsulinemic hypoglycemia

	Serum insulin at hypoglycemia, pmol/L ( $\mu$ U/mL)	Glucose infusion rate to maintain euglycemia (mg/kg/min)	Glycemic response to glucagon, mmol/L (mg/dL)	Free fatty acid/ketone bodies	Ref.
1	ND	>8	>1.5 (27)	ND	10
2	Any detectable level	>10 (neonate) >7 (5 years old), >4 (adults)	>1.7 (30)	Inappropriately low fatty acids and ketones	9
3	>6.95 (1)	ND	>2–3 (36–54)	Negative ketone bodies in urine/plasma	13
4	Any detectable level	>8	ND	ND	8
Proposed criteria	>20.84 (3)	>8 (neonates), >3 (adults), in-between (children)	>2 (36)	3-hydroxybutyrate < 1.3 mmol/L, FFA < 1 mmol/L	

ND, not described; FFA, free fatty acid.

hypoglycemia after an overnight fast. When the patient has hyperinsulinemia, hypoglycemic episodes can occur at any time point, sometimes even at 2 hours after the last meal.

#### (2) Does the patient respond to glucagon injection?

When hypoglycemia is caused by a defect in glycogenolysis, the patient does not respond to intramuscular/intravenous injection of glucagon, which stimulates glycogenolysis. Similarly, when a patient has a defect in gluconeogenesis, by the time the patient becomes hypoglycemic, the glycogen storage in the liver should have been exhausted; therefore the patient does not respond to glucagon either. Only when the patient has hyperinsulinemia can hepatic glycogen be mobilized by glucagon, and glycemic response (>1.7–2.0 mmol/L) is seen.

#### (3) What is the amount of glucose infusion needed to keep the patient euglycemic?

When hypoglycemia is caused by etiologies other than hyperinsulinism, euglycemia should be maintained by providing the amount of intravenous glucose that corresponds to the normal hepatic (or possibly renal) glucose output: 4–6 mg/kg/min in neonates, 1–2 mg/kg/min in adults, and intermediate values in older children. When euglycemia cannot be maintained by these amounts of continuous glucose infusion, clinicians may suspect the presence of hyperinsulinemia.

#### 2) Laboratory evidence of hyperinsulinism

##### (1) Insulin at hypoglycemia

HI is diagnosed by demonstrating inappropriately elevated insulin in the presence of hypoglycemia (<2.5 mmol/L, 45 mg/dL). However, it is often difficult to prove hyperinsulinemia by a critical sample taken during a hypoglycemic event<sup>11,12</sup>. In addition, the term “inappropriately elevated” insulin level is not precisely defined: some authors suggest that any detectable level of insulin is abnormal<sup>8,9</sup>, whereas others propose different cutoffs<sup>13</sup>. With regard to the insulin levels during hypoglycemia, “any detectable level” is probably an overstatement because it may suggest but not prove HI. The cutoffs depend on the sensitivity of the particular assay as well as on the insulin sensitivity of each patient. In our own series of 94 confirmed Asian patients with CHI, the insulin at hypoglycemia ranged

8.75–1,250 pmol/L (1.26–180  $\mu$ U/mL) with a median of 73.3 pmol/L (10.55  $\mu$ U/mL; unpublished data). In contrast, insulin levels during hypoglycemic events in patients without HI ranged from undetectable to 43.1 pmol/L (6.2  $\mu$ U/mL) while the detection limit was  $>2.1$  pmol/L (0.3  $\mu$ U/mL). Clearly, these values overlap.

(2) Relatively low free fatty acid and ketone bodies at hypoglycemia

Insulin inhibits lipolysis; therefore, low free fatty acid and ketone bodies during hypoglycemia are also used as diagnostic adjuncts. In normal infants (0–24 months of age), blood 3-hydroxybutyrate and free fatty acid levels after a 20-hour fast are 3.11 mmol/L (range, 1.29–4.34 mmol/L) and 2.15 mmol/L (range, 1.03–3.24 mmol/L), respectively<sup>14</sup>. These values can be used to set cutoffs for HI. In our series of 207 cases with confirmed CHI, the highest 3-hydroxybutyrate at hypoglycemia was 0.44 mmol/L (unpublished data).

3) Proposed diagnostic criteria to suspect HI

It is difficult to set definitive diagnostic criteria for HI. Several authors propose different cutoff values to diagnose HI (Table 1), but these criteria should be regarded as suggestive and not necessarily diagnostic. The bottom of Table 1 includes proposed criteria that strongly suggest the presence of HI in children.

## 2. Treatment goals of CHI

The goals of HI treatment are to prevent neurological sequelae of hypoglycemia. Factors that could affect neurological outcomes include age, comorbid conditions, severity of the initial episode, and duration and frequency of subsequent

hypoglycemic episodes<sup>15–18</sup>. Therefore, the treatment goals should be individualized. Currently, blood glucose  $>3.33$ – $3.89$  mmol/L (60–70 mg/dL) is the most commonly recommended target for HI treatment<sup>10,11,13</sup>.


## Insulin secretion and the ATP-sensitive potassium channel

### 1. Glucose induced insulin secretion (the GSIS pathway) (Fig. 2)

In pancreatic  $\beta$ -cells, extracellular glucose is transported into the cytoplasm by the action of glucose transporter (GLUT2). The glucose is then phosphorylated by glucokinase. Glucokinase is not easily saturated by the physiological range of intracellular glucose, and is not inhibited by its end product glucose-6-phosphate. Therefore, it serves as the fuel gauge of the  $\beta$ -cells. Glucose-6-phosphate is then metabolized via glycolysis, the Krebs cycle, and oxidative phosphorylation to generate ATP. An increased ATP/ADP ratio within  $\beta$ -cells leads to the closure of the ATP-sensitive potassium channel ( $K_{ATP}$  channel), which causes depolarization of the cell membrane and opening of the voltage-gated calcium channel. The resulting influx of calcium ions then causes fusion of the insulin secretory granules with the cell membrane and secretion of insulin<sup>19</sup>.

### 2. The $K_{ATP}$ channel

The  $K_{ATP}$  channel is an octameric structure composed of four molecules of pore-forming Kir6.2 and four molecules of SUR1 that surround the pore and regulate the channel activity.



**Fig. 2.** The glucose-induced insulin secretion pathway. GLUT2, glucose transporter 2; GCK, glucokinase; G6P, glucose 6-phosphate; MCT1, monocarboxylate transporter 1; GDH, glutamate dehydrogenase; HADH, L-3-hydroxyacyl-coenzyme A dehydrogenase;  $\alpha$ -KG,  $\alpha$ -ketoglutarate; Ins, insulin.

Intracellular ATP binds to Kir6.2 molecules to inhibit the channel activity, whereas MgADP binds to SUR1 to activate the channel. Therefore, channel activity is controlled by the ATP/ADP ratio within the cells.

In the endoplasmic reticulum, Kir6.2 and SUR1 associate with each other to form the channel that is transferred to the Golgi apparatus and to the cell surface. If Kir6.2 and SUR1 do not associate with each other, they cannot escape the endoplasmic reticulum and are degraded there.

Kir6.2 and SUR1 are encoded by *KCNJ11* (1 exon) and *ABCC8* (39 exons), respectively. These genes are located side-by-side with close proximity on 11p15.1.

## Known etiologies of CHI (Table 2)

### 1. Transient and persistent CHI

There are two main types of CHI: transient CHI, which usually develops soon after birth and resolves spontaneously within the first 3–4 weeks of life and persistent CHI, which can develop later in life as well as in the neonatal period, and lasts longer. The distinction between transient and persistent CHI is not possible on the basis of laboratory test results. In our national survey in Japan, only shorter gestational age and lighter birth weight were predictors of transient CHI<sup>20</sup>. The incidence of persistent CHI is generally estimated as 1 in 50,000 live births<sup>9</sup> although the incidence could be higher in certain populations (e.g., 1 in 2,500 births in Saudi Arabia). On the contrary, the incidence of transient CHI is much higher. In the national survey in Japan, the incidence of transient CHI (1 in 17,000 births) was approximately twice as high as that for persistent CHI (1 in 35,400 births)<sup>20</sup>.

#### 1) Transient CHI

Transient CHI is believed to be caused mainly by nongenetic factors, e.g., small size for the infant's gestational age or stressful perinatal conditions such as cardiopulmonary disorders. An important exception is the monoallelic inactivating mutation in

*HNF4A*<sup>21–23</sup>. Unlike other patients with transient CHI, patients with *HNF4A* mutations are often born large for gestational age. Importantly, a fraction of these patients develop a form of dominantly inherited diabetes, maturity-onset diabetes of the young type 1 (MODY1), later in life and therefore should be followed up after resolution of CHI (21–23). Because *HNF1A* is in the same pathway with *HNF4A* and its mutation is the cause of MODY3, researchers checked for mutations in *HNF1A* in patients with transient CHI, and indeed found some patients with mutations in *HNF1A*<sup>24,25</sup>.

#### 2) Persistent CHI

In contrast, persistent CHI is believed to have genetic etiologies. However, even with the most comprehensive analysis, the responsible genes can be identified in only 53% of diazoxide-responsive CHI patients<sup>26</sup> although in unresponsive patients,  $K_{ATP}$  channel mutations could be identified in most (87.6%–88%) cases<sup>26,27</sup>.

### 2. Causes of persistent CHI

Table 2 lists known causes of CHI. The most common of these are inactivating mutations in one of the  $K_{ATP}$  channel genes, *ABCC8* or *KCNJ11* ( $K_{ATP}$ -CHI). The second most common is an activating mutation of glutamate dehydrogenase (GDH-CHI). Others are relatively rare. When confined to families with consanguinity, inactivating mutations in L-3-hydroxyacyl-coenzyme A dehydrogenase (HADH-CHI) are the most common cause<sup>26,27</sup>.

#### 1) $K_{ATP}$ -CHI

Three distinct subtypes of  $K_{ATP}$ -CHI are known:

(1) Recessively-inherited  $K_{ATP}$ -CHI: Recessive  $K_{ATP}$ -CHI is caused by biallelic mutations in one of the  $K_{ATP}$  channel genes. This is the most severe form of  $K_{ATP}$ -CHI, and all  $\beta$ -cells in the pancreas present in abnormal (diffuse) form. Pathologically, recessive  $K_{ATP}$ -CHI is characterized by large  $\beta$ -cells with abnormally enlarged nuclei<sup>28</sup>.

**Table 2. Genetic causes of congenital hyperinsulinism**

Gene	Protein	Chromosome	Inheritance	Note
$K_{ATP}$ channel			AR, AD, Focal	Diazoxide unresponsive Usher CHI (contiguous deletion)
<i>ABCC8</i>	SUR1	11p15.1		
<i>KCNJ11</i>	Kir6.2	11p15.1		
<i>GLUD1</i>	Glutamate dehydrogenase	10q23.3	AD	Hyperammonemia
<i>GCK</i>	Glucokinase	7p15	AD	Diffuse/focal?
<i>HADH</i>	L-3-hydroxyacyl-coenzyme A dehydrogenase	4q22–q26	AR	
<i>UCP2</i>	Uncoupling protein 2	11q13	AD	
<i>SLC16A1</i>	Monocarboxylate transporter 1	1p12	AD	Exercise induced HI
<i>HNF4A</i>	Hepatocyte nuclear factor 4a	20q13.12	AD	Transient/persistent macrosomia
<i>HNF1A</i>	Hepatocyte nuclear factor 1a	12q24.2	AD	Variable onset glycogenosis renal tubular dysfunction

AR, autosomal recessive; AD, autosomal dominant; CHI, congenital hyperinsulinism; HI, hyperinsulinism

(2) Dominantly inherited  $K_{ATP}$ -CHI: Dominant  $K_{ATP}$ -CHI is caused by a monoallelic mutation in the  $K_{ATP}$  channel genes. The presentation is usually relatively milder, and patients often respond to diazoxide<sup>29)</sup> although there are some refractory cases<sup>30)</sup>.

### (3) Focal $K_{ATP}$ -CHI:

#### i) Pathogenesis

In patients with focal  $K_{ATP}$ -CHI, abnormal  $\beta$ -cells are confined to a restricted region in the pancreas. In close proximity with the  $K_{ATP}$  channel genes at chromosome 11p15.1, an imprinted region at 11p15.5 harbors maternally expressed tumor suppressors, *H19* and *CDKN1c*, and a paternally expressed growth factor gene, *IGF2*. The focal lesion arises in an individual with a paternally inherited, monoallelic mutation in one of the  $K_{ATP}$  channel genes. When segmental paternal uniparental disomy occurs as a somatic mutation during the development of the pancreas, that particular cell loses  $K_{ATP}$  channel activity. At the same time, the tumor-suppressor activities of *H19* and *CDKN1c* are lost, and the activity of *IGF2* is doubled. This leads to a growth advantage for the abnormal  $\beta$ -cells and eventually leads to formation of a focal lesion<sup>31-34)</sup>. Histologically, the focal lesion is characterized by the presence of large  $\beta$ -cells with enlarged nuclei similar to those of the diffuse lesion, and  $\beta$ -cells outside the focus have normal histology<sup>35-37)</sup>.

#### ii) Clinical implication

Although 96.2% of focal lesions are unresponsive to diazoxide<sup>3)</sup>, when the focal lesion is identified preoperatively, partial pancreatectomy can cure the patient without postoperative complications. Therefore, the identification and localization of focal lesions are extremely important. However, because they are generated during the normal organogenesis of the pancreas, they cannot usually be detected using conventional imaging modalities such as computed tomography (CT), magnetic resonance imaging, and angiography. The focal lesions can be preoperatively identified using molecular analysis and <sup>18</sup>F-fluoro-L-DOPA positron emission tomography (<sup>18</sup>F-DOPA PET) scans, thereby enabling surgeons to plan the surgical procedure and to find the lesion intraoperatively.

#### iii) <sup>18</sup>F-DOPA PET scan

<sup>18</sup>F-DOPA is incorporated into  $\beta$ -cells by DOPA-decarboxylase, which is abundant in  $\beta$ -cells. Following the initial description of its role in identifying the focal lesion<sup>38)</sup>, its usefulness has been reported in a number of publications<sup>39,40)</sup>. <sup>18</sup>F-DOPA PET detects focal lesions as small as 5 mm and is better preformed as PET-CT. However, there are some challenges in interpreting the results. First, artifact uptakes tend to be found in the head of the pancreas because the head has a larger mass than the rest of the pancreas and because <sup>18</sup>F-DOPA is excreted into the bile duct. Second, <sup>18</sup>F-DOPA PET does not necessarily show the exact size of the lesion, especially when the lesion extends so-called tentacles out of the central lesion. These problems appeared more pronounced in our experience in Japan<sup>41)</sup>.

#### iv) Epidemiology of focal $K_{ATP}$ -CHI

Previously, it was reported that approximately 40%–60% of surgically treated patients had focal CHI<sup>31,42,43)</sup>. However, recent molecular analysis has revealed a racial disparity in the frequency of paternally inherited monoallelic mutation in  $K_{ATP}$ -CHI patients. For example, the percentage of patients with a paternally inherited monoallelic  $K_{ATP}$ -channel mutation is 20.6% in Spain<sup>44)</sup>, 33% in Norway<sup>45)</sup>, 84.2% in Japan<sup>46)</sup>, 37.7% in Germany<sup>47)</sup>, 58% in China<sup>48)</sup>, and 25% in the UK<sup>27)</sup>. Obviously, these figures could be affected by ascertainment biases and by small sample sizes. In Japan, we have presently identified 46 patients with  $K_{ATP}$ -CHI, and 37 (80.4%) have paternal mutations. Therefore, combined with the report from China<sup>48)</sup>, it appears that Asians tend to have a higher frequency of paternally inherited mutation, suggesting the presence of focal CHI. The identification of focal CHI, therefore, is even more important for Asian patients.

#### v) Discordant molecular and <sup>18</sup>F-DOPA PET results

Not all patients with a paternally inherited  $K_{ATP}$  channel mutation have focal uptake by <sup>18</sup>F-DOPA PET, and some of these actually show diffuse histology. For example, Banerjee et al.<sup>49)</sup> reported that 31% of patients with a paternal monoallelic mutation showed diffuse uptake on <sup>18</sup>F-DOPA PET. Similarly, Kapoor et al.<sup>27)</sup> reported that 26% of patients with paternal mutation appeared to have a diffuse lesion on PET. These results may indicate that not all maternal mutations were identified by the molecular analysis. However, when monoallelic mutations were identified in patients with  $K_{ATP}$ -CHI, the majority were paternal mutations: 79.3% and 65% in two UK series<sup>27,49)</sup>, 84.2% in Japan<sup>46)</sup>, and 83.3% in Norway<sup>45)</sup>. Selectively missing the maternal allele during the molecular analysis is statistically highly unlikely. Another possibility, therefore, is that these patients have unusually scattered focal CHI that resembles true diffuse CHI. Further analysis is necessary to address this problem.

#### 2) Non- $K_{ATP}$ channel CHI

Most other persistent CHI are caused by excessive anaplerosis (replenishment of metabolic intermediate) into the GSIS pathway. With the exception of *GCK* and *SLC16A1* mutations, these non- $K_{ATP}$ -channel CHI are usually responsive to diazoxide.

#### (1) GDH

GDH is encoded by *GLUD1* at chromosome 10q23.3. GDH mediates conversion of glutamate to  $\alpha$ -ketoglutarate and ammonia, which is one of the major anaplerotic pathways in the Krebs cycle. An activating mutation in the *GLUD1* gene then supplies excess  $\alpha$ -ketoglutarate into the Krebs cycle. The resulting overproduction of ATP causes CHI associated with hyperammonemia (hyperinsulinism–hyperammonemia syndrome)<sup>50)</sup>. Both dominantly inherited and sporadic cases have been reported. Hypoglycemia is responsive to diazoxide, but hyperammonemia is resistant to conventional treatment.

Because GDH is allosterically activated by leucine, GDH-CHI presents with the typical leucine-sensitive CHI.

### 3) HADH

HADH—previously known as short-chain hydroxyacyl CoA dehydrogenase—is encoded by *HADH* at 4q22–q26. HADH functions in the mitochondrial matrix to catalyze the oxidation of straight-chain 3-hydroxyacyl-CoAs as part of the  $\beta$ -oxidation pathway. Unlike other proteins in the  $\beta$ -oxidation pathway, HADH is abundant in pancreatic  $\beta$ -cells and inhibits the activity of GDH. Biallelic HADH mutation then causes activation of GDH and hyperinsulinemia<sup>51–53</sup>.

### 4) Glucokinase

Glucokinase is encoded by *GCK* at chromosome 7p15. Patients with GCK-HI have an activating mutation in *GCK*. This leads to overactivity in the GSIS pathway and oversecretion of insulin<sup>54–56</sup>. On the contrary, inactivating monoallelic mutation is a cause of MODY2 or GCK-MODY<sup>57</sup>. Recently, a somatic activating mutation in *GCK* has been proposed as a cause of a novel form of diazoxide-responsive focal CHI<sup>58</sup>.

### 5) Uncoupling protein 2

Mitochondrial uncoupling protein 2 (UCP2) is encoded by *UCP2* at chromosome 11q13. UCP2 protein leaks protons across the inner mitochondrial membrane, thereby uncoupling the oxidative phosphorylation from ATP generation. Patients with a monoallelic mutation in *UCP2* have excessive ATP production leading to HI<sup>59</sup>.

### 6) Monocarboxylate transporter 1

Monocarboxylate transporter 1 (MCT1) encoded by *SLC16A1* at 1p12 is in the cell membrane and transports extracellular lactate and pyruvate into the cells. In pancreatic  $\beta$ -cells, the activity of MCT1 is normally suppressed to prevent lactate influx during exercise. In patients with exercise-induced HI, researchers identified mutations in the promoter of *SLC16A1* that activate the transporter. During exercise, extracellular lactate is fluxed into the  $\beta$ -cells and is converted to pyruvate, which fuels the GSIS pathway. The resulting increase in ATP production leads to HI<sup>60</sup>.

## 3. Syndromic CHI

A variety of syndromes are reportedly associated with CHI. Because CHI is not a common feature of these syndromes, some of these associations may be coincidental. Nevertheless, CHI is frequently associated with Beckwith-Wiedemann syndrome, Sotos syndrome, Kabuki syndrome, Costello syndrome, mosaic Turner syndrome, or congenital deficiency of glycosylation<sup>61</sup>. In these syndromes, the association may have biological implications more than by-chance association. Of note is the Usher-CHI syndrome<sup>62,63</sup> in which CHI is associated with the symptoms of Usher syndrome, i.e., hearing loss and retinitis pigmentosa. This association is caused by biallelic deletions

encompassing the  $K_{ATP}$  channel genes at 15p11 and the adjacent *USH1C* gene at 11p14.3, which is responsible for Usher syndrome.

## Current treatment strategies

Current treatment strategies are summarized in Table 3 and are reviewed below.

### 1. Diazoxide

Diazoxide is a benzothiazine derivative that acts on the SUR1 subunit of the  $K_{ATP}$  channel, activating it. Diazoxide is used orally in three divided doses (5–15 mg/kg/day) and is effective for a variety of CHI subtypes<sup>64</sup>. However, it is generally ineffective for the most severe, neonatal-onset, recessive, and focal forms of  $K_{ATP}$ -CHI. Dominant  $K_{ATP}$  channel CHI often responds to diazoxide, although some unresponsive cases have been reported<sup>32</sup>. Hypertrichosis occurs in most patients and could be a serious concern. Other side effects include water retention, which could cause serious problems such as congestive heart failure or reopening of the ductus arteriosus<sup>65,66</sup>. This side effect may be of particular concern in patients with a low birth weight, as in transient CHI. Routine coadministration of diuretics is advised.

### 2. Octreotide

Octreotide is a somatostatin analog that acts on the somatostatin receptors SSTR2 and SSTR5 and inhibits secretion of a variety of hormones, including gastrin, cholecystokinin, glucagon, growth hormone, secretin, pancreatic polypeptide, thyroid stimulating hormone (TSH) vasoactive intestinal peptide, and insulin. Although its use for CHI has not been licensed in any country, it has been used for nearly 20 years for both short- and long-term control of diazoxide-unresponsive CHI<sup>67,68</sup>. It is administered as multiple daily subcutaneous injections (3–4 times/day) or by continuous subcutaneous infusions using an insulin pump. In our experience, many

**Table 3. Treatment for congenital hyperinsulinism**

Nutritional
Hypertonic glucose infusion
Cornstarch
Glycogen storage disorder formula
Enteral feeding (nasogastric tube feeding, gastrostomy)
Medical
Diazoxide, 5–20 mg/kg/day, po
Nifedipine, 0.25–2.5 mg/kg/day, po
Octreotide, 5–25 $\mu$ g/kg/day, sc
Glucagon, 1–20 $\mu$ g/kg/hr, sc, iv, im
Surgical
Pancreatectomy (partial, subtotal, neartotal)

po, per oral; sc, subcutaneous; iv, intravenous; im, intramuscular.

patients with  $K_{ATP}$  channel CHI can be maintained on long-term treatment until spontaneous remission at 2–5 years of age<sup>69</sup>. Common adverse events include gastrointestinal symptoms, white stool, dilated gall bladder with or without gall stones, and growth deceleration after 2 years of age. Rarer, but more serious side effects, include hepatitis<sup>70</sup>, necrotizing enterocolitis<sup>71</sup> and long QT syndrome<sup>72</sup>.

### 3. Glucagon

Glucagon stimulates glycogenolysis and gluconeogenesis to increase hepatic glucose output. It is administered by intravenous, subcutaneous, or intramuscular routes, and has been used mainly for short-term control of diazoxide-unresponsive patients who are not adequately controlled by other means. However, as is the case for octreotide, its long-term use until spontaneous remission has been reported<sup>73,74</sup>. Apart from its gastrointestinal side effects, its crystallization in the infusion tubes has been a major practical problem during long-term use. Part of this problem may be ameliorated by the development of a water-soluble formulation that is in a phase 2 clinical trial (ClinicalTrials.gov Identifier: NCT01972152).

### 4. Pancreatectomy

When patients are not responsive to medical treatment and cannot be weaned off treatment with intravenous glucose infusions, pancreatectomy should be considered. When a focal lesion is identified preoperatively, partial pancreatectomy is the treatment of choice. However, the lesion is not always visible or palpable at the site indicated by <sup>18</sup>F-DOPA PET. Although intraoperative sonography can aid in identification<sup>75</sup>, repeated intraoperative biopsy may be necessary to help surgeons determine the extent of pancreatectomy. Such a treatment is made possible only by a multidisciplinary team composed of surgeons, radiologists, pediatric endocrinologists, and pathologists who are well experienced in the treatment of CHI<sup>76,77</sup>. When a focal lesion is identified in the body or tail of the pancreas using <sup>18</sup>F-DOPA PET, resection is relatively easy; even if the exact location of the focal lesion cannot be identified, distal pancreatectomy of <70% typically cures the patient without a risk of postoperative diabetes. On the contrary, when the lesion is identified in the head of the pancreas, resection may be difficult without damaging important adjacent structures such as the main pancreatic duct or the common bile duct. In those cases, pancreatic head resection with Roux-en-Y reconstruction of distal pancreatojejunostomy has been proposed<sup>78</sup>. For patients with diazoxide-unresponsive diffuse CHI, extended resection of the pancreas is still needed. Even in these cases, near-total pancreatectomy should be avoided as much as possible in order to reduce the development of postsurgical diabetes<sup>79,80</sup>. A 70%–90% resection may be considered to reduce the pancreatic mass and to facilitate medical management.

## Other unsolved questions and future perspectives

### 1. Causes of the remaining 50% of persistent CHI

At present, even with the most comprehensive molecular analysis, mutations in known causative genes cannot be identified in 21.3% of patients<sup>26</sup>. When confined to diazoxide-responsive cases, mutations are not identified in 53%. Therefore, if we assume that all persistent CHI is genetic in origin, there must be unidentified causative genes. In an effort to address this issue, Proverbio et al.<sup>81</sup> analyzed 17 families with CHI who lacked mutations in *ABCC8/KCNJ11* using a combination of transmission disequilibrium tests and whole-exome sequencing and reported 21 novel genes as possible candidates. Although none of these have been confirmed as causative, further efforts employing next-generation sequencing may answer these questions.

Using next-generation sequencing, Flanagan et al.<sup>82</sup> took a different approach of sequencing the entire genomic region of the *ABCC8* and *HADH* genes<sup>82</sup>. By this strategy, they identified deep intronic mutations of both genes causing CHI, c.1333-1013A>G in *ABCC8* and c.636p471G>T *HADH*. Surprisingly, these mutations were common in the Irish and Turkish populations, accounting for 14% of focal hyperinsulinism cases and 32% of subjects with *HADH* mutations.

### 2. Causes of transient CHI (a hypothesis)

Transient CHI is common in infants who were born small for their gestational ages (SGA) or in those with perinatal stress. However, little is known about its cause. SGA infants are in a hypoxemic condition in utero<sup>83</sup>. Because  $\beta$ -cell function is inhibited by hypoxia-inducible factor 1 $\alpha$  (HIF1 $\alpha$ )<sup>84,85</sup>, a sudden increase in the oxygen tension at delivery may downregulate HIF1 $\alpha$  leading to hyperinsulinemia. In line with the observation that oxygenation of fetal blood improves with gestational ages<sup>83</sup>, it has been reported that blood insulin levels at birth correlate with the gestational age of the infants: 9.2  $\mu$ IU/mL for full term; 10.3  $\mu$ IU/mL for early term; 13.2  $\mu$ IU/mL for late preterm; and 18.9  $\mu$ IU/mL for early preterm<sup>86</sup>. Hyperinsulinemia at birth, therefore, is a common finding in newborns with a lower birth weight.

### 3. Mechanism of spontaneous remission of CHI

Both diffuse and focal HI resolve spontaneously over time<sup>87</sup>. A previously proposed mechanism for spontaneous remission of CHI is apoptotic death of insulin-oversecreting  $\beta$ -cells<sup>88</sup>. However, the initial event could be functional shutdown of insulin secretion rather than apoptotic cell death because the abnormal  $\beta$ -cells could still be observed by <sup>18</sup>F-DOPA PET at an early stage of the spontaneous remission of focal  $K_{ATP}$ -CHI<sup>89</sup>. Manipulating the process of functional shutoff could be an

attractive treatment option for CHI.

#### 4. Novel medications for diazoxide-unresponsive CHI

##### 1) Novel somatostatin analogues

Novel somatostatin analogues have been successfully used for CHI or other forms of HI, including lanreotide<sup>90,91</sup> or long-acting octreotide<sup>92</sup> for CHI. In addition, pasireotide has been tested for severe postgastric bypass HI<sup>93</sup>. Although octreotide and lanreotide have affinities for somatostatin receptors SSTR2 and SSTR5, pasireotide has a broader spectrum of activity for other types of SSTRs<sup>94</sup>.

##### 2) Small molecule corrector of $K_{ATP}$ -channel CHI

The search for small molecules to treat CHI is fueled by previous efforts to correct the trafficking defect of the cystic fibrosis transmembrane conductance regulator, which is deficient in patients with cystic fibrosis. The idea is to use small molecules as pharmacological chaperones to correct the trafficking defect and help their expression to the cell surface<sup>95</sup>. This strategy is applicable to certain mutations of the  $K_{ATP}$ -channel genes. Thus far, sulfonylureas<sup>96</sup> and carbamazepine<sup>97</sup> have been successfully used to correct the trafficking defects of mutations within the transmembrane domains 0 and 1 (TMD0, TMD1) of ABCC8.

##### 3) Glucagon-like peptide 1 (GLP1) receptor antagonist

GLP1 is secreted from the L-cells of the small intestine and binds to the GLP1 receptors in pancreatic  $\beta$ -cells, thereby stimulating the secretion of insulin (the incretin pathway). This pathway has a role in the amplification of postprandial insulin secretion and has been the target of novel treatments for type 2 diabetes. An antagonist of the GLP1 receptor, exendin<sup>9,39</sup>, has been shown to be effective for the treatment of CHI in an open-label, randomized clinical trial<sup>98</sup>. Although the effect was not complete, and monotherapy of CHI with this class of medication appears impractical, GLP1 receptor antagonists may have a role in adjunctive treatment of CHI.

##### 4) Mammalian target of rapamycin (mTOR) inhibitors

mTOR is a member of the serine/threonine kinase family and is induced by amino acids (arginine and branched-chain amino acids), stress, high-energy status, oxygen, and growth factors. mTOR is complexed with regulatory-associated protein of mTOR (Raptor), mammalian LST8/G-protein  $\beta$ -subunit-like protein (mLST8/G $\beta$ L), PRAS40, and DEPTOR to form the mTORC1 complex. Alternatively, mTOR is complexed with mLST8/G $\beta$ , rapamycin-insensitive companion of mTOR (Rictor), and mammalian stress-activated protein kinase-interacting protein 1 (mSIN1) to form mTORC2 and is active in a variety of cellular mechanisms, including protein synthesis, cell proliferation, or cell survival. Therefore, mTOR inhibitors have been widely used to treat neoplasms. In terms of glucose metabolism, activation of mTORC1 is known to cause increased

glucose uptake and glycolysis via HIF1. In addition, mTORC2 is known to play an important role in maintaining the  $\beta$ -cell mass through the phosphatidylinositol-3-kinase/mTORC2/AKT signaling pathway<sup>99</sup>. Sirolimus, one of the mTOR inhibitors, was successfully used to treat patients with diazoxide unresponsive CHI<sup>100</sup>. These classes of medications therefore may have a role in the treatment of CHI as well.

#### Conflict of interest

No potential conflict of interest relevant to this article was reported.

#### Acknowledgments

This work was supported by a Grant-in-aid for Scientific Research from the Ministry of Health, Labour and Welfare of Japan (Research on Measures for Intractable Diseases 2012-070).

#### References

1. Menni F, de Lonlay P, Sevin C, Touati G, Peigne C, Barbier V, et al. Neurologic outcomes of 90 neonates and infants with persistent hyperinsulinemic hypoglycemia. *Pediatrics* 2001;107:476-9.
2. Steinkrauss L, Lipman TH, Hendell CD, Gerdes M, Thornton PS, Stanley CA. Effects of hypoglycemia on developmental outcome in children with congenital hyperinsulinism. *J Pediatr Nurs* 2005;20:109-18.
3. Ludwig A, Ziegenhorn K, Empting S, Meissner T, Marquard J, Holl R, et al. Glucose metabolism and neurological outcome in congenital hyperinsulinism. *Semin Pediatr Surg* 2011;20:45-9.
4. Avatapalle HB, Banerjee I, Shah S, Pryce M, Nicholson J, Rigby L, et al. Abnormal neurodevelopmental outcomes are common in children with transient congenital hyperinsulinism. *Front Endocrinol (Lausanne)* 2013;4:60.
5. Shilyansky J, Fisher S, Cutz E, Perlman K, Filler RM. Is 95% pancreatectomy the procedure of choice for treatment of persistent hyperinsulinemic hypoglycemia of the neonate? *J Pediatr Surg* 1997;32:342-6.
6. Meissner T, Wendel U, Burgard P, Schaetzle S, Mayatepek E. Long-term follow-up of 114 patients with congenital hyperinsulinism. *Eur J Endocrinol* 2003;149:43-51.
7. Levy-Shraga Y, Pinhas-Hamiel O, Kraus-Houminer E, Landau H, Mazor-Aronovitch K, Modan-Moses D, et al. Cognitive and developmental outcome of conservatively treated children with congenital hyperinsulinism. *J Pediatr Endocrinol Metab* 2013;26:301-8.
8. Hussain K. Diagnosis and management of hyperinsulinaemic hypoglycaemia of infancy. *Horm Res* 2008;69:2-13.
9. Arnoux JB, Verkarre V, Saint-Martin C, Montravers



- F, Brassier A, Valayannopoulos V, et al. Congenital hyperinsulinism: current trends in diagnosis and therapy. *Orphanet J Rare Dis* 2011;6:63.
10. Mohamed Z, Arya VB, Hussain K. Hyperinsulinaemic hypoglycaemia: genetic mechanisms, diagnosis and management. *J Clin Res Pediatr Endocrinol* 2012;4:169-81.
  11. Palladino AA, Bennett MJ, Stanley CA. Hyperinsulinism in infancy and childhood: when an insulin level is not always enough. *Clin Chem* 2008;54:256-63.
  12. De Leon DD, Stanley CA. Determination of insulin for the diagnosis of hyperinsulinemic hypoglycemia. *Best Pract Res Clin Endocrinol Metab* 2013;27:763-9.
  13. Arnoux JB, de Lonlay P, Ribeiro MJ, Hussain K, Blankenstein O, Mohnike K, et al. Congenital hyperinsulinism. *Early Hum Dev* 2010;86:287-94.
  14. van Veen MR, van Hasselt PM, de Sain-van der Velden MG, Verhoeven N, Hofstede FC, de Koning TJ, et al. Metabolic profiles in children during fasting. *Pediatrics* 2011;127:e1021-7.
  15. Vannucci RC, Vannucci SJ. Hypoglycemic brain injury. *Semin Neonatol* 2001;6:147-55.
  16. Salhab WA, Wyckoff MH, Laptook AR, Perlman JM. Initial hypoglycemia and neonatal brain injury in term infants with severe fetal acidemia. *Pediatrics* 2004;114:361-6.
  17. Rozance PJ, Hay WW. Hypoglycemia in newborn infants: Features associated with adverse outcomes. *Biol Neonate* 2006;90:74-86.
  18. Gataullina S, Dellatolas G, Perdry H, Robert JJ, Valayannopoulos V, Touati G, et al. Comorbidity and metabolic context are crucial factors determining neurological sequelae of hypoglycaemia. *Dev Med Child Neurol* 2012;54:1012-7.
  19. Ashcroft FM, Rorsman P. K(ATP) channels and islet hormone secretion: new insights and controversies. *Nat Rev Endocrinol* 2013;9:660-9.
  20. Yorifuji T, Masue M, Nishibori H. Congenital hyperinsulinism: global and Japanese perspectives. *Pediatr Int* Forthcoming 2014. <http://dx.doi.org/10.1111/ped.12390>.
  21. Pearson ER, Boj SF, Steele AM, Barrett T, Stals K, Shield JP, et al. Macrosomia and hyperinsulinaemic hypoglycaemia in patients with heterozygous mutations in the HNF4A gene. *PLoS Med* 2007;4:e118.
  22. Kapoor RR, Locke J, Colclough K, Wales J, Conn JJ, Hattersley AT, et al. Persistent hyperinsulinemic hypoglycemia and maturity-onset diabetes of the young due to heterozygous HNF4A mutations. *Diabetes* 2008;57:1659-63.
  23. Flanagan SE, Kapoor RR, Mali G, Cody D, Murphy N, Schwahn B, et al. Diazoxide-responsive hyperinsulinemic hypoglycemia caused by HNF4A gene mutations. *Eur J Endocrinol* 2010;162:987-92.
  24. Stanescu DE, Hughes N, Kaplan B, Stanley CA, De Leon DD. Novel presentations of congenital hyperinsulinism due to mutations in the MODY genes: HNF1A and HNF4A. *J Clin Endocrinol Metab* 2012;97:E2026-30.
  25. Colclough K, Bellanne-Chantelot C, Saint-Martin C, Flanagan SE, Ellard S. Mutations in the genes encoding the transcription factors hepatocyte nuclear factor 1 alpha and 4 alpha in maturity-onset diabetes of the young and hyperinsulinemic hypoglycemia. *Hum Mutat* 2013;34:669-85.
  26. Snider KE, Becker S, Boyajian L, Shyng SL, MacMullen C, Hughes N, et al. Genotype and phenotype correlations in 417 children with congenital hyperinsulinism. *J Clin Endocrinol Metab* 2013;98:E355-63.
  27. Kapoor RR, Flanagan SE, Arya VB, Shield JP, Ellard S, Hussain K. Clinical and molecular characterisation of 300 patients with congenital hyperinsulinism. *Eur J Endocrinol* 2013;168:557-64.
  28. Sempoux C, Guiot Y, Jaubert F, Rahier J. Focal and diffuse forms of congenital hyperinsulinism: the keys for differential diagnosis. *Endocr Pathol* 2004;15:241-6.
  29. Pinney SE, MacMullen C, Becker S, Lin YW, Hanna C, Thornton P, et al. Clinical characteristics and biochemical mechanisms of congenital hyperinsulinism associated with dominant KATP channel mutations. *J Clin Invest* 2008;118:2877-86.
  30. Macmullen CM, Zhou Q, Snider KE, Tewson PH, Becker SA, Aziz AR, et al. Diazoxide-unresponsive congenital hyperinsulinism in children with dominant mutations of the  $\beta$ -cell sulfonylurea receptor SUR1. *Diabetes* 2011;60:1797-804.
  31. de Lonlay P, Fournet JC, Rahier J, Gross-Morand MS, Poggi-Travert F, Foussier V, et al. Somatic deletion of the imprinted 11p15 region in sporadic persistent hyperinsulinemic hypoglycemia of infancy is specific of focal adenomatous hyperplasia and endorses partial pancreatectomy. *J Clin Invest* 1997;100:802-7.
  32. Verkarre V, Fournet JC, de Lonlay P, Gross-Morand MS, Devillers M, Rahier J, et al. Paternal mutation of the sulfonylurea receptor (SUR1) gene and maternal loss of 11p15 imprinted genes lead to persistent hyperinsulinism in focal adenomatous hyperplasia. *J Clin Invest* 1998;102:1286-91.
  33. Fournet JC, Mayaud C, de Lonlay P, Gross-Morand MS, Verkarre V, Castanet M, et al. Unbalanced expression of 11p15 imprinted genes in focal forms of congenital hyperinsulinism: association with a reduction to homozygosity of a mutation in ABCC8 or KCNJ11. *Am J Pathol* 2001;158:2177-84.
  34. Damaj L, le Lorch M, Verkarre V, Werl C, Hubert L, Nihoul-Fekete C, et al. Chromosome 11p15 paternal isodisomy in focal forms of neonatal hyperinsulinism. *J Clin Endocrinol Metab* 2008;93:4941-7.
  35. Suchi M, Thornton PS, Adzick NS, MacMullen C, Ganguly A, Stanley CA, et al. Congenital hyperinsulinism: intraoperative biopsy interpretation can direct the extent of pancreatectomy. *Am J Surg Pathol* 2004;28:1326-35.
  36. Suchi M, MacMullen CM, Thornton PS, Adzick NS, Ganguly A, Ruchelli ED, et al. Molecular and immunohistochemical analyses of the focal form of congenital hyperinsulinism.

- Mod Pathol 2006;19:122-9.
37. Rahier J, Guiot Y, Sempoux C. Morphologic analysis of focal and diffuse forms of congenital hyperinsulinism. *Semin Pediatr Surg* 2011;20:3-12.
  38. Otonkoski T, Nanto-Salonen K, Seppanen M, Veijola R, Huopio H, Hussain K, et al. Noninvasive diagnosis of focal hyperinsulinism of infancy with [18F]-DOPA positron emission tomography. *Diabetes* 2006;55:13-8.
  39. Blomberg BA, Moghbel MC, Saboury B, Stanley CA, Alavi A. The value of radiologic interventions and (18) F-DOPA PET in diagnosing and localizing focal congenital hyperinsulinism: systematic review and meta-analysis. *Mol Imaging Biol* 2013;15:97-105.
  40. Yang J, Hao R, Zhu X. Diagnostic role of 18F-dihydroxyphenylalanine positron emission tomography in patients with congenital hyperinsulinism: a meta-analysis. *Nucl Med Commun* 2013;34:347-53.
  41. Masue M, Nishibori H, Fukuyama S, Yoshizawa A, Okamoto S, Doi R, et al. Diagnostic accuracy of [<sup>18</sup>F]-fluoro-L-dihydroxyphenylalanine positron emission tomography scan for persistent congenital hyperinsulinism in Japan. *Clin Endocrinol (Oxf)* 2011;75:342-6.
  42. Glaser B, Ryan F, Donath M, Landau H, Stanley CA, Baker L, et al. Hyperinsulinism caused by paternal-specific inheritance of a recessive mutation in the sulfonylurea-receptor gene. *Diabetes* 1999;48:1652-7.
  43. De Leon DD, Stanley CA. Mechanisms of disease: advances in diagnosis and treatment of hyperinsulinism in neonates. *Nat Clin Pract Endocrinol Metab* 2007;3:57-68.
  44. Fernandez-Marmiesse A, Salas A, Vega A, Fernandez-Lorenzo JR, Barreiro J, Carracedo A. Mutation spectra of ABCC8 gene in Spanish patients with Hyperinsulinism of Infancy (HI). *Hum Mutat* 2006;27:214.
  45. Sandal T, Laborie LB, Brusgaard K, Eide SA, Christesen HB, Sovik O, et al. The spectrum of ABCC8 mutations in Norwegian patients with congenital hyperinsulinism of infancy. *Clin Genet* 2009;75:440-8.
  46. Yorifuji T, Kawakita R, Nagai S, Sugimine A, Doi H, Nomura A, et al. Molecular and clinical analysis of Japanese patients with persistent congenital hyperinsulinism: predominance of paternally inherited monoallelic mutations in the KATP channel genes. *J Clin Endocrinol Metab* 2011;96:E141-5.
  47. Mohnike K, Wieland I, Barthlen W, Vogelgesang S, Empting S, Mohnike W, et al. Clinical and genetic evaluation of patients with KATP channel mutations from the German registry for congenital hyperinsulinism. *Horm Res Paediatr* 2014;81:156-68.
  48. Su C, Gong C, Sanger P, Li W, Wu D, Gu Y, et al. Long-term follow-up and mutation analysis of 27 chinese cases of congenital hyperinsulinism. *Horm Res Paediatr* 2014;81:169-76.
  49. Banerjee I, Skae M, Flanagan SE, Rigby L, Patel L, Didi M, et al. The contribution of rapid KATP channel gene mutation analysis to the clinical management of children with congenital hyperinsulinism. *Eur J Endocrinol* 2011;164:733-40.
  50. Stanley CA, Lieu YK, Hsu BY, Burlina AB, Greenberg CR, Hopwood NJ, et al. Hyperinsulinism and hyperammonemia in infants with regulatory mutations of the glutamate dehydrogenase gene. *N Engl J Med* 1998;338:1352-7.
  51. Molven A, Matre GE, Duran M, Wanders RJ, Rishaug U, Njolstad PR, et al. Familial hyperinsulinemic hypoglycemia caused by a defect in the SCHAD enzyme of mitochondrial fatty acid oxidation. *Diabetes* 2004;53:221-7.
  52. Flanagan SE, Patch AM, Locke JM, Akcay T, Simsek E, Alaei M, et al. Genome-wide homozygosity analysis reveals HADH mutations as a common cause of diazoxide-responsive hyperinsulinemic-hypoglycemia in consanguineous pedigrees. *J Clin Endocrinol Metab* 2011;96:E498-502.
  53. Heslegrave AJ, Hussain K. Novel insights into fatty acid oxidation, amino acid metabolism, and insulin secretion from studying patients with loss of function mutations in 3-hydroxyacyl-CoA dehydrogenase. *J Clin Endocrinol Metab* 2013;98:496-501.
  54. Kassem S, Bhandari S, Rodríguez-Bada P, Motaghedi R, Heyman M, Garcia-Gimeno MA, et al. Large islets, beta-cell proliferation, and a glucokinase mutation. *N Engl J Med* 2010;362:1348-50.
  55. Barbetti F, Cobo-Vuilleumier N, Dionisi-Vici C, Toni S, Ciampalini P, Massa O, et al. Opposite clinical phenotypes of glucokinase disease: description of a novel activating mutation and contiguous inactivating mutations in human glucokinase (GCK) gene. *Mol Endocrinol* 2009;23:1983-9.
  56. Christesen HB, Tribble ND, Molven A, Siddiqui J, Sandal T, Brusgaard K, et al. Activating glucokinase (GCK) mutations as a cause of medically responsive congenital hyperinsulinism: prevalence in children and characterisation of a novel GCK mutation. *Eur J Endocrinol* 2008;159:27-34.
  57. Osbak KK, Colclough K, Saint-Martin C, Beer NL, Bellanne-Chantelot C, Ellard S, et al. Update on mutations in glucokinase (GCK), which cause maturity-onset diabetes of the young, permanent neonatal diabetes, and hyperinsulinemic hypoglycemia. *Hum Mutat* 2009;30:1512-26.
  58. Henquin JC, Sempoux C, Marchandise J, Godecharles S, Guiot Y, Nenquin M, et al. Congenital hyperinsulinism caused by hexokinase I expression or glucokinase-activating mutation in a subset of  $\beta$ -cells. *Diabetes* 2013;62:1689-96.
  59. Gonzalez-Barroso MM, Giurgea I, Bouillaud F, Anedda A, Bellanne-Chantelot C, Hubert L, et al. Mutations in UCP2 in congenital hyperinsulinism reveal a role for regulation of insulin secretion. *PLoS One* 2008;3:e3850.
  60. Otonkoski T, Kaminen N, Ustinov J, Lapatto R, Meissner T, Mayatepek E, et al. Physical exercise-induced hyperinsulinemic hypoglycemia is an autosomal-dominant trait characterized by abnormal pyruvate-induced insulin release. *Diabetes* 2003;52:199-204.
  61. Kapoor RR, James C, Hussain K. Hyperinsulinism in

- developmental syndromes. *Endocr Dev* 2009;14:95-113.
62. Bitner-Glindzicz M, Lindley KJ, Rutland P, Blyndon D, Smith VV, Milla PJ, et al. A recessive contiguous gene deletion causing infantile hyperinsulinism, enteropathy and deafness identifies the Usher type 1C gene. *Nat Genet* 2000;26:56-60.
  63. Al Mutair AN, Brusgaard K, Bin-Abbas B, Hussain K, Felimban N, Al Shaikh A, et al. Heterogeneity in phenotype of usher-congenital hyperinsulinism syndrome: hearing loss, retinitis pigmentosa, and hyperinsulinemic hypoglycemia ranging from severe to mild with conversion to diabetes. *Diabetes Care* 2013;36:557-61.
  64. Touati G, Poggi-Travert F, Ogier de Baulny H, Rahier J, Brunelle F, Nihoul-Fekete C, et al. Long-term treatment of persistent hyperinsulinaemic hypoglycaemia of infancy with diazoxide: a retrospective review of 77 cases and analysis of efficacy-predicting criteria. *Eur J Pediatr* 1998;157:628-33.
  65. Demirel F, Unal S, Cetin II, Esen I, Arasli A. Pulmonary hypertension and reopening of the ductus arteriosus in an infant treated with diazoxide. *J Pediatr Endocrinol Metab* 2011;24:603-5.
  66. Yoshida K, Kawai M, Marumo C, Kanazawa H, Matsukura T, Kusuda S, et al. High prevalence of severe circulatory complications with diazoxide in premature infants. *Neonatology* 2014;105:166-71.
  67. Glaser B, Hirsch HJ, Landau H. Persistent hyperinsulinemic hypoglycemia of infancy: long-term octreotide treatment without pancreatectomy. *J Pediatr* 1993;123:644-50.
  68. Thornton PS, Alter CA, Katz LE, Baker L, Stanley CA. Short- and long-term use of octreotide in the treatment of congenital hyperinsulinism. *J Pediatr* 1993;123:637-43.
  69. Yorifuji T, Kawakita R, Hosokawa Y, Fujimaru R, Matsubara K, Aizu K, et al. Efficacy and safety of long-term, continuous subcutaneous octreotide infusion for patients with different subtypes of KATP-channel hyperinsulinism. *Clin Endocrinol (Oxf)* 2013;78:891-7.
  70. Koren I, Riskin A, Barthlen W, Gillis D. Hepatitis in an infant treated with octreotide for congenital hyperinsulinism. *J Pediatr Endocrinol Metab* 2013;26:183-5.
  71. Laje P, Halaby L, Adzick NS, Stanley CA. Necrotizing enterocolitis in neonates receiving octreotide for the management of congenital hyperinsulinism. *Pediatr Diabetes* 2010;11:142-7.
  72. Celik N, Cinaz P, Emeksiz HC, Hussain K, Camurdan O, Bideci A, et al. Octreotide-induced long QT syndrome in a child with congenital hyperinsulinemia and a novel missense mutation (p.Met115Val) in the ABCC8 gene. *Horm Res Paediatr* 2013;80:299-303.
  73. Mohnike K, Blankenstein O, Pfuetzner A, Potzsch S, Schober E, Steiner S, et al. Long-term non-surgical therapy of severe persistent congenital hyperinsulinism with glucagon. *Horm Res* 2008;70:59-64.
  74. Neylon OM, Moran MM, Pellicano A, Nightingale M, O'Connell MA. Successful subcutaneous glucagon use for persistent hypoglycaemia in congenital hyperinsulinism. *J Pediatr Endocrinol Metab* 2013;26:1157-61.
  75. von Rohden L, Mohnike K, Mau H, Eberhard T, Mohnike W, Blankenstein O, et al. Visualization of the focus in congenital hyperinsulinism by intraoperative sonography. *Semin Pediatr Surg* 2011;20:28-31.
  76. Adzick NS, Thornton PS, Stanley CA, Kaye RD, Ruchelli E. A multidisciplinary approach to the focal form of congenital hyperinsulinism leads to successful treatment by partial pancreatectomy. *J Pediatr Surg* 2004;39:270-5.
  77. Palladino AA, Stanley CA. A specialized team approach to diagnosis and medical versus surgical treatment of infants with congenital hyperinsulinism. *Semin Pediatr Surg* 2011;20:32-7.
  78. Laje P, Stanley CA, Palladino AA, Becker SA, Adzick NS. Pancreatic head resection and Roux-en-Y pancreaticojejunostomy for the treatment of the focal form of congenital hyperinsulinism. *J Pediatr Surg* 2012;47:130-5.
  79. Barthlen W, de Lonlay P. Congenital hyperinsulinism. *Semin Pediatr Surg* 2011;20:1-2.
  80. Barthlen W. Surgery in congenital hyperinsulinism-tips and tricks not only for surgeons: a practical guide. *Semin Pediatr Surg* 2011;20:56-9.
  81. Proverbio MC, Mangano E, Gessi A, Bordoni R, Spinelli R, Asselta R, et al. Whole genome SNP genotyping and exome sequencing reveal novel genetic variants and putative causative genes in congenital hyperinsulinism. *PLoS One* 2013;8:e68740.
  82. Flanagan SE, Xie W, Caswell R, Damhuis A, Vianey-Saban C, Akcay T, et al. Next-generation sequencing reveals deep intronic cryptic ABCC8 and HADH splicing founder mutations causing hyperinsulinism by pseudoexon activation. *Am J Hum Genet* 2013;92:131-6.
  83. Thompson LP, Al-Hasan Y. Impact of oxidative stress in fetal programming. *J Pregnancy* 2012;2012:582748.
  84. Heinis M, Simon MT, Ilc K, Mazure NM, Pouyssegur J, Scharfmann R, et al. Oxygen tension regulates pancreatic beta-cell differentiation through hypoxia-inducible factor 1alpha. *Diabetes* 2010;59:662-9.
  85. Hakim F, Kaitsuka T, Raed JM, Wei FY, Shiraki N, Akagi T, et al. High oxygen condition facilitates the differentiation of mouse and human pluripotent stem cells into pancreatic progenitors and insulin-producing cells. *J Biol Chem* 2014;289:9623-38.
  86. Wang G, Divall S, Radovick S, Paige D, Ning Y, Chen Z, et al. Preterm birth and random plasma insulin levels at birth and in early childhood. *JAMA* 2014;311:587-96.
  87. Mazor-Aronovitch K, Landau H, Gillis D. Surgical versus non-surgical treatment of congenital hyperinsulinism. *Pediatr Endocrinol Rev* 2009;6:424-30.
  88. Kassem SA, Ariel I, Thornton PS, Scheimberg I, Glaser B. Beta-cell proliferation and apoptosis in the developing normal human pancreas and in hyperinsulinism of infancy. *Diabetes* 2000;49:1325-33.
  89. Yorifuji T, Hosokawa Y, Fujimaru R, Kawakita R, Doi

- H, Matsumoto T, et al. Lasting 18F-DOPA PET uptake after clinical remission of the focal form of congenital hyperinsulinism. *Horm Res Paediatr* 2011;76:286-90.
90. Modan-Moses D, Koren I, Mazor-Aronovitch K, Pinhas-Hamiel O, Landau H. Treatment of congenital hyperinsulinism with lanreotide acetate (Somatuline Autogel). *J Clin Endocrinol Metab* 2011;96:2312-7.
91. Kuhnen P, Marquard J, Ernert A, Meissner T, Raile K, Wannemacher G, et al. Long-term lanreotide treatment in six patients with congenital hyperinsulinism. *Horm Res Paediatr* 2012;78:106-12.
92. Le Quan Sang KH, Arnoux JB, Mamoune A, Saint-Martin C, Bellanne-Chantelot C, Valayannopoulos V, et al. Successful treatment of congenital hyperinsulinism with long-acting release octreotide. *Eur J Endocrinol* 2012;166:333-9.
93. de Heide LJ, Laskewitz AJ, Apers JA. Treatment of severe postRYGB hyperinsulinemic hypoglycemia with pasireotide: a comparison with octreotide on insulin, glucagon, and GLP-1. *Surg Obes Relat Dis* 2013 Dec 4 [Epub]. <http://dx.doi.org/10.1016/j.soard.2013.11.006>.
94. Feelders RA, de Herder WW, Neggers SJ, van der Lely AJ, Hofland LJ. Pasireotide, a multi-somatostatin receptor ligand with potential efficacy for treatment of pituitary and neuroendocrine tumors. *Drugs Today (Barc)* 2013;49:89-103.
95. Martin GM, Chen PC, Devaraneni P, Shyng SL. Pharmacological rescue of trafficking-impaired ATP-sensitive potassium channels. *Front Physiol* 2013;4:386.
96. Yan FF, Casey J, Shyng SL. Sulfonylureas correct trafficking defects of disease-causing ATP-sensitive potassium channels by binding to the channel complex. *J Biol Chem* 2006;281:33403-13.
97. Chen PC, Olson EM, Zhou Q, Kryukova Y, Sampson HM, Thomas DY, et al. Carbamazepine as a novel small molecule corrector of trafficking-impaired ATP-sensitive potassium channels identified in congenital hyperinsulinism. *J Biol Chem* 2013;288:20942-54.
98. Calabria AC, Li C, Gallagher PR, Stanley CA, De Leon DD. GLP-1 receptor antagonist exendin-(9-39) elevates fasting blood glucose levels in congenital hyperinsulinism owing to inactivating mutations in the ATP-sensitive K<sup>+</sup> channel. *Diabetes* 2012;61:2585-91.
99. Gu Y, Lindner J, Kumar A, Yuan W, Magnuson MA. Rictor/mTORC2 is essential for maintaining a balance between beta-cell proliferation and cell size. *Diabetes* 2011;60:827-37.
100. Senniappan S, Alexandrescu S, Tatevian N, Shah P, Arya V, Flanagan S, et al. Sirolimus therapy in infants with severe hyperinsulinemic hypoglycemia. *N Engl J Med* 2014;370:1131-7.