



Published in final edited form as:

Cell. 2014 July 31; 158(3): 607–619. doi:10.1016/j.cell.2014.05.047.

MicroRNA Directly Enhances Mitochondrial Translation during Muscle Differentiation

Xiaorong Zhang¹, Xinxin Zuo¹, Bo Yang¹, Zongran Li¹, Yuanchao Xue^{1,2}, Yu Zhou², Jie Huang¹, Xiaolu Zhao¹, Jie Zhou¹, Yun Yan¹, Huiqiong Zhang¹, Peipei Guo¹, Hui Sun¹, Lin Guo¹, Yi Zhang^{1,*}, and Xiang-Dong Fu^{1,2,3}

¹State Key Laboratory of Virology, College of Life Sciences, Wuhan University, Wuhan, Hubei 430072, China

²Department of Cellular and Molecular Medicine, University of California, San Diego, La Jolla, CA 92093-0651, USA

³Institute of Genomic Medicine, University of California, San Diego, La Jolla, CA 92093-0651, USA

Abstract

MicroRNAs are well known to mediate translational repression and mRNA degradation in the cytoplasm. Various microRNAs have also been detected in membrane-compartmentalized organelles, but the functional significance has remained elusive. Here we report that miR-1, a microRNA specifically induced during myogenesis, efficiently enters the mitochondria where it unexpectedly stimulates, rather than represses, the translation of specific mitochondrial genome-encoded transcripts. We show that this positive effect requires specific miR:mRNA base-pairing and Ago2, but not its functional partner GW182, which is excluded from the mitochondria. We provide evidence for the direct action of Ago2 in mitochondrial translation by CLIP-seq, functional rescue with mitochondria-targeted Ago2, and selective inhibition of the microRNA machinery in the cytoplasm. These findings unveil a positive function of microRNA in mitochondrial translation and suggest a highly coordinated myogenic program via miR-1 mediated translational stimulation in the mitochondria and repression in the cytoplasm.

© 2014 Elsevier Inc. All rights reserved.

Correspondence: Xiang-Dong Fu, Tel: 858-534-4937, Fax: 858-822-6920, xdfu@ucsd.edu.

*Current Address: Center for Genome Analysis, ABLifeInc, Novonest Building 135, 8 Nanhu Avenue, East Lake Hi-Tech Development Zone, Wuhan, Hubei 430079, China

Accession Numbers

The Ago2 CLIP-seq data performed on C2C12 myoblasts and myotubes are available at the Gene Expression Omnibus under the accession number GSE57596

Author contribution: X-R.Z., X-X.Z., and X-D.F. designed the experiments; Y.Z. contributed to experimental design in the initial phase of this project; X-R.Z., X-X.Z., L.Z. carried out most of the experiments; Y.X. performed Ago2 CLIP-seq; B.Y.; Y.Z., J.H., J.Z., and Y.Y. analyzed the data; X-L.Z. and L.G. performed mass spectrometric analysis of IPed Ago2; H.Z., P.G., and H.S. contributed to data collection; X-R.Z., X-X.Z., and X-D.F. wrote the paper.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

The function and mechanism of microRNAs (miRNAs) have been well studied in higher eukaryotic cells. miRNAs bind their target mRNAs via partial base-pairing within the RNA-Induced Silencing Complexes (RISC) and regulate both mRNA stability and translation (Carthew and Sontheimer, 2009; Chekulaeva and Filipowicz, 2009). miRNA-mediated RNA degradation can be accomplished via a slicing-competent Ago protein (Ago2 in mammals) to cleave target mRNA in the mRNA:miRNA duplex. The broad function of RISC in translational control is mediated by the key Ago partner GW182 to recruit deadenylating/decapping enzymes to allow exonucleases to attack unprotected mRNA and repress translation through competing with the cap binding protein eIF4E and/or interfering with ribosome scanning (Czech and Hannon, 2011).

Besides their primary functions in translational repression, miRNAs have also been implicated in enhancing translation under specific cellular conditions (Vasudevan, 2012). This was first observed in serum-starved cells (Vasudevan et al., 2007), but such an opposite function of miRNAs has not been widely appreciated, partly because the potential mechanism involved has remained largely unclear. Biochemical experiments with fly extracts provided some initial hint to this unconventional miRNA function, showing that a target mRNA lacking both cap and typical poly (A) tail permits enhanced translation when the miRNA is assembled into a complex with a GW182-detached Ago protein (Iwasaki and Tomari, 2009). Related phenomena have also been observed in immature oocytes where the miRNA machinery is either in an inactive state (Suh et al., 2010) or even stimulates translation of target transcripts (Mortensen et al., 2011).

The miRNA machinery is known to primarily act in the cytoplasm. However, growing evidence suggests that both miRNAs and proteins of the Argonaute family also play important roles in transcriptional control (Cernilogar et al., 2011; Guang et al., 2008) and DNA repair (Wei et al., 2012). miRNAs have also been detected in membrane-bound compartments, such as secreted vesicles (Zhang et al., 2010) and mitochondria (Bandiera et al., 2011; Barrey et al., 2011; Das et al., 2012; Kren et al., 2009; Sripada et al., 2012). The presence of miRNAs in the mitochondria is somewhat surprising because mitochondria maintain their own genome, and in many aspects, resemble bacteria (Taanman, 1999); however, the functional significance of miRNAs in the mitochondria has remained largely unknown. One report indicates that miR-181c could repress the translation of a mitochondrial transcript, but paradoxically, reduced translation led to an overall enhancement of mitochondrial activities in ventricular myocytes (Das et al., 2012).

Our current study was initially geared to investigate a puzzle that mitochondrial DNA (mtDNA)-encoded proteins were greatly elevated during muscle differentiation, but without a significant increase in mtDNA copy number or transcription. This leads to an unexpected finding that the muscle-specific miR-1 is able to stimulate mitochondrial translation of multiple mtDNA-encoded transcripts, while repressing its nuclear DNA-encoded targets in the cytoplasm. Such a miRNA-dependent effect requires Ago2, but not its normal functional partner GW182, consistent with the presence of a significant amount of Ago2, but not GW182, in the mitochondria. These findings provide critical insights into miRNA action

mechanisms, functional conservation of the Argonaute family of proteins in diverse organisms, change in bioenergetics mechanisms during cell differentiation, and the regulation of mitochondrial translation in higher eukaryotic cells.

Results

Marked increase in mitochondrial protein synthesis during muscle differentiation

Mitochondria generate ATP through oxidative phosphorylation to provide the chemical form of energy for cellular activities (Saraste, 1999). The energy demand is especially high in cardiac and skeletal muscle (Lopaschuk et al., 2010; Moyes et al., 1997). Consistently, we detected much higher levels of mtDNA-encoded proteins in muscle cells relative to constant amounts of nuclear DNA-encoded histone H3 protein and 5S rRNA in different mouse tissues (Figure 1A). Such a marked increase was also evident during C2C12 cell differentiation from myoblasts to myotubes, characterized by the induction of the Myosin Heavy Chain (MHC) and the muscle-specific miRNA miR-1 (Chen et al., 2006; Rao et al., 2006). However, both actin and the nuclear DNA-encoded, mitochondrial residential protein VDAC were constant (Figure 1B). Quantitative analysis indicates that the mitochondrial ND1 and COX1 proteins were increased by more than 15-fold during C2C12 cell differentiation (Figure 1C).

Significantly, such a dramatic elevation in protein synthesis was not accompanied by mtDNA replication during C2C12 cell differentiation (Figure 1D), as reported earlier (Franko et al., 2008). The mRNA levels for both COX1 and ND1 were also invariant (Figure 1E), which was further confirmed by RT-qPCR with three distinct primer sets (Figure S1). This is consistent with little change in TFAM expression (Figure 1B), a vital transcriptional activator in the mitochondria (Bestwick and Shadel, 2013). Since we did not detect any induction of TACO1 (Figure 1B), a major translational activator for COX1 (Weraarpachai et al., 2009), these data suggest a previously unknown mechanism that may account for induced mitochondrial translation during muscle differentiation.

Evidence for the presence of Ago2 within the mitochondria

Given the recent observations that various miRNAs were detected in isolated mitochondria from rat liver (Kren et al., 2009), rat ventricular myocytes (Das et al., 2012), and several common human cell lines (Sripada et al., 2012), we reasoned that reduced miRNA targeting to the mitochondria might account for increased translation in the mitochondria during muscle differentiation. Alternatively, enhanced mitochondrial translation may indirectly result from repression of some nuclear DNA-encoded mitochondrial translational repressors, in line with the induction of Ago2 expression during C12C12 differentiation (Figure 1B).

To begin to explore these possibilities, we first used highly purified mitochondria to rigorously demonstrate that a fraction of Ago2 is indeed present within the mitochondria. Because purified mitochondria are often contaminated with endoplasmic reticulum (ER) proteins due to the intimate association between ER and the mitochondria (de Brito and Scorrano, 2008; Rizzuto et al., 1998), we selectively removed the outer membrane by treating purified mitochondria (MT) with digitonin to prepare mitoplasts (MP) from mouse

heart or C2C12 cells before and after differentiation (Bosmann et al., 1972; Schnaitman and Greenawalt, 1968). We found that our initially purified mitochondria were largely free of ER contamination as indicated by the absence of the ER-specific protein ERp60 or ERp72 (Figure 2A). Upon digitonin treatment followed by sedimentation or fractionation on a sucrose gradient, the outer membrane was removed, as evidenced by the lack of Tom20, but the inner membrane was largely intact, as indicated by the presence of the inner membrane marker NDUFB8 and the mitochondrial matrix protein HSP60 (Figures 2A and S2A).

Under these rigorously controlled conditions, we detected Ago2 in both highly purified MT and MP, which became Trypsin sensitive only when the inner membrane was further permeabilized with Triton X-100. Neither Ago1 nor Ago3 was detectable in MT and MP (Figure S2A), indicating that Ago2 may be selectively imported into the mitochondria. Tandem mass spectrometric analysis confirmed the identity of Ago2 immunoprecipitated from MP (Figure S2B). These data agree with the recent confocal microscopic result (Bandiera et al., 2011) and provide the strongest biochemical evidence to date for the presence of Ago2 inside the mitochondria.

Quantification of Ago2 in the mitochondria during muscle differentiation

We next wished to quantitatively determine the levels of Ago2 in the mitochondria during C2C12 cell differentiation by using the standard curve generated with Ago2 expressed in insect cells (Figure S2C and S2D). Because induced myotubes contain more than one nucleus, we determined the copy number of nuclear DNA by qPCR to calculate the number of Ago2 molecules per nuclear genome. This revealed 1.34×10^5 and 7.35×10^5 Ago2 molecules in C2C12 myoblasts and myotubes, respectively (Figure 2B), which is similar to the recent estimate of 1.4×10^5 – 1.7×10^5 Ago2 per cell on a human melanoma cell line (Wang et al., 2012).

We then determined the amount of Ago2 within the mitochondria by using mtDNA as an internal control (the copy number of mtDNA varies from 0 to 11 per mitochondria with the average=2, see (Cavelier et al., 2000)), which is necessary because of potential damage during MP preparation. This analysis revealed 31 and 314 Ago2 molecules, respectively, per mtDNA in C2C12 cells before and after differentiation (Figure 2B). We also determined the ratio of mitochondrial and nuclear DNA in our cellular system, revealing a ratio of 554 in myoblasts and 774 in myotubes, thus suggesting the presence of $31 \times 554 / 1.34 \times 10^5 = 12.8\%$ and $314 \times 774 / 7.35 \times 10^5 = 33.1\%$ of total cellular Ago2 in the mitochondria of C2C12 cells before and after differentiation, respectively. These quantitative data suggest that more Ago2 is targeted to the mitochondria upon C2C12 differentiation into myotubes.

Efficient entry of induced miR-1 into the mitochondria of muscle cells

We similarly analyzed the quantity of miR-1 in the mitochondria by using highly purified mitoplasts, finding that a fraction of the newly induced miR-1 was indeed detectable inside the mitochondria of myotubes, which became sensitive to nuclease only in the presence of Triton X-100 (Figure 2C). The nuclease sensitivity of miR-1 was similar to the mitochondrial 16S rRNA. In contrast, both nuclear U6 snRNA and cytoplasmic GAPDH mRNA were largely removed from purified MT and MP, and any residual amounts could be

fully degraded by a combination of RNaseT1 (RN) and Micrococcal nuclease (MN) in the absence of detergent (Figure 2C). We further showed that our mitoplasts were highly purified away from the abundant 7SK non-coding RNA and multiple nuclear DNA-transcribed mRNAs for mitochondrial residential proteins (e.g. NDUFB8, COX4V1, and NDUFV1) (Figure S2E).

Using the same quantification strategy for Ago2, we determined the amount of miR-1 in the mitochondria, revealing 5.9×10^4 miR-1 per nuclear genome in myotubes (note that miR-1 was not detectable in myoblasts), 31.5 per mtDNA, and a total amount of $31.5 \times 775 / 5.9 \times 10^4 = 41.4\%$ in the mitochondria. Therefore, similar to Ago2, miR-1 is not only present in the mitochondria in a substantial quantity, but also increased, rather than decreased, during muscle cell differentiation.

Having thoroughly demonstrated the presence of both Ago2 and miR-1 in the mitochondria, we next performed Ribo-IP to determine whether Ago2 is associated with mitochondrial transcripts. Using consecutively purified MT and MP (characterized in the lower panel of Figure 2D) from the same cell population, we found that anti-Ago2, but not control IgG, efficiently captured multiple mitochondrial transcripts (Figure 2D, upper panel). These semi-quantitative results were further conformed by RT-qPCR with RNA recovered from purified mitoplasts (Figure S2F). Therefore, in contrast to our initial expectation, the miRNA machinery appears to become increased, rather than decreased, in the mitochondria of differentiated C2C12 cells.

Identification of specific miR-1 targets in C2C12 cells

To determine specific targets for Ago2 and miR-1, we first designed a miRACE strategy to experimentally identify specific miRNA targets (Figure S3A), instead of relying on computational prediction, which is analogous to a published method (Easow et al., 2007). In this procedure, miRNA:mRNA duplexes were captured from whole cell extracts with anti-Ago2 followed by in situ extension of endogenous miRNA into target mRNA with reverse transcriptase. The extended products were poly(C)-tailed at the 3' end and PCR-amplified using a common primer linked to a poly(G) sequence and a miRNA-specific primer. While we are still refining this method for high throughput analysis, we initially focused on identifying miR-1 targets by cloning and sequencing a sizable number of PCR products. This revealed that several cytoplasmic targets for miR-1, one of which (ELL2) was further validated (see below). Importantly, we identified multiple mitochondrial transcripts for this miRNA (Figure S3B).

The second approach we took was to map specific Ago2 binding events by crosslinking immuno precipitation coupled with deep sequencing (CLIP-seq). We performed duplicated CLIP-seq on C2C12 myoblasts and myotubes, obtaining ~10 million uniquely mapped tags from each CLIP-seq library, which showed high concordance within the same cell types and large differences between the two cell types (Figure S3C). Compiling the normalized Ago2 binding events (to 1 million) on potential miR-1 cytoplasmic targets predicted by the miRNA target prediction program PITA (Kertesz et al., 2007), we observed no enrichment on myoblasts, but significant enrichment on myotubes, consistent with induced miR-1 only in myotubes (Figure 3A). Of five most characterized cytoplasmic miR-1 targets reported in

the literature, four showed a dramatic increase in Ago2 binding on the target sites (Figure 3B). HDAC4 exhibited an Ago2 peak before C2C12 differentiation, indicative of the involvement of another miRNA(s), but the Ago2 level was further enhanced after C2C12 differentiation (Figure 3B). Together, these data demonstrated the robustness of our CLIP-seq data in pinpointing specific miRNA targeting sites in the nuclear genome.

Critical insights from mapped Ago2-RNA interactions in the mitochondria

We then turned our attention to Ago2 interaction with the mitochondrial genome. Because the nuclear genome is populated with multiple mitochondrial pseudogenes, we took extra caution in mapping CLIP-seq tags to mtDNA by dividing the CLIP-seq tags into three classes that were (1) uniquely mapped to mitochondrial transcripts from mtDNA, (2) uniquely mapped to mitochondrial pseudogenes in the nuclear genome, and (3) commonly mapped to both nuclear and mitochondrial genomes (Table S1). This analysis revealed that the number of tags uniquely mapped to the nuclear mitochondrial pseudogenes is largely negligible. We therefore combined tags from class 1 and 3 (Table S2) to generate an Ago2 interaction map in the mitochondria. Significantly, we observed dramatic induction of Ago2 binding to multiple specific mitochondrial transcripts upon C2C12 differentiation (Figure 3C).

A major technical concern is potential contamination of abundant mitochondrial transcripts during immuno precipitation, which might be the reason that most published CLIP-seq studies ignored tags that were mapped to mitochondrial transcripts. To address this problem, we analyzed crosslink-induced mutation sites (CIMS), which is known to occur in internal regions, while insertions and point mutations due to sequencing errors are typically distributed at both ends (Zhang and Darnell, 2011) (Figure S3D). The CIMSs (blue tags on the opposite side of red CLIP tags) were largely concordant with the distribution of the CLIP tags mapped on mitochondrial transcripts, thus confirming direct interactions (Figure 3C).

Importantly, using the PITA program, we identified potential target regions for the two most highly induced miR-1 (green bars) and miR-206 (red bars), which shared related seed sequences. Two of these predicated sites exactly match the miRACE-identified target sites for miR-1 in ND1 and COX1 (blue bars in Figure 3C). These candidate miR-1 target sites correspond to multiple induced Ago2 CLIP-seq peaks. Specifically, we noted a major Ago2 CLIP-seq peak on ND1 (peak 1), another major peak on ATP8 (peak 3), multiple peaks on COX3 (peak 4), two small peaks on ND5 (peak 5 and 6), and one peak on cytb (peak 7) (Figure 3C). Interestingly, Ago2 seems to prevalently interact with 12S mitochondria-specific rRNA (Rrn1) in myoblasts, which was further induced after C2C12 differentiation to myotubes. Furthermore, Ago2 also shows broad interactions with multiple mitochondrial transcripts, including COX1, which contains a miR-1 target that had been both predicted and experimentally mapped. This broad binding pattern implies either multiple miRNA:mRNA interactions and/or some distinct functional properties of Ago2 in the mitochondria (see Discussion).

Polysome profiling of mitochondrial translation during muscle differentiation

A gold standard for studying translational control is through polysome profiling, a routine for investigating cytoplasmic translation, but a major challenge for detecting potential polysomes from the mitochondria partly because of coupled transcription/translation on the inner membrane (Bestwick and Shadel, 2013; Bonawitz et al., 2006). By optimizing the protocol for mitoribosome fractionation, we detected both small (28S) and large (39S) ribosome fractions as well as monosomes (55S) on a 10–30% sucrose gradient, which is characterized by the distribution of 12S and 16S rRNAs and their associated ribosome proteins (MRPS27 and MRPL24) (Figure 3D). These profiles resemble closely to those reported in the literature (Antonicka et al., 2013; Surovtseva et al., 2011). Interestingly, we also detected putative polysomes near the bottom of the gradient (fractions 12 and 13) on both C2C12 myoblasts and myotubes, which could be converted to monosomes by RNase I treatment (Figure 3D).

Importantly, we observed that Ago2 co-sediments with 28S and 39S ribosomes, and to a less extent, with monosomes or potential polysomes on myoblasts, and such distribution became broadened on myotubes, indicative of additional induced interactions (Figure 3D). This is consistent with Ago2 CLIP-seq signals on 12S rRNA in myoblasts, and to less extent, on 16S rRNA in myotubes (see Figure 3C). Significantly, by RT-qPCR quantification of ND1 and COX1 mRNAs and plotting the data according to percent distribution on the gradient, we found that both mitochondrial transcripts showed increased association with the putative polysome fractions from myotubes relative to myoblasts. These data provide initial support to enhanced mitochondrial translation upon C2C12 cell differentiation.

Opposite functions of miR-1 in the cytoplasm versus mitochondria

To determine the precise role of miR-1 in mitochondrial translation, we first validated the expected effect of miR-1 on repressing translation of its cytoplasmic target HDAC4 (Chen et al., 2006), and the newly identified target ELL2 (Figure 4A). Interestingly, we detected the opposite effects of miR-1 on mitochondrial ND1 and COX1 transcripts in undifferentiated C2C12 cells, which showed increased, rather than decreased, translation, while the level of COX2 was unaffected (Figure 4B). Using Ago2 knockout mouse embryonic fibroblasts (MEFs), we demonstrated that the miR-1 effects on both COX1 and ND1 were Ago2 dependent (Figure 4C).

To further characterize this positive effect, we performed a converse experiment using an antagomir to block miR-1 in C2C12 myotubes (Figure 4D). While antagomir-1 had little effect on ND1 and COX1 proteins before miR-1 induction in C2C12 myoblasts, the miR-1 inhibitor diminished miR-1 expression detected by Northern and blocked enhanced ND1 and COX1 translation measured by Western (Figure 4D). Consistent with the regulation at the translational level, little difference was detected on the mRNA level for both of these mitochondrial transcripts (Figure 4D, lower panel). We could also prevent miR-1 dependent translational enhancement by co-expressing a miR-1 sponge in the cytoplasm (Figure S4A).

As expected, the activities of ND1-containing Complex I and COX1-containing Complex IV were both elevated in miR-1 transfected myoblasts (Figure 4E). ATP production was also

enhanced upon C2C12 differentiation, which was significantly attenuated by antagomir-1 (Figure 4F). Despite relatively modest enhancement in these assays, which might be due to uncoordinated induction of other respiratory chain components in miR-1 transfected cells, these results linked miR-1 enhanced translation to increased respiratory activities.

Base-pairing requirement for miR-1 induced mitochondrial translation

To understand how miR-1 enhances mitochondrial translation, we first asked if there is anything unusual about the miR-1 targeting site in ND1 or COX1. We found that, as expected, the target sites cloned in the 3' UTR of a luciferase reporter mediated translational repression in the cytoplasm, and a mutant miR-1 lost the effect, which could be reversed by reverting the target sequences in the reporter to restore base pairing (Figure S4B and S4C). The stimulating effect of miR-1 did not seem to result from induced myogenesis or increased mtDNA replication (Figure S4D to S4F). We also found that miR-1 was able to enhance ND1 translation (Figure 4G) and ATP production (Figure 4H) in non-myogenic HeLa cells. These data argue against enhanced mitochondrial translation as an indirect result of the induced myogenic program.

The mapped miR-1 target sites in both ND1 and COX1 fulfill the established base-pairing rule for miRNA targeting (Figure 5A). The seed regions in both COX1 and ND1 appear to be more conserved during evolution compared to those in the middle and the 3' side (Figure S5A). We mutated either the seed region or the 3' sequence and found that both mutations abolished miR-1 induced ND1 and COX1 translation (Figure 5A and 5B). Again, the mRNA levels remained unaltered, and neither wild-type miR-1 nor its mutants exhibited detectable effect on the nuclear DNA-encoded mitochondrial protein COX4 (Figure 5B).

Direct action of miR-1 in mitochondrial translation

We took two additional strategies to further demonstrate the direct act of miR-1 within the mitochondria. We designed two separate miRNA mimics with altered targeting specificities. miR-COX1 would more efficiently target COX1, with reduced potential in targeting ND1, while miR-ND1 would only target ND1 (Figure 5C). The specificity of these designer miRNA mimics was confirmed on their cytoplasmic reporters (Figure S5B and S5C). In transfected C2C12 myoblasts, we found that miR-COX1 elevated COX1, with less effect on ND1, and as expected, miR-ND1 only stimulated ND1 (Figure 5D). We also tested another miRNA miR-101, which targets a specific region in ND1, and miR-499-5p, which targets a different region in ND1 (Figure 5E). We found that both exclusively enhanced ND1 translation, while several non-targeting miRNAs tested lacked the effect (Figure 5F). Because it is unlikely that specific miRNA mimics all caused the same indirect effects in the cytoplasm to stimulate mitochondrial translation, these data support the direct act of multiple miRNAs in enhancing mitochondrial translation.

To provide decisive evidence, we took advantage of Ago2 knockout MEFs to functionally rescue miR-1 dependent mitochondrial translation with either HA-tagged Ago2 or mitochondria-targeted Ago2 that carries a specific mitochondrial targeting signal peptide (su9). Confocal microscopy confirmed the colocalization of Su9-HA-Ago2 with Mito-GFP, a well-established marker for the mitochondria (Chen et al., 2003), while HA-Ago2 was

diffusely localized in the cytoplasm (Figure 5G). Using an RNAi-based luciferase reporter containing a perfect complementary sequence to miR-1, we found that the reporter was potently suppressed in wild-type, but not in Ago2 knockout MEFs (Figure 5H, upper panel). Exogenous HA-Ago2 rescued the RNAi effect and restored miR-1-enhanced translation of ND1 and COX1, and in sharp contrast, mitochondria-targeted Su9-HA-Ago2 could not rescue the RNAi effect in the cytoplasm, but potently restored the effect of miR-1 in the mitochondria (Figure 5H). Together with the specificity switch experiments, these data demonstrate the direct effect of miR-1 in enhancing mitochondrial translation.

Differential sensitivity to cytoplasmic and mitochondrial translational inhibitors

Thus far, we had measured the steady state levels of mitochondrial proteins to infer miR-1 regulated translation. To directly demonstrate enhanced mitochondrial translation, we used the methionine analog AHA to metabolically label nascent proteins in the mitochondria while blocking cytoplasmic translation by Emetine (Weraarpachai et al., 2009). In this experimental setting, C2C12 myoblasts were first transfected with miR-1 and then switched to the media containing Emetine. AHA was next added to allow its incorporation into nascent polypeptides, which were subsequently labeled by the reaction with fluorescent alkyne and resolved on SDS-PAGE. As expected, Emetine plus the mitochondrial translation inhibitor Chloramphenicol (INN) potently blocked all translation activities in the cell (Figure 6A, lanes 3 and 4).

In the presence of Emetine, but absence of INN, we detected translation products of all mtDNA-encoded transcripts. Compared to loading controls, we noted some general enhancement of mitochondrial translation by miR-1. In addition, miR-1 markedly enhanced COX1 translation (Figure 6A, compare lanes 5 and 6). We also reproducibly detected enhanced (although to less degrees) translation of ND1 (note that ND1 was less efficiently labeled, which is a well-known fact) as well as cytb, COX3, and ATP8 (red arrows in Figure 6A), all of which contain at least one predicted miR-1 targeting site each underlying a specific Ago2 CLIP-seq peak (see Figure 3C). In contrast, overexpressed miR-499-5p had no effect on COX1, but enhanced ND1 translation (Figure 6A, lanes 7 and 8), consistent with its targeting specificity. We also noted its ability to enhance ND4L, which also contains a predicted miR-499-5p targeting site (data not shown). Finally, using INN, we examined and ruled out the potential effect of miR-1 on the protein stability of ND1 (Figure 6B). Together, these data provide concrete evidence for miRNA-augmented mitochondrial translation.

Potential mechanism for miRNA-enhanced translation in the mitochondria

To gain mechanistic insights into miRNA-enhanced mitochondrial translation, we followed the clue from a *Drosophila* study, which suggests three critical requirements for converting miRNA-dependent translational repression to activation: (1) lack of the cap at the 5' end, (2) lack of a typical poly (A) tail at the 3' end, and (3) detachment of GW182 from an Ago protein (Iwasaki and Tomari, 2009; Vasudevan, 2012). Interestingly, all mitochondrial transcripts automatically fulfill the first two requirements because mitochondrial transcripts resemble mRNAs in prokaryotic cells by having no cap nor typical long poly(A) tail (Gagliardi et al., 2004; Temperley et al., 2010). This prompted us to experimentally test the

third requirement by determining whether GW182 maintains its partnership with Ago2 within the mitochondria.

We found that, compared to Ago2, GW182 was only detected in the cytoplasm (Figure 6C). Three GW182 paralogs are encoded in mammalian genomes, including TNRC6A (which corresponds to GW182), TNRC6B and TNRC6C (Ding and Han, 2007; Jakymiw et al., 2007) and published works indicate that all function in the miRNA pathway (Liu et al., 2005; Meister et al., 2005; Zipprich et al., 2009). We found that all three GW182 family members were excluded from the mitochondria (Figure 6C), suggesting that the miRNA machinery is rearranged in the mitochondria.

We took advantage of this mechanistic insight to further test the selective requirement for GW182 in miRNA-mediated functions in the cytoplasm versus the mitochondria. GW182 has been shown to be required for miRNA-mediated translational repression in the cytoplasm (Eulalio et al., 2008; Zipprich et al., 2009). As expected, GW182 knockdown prevented miR-1 mediated translational repression of its two cytoplasmic targets HDAC4 and Hand2 (Figure 6E, upper panel) as well as the cytoplasmic reporter containing the miRNA target site from ND1 (Figure 6E, low panel). Under these conditions, GW182 RNAi-treated C2C12 myoblasts continued to show enhanced ND1 translation by miR-1 (Figure 6D, upper panel). Based on these results, we propose that the rearranged miRNA machinery, coupled with the prokaryotic characteristics of the mitochondrial transcripts, is responsible for translational enhancement in the mitochondria.

Discussion

The findings presented in this report have important implications in diverse functions of the miRNA machinery in mammalian cells, miRNA-mediated switch of energy sources during cellular differentiation and reprogramming, and initiation of mitochondrial translation, which has been poorly understood despite intensive efforts in past decades. Below we discuss critical insights into each of these functional aspects.

Selective targeting of Ago2 and specific miRNAs to the mitochondria

The presence of the miRNA machinery in the mitochondria was only recently recognized (Bandiera et al., 2011; Barrey et al., 2011; Bian et al., 2010; Das et al., 2012; Kren et al., 2009; Sripada et al., 2012). In the present study, we carefully examined this critical issue by using highly purified mitoplasts to perform protease and nuclease protection assays and employed quantitative data to demonstrate that a significant fraction of Ago2 and miR-1 indeed reside within the mitochondria, especially when C2C12 cells are differentiated into myotubes.

An important question is how Ago2 is able to selectively enter the mitochondria. A previous study predicted a mitochondrial targeting sequence near the N-terminus of the Ago2 protein (Bandiera et al., 2011). We fused this sequence to GFP, but failed to detect its ability to target GFP to the mitochondria (data not shown). While still a mystery, it is important to note that most nuclear DNA-encoded mitochondrial proteins do not contain any identifiable mitochondrial targeting sequences, indicating the existence of other mechanisms for

importing cytoplasmic proteins into the mitochondria, such as communication between mitochondria and ER where Ago2 was initially identified (Cikaluk et al., 1999; Kornmann et al., 2009).

Although 5SrRNA and tRNA can be actively imported into mitochondria (Mahata et al., 2006; Wang et al., 2010), we do not know how exactly miRNAs enter the mitochondria. Our preliminary mitochondrial import assay indicates that multiple specific and non-specific miRNAs, but not their precursors, were able to enter purified mitochondria without the need for ATP hydrolysis (data not shown). We thus suspect that, while most miRNAs may be tied up on their cytoplasmic targets, thus preventing them from free diffusion, a subset of “free” miRNAs, including those newly induced, might enter the mitochondria. Clearly, the mechanism for targeting a subset of miRNAs and specific components of the miRNA machinery to the mitochondria awaits future studies.

Enhanced translation by the miRNA/Ago2 complex in the mitochondria

We provided a series of evidence for direct action of the miRNA/Ago2 complex in enhancing mitochondrial translation, including (1) direct Ago2 binding to specific mtDNA-encoded transcripts, (2) the requirement for specific base-pairing interactions to elicit the effect, (3) functional rescue of Ago2 knockout MEFs with mitochondria-targeted Ago2, and (4) selective inactivation of the miRNA machinery in the cytoplasm by GW182 knockdown. The latter two pieces of information demonstrate that miR-1 is able to stimulate mitochondrial translation when the miRNA machinery is selectively inactivated in the cytoplasm.

It is currently unclear how the miRNA/Ago2 complex precisely augments mitochondrial translation. In fact, this is related to a fundamental question with respect to translation initiation in the mitochondria, despite of extensive efforts in the past decades. This is because the fully processed mitochondrial transcripts do not contain a cap structure (a key feature of eukaryotic cytoplasmic mRNAs) nor contain any identifiable ribosomal binding site (a mechanism used by bacteria). Based on our Ago2 mapping data, it is curious to note extensive interactions of Ago2 with the 12S rRNA. We also observed extensive association of Ago2 with mitochondrial transcripts, some with more specific interactions while others in a more continuous fashion as if those transcripts are “wrapped” around the Ago2-containing complex. As such, Ago2 may function as a key mitochondrial translation initiation factor to facilitate ribosome-mRNA interactions.

A coordinated myogenic program regulated by miR-1

Our findings also extend the physiological function of miR-1 in skeletal and cardiac muscle. Genetic ablation of this miRNA demonstrated a profound effect on heart functions in the mouse (Zhao et al., 2007). However, as the heart contains fully differentiated cardiomyocytes in adult, it has been unclear how this miRNA might contribute to muscle physiology. Our data demonstrate that this miRNA is able to boost ATP production in the mitochondria, which is in high demand for heart contraction, thus providing a plausible explanation for its functional requirement in the heart.

Published studies have demonstrated a vital role of miR-1 in inhibiting HDAC4 (Chen et al., 2006), which suppresses Mef2 (Arnold et al., 2007), and elevated Mef2 further activates the expression of miR-1 (Liu et al., 2007), thereby forming an auto-regulatory loop to maintain the myogenic program. Our data add a further role of miR-1 in enhancing protein synthesis and ATP production in the mitochondria of differentiated muscle cells and suggest miR-1 in coordinating regulatory networks in both the cytoplasm and the mitochondria during and after muscle differentiation, as illustrated in Figure 6F. Therefore, the newly elucidated role of the miRNA complex in the mitochondria suggests a new layer of crosstalk between nuclear and mitochondrial genomes in eukaryotic cells.

Experimental Procedures

Isolation of mitochondria and purification of mitoplasts

Cardiomyocytes from mouse heart, C2C12-derived myoblasts or myotubes were collected by centrifugation, homogenized in a pre-chilled Dounce homogenizer (Kontes), and post-nuclear supernatant collected. Mitochondria were sedimented at 13,000g for 10 min, washed once in the same buffer, and further purified by centrifugation at 40,000g for 1 hr at 4°C on a sucrose gradient (17, 31, 42, 50%) in T₁₀E₂₀buffer (10 mM Tris-HCl pH 7.4, 20 mM EDTA). Purification of mitoplasts was according to the previous procedure (Schnaitman and Greenawalt, 1968).

Trypsin/Nuclease protection assays

For trypsin protection, isolated mitochondria and mitoplasts were either kept on ice as control or treated with Trypsin (100 µg/ml) alone or with Trypsin (100 µg/ml) plus Triton X-100 (1%). Treated mitochondria or mitoplasts were collected by centrifugation at 18,000g for 3 min and the pellets were re-suspended in SDS-PAGE sample buffer for Western blotting. The nuclease protection was similarly performed on isolated mitochondria and mitoplasts with RNaseT1 (3000 U/mL) plus Micrococcal nuclease (1000 U/mL) digestion on ice for 30 min in the presence or absence of Triton X-100. RNA was extracted with Trizol from pelleted mitochondria or mitoplasts for RT-qPCR.

Analysis of mitochondrial translation by polysome profiling

Sucrose gradient sedimentation of mitochondrial ribosome/polysome was carried out as described (Antonicka et al., 2013). Briefly, mitochondria purified from C2C12 myoblasts or myotubes were suspended in the lysis buffer (260 mM sucrose, 100 mM KCl, 20 mM MgCl₂, 10 mM Tris-pH 7.5, 1% Triton X-100, 5 mM β-mercaptoethanol, and a cocktail of protease inhibitors from Roche) on ice for 20 min. The lysate was centrifuged at 9000g for 30 min to remove particles, loaded on an 8 ml 10–30% sucrose gradient, and centrifuged at 180,000g for 260 min in a Beckman SW41-Ti rotor. To characterize putative polysomes, the lysate was treated with 5 U/µL RNase I for 40 min at 25°C. After centrifugation, 13 fractions were collected for RNA and protein analysis.

Detection of nascent protein synthesis

Click-iT reagents (Life Technology) were used to label nascent polypeptides. C2C12 myoblasts were first transfected with miRNA or control RNA at 30–40% confluence in 24-

well plates. After culturing overnight, the plate was washed 3 times with pre-warmed PBS. The cells were incubated in methionine-free DMEM for 45 min followed by the addition of 20 μ M of Emetine, and incubated for another 15 min. The media were replaced with methionine-free DMEM containing 50 μ M methionine analog AHA and 20 μ M Emetine. After incubation for 4 hrs, AHA-containing nascent proteins were labeled with TAMRA (Life Technology), resolved in 4–12% gradient SDS-PAGE, and analyzed by Typhoon 9400. These and other methods are detailed in Extended Experimental Procedures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Drs. ImmoScheffler, Amy Pasquinelli, Shobha Vasudevan, and David Chan for insightful discussion during the course of this investigation, Dr. Xiaoyan Ding for providing the C2C12 cell line, Dr. Anna Lombes for the gift of anti-ND1 antibody. This work was supported by the China 973 program (2011CB811300, 2012CB910800), the Chinese 111 program grant (B06018) and NIH grants (GM049369 and GM052872).

References

- Antonicka H, Sasarman F, Nishimura T, Paupe V, Shoubbridge EA. The mitochondrial RNA-binding protein GRSF1 localizes to RNA granules and is required for posttranscriptional mitochondrial gene expression. *Cell metabolism*. 2013; 17:386–398. [PubMed: 23473033]
- Arnold MA, Kim Y, Czubryt MP, Phan D, McAnally J, Qi X, Shelton JM, Richardson JA, Bassel-Duby R, Olson EN. MEF2C transcription factor controls chondrocyte hypertrophy and bone development. *Developmental cell*. 2007; 12:377–389. [PubMed: 17336904]
- Bandiera S, Ruberg S, Girard M, Cagnard N, Hanein S, Chretien D, Munnich A, Lyonnet S, Henrion-Caude A. Nuclear outsourcing of RNA interference components to human mitochondria. *PloS one*. 2011; 6:e20746. [PubMed: 21695135]
- Barrey E, Saint-Auret G, Bonnamy B, Damas D, Boyer O, Gidrol X. Pre-microRNA and mature microRNA in human mitochondria. *PloS one*. 2011; 6:e20220. [PubMed: 21637849]
- Bestwick ML, Shadel GS. Accessorizing the human mitochondrial transcription machinery. *Trends in biochemical sciences*. 2013; 38:283–291. [PubMed: 23632312]
- Bian Z, Li LM, Tang R, Hou DX, Chen X, Zhang CY, Zen K. Identification of mouse liver mitochondria-associated miRNAs and their potential biological functions. *Cell Res*. 2010; 20:1076–1078. [PubMed: 20733615]
- Bonawitz ND, Clayton DA, Shadel GS. Initiation and beyond: multiple functions of the human mitochondrial transcription machinery. *Molecular cell*. 2006; 24:813–825. [PubMed: 17189185]
- Bosmann HB, Myers MW, Dehond D, Ball R, Case KR. Mitochondrial autonomy. Sialic acid residues on the surface of isolated rat cerebral cortex and liver mitochondria. *J Cell Biol*. 1972; 55:147–160. [PubMed: 4653414]
- Carthew RW, Sontheimer EJ. Origins and Mechanisms of miRNAs and siRNAs. *Cell*. 2009; 136:642–655. [PubMed: 19239886]
- Cavelier L, Johannisson A, Gyllensten U. Analysis of mtDNA copy number and composition of single mitochondrial particles using flow cytometry and PCR. *Exp Cell Res*. 2000; 259:79–85. [PubMed: 10942580]
- Cernilogar FM, Onorati MC, Kothe GO, Burroughs AM, Parsi KM, Breiling A, Lo Sardo F, Saxena A, Miyoshi K, Siomi H, et al. Chromatin-associated RNA interference components contribute to transcriptional regulation in *Drosophila*. *Nature*. 2011; 480:391–395. [PubMed: 22056986]
- Chekulaeva M, Filipowicz W. Mechanisms of miRNA-mediated post-transcriptional regulation in animal cells. *Current opinion in cell biology*. 2009; 21:452–460. [PubMed: 19450959]

- Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE, Chan DC. Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic development. *The Journal of cell biology*. 2003; 160:189–200. [PubMed: 12527753]
- Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, Conlon FL, Wang DZ. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nature genetics*. 2006; 38:228–233. [PubMed: 16380711]
- Cikaluk DE, Tahbaz N, Hendricks LC, DiMattia GE, Hansen D, Pilgrim D, Hobman TC. GERp95, a membrane-associated protein that belongs to a family of proteins involved in stem cell differentiation. *Molecular biology of the cell*. 1999; 10:3357–3372. [PubMed: 10512872]
- Czech B, Hannon GJ. Small RNA sorting: matchmaking for Argonautes. *Nature reviews Genetics*. 2011; 12:19–31.
- Das S, Ferlito M, Kent OA, Fox-Talbot K, Wang R, Liu D, Raghavachari N, Yang Y, Wheelan SJ, Murphy E, et al. Nuclear miRNA regulates the mitochondrial genome in the heart. *Circulation research*. 2012; 110:1596–1603. [PubMed: 22518031]
- de Brito OM, Scorrano L. Mitofusin 2 tethers endoplasmic reticulum to mitochondria. *Nature*. 2008; 456:605–610. [PubMed: 19052620]
- Ding L, Han M. GW182 family proteins are crucial for microRNA-mediated gene silencing. *Trends in cell biology*. 2007; 17:411–416. [PubMed: 17766119]
- Easow G, Teleman AA, Cohen SM. Isolation of microRNA targets by miRNP immuno purification. *Rna*. 2007; 13:1198–1204. [PubMed: 17592038]
- Eulalio A, Huntzinger E, Izaurralde E. GW182 interaction with Argonaute is essential for miRNA-mediated translational repression and mRNA decay. *Nat Struct Mol Biol*. 2008; 15:346–353. [PubMed: 18345015]
- Franko A, Mayer S, Thiel G, Mercy L, Arnould T, Hornig-Do H, Wiesner R, Goffart S. CREB-1a Is Recruited to and Mediates Upregulation of the Cytochrome c Promoter during Enhanced Mitochondrial Biogenesis Accompanying Skeletal Muscle Differentiation. *Molecular and Cellular Biology*. 2008; 28:2446–2459. [PubMed: 18227154]
- Gagliardi D, Stepien PP, Temperley RJ, Lightowlers RN, Chrzanowska-Lightowlers ZM. Messenger RNA stability in mitochondria: different means to an end. *Trends in genetics : TIG*. 2004; 20:260–267. [PubMed: 15145579]
- Guang S, Bochner AF, Pavelec DM, Burkhardt KB, Harding S, Lachowiec J, Kennedy S. An Argonaute transports siRNAs from the cytoplasm to the nucleus. *Science*. 2008; 321:537–541. [PubMed: 18653886]
- Iwasaki S, Tomari Y. Argonaute-mediated translational repression (and activation). *Fly*. 2009; 3:204–206. [PubMed: 19556851]
- Jakymiw A, Pauley KM, Li S, Ikeda K, Lian S, Eystathioy T, Satoh M, Fritzler MJ, Chan EK. The role of GW/P-bodies in RNA processing and silencing. *Journal of cell science*. 2007; 120:1317–1323. [PubMed: 17401112]
- Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E. The role of site accessibility in microRNA target recognition. *Nature genetics*. 2007; 39:1278–1284. [PubMed: 17893677]
- Kornmann B, Currie E, Collins SR, Schuldiner M, Nunnari J, Weissman JS, Walter P. An ER-mitochondria tethering complex revealed by a synthetic biology screen. *Science*. 2009; 325:477–481. [PubMed: 19556461]
- Kren BT, Wong PY, Sarver A, Zhang X, Zeng Y, Steer CJ. MicroRNAs identified in highly purified liver-derived mitochondria may play a role in apoptosis. *RNA Biol*. 2009; 6:65–72. [PubMed: 19106625]
- Liu J, Rivas FV, Wohlschlegel J, Yates JR 3rd, Parker R, Hannon GJ. A role for the P-body component GW182 in microRNA function. *Nat Cell Biol*. 2005; 7:1261–1266. [PubMed: 16284623]
- Liu N, Williams AH, Kim Y, McAnally J, Bezprozvannaya S, Sutherland LB, Richardson JA, Bassel-Duby R, Olson EN. An intragenic MEF2-dependent enhancer directs muscle-specific expression of microRNAs 1 and 133. *Proc Natl Acad Sci U S A*. 2007; 104:20844–20849. [PubMed: 18093911]

- Lopaschuk GD, Ussher JR, Folmes CD, Jaswal JS, Stanley WC. Myocardial fatty acid metabolism in health and disease. *Physiol Rev.* 2010; 90:207–258. [PubMed: 20086077]
- Mahata B, Mukherjee S, Mishra S, Bandyopadhyay A, Adhya S. Functional delivery of a cytosolic tRNA into mutant mitochondria of human cells. *Science.* 2006; 314:471–474. [PubMed: 17053148]
- Meister G, Landthaler M, Peters L, Chen PY, Urlaub H, Luhrmann R, Tuschl T. Identification of novel argonaute-associated proteins. *Current biology : CB.* 2005; 15:2149–2155. [PubMed: 16289642]
- Mortensen RD, Serra M, Steitz JA, Vasudevan S. Posttranscriptional activation of gene expression in *Xenopus laevis* oocytes by microRNA-protein complexes (microRNPs). *Proc Natl Acad Sci U S A.* 2011; 108:8281–8286. [PubMed: 21536868]
- Moyes C, Mathieu-Costello O, Tsuchiya N, Filburn C, Hansford R. Mitochondrial biogenesis during cellular differentiation. *American Journal of Physiology- Cell Physiology.* 1997; 272:1345–1351.
- Rao PK, Kumar RM, Farkhondeh M, Baskerville S, Lodish HF. Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc Natl Acad Sci U S A.* 2006; 103:8721–8726. [PubMed: 16731620]
- Rizzuto R, Pinton P, Carrington W, Fay FS, Fogarty KE, Lifshitz LM, Tuft RA, Pozzan T. Close contacts with the endoplasmic reticulum as determinants of mitochondrial Ca²⁺ responses. *Science.* 1998; 280:1763–1766. [PubMed: 9624056]
- Saraste M. Oxidative phosphorylation at the fin de siecle. *Science.* 1999; 283:1488–1493. [PubMed: 10066163]
- Schnaitman C, Greenawalt JW. Enzymatic properties of the inner and outer membranes of rat liver mitochondria. *J Cell Biol.* 1968; 38:158–175. [PubMed: 5691970]
- Sripada L, Tomar D, Prajapati P, Singh R, Singh AK, Singh R. Systematic analysis of small RNAs associated with human mitochondria by deep sequencing: detailed analysis of mitochondrial associated miRNA. *PLoS one.* 2012; 7:e44873. [PubMed: 22984580]
- Suh N, Baehner L, Moltzahn F, Melton C, Shenoy A, Chen J, Blelloch R. MicroRNA function is globally suppressed in mouse oocytes and early embryos. *Current biology : CB.* 2010; 20:271–277. [PubMed: 20116247]
- Surovtseva YV, Shutt TE, Cotney J, Cimen H, Chen SY, Koc EC, Shadel GS. Mitochondrial ribosomal protein L12 selectively associates with human mitochondrial RNA polymerase to activate transcription. *Proc Natl Acad Sci U S A.* 2011; 108:17921–17926. [PubMed: 22003127]
- Taanman JW. The mitochondrial genome: structure, transcription, translation and replication. *Biochimica et biophysica acta.* 1999; 1410:103–123. [PubMed: 10076021]
- Temperley RJ, Wydro M, Lightowlers RN, Chrzanoska-Lightowlers ZM. Human mitochondrial mRNAs—like members of all families, similar but different. *Biochimica et biophysica acta.* 2010; 1797:1081–1085. [PubMed: 20211597]
- Vasudevan S. Posttranscriptional upregulation by microRNAs. *Wiley interdisciplinary reviews RNA.* 2012; 3:311–330. [PubMed: 22072587]
- Vasudevan S, Tong Y, Steitz J. Switching from Repression to Activation: MicroRNAs Can Up-Regulate Translation. *Science.* 2007; 318:1931. [PubMed: 18048652]
- Wang D, Zhang Z, O’Loughlin E, Lee T, Houel S, O’Carroll D, Tarakhovsky A, Ahn NG, Yi R. Quantitative functions of Argonaute proteins in mammalian development. *Genes & development.* 2012; 26:693–704. [PubMed: 22474261]
- Wang G, Chen HW, Oktay Y, Zhang J, Allen EL, Smith GM, Fan KC, Hong JS, French SW, McCaffery JM, et al. PNPASE regulates RNA import into mitochondria. *Cell.* 2010; 142:456–467. [PubMed: 20691904]
- Wei W, Ba Z, Gao M, Wu Y, Ma Y, Amiard S, White CI, Rendtlew Danielsen JM, Yang YG, Qi Y. A role for small RNAs in DNA double-strand break repair. *Cell.* 2012; 149:101–112. [PubMed: 22445173]
- Weraarpachai W, Antonicka H, Sasarman F, Seeger J, Schrank B, Kolesar JE, Lochmuller H, Chevrette M, Kaufman BA, Horvath R, et al. Mutation in TACO1, encoding a translational activator of COX I, results in cytochrome c oxidase deficiency and late-onset Leigh syndrome. *Nature genetics.* 2009; 41:833–837. [PubMed: 19503089]

- Zhang C, Darnell RB. Mapping in vivo protein-RNA interactions at single-nucleotide resolution from HITS-CLIP data. *Nature biotechnology*. 2011; 29:607–614.
- Zhang Y, Liu D, Chen X, Li J, Li L, Bian Z, Sun F, Lu J, Yin Y, Cai X, et al. Secreted monocytic miR-150 enhances targeted endothelial cell migration. *Mol Cell*. 2010; 39:133–144. [PubMed: 20603081]
- Zhao Y, Ransom JF, Li A, Vedantham V, von Drehle M, Muth AN, Tsuchihashi T, McManus MT, Schwartz RJ, Srivastava D. Dysregulation of cardiogenesis, cardiac conduction, and cell cycle in mice lacking miRNA-1-2. *Cell*. 2007; 129:303–317. [PubMed: 17397913]
- Zipprich JT, Bhattacharyya S, Mathys H, Filipowicz W. Importance of the C-terminal domain of the human GW182 protein TNRC6C for translational repression. *Rna*. 2009; 15:781–793. [PubMed: 19304925]

Highlights

- miR-1, Argonaute 2, but not GW182, are localized in the mitochondria
- Argonaute 2 binds mitochondrial transcripts and ribosomal RNAs
- microRNAs are able to directly enhance mitochondrial translation
- miR-1 mediates a coordinated myogenic program for muscle differentiation




Figure 1. Increased mitochondrial proteins without a significant elevation in mtDNA copy number or transcription

(A) Levels of COX1 and ND1 proteins detected by Western blotting and miR-1 by Northern blotting in different adult mouse tissues. Levels of Histone H3 and 5S rRNA served as protein and RNA loading controls, respectively.

(B) Western and TaqMan PCR analysis of COX1, ND1 and miR-1 in C2C12 cells before and after differentiation. Panactin and VDAC served as protein loading controls and 28S rRNA as an RNA loading control. MHC was analyzed as a control for the induction of the myogenic program. Tfam and TACO1 were analyzed to detect potential regulation of critical mitochondrial transcriptional and translational regulators.

(C, D, E) Quantification of proteins normalized to pan actin (C), mtDNA normalized to nuclear DNA (D), and RNA normalized to GAPDH, (E) during C2C12 cell differentiation. See also Fig. S1. Data in C and E are based on three biological repeats and data in D on ten biological repeats. Errors bars are mean \pm SD.




Figure 2. Detection and quantitative analysis of Ago2 and miR-1 in the mitochondria

(A) Trypsin protection analysis of purified mitochondria (MT) and mitoplasts (MP) from adult mouse heart, C2C12 myoblasts, and C2C12 myotubes. See also Fig. S2A.

(B) Quantification of Ago2 and miR-1 per C2C12 nucleus or per mitochondrial genome. The average number is based on two independent experiments. See also Fig. S2C, D.

(C) Nuclease protection analysis. Three aliquots of purified mitochondria and mitoplasts were treated with isolation buffer, with RNaseT1 (RN) plus MNase (MN), or with the combination of nucleases in the presence of Triton X-100.

(D) Purified mitochondria and mitoplasts from the same fractionation experiment were first characterized by the lack of the ER marker ERp72, but the presence of outer membrane protein marker Tom20 in the mitochondria and the inner membrane marker NDUFB8 in both mitochondria and mitoplasts (lower panel). Equal amounts of purified mitochondria and mitoplasts based on the levels of 16S rRNA were subjected to ribo-IP with anti-Ago2 or control IgG (upper panel). Western blotting data showed specific anti-Ago2 IP. Individual mitochondrial transcripts were examined by semi-quantitative RT-PCR. See also Fig. S2E, F for real-time RT-PCR analysis of Ago2-associated mitochondrial transcripts from purified mitoplasts.




Figure 3. Mapping of Ago2-RNA interactions by CLIP-seq and analysis of mitochondrial translation by polysome profiling

(A) Ago2 CLIP-seq profiles on predicted miR-1 targets before and after C2C12 differentiation. Duplicated data in each cell type were individually plotted against the center of PITA predicted miR-1 target sites.

(B) Ago2 CLIP tags on known cytoplasmic targets for miR-1 before (upper panels) and after (low panels) C2C12 differentiation. The chromosomal location of each target site is shown on the top of each panel. Red: CLIP tags; Blue: CIMS tags.

(C) The distribution of Ago2 CLIP-seq tags in the mitochondrial genome before and after C2C12 cell differentiation. The annotated genes are displayed on the top. PITA predicted target sites for miR-1 (green) and miR-206 (red) are indicated right below the gene track. The CLIP tag distribution (red) and corresponding CIMS (dark blue) are separately mapped to the mitochondrial genome in C2C12 myoblasts cultured in growth media (GM) and in

C2C12 myotubes maintained in differentiation media (DM) for 4 days. The locations of miRACE identified miR-1 target sites are indicated between the two CLIP panels.

(D) Analysis of mitochondrial translation by polysome profiling. Purified mitochondria were used for fractionation on a sucrose gradient, which was free of cytoplasmic ribosomes, as indicated by the lack of Western blotting for representative cytoplasmic ribosomal proteins (RPS3 and RPL4). Representative mitochondrial ribosomal proteins (MRPS27 and MRPL24) and Ago2 on individual gradient fractions were detected by Western blotting and specific rRNA, mRNA and miRNA transcripts were quantified by real-time RT-PCR (Taqman for miR-1). The assignment of small and large ribosomal subunits, monosomes, and putative polysomes was based on the distribution of both rRNAs and proteins and comparison with published mitochondrial polysome profiles. The putative polysome fractions were characterized by RNase I treatment. The relative abundance of individual transcripts in each fraction was presented as the percentage of the total fraction. The increased polysome association of ND1 and COX1 transcripts is highlighted with dark blue bars proportional to their enhancement from myoblasts to myotubes. See also Fig. S3.




Figure 4. Opposite effects of miRNAs on translation in the cytoplasm versus mitochondria
 (A) Western blotting of HDAC4 and ELL2 in C2C12 myoblasts after transfection with miR-1 or control siRNA.
 (B) Western blotting of ND1 and COX1 in C2C12 myoblasts 48 hrs after transfection with control RNA or miR-1.
 (C) ND1 and COX1 protein levels in wild-type and Ago2 knockout MEFs transfected with either control RNA or miR-1.
 (D) Antagomir-1 blocked miR-1 expression and diminished enhanced ND1 and COX1 translation in myotubes. In this experiment, C2C12 myoblasts were first transfected with control RNA or antagomir-1 and subsequently transferred to differentiation media after 24 hrs. Western blotting of ND1 and COX1 was performed after culturing the cells for another 3 days. Actin and 5S rRNA served as loading controls for protein and RNA. The lower panel showed that antagomir-1 had no effect on the mRNA levels of both ND1 and COX1.
 (E) Enhanced activities of Complex I and IV by miR-1 in transfected C2C12 myoblasts.
 (F) Antagomir-1 blocked enhanced ATP production in C2C12 myotubes.
 (G) miR-1 enhanced ND1 protein synthesis without affect its mRNA levels in transfected HeLa cells.
 (H) miR-1 induced ATP production in transfected HeLa cells.
 Data in D, E, F, G, H are based on three biological repeats and shown as mean \pm SD; p-values are indicated in each panel. See also Fig. S4.




Figure 6. miR-1 enhanced protein synthesis in the mitochondria

(A) Detection of nascent protein synthesis in the mitochondria when cytoplasmic translation was blocked by Emetine. C2C12 myoblasts were first transfected with a control RNA or miR-1 or miR-499-5p and protein synthesis was monitored by AHA incorporation in the presence of Emetine. Cytoplasmic Actin and mitochondrial VDAC served as loading controls. Emetine blocked all cytoplasmic translation to allow detection of mitochondrial translational activities. Red arrows highlight individual elevated bands by miR-1 or miR-499-5p.

(B) Lack of effect of miR-1 on ND1 protein stability. The protein levels were determined in the presence of INN in control and miR-1 transfected C2C12 cells and the data were normalized against the protein level of ND1 at the 0 time point.

(C) Restriction of GW182 and its paralogs TNRC6B and TNRC6C from the mitochondria. Lamin A/C was purified away from the mitochondria. ND1 and Tom20 served as controls for proteins localized within the mitochondria or associated with the outer membrane of purified mitochondria.

(D) miR-1 enhanced mitochondrial translation when the miRNA machinery was selectively inactivated by knocking down GW182 in the cytoplasm. GW182 showed the expected requirement for miR-1 to act on its cytoplasmic targets HDAC4 and Hand2 (upper panel) and on a cytoplasmic luciferase reporter containing the miR-1 targeting site in ND1 (lower panel). When the function of the miRNA machinery was largely compromised in the cytoplasm of GW182 knockdown cells, transfected miR-1 continued to show the ability to enhance ND1 translation in the mitochondria.

(E) Expanded roles of miR-1 in muscle differentiation. miR-1 is known to be induced by SRF, MyoD and Mef2 during differentiation. Induced miR-1 has been thought to act on its cytoplasmic targets to inhibit cell proliferation via Hand2 and promote cell differentiation through HDAC4 and Mef2. The newly elucidated function of miR-1 contributes to efficient protein synthesis in the mitochondria, resulting in boosted ATP production to meet the increasing energy demand in differentiating muscle cells. Therefore, miR-1 mediated translational repression in the cytoplasm and translational enhancement in the mitochondria

may constitute a highly coordinated myogenic program for muscle differentiation and function.

Data in B and D are based on three biological repeats and shown as mean \pm SD with p-values indicated in D.