



Published in final edited form as:

Cell Rep. 2014 July 24; 8(2): 382–392. doi:10.1016/j.celrep.2014.06.027.

## Appropriate crypt formation in the uterus for embryo homing and implantation requires Wnt5a-ROR signaling

Jeeyeon Cha<sup>1,2</sup>, Amanda Bartos<sup>1</sup>, Craig Park<sup>1</sup>, Xiaofei Sun<sup>1</sup>, Yingju Li<sup>1</sup>, Sang-Wook Cha<sup>2</sup>, Rieko Ajima<sup>3,4</sup>, Hsin-Yi Henry Ho<sup>5,6</sup>, Terry P. Yamaguchi<sup>3</sup>, and Sudhansu K. Dey<sup>1,2,\*</sup>

<sup>1</sup>Division of Reproductive Sciences, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA

<sup>2</sup>Division of Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA

<sup>3</sup>Cancer and Developmental Biology Laboratory, Center for Cancer Research, National Cancer Institute-Frederick, NIH, Frederick, MD 21702, USA

<sup>4</sup>Division of Mammalian Development, National Institute of Genetics, Yata 1111, Mishima 411-8540, Japan

<sup>5</sup>Department of Neurobiology, Harvard Medical School, Boston, MA 02115

<sup>6</sup>Department of Cell Biology and Human Anatomy, UC Davis School of Medicine, Davis, CA 95616, USA

### Summary

Embryo homing and implantation occur within a crypt (implantation chamber) at the antimesometrial (AM) pole along the uterus. The mechanism by which this is achieved is not known. Here we show that villi-like epithelial projections from the main uterine lumen towards the AM pole at regularly spaced intervals to form crypts for embryo implantation were disrupted in mice with uterine loss or gain of function of *Wnt5a*, or loss of function of both *Ror1* and *Ror2*. This disruption of Wnt5a-ROR signaling resulted in disorderly epithelial projections, crypt formation, embryo spacing, and impaired implantation. These early disturbances under abnormal Wnt5a-ROR signaling were reflected in adverse late pregnancy events, including defective decidualization and placentation, ultimately leading to compromised pregnancy outcome. This

© 2014 The Authors. Published by Elsevier Inc. All rights reserved.

\*Address for correspondence: SK Dey, Cincinnati Children's Hospital Medical Center, Division of Reproductive Sciences, MLC 7045, 3333 Burnet Avenue, Cincinnati, Ohio 45229, USA; Phone: 513-803-1158; sk.dey@cchmc.org.

### Author contributions

J.C., C.P., X.S., S.-W.C., R.A., H.-Y.H.H., T.P.Y. and S.K.D. designed experiments. J.C., A.B., C.P., X.S., Y.L. and R.A. performed all animal, biochemical, histological, molecular and cell biological experiments. S.-W.C., H.-Y.H.H., R.A. and T.P.Y. provided reagents, mouse lines, and technical advice. J.C., A.B., C.P., X.S., Y.L., S.-W.C., R.A., H.-Y.H.H., T.P.Y. and S.K.D. analyzed data. J.C., C.P., X.S., H.-Y.H.H., T.P.Y. and S.K.D. wrote the manuscript.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

study presents deeper insight regarding the formation of organized epithelial projections for crypt formation and embryo implantation for pregnancy success.

---

## Introduction

Reciprocal interactions between the receptive uterus and blastocyst are critical to implantation. The uterus undergoes morphological, cellular and molecular changes during pregnancy. This plasticity is reflected in its requirement for transcription factors, homeotic proteins, growth factors, morphogens and signaling molecules (Cha et al., 2012). Genetic intervention of these signaling pathways results in defective or deferred implantation, which propagates adverse ripple effects throughout the remainder of gestation, compromising pregnancy success (Song et al., 2002; Ye et al., 2005; Sun et al., 2012).

Blood vessels enter the uterus through the mesometrium, positioning the uterus along a mesometrial-antimesometrial (**M-AM**) axis. Normally, implantation occurs within a crypt (implantation chamber) at the AM pole as evident from histology of uterine cross-sections (Daikoku et al., 2011). Blastocyst attachment to the luminal epithelium (**LE**) occurs in the evening of day 4 of pregnancy (1800–2000 h) and is coincident with increased endometrial vascular permeability at the site of blastocyst attachment. In mice, this vascular permeability can be monitored by intravenous injection of a blue dye solution, which marks the implantation sites (**IS**) as distinct blue bands (Psychoyos, 1973). With progression of implantation on day 5, stromal cells around the implantation chamber undergo proliferation and differentiation into specialized decidual cells (decidualization). Decidualization supports embryonic growth and development and later directs placentation to establish maternal-fetal vascular connection.

Wnt signaling is a conserved pathway with roles in organogenesis and cell fate determination (Angers and Moon, 2009; MacDonald et al., 2009; van Amerongen and Nusse, 2009). In canonical signaling, Wnt ligands bind Frizzled (Fzd) and LRP5/6 receptors to stabilize  $\beta$ -catenin for nuclear translocation;  $\beta$ -catenin then binds to TCF/LEF transcription factors and activates gene transcription. In non-canonical signaling, Wnts function independently of  $\beta$ -catenin-mediated transcription to direct cell movements, cell shape and polarity (Kikuchi et al., 2011; van Amerongen and Nusse, 2009; Veeman et al., 2003). Wnt5a is considered a non-canonical Wnt ligand, and *Wnt5a* deletion in *Xenopus*, zebrafish, and mice disturbs directed cellular movements and polarity (Gao et al., 2011; Miyoshi et al., 2012; Schambony and Wedlich, 2007; Yamaguchi et al., 1999).

Several receptors including Frizzled, ROR, and Ryk were shown to mediate non-canonical Wnt5a signaling in various systems (Green et al., 2008; Macheda et al., 2012; Minami et al., 2010; Niehrs, 2012). Significance of Wnt5a-ROR signaling in morphogenetic movement was recently reported in mouse embryogenesis (Ho et al., 2012). However, its role in the adult female reproductive tract during pregnancy is unknown.

Utilizing *LoxP-Cre* system, we show here that an appropriate dosage of Wnt5a is required for normal blastocyst attachment and implantation. Females with either uterine inactivation (*Wnt5a*<sup>d/d</sup>) or overexpression (*Wnt5a*<sup>GOF</sup>) of *Wnt5a* are sterile or give birth to small litters.

This subfertility phenotype is due to aberrant blastocyst attachment arising from defective LE organization, crypt formation and embryo positioning in the uterus. These aberrations led to defective implantation, embryo crowding, poor decidualization and placentation, leading to increased resorption rate and subfertility. Mice with uterine inactivation of both *Ror1* and *Ror2* (*Ror1<sup>d/d</sup>/Ror2<sup>d/d</sup>*) also showed similar phenotypes. These results suggest that uterine Wnt5a-ROR signaling contributes to appropriate uterine LE organization and crypt formation for blastocyst attachment.

## Results

### **Wnt5a is expressed in a spatiotemporal manner in the periimplantation uterus**

To better understand the function of Wnt5a in early pregnancy, we assessed the spatiotemporal uterine *Wnt5a* expression. While *Wnt5a* expression was low to undetectable on days 1 and 2 of pregnancy, the expression was localized to the LE and subepithelial stroma on days 3 and 4, prior to and during the receptive phase. With implantation, the expression was highly localized in stromal cells at the M pole on day 5 (Figure 1A). On day 8, uterine *Wnt5a* expression was lower, indicating downregulation with pregnancy progression (Figure 1A). These results suggest that while *Wnt5a* expression on days 3 and 4 is associated with uterine receptivity, its localized expression at the M pole on day 5 shows a Wnt5a gradient along the M-AM axis.

### **Mice with uterine inactivation of *Wnt5a* show severely compromised fertility**

To explore uterine Wnt5a's role in pregnancy, we generated females with uterine inactivation of *Wnt5a* (*Wnt5a<sup>d/d</sup>*). *Wnt5a<sup>d/d</sup>* females and their floxed littermates (*Wnt5a<sup>f/f</sup>*) were mated with WT males. We found that >70% of plug-positive *Wnt5a<sup>d/d</sup>* females failed to produce any litters after multiple matings (Figure 1B). Moreover, those 33% of plug-positive *Wnt5a<sup>d/d</sup>* females that delivered pups had significantly small litter sizes (Figure 1B). Females with deletion of one allele of *Wnt5a* showed a modest reduction in litter size (*Wnt5a<sup>d/+</sup>*: 5.0 ± 1.4 pups/litter, n=4; *Wnt5a<sup>f/f</sup>*: 8.0 ± 0.6, n=6; mean ± SEM). This suggests that regulated Wnt5a levels are critical for pregnancy success. Wnt5a was efficiently deleted in *Wnt5a<sup>d/d</sup>* uteri, but not completely absent on days 4 or 5 of pregnancy (Figure 1C–D). The residual Wnt5a may explain retention of some fertility in *Wnt5a<sup>d/d</sup>* females.

### **Mice with uterine overexpression of *Wnt5a* show reduced fertility**

To further assess Wnt5a's role, we generated females with uterine overexpression of *Wnt5a* (*Wnt5a<sup>GOF</sup>*) (Figure S1A–E). *Wnt5a<sup>GOF</sup>* females mated with WT males had a significant reduction in litter size (Figure 1B). We confirmed upregulation of *Wnt5a* in *Wnt5a<sup>GOF</sup>* uteri on the mornings of days 4 and 5. In situ hybridization results showed ectopic *Wnt5a* expression in all major uterine compartments (Figure 1C), and Wnt5a protein levels were 2–3 fold higher in *Wnt5a<sup>GOF</sup>* uteri (Figure 1D). Interestingly, females with one allele of upregulated *Wnt5a* showed normal litter size (*Wnt5a<sup>GOF/+</sup>*: 7.0 ± 1.4 pups/litter; *Wnt5a<sup>f/f</sup>*: 8.3 ± 1.3; n=5, mean ± SEM).

Approaching blastocyst attachment, the receptive uterine LE shows a decrease in apicobasal cell polarity with the loss of surface microvilli to facilitate trophectoderm adhesion to the

apical surface of the LE (Daikoku et al., 2011; Schlafke and Enders, 1975). EM analysis showed persistent apicobasal polarity of the LE in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice with higher cell heights and retention of glycosylated microvilli on the apical surface (Figure 1E). These results suggest that aberrant uterine *Wnt5a* levels interfered with critical aspects of the attachment reaction.

### ***Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice show defective implantation**

To address the severe subfertility phenotypes in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females, we examined implantation status on day 5 of pregnancy. We found that *Wnt5a<sup>d/d</sup>* or *Wnt5a<sup>GOF</sup>* uteri either failed to show blue bands or had very weak bands compared to those in floxed littermates (Figure 1F), and unattached blastocysts were recovered from their uteri. In addition, implantation sites in these females were often irregularly spaced. (Figure 1F). These characteristics were reflected in cross-sections of day 5 IS with embryo implantation in the middle of the lumen (Figure S1F). We next asked whether decidualization and continued embryonic growth are affected under aberrant *Wnt5a* signaling.

### **Decidual development is aberrant in females with dysregulated uterine *Wnt5a* levels**

In mice, stromal cells surrounding the blastocyst undergo differentiation to form the primary decidual zone (PDZ) beginning on day 5 afternoon. PDZ is avascular with stromal cells assuming organized epithelioid characteristics with the acquisition of epithelial cell markers (Paria et al., 1999). The presumed functions of the PDZ are to support the early developing embryo and protect it from noxious substances from the maternal circulation by creating a selective barrier (Paria et al., 1999). By day 6, the PDZ is well developed and a secondary decidual zone (SDZ) begins to form at the periphery of the PDZ (Cha et al., 2013). Decidual growth peaks between days 7 and 8, after which the decidua progressively regresses to accommodate embryonic growth and placentation (Cha et al., 2012; Dey et al., 2004). We asked whether PDZ formation is disturbed in mice with aberrant *Wnt5a* expression. Histology of cross-sections of day 6 IS showed atypical implantation with embryos implanted close to the M pole rather than the AM pole (Figure 2A). Abnormal PDZ development was associated with atypical distribution of cyokeratin 8 (CK8), E-cadherin and ZO-1: ZO-1 distribution was reduced with retention of CK8/E-cadherin in the LE of *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* uteri, suggesting poor trophoblast penetration through the LE (Figure 2B). Notably, the PDZ was absent in some IS with the embryo entrapped within the intact epithelium (Figure 2B). These results suggest that disturbances in periimplantation *Wnt5a* signaling contribute to abnormal PDZ formation.

### **Implantation sites in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice show aberrant M-AM orientation**

To gain further insight in decidualization under dysregulated *Wnt5a* levels, we assessed decidual growth on day 8. We found abnormal embryo spacing and different sizes of IS in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females (Figure 2C), indicating asynchronous development. Furthermore, *Wnt5a<sup>d/d</sup>* IS assumed a more spherical morphology with reduced *Cox2* expression in contrast to the elliptical appearance in floxed littermates with appropriate M-AM axis formation and *Cox2* expression at the M pole (Figure 2D). Thus appropriate M-AM orientation in *Wnt5a<sup>d/d</sup>* IS was not realized. Ratios of the length along the M-AM axis

respective to the anterior-posterior (A-P) axis in *Wnt5a*<sup>d/d</sup> IS were ~25% smaller than *Wnt5a*<sup>f/f</sup> sites (Figure S2A). Similar phenotypes were noted in *Wnt5a*<sup>GOF</sup> females (Figure 2C, Figure S2A). Since *Wnt5a* levels are normally low from day 8 onward, we attribute these defects to adverse effects originating at the time of implantation. In several *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females, some decidual growth was noted on day 8 even with inappropriate M-AM orientation of IS (Figure 2C, Figure S2A). This observation resembles the results of previous studies that decidualization initiated from aberrant implantation can persist for a limited time (Song et al., 2002; Sun et al., 2012).

### ***Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females show increased resorption rates**

We speculated that early defects would lead to abnormalities at later stages. Indeed, increasing numbers of resorption sites on days 10 and 12 were noted in *Wnt5a*-dysregulated females (Figure 2E–G). Placental development was disrupted and embryos were substantially smaller or absent in many IS from day 10 onwards. On day 10, the trophoblast giant cell (TGC) layer, as marked by *Prl3d1* expression, normally lies between the maternal decidua basalis and developing placenta. However, clusters of *Prl3d1*-positive TGCs were present in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> IS in lieu of the embryo proper (Figure S2B). Implantation sites that contained placentae had smaller spongiotrophoblast compartments as indicated by reduced *Tpbpa* expression (Figure S2B). Many *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> IS underwent resorption by day 12, but those few IS that persisted showed abnormal placentation (Figure S2C). *Gcm1* positive syncytiotrophoblasts in the labyrinth failed to elongate, suggesting disrupted secondary branching morphogenesis of placental villi (Figure S2C). In summary, both *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> mice showed more or less similar phenotypes, suggesting that regulated *Wnt5a* levels are critical for normal implantation, embryo spacing, decidualization, placentation.

### **Expression of uterine receptivity marker genes are not altered under dysregulated *Wnt5a* levels**

To examine *Wnt5a*'s role in uterine receptivity, we examined the expression of estrogen- and progesterone (P<sub>4</sub>)-responsive markers of uterine receptivity in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> mice on day 4 (day of receptivity) (Lee et al., 2006; Lim et al., 1997; Matsumoto et al., 2002; Song et al., 2000; Stewart et al., 1992). Expression patterns of P<sub>4</sub>-responsive genes *Hoxa10* and *Ihh* in stromal and epithelial cells, respectively, and estrogen-responsive gene *Lif* in glandular epithelial cells in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> mice were similar to those in floxed uteri (Figure S3A & B). Expression of *Msx1*, another uterine receptivity marker (Daikoku et al., 2011), was also comparable in these females (Figure S3A & B; S3A–C). In addition, apicobasal polarity of LE as seen by immunofluorescence (IF) of αPKC and ZO1 was comparable (Figure S3D). Uterine receptivity is characterized by differentiation of LE with increased stromal cell proliferation. We found that the population of Ki67 positive cells in the LE and stroma of *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females were nearly comparable to that of floxed littermates on day 4 (Figure S3E). These results suggest that defective attachment and implantation seen in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> mice are not due to defects in uterine receptivity on day 4 morning. *Wnt11*, another non-canonical ligand and modestly expressed in the LE, was not altered in *Wnt5a*<sup>d/d</sup> or *Wnt5a*<sup>GOF</sup> day 4 uteri (Figure S3F). Expression of *Wnt7a*, reported to be critical for uterine gland formation (Dunlap et al., 2011), was also

similar in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice with no apparent defects in gland formation (Figure S3F). These results suggest that uterine *Wnt11* and *Wnt7a* expression do not respond to fluctuating uterine *Wnt5a* levels and may not intersect with uterine *Wnt5a* signaling.

### ***Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females display aberrant crypt formation and *Hbegf* expression prior to blastocyst attachment**

In mice, blastocysts normally implant within specialized crypts at the AM pole (Daikoku et al., 2011; Kirby, 1971). Our observation that blastocysts implanted closer to the M-pole in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice on days 5 and 6 suggested that blastocyst attachment failed to occur towards the AM pole (Figure 2A & Figure S1F). In support of this, we performed histological analysis of longitudinal uterine sections on day 4 evening prior to blastocyst attachment with appropriate orientation of the M-AM axis confirmed by the positioning of glands at the AM pole. We found that regularly-spaced villi-like epithelial projections extended from the primary uterine lumen into the stroma towards the AM pole on day 4 evening in floxed females (Figure 3A). In contrast, these projections were randomly present in both M and AM directions in *Wnt5a<sup>d/d</sup>* females with presence of glands even at the M pole. Furthermore, crypt formation was frequently absent in these females with blastocysts situated near the M pole (Figure 3A). *Wnt5a<sup>GOF</sup>* mice also showed defects in crypt formation, embryo placement, including blastocysts situated within the primary lumen rather than in a crypt or two embryos closely apposed within one site (Figure 3A). Intriguingly, directional villi-like projections towards the AM pole in floxed mice were not seen on day 3 but became evident on day 4, coincident with the initiation of a *Wnt5a* expression gradient along the M-AM axis in longitudinal sections (Figure 3B & Figure S4A). The loss of the *Wnt5a* gradient is consistent with our observations of aberrant embryo spacing and crowding on day 5 under dysregulated uterine *Wnt5a* levels (Figure 1F).

The altered LE architecture in these mice led us to hypothesize that on-time embryo-uterine interactions were disturbed. *Hbegf* is first expressed in the LE around the blastocyst several hours before the attachment reaction and is the first known molecular link mediating blastocyst-uterine interactions in a juxtacrine and paracrine fashion (Das et al., 1994). We found that *Hbegf* expression was undetectable in the LE around the misplaced blastocyst in *Wnt5a<sup>d/d</sup>* or *Wnt5a<sup>GOF</sup>* uteri on day 4 evening (Figure 3C), suggesting that the bidirectional communication was not realized under aberrant *Wnt5a* signaling.

### ***Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice have normal ovarian function**

Since *Pgr<sup>Cre</sup>* is also expressed in ovary, we examined whether infertility/subfertility in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females is due to ovarian dysfunction. *Wnt5a* is mainly expressed in corpora lutea. Its expression was slightly lower in *Wnt5a<sup>d/d</sup>* ovaries but higher in *Wnt5a<sup>GOF</sup>* ovaries (Figure S4B), although qPCR results showed no significant changes in transcript levels in these ovaries (Figure S4C). More importantly, we assessed ovarian function with respect to ovulation, fertilization, preimplantation embryo development, and hormone production. The number of IS plus recovered blastocysts in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females on day 5 were comparable to the number of blue bands in floxed littermates (Figure S4D), providing evidence that these processes were comparable among these three genotypes. Serum levels of estradiol-17 $\beta$  ( $E_2$ ) and  $P_4$  on days 4, 8 and 12 of

pregnancy in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females were comparable to their floxed counterparts (Figure S4E & F). These results suggest normal ovarian function and hormonal milieu in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females.

### Uterine *Wnt5a* promotes ROR2 phosphorylation

*Wnt5a* can bind to ROR receptors to mediate non-canonical Wnt signaling (Green et al., 2008; Grumolato et al., 2010; Minami et al., 2010; Niehrs, 2012). ROR2 levels were upregulated in *Wnt5a<sup>d/d</sup>* uteri but downregulated in *Wnt5a<sup>GOF</sup>* uteri (Figure 4A); ROR1 levels did not show a similar pattern (Figure 4A). This observation suggests that *Wnt5a* and ROR2 interact. ROR2 was primarily localized in epithelia of floxed uteri with seemingly higher intensity in *Wnt5a<sup>d/d</sup>* mice on day 4, although some signals were noted in the underlying stroma (Figure S5A). This result corroborates enriched ROR2 levels in isolated primary uterine epithelial cells compared to stromal cells (Figure S5B). ROR1 was also enriched in epithelial cells rather than stromal cells (Figure S5B).

These results led us to explore if ROR2 responds to *Wnt5a* stimulation. *Wnt5a* can modulate ROR2 receptor function via tyrosine kinase activity and ROR2 autophosphorylation (Mikels et al., 2009). IP-kinase assay in a uterine epithelial cell line expressing ROR2 showed that ROR2 undergoes tyrosine phosphorylation after *Wnt5a* stimulation (Figure 4B). IP assays also showed increased tyrosine phosphorylation of ROR2 in *Wnt5a<sup>GOF</sup>* uterine extracts (Figure 4C). Furthermore, gel-shift assays for ROR1 and ROR2 phosphorylation showed downward and upward shifts in day 5 uterine extracts of *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females, respectively (Figure 4D). These results intrigued us to determine the significance of ROR1 and ROR2 in female fertility.

### Mice with uterine deletion of both *Ror1* and *Ror2* show severe subfertility

We first examined if mice with uterine inactivation of ROR2 (*Ror2<sup>d/d</sup>*) have adverse effects on fertility. Despite efficient depletion of ROR2 (Figure 4E), *Ror2<sup>d/d</sup>* females had normal fertility (Figure 5A). We speculated that ROR1 compensated for ROR2 deficiency because of their redundant functions (Ho et al., 2012; Lyashenko et al., 2010; Nomi et al., 2001). Indeed *Ror2<sup>d/d</sup>* uteri had increased levels of ROR1 (Figure 4E). Females with uterine deletion of *Ror1* showed normal fertility (*Ror1<sup>d/d</sup>*: 8.50 ± 0.56 pups/litter, n=6; *Ror1<sup>ff/ff</sup>*: 9.0 ± 0.81, n=6). Mice with uterine deletion of both alleles of *Ror1* and one allele of *Ror2* also had fertility comparable to floxed littermates (*Ror1<sup>d/d</sup>Ror2<sup>d/+</sup>*: 7.0 ± 1.0 pups/litter, n=3; *Ror1<sup>ff/ff</sup>Ror2<sup>ff/ff</sup>*: 8.17 ± 0.48, n=6; mean ± SEM), suggesting that one allele of uterine *Ror2* is sufficient for normal fertility. We then examined fertility in mice with uterine deletion of both alleles of *Ror1* and *Ror2* (*Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>*). *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females showed severe subfertility with only 29% of plug-positive females producing a small number of pups (Figure 5B) and high resorption rates on day 12 (Figure 5C & D).

On day 5, defective implantation was marked by very weak or absence of blue bands in *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* uteri with efficient deletion of both genes; unattached blastocysts were recovered from these mice (Figure 5E & F). Again, we found defective crypt formation with embryos entrapped within the highly branched LE on day 4 evening prior to blastocyst attachment (Figure 5G). These findings suggest that blastocyst attachment was defective

leading to late stage defects. Defective implantation seemingly did not result from aberrant expression of uterine receptivity genes *Lif* or *Ihh*; their expression was normal (Figure S5C–D). Ovarian levels of ROR1 or ROR2 and serum P<sub>4</sub> and E<sub>2</sub> levels were also comparable between *Ror1<sup>ff</sup>Ror2<sup>ff</sup>* and *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females (Figure S5E & F), suggesting normal ovarian function and responsiveness.

On day 6, the formation of the PDZ as determined by histology and localization of ZO-1/CK8 showed embryos entrapped within the intact LE (Figure S5G & H). These changes are strikingly similar to those observed in females with dysregulated Wnt5a signaling, thus reinforcing the critical role of Wnt5a-ROR signaling in blastocyst attachment, implantation and decidualization.

### Wnt5a-ROR signaling influences *Bmp2* signaling during implantation

Normally, *Bmp2* is expressed in the stroma surrounding the blastocyst at the initiation of implantation and markedly increases in decidualization. Since *Bmp2* is critical for these events (Lee et al., 2007; Paria et al., 2001), we assessed uterine *Bmp2* expression in *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* females. We found reduced *Bmp2* expression at the IS of these mice on day 5 with attenuated nuclear localization of pSmad1/5/8 (Figure 5H & I). *Bmp2* signaling is mediated by phosphorylation of Smad proteins, and Wnt5a was shown to intersect with Smad signaling in other systems (Miyoshi et al., 2012). These results suggest that *Bmp2* signaling is compromised with altered Wnt5a levels. In contrast, *Ptgs2* (encoding Cox2) expression which is normally expressed both in the LE and subepithelial stroma at the IS and critical for implantation (Lim et al., 1997), was comparable among *Wnt5a<sup>ff</sup>*, *Wnt5a<sup>d/d</sup>*, *Wnt5a<sup>GOF</sup>* and *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females (Figure 5H & I). These results suggest that specific signaling pathways are affected under altered Wnt5a-ROR signaling.

## Discussion

This study provides evidence for the requirement of Wnt5a-ROR signaling in promoting villi-like epithelial projections towards the AM pole for crypt formation and implantation. Although the presence of crypts were noted in uterine cross-sections in mice (Daikoku et al., 2011; Kirby, 1971), the underlying mechanism regulating the formation of regularly spaced villi-like epithelial projections from the main uterine lumen to form crypts at the AM pole remained unknown. Wnt5a-ROR signaling has been reported to influence tissue patterning, cell polarity and directional cell movement during organogenesis (Angers and Moon, 2009; Bayly and Axelrod, 2011). Our findings that Wnt5a-ROR signaling is active in the adult uterus to confer appropriate LE organization for implantation and that dysregulation of this pathway leads to aberrant crypt formation and subfertility reveals an unrecognized critical event in implantation.

Defective implantation and severe subfertility in both *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* mice suggest that regulated Wnt5a levels are critical for early pregnancy events. Localized *Wnt5a* expression at the M pole prior to and following implantation suggests the presence of a Wnt5a gradient that directs uterine decidual growth patterning along the M-AM axis, conferring an elliptical shape and proper embryonic growth. Appropriate directional decidual



formation is essential for normal embryonic growth and placentation. Our results suggest that aberrant Wnt5a-ROR signaling leads to abnormal decidual growth.

One major function of the decidua is to regulate trophoblast invasion into the stroma to direct placentation at the M pole. Our finding of defective implantation and decidual growth restriction under aberrant uterine Wnt5a levels suggests that some decidualization could be initiated, but not sustained to support placentation and term pregnancy. This explains the abnormal placentation seen on day 10 onwards in females with dysregulated Wnt5a-ROR signaling. Notably, *Wnt5a* expression is appreciably lower during decidualization in floxed females, suggesting that this defect originated from aberrations in the implantation process. This resembles previous work by us and others (Song et al., 2002; Sun et al., 2012; Ye et al., 2005).

The epithelial-mesenchymal interaction during implantation constitutes a reciprocal signaling circuitry between the LE and stroma. Our study suggests that regulated Wnt5a-ROR2 signaling participates in this transition. Reduced *Bmp2* and pSMAD1/5/8 expression with disrupted PDZ formation under aberrant Wnt5a-ROR signaling suggests that this pathway influences *Bmp2* signaling critical for decidualization (Lee et al., 2007).

Alternatively, the defective LE architecture, conferring poor blastocyst attachment under abnormal-Wnt5a-ROR signaling perhaps resulted in defective decidualization and *Bmp2* signaling. This is a possibility since LE architecture had already become aberrant in day 4 evening preceding *Bmp2* expression. These results indicate a relationship between Wnt5a-ROR signaling, appropriate LE architecture, and *Bmp2* signaling for implantation and decidualization. In this context, Wnt5a was shown to aid in intestinal stem cell niche regeneration after injury via BMP-Smad signaling (Miyoshi et al., 2012).

Wnt5a-ROR signaling is active in embryogenesis (Ho et al., 2012; Yamaguchi et al., 1999). However, subfertility phenotypes in *Wnt5a<sup>d/d</sup>* or *Wnt5a<sup>GOF</sup>* females in our studies did not stem from embryogenesis defects, since females were mated with WT males. Furthermore, uterine deletion of *Ror1/Ror2* resulted in similar fertility defects seen in mice with dysregulated uterine Wnt5a levels, suggesting that the defects primarily originated from uterine aberrations. This is further reinforced by similar changes in LE morphology in mice with dysregulated Wnt5a or *Ror1/Ror2* levels. Since ovulation, fertilization, and preimplantation embryo development were not disturbed with comparable levels of E<sub>2</sub> and P<sub>4</sub> under these experimental conditions, we believe that ovarian dysfunction did not contribute to fertility defects in these mice.

In WT females, invariable orientation of villous-like epithelial projections from the primary lumen towards the AM pole with regular spacing is reminiscent of the patterning effect of PCP signaling in organogenesis. Whether Wnt5a-ROR signaling involves PCP signaling would require functional studies using mice with conditional uterine deletion of PCP components.

Although certain aspects of implantation are similar in mice and humans, there are species-specific differences. Nonetheless, placentation in both mice and humans is hemochorial. Regarding the relevance of embryo spacing, abnormal implantation can occur close to or on

the cervix in placenta previa which causes extensive bleeding with increased mortality and/or morbidity of the mother and fetus. Aberrant embryo spacing can also contribute to complications in multiple gestation pregnancy. Since Wnt signaling is highly conserved across species, it would be interesting to know whether Wnt5a-ROR signaling is involved in human implantation.

Collectively, our studies show that altered Wnt5a-ROR signaling leads to defects prior to and during blastocyst attachment, poor decidual growth, placentation and pregnancy outcome. This study has revealed a critical role for Wnt5a-ROR signaling in LE remodeling for blastocyst attachment and implantation and has expanded our understanding of implantation biology.

## Experimental Procedures

For detailed methods, see Supplemental Experimental Procedures.

### Mice

*Pgr<sup>Cre/+</sup>*, *Wnt5a<sup>loxP/loxP</sup>* and/or *Ror1<sup>loxP/loxP</sup>/Ror2<sup>loxP/loxP</sup>* floxed lines were generated as described (Ho et al., 2012; Miyoshi et al., 2012; Soyal et al., 2005). Details of *Wnt5a* conditionally overexpressing mouse line generation are described in Supplemental Experimental Procedures. *Wnt5a<sup>d/d</sup>*, *Wnt5a<sup>GOF</sup>*, and *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* mice were generated by mating floxed females with *Pgr<sup>Cre/+</sup>* males, as described (Daikoku et al., 2011). All mice used in this study were housed in the Cincinnati Children's Animal Care Facility according to NIH and institutional guidelines for the use of laboratory animals. All protocols were approved by the Institutional Animal Care and Use Committee.

### Analysis of Pregnancy Events

Ovulation, fertilization, preimplantation embryo development, and implantation were assessed as described (Daikoku et al., 2011).

### Immunostaining and histology

Tissue sections from control and experimental groups were processed onto the same slide for immunohistochemistry and IF (Daikoku et al., 2011).

### RNA isolation and qRT-PCR

qRT-PCR was performed as described (Daikoku et al., 2011).

### In Situ Hybridization

Paraformaldehyde-fixed frozen sections from control and experimental groups were processed onto the same slide and hybridized with <sup>35</sup>S-labeled or DIG-labeled cRNA probes (Daikoku et al., 2011; Simmons et al., 2008).

### Primary uterine cell culture

Primary epithelial and stromal cells were isolated and cultured with 95% purity as described (Daikoku et al., 2011).

## Immunoblotting

Western blotting was performed as described (Daikoku et al., 2011). The same blots were used for quantitative analysis of each protein. Bands were visualized using an ECL kit (GE Healthcare). Actin served as a loading control.

## Immunoprecipitation

IP was performed as described (Daikoku et al., 2011).

## IP-Kinase assay

Ishikawa cells (ATCC) expressing endogenous ROR2 were lysed and subjected to IP-kinase assay as previously described with modification (Paria et al., 1993).

**Electron microscopy**—Sections (80 nm) of *Wnt5a<sup>f/f</sup>* and *Wnt5a<sup>GOF</sup>* uteri on day 5 of pseudopregnancy were cut with a Leica UC-7 ultramicrotome, stained with 3% uranyl acetate and Sato's Lead, and examined under a JEOL JEM-1400 Transmission Electron Microscope at 80KV.

**Assay of serum E<sub>2</sub> and P<sub>4</sub> Levels**—Sera were collected on days 4, 8, and 12 of pregnancy, and E<sub>2</sub> and P<sub>4</sub> levels were measured by EIA kits (Cayman) (Daikoku et al., 2011).

## Statistics

Statistical analyses were performed, using two-tailed Student's t-test. A value of  $p < 0.05$  was considered statistically significant.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank Michael Greenberg (Harvard) for generously providing the *Ror1/Ror2* floxed mice, and Francesco DeMayo and John B. Lydon (Baylor) for *Pgr-Cre* mice. The ROR2 antibody was kindly provided by Yasuhiro Minami (Kyoto), and the phospho-Smad1/5/8 antibody by Dan Vasilias, Susan Morton, Tom Jessell and Ed Laufer (Columbia). We are thankful to Barbara Fegley and the Electron Microscopy Research Lab (University of Kansas Medical Center) for assistance with electron microscopy experiments (9P20GM104936). This work was supported in part by grants from the NIH (HD068524 and DA06668) and March of Dimes (21-FY12-127 and 22-FY13-543) (to S.K.D.) and by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research (T.P.Y.). J.C. is a National Research Service Award fellow (NIA/NIH F30AG040858) of the University of Cincinnati MSTP, and X.S. was supported by a Lalor Foundation Fellowship.

## References


- Angers S, Moon RT. Proximal events in Wnt signal transduction. *Nat Rev Mol Cell Biol.* 2009; 10:468–477. [PubMed: 19536106]
- Bayly R, Axelrod JD. Pointing in the right direction: new developments in the field of planar cell polarity. *Nat Rev Genet.* 2011; 12:385–391. [PubMed: 21502960]
- Cha J, Sun X, Dey SK. Mechanisms of implantation: strategies for successful pregnancy. *Nat Med.* 2012; 18:1754–1767. [PubMed: 23223073]

- Daikoku T, Cha J, Sun X, Tranguch S, Xie H, Fujita T, Hirota Y, Lydon J, DeMayo F, Maxson R, et al. Conditional deletion of *Msx* homeobox genes in the uterus inhibits blastocyst implantation by altering uterine receptivity. *Dev Cell*. 2011; 21:1014–1025. [PubMed: 22100262]
- Das SK, Wang XN, Paria BC, Damm D, Abraham JA, Klagsbrun M, Andrews GK, Dey SK. Heparin-binding EGF-like growth factor gene is induced in the mouse uterus temporally by the blastocyst solely at the site of its apposition: a possible ligand for interaction with blastocyst EGF-receptor in implantation. *Development*. 1994; 120:1071–1083. [PubMed: 8026321]
- Dey SK, Lim H, Das SK, Reese J, Paria BC, Daikoku T, Wang H. Molecular cues to implantation. *Endocr Rev*. 2004; 25:341–373. [PubMed: 15180948]
- Dunlap KA, Filant J, Hayashi K, Rucker EB 3rd, Song G, Deng JM, Behringer RR, DeMayo FJ, Lydon J, Jeong JW, et al. Postnatal deletion of *Wnt7a* inhibits uterine gland morphogenesis and compromises adult fertility in mice. *Biol Reprod*. 2011; 85:386–396. [PubMed: 21508348]
- Gao B, Song H, Bishop K, Elliot G, Garrett L, English MA, Andre P, Robinson J, Sood R, Minami Y, et al. Wnt signaling gradients establish planar cell polarity by inducing *Vangl2* phosphorylation through *Ror2*. *Dev Cell*. 2011; 20:163–176. [PubMed: 21316585]
- Green JL, Kuntz SG, Sternberg PW. Ror receptor tyrosine kinases: orphans no more. *Trends Cell Biol*. 2008; 18:536–544. [PubMed: 18848778]
- Grumolato L, Liu G, Mong P, Mudbhary R, Biswas R, Arroyave R, Vijayakumar S, Economides AN, Aaronson SA. Canonical and noncanonical Wnts use a common mechanism to activate completely unrelated coreceptors. *Genes Dev*. 2010; 24:2517–2530. [PubMed: 21078818]
- Ho HY, Susman MW, Bikoff JB, Ryu YK, Jonas AM, Hu L, Kuruvilla R, Greenberg ME. Wnt5a-Ror-Dishevelled signaling constitutes a core developmental pathway that controls tissue morphogenesis. *Proc Natl Acad Sci U S A*. 2012; 109:4044–4051. [PubMed: 22343533]
- Kikuchi A, Yamamoto H, Sato A, Matsumoto S. New insights into the mechanism of Wnt signaling pathway activation. *Int Rev Cell Mol Biol*. 2011; 291:21–71. [PubMed: 22017973]
- Kirby, DRS. Blastocyst-Uterine Relationship before and during Implantation. In: Blandau, RJ., editor. *The Biology of the Blastocyst*. Chicago, IL: The University of Chicago Press; 1971. p. 393-412.
- Lee K, Jeong J, Kwak I, Yu CT, Lanske B, Soegiarto DW, Toftgard R, Tsai MJ, Tsai S, Lydon JP, et al. Indian hedgehog is a major mediator of progesterone signaling in the mouse uterus. *Nat Genet*. 2006; 38:1204–1209. [PubMed: 16951680]
- Lee KY, Jeong JW, Wang J, Ma L, Martin JF, Tsai SY, Lydon JP, DeMayo FJ. *Bmp2* is critical for the murine uterine decidual response. *Mol Cell Biol*. 2007; 27:5468–5478. [PubMed: 17515606]
- Lim H, Paria BC, Das SK, Dinchuk JE, Langenbach R, Trzaskos JM, Dey SK. Multiple female reproductive failures in cyclooxygenase 2-deficient mice. *Cell*. 1997; 91:197–208. [PubMed: 9346237]
- Lyashenko N, Weissenbock M, Sharir A, Erben RG, Minami Y, Hartmann C. Mice lacking the orphan receptor *ror1* have distinct skeletal abnormalities and are growth retarded. *Dev Dyn*. 2010; 239:2266–2277. [PubMed: 20593419]
- MacDonald BT, Tamai K, He X. Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Dev Cell*. 2009; 17:9–26. [PubMed: 19619488]
- Macheda ML, Sun WW, Kugathasan K, Hogan BM, Bower NI, Halford MM, Zhang YF, Jacques BE, Lieschke GJ, Dabdoub A, et al. The Wnt receptor *Ryk* plays a role in mammalian planar cell polarity signaling. *J Biol Chem*. 2012; 287:29312–29323. [PubMed: 22773843]
- Matsumoto H, Zhao X, Das SK, Hogan BL, Dey SK. Indian hedgehog as a progesterone-responsive factor mediating epithelial-mesenchymal interactions in the mouse uterus. *Dev Biol*. 2002; 245:280–290. [PubMed: 11977981]
- Mikels A, Minami Y, Nusse R. *Ror2* receptor requires tyrosine kinase activity to mediate *Wnt5A* signaling. *J Biol Chem*. 2009; 284:30167–30176. [PubMed: 19720827]
- Minami Y, Oishi I, Endo M, Nishita M. Ror-family receptor tyrosine kinases in noncanonical Wnt signaling: their implications in developmental morphogenesis and human diseases. *Dev Dyn*. 2010; 239:1–15. [PubMed: 19530173]
- Miyoshi H, Ajima R, Luo CT, Yamaguchi TP, Stappenbeck TS. *Wnt5a* potentiates TGF-beta signaling to promote colonic crypt regeneration after tissue injury. *Science*. 2012; 338:108–113. [PubMed: 22956684]

- Niehrs C. The complex world of WNT receptor signalling. *Nat Rev Mol Cell Biol.* 2012; 13:767–779. [PubMed: 23151663]
- Nomi M, Oishi I, Kani S, Suzuki H, Matsuda T, Yoda A, Kitamura M, Itoh K, Takeuchi S, Takeda K, et al. Loss of mRor1 enhances the heart and skeletal abnormalities in mRor2-deficient mice: redundant and pleiotropic functions of mRor1 and mRor2 receptor tyrosine kinases. *Mol Cell Biol.* 2001; 21:8329–8335. [PubMed: 11713269]
- Paria BC, Das SK, Andrews GK, Dey SK. Expression of the epidermal growth factor receptor gene is regulated in mouse blastocysts during delayed implantation. *Proc Natl Acad Sci U S A.* 1993; 90:55–59. [PubMed: 7678348]
- Paria BC, Ma W, Tan J, Raja S, Das SK, Dey SK, Hogan BL. Cellular and molecular responses of the uterus to embryo implantation can be elicited by locally applied growth factors. *Proc Natl Acad Sci U S A.* 2001; 98:1047–1052. [PubMed: 11158592]
- Paria BC, Zhao X, Das SK, Dey SK, Yoshinaga K. Zonula occludens-1 and E-cadherin are coordinately expressed in the mouse uterus with the initiation of implantation and decidualization. *Dev Biol.* 1999; 208:488–501. [PubMed: 10191061]
- Psychoyos, A. Endocrine control of egg implantation. In: Greep, EGARO.; Geiger, SR., editors. *Handbook of Physiology.* Washington, D.C: American Physiology Society; 1973. p. 187-215.
- Schambony A, Wedlich D. Wnt-5A/Ror2 regulate expression of XPAPC through an alternative noncanonical signaling pathway. *Dev Cell.* 2007; 12:779–792. [PubMed: 17488628]
- Schlafke S, Enders AC. Cellular basis of interaction between trophoblast and uterus at implantation. *Biol Reprod.* 1975; 12:41–65. [PubMed: 1095088]
- Simmons DG, Rawn S, Davies A, Hughes M, Cross JC. Spatial and temporal expression of the 23 murine Prolactin/Placental Lactogen-related genes is not associated with their position in the locus. *BMC Genomics.* 2008; 9:352. [PubMed: 18662396]
- Song H, Lim H, Das SK, Paria BC, Dey SK. Dysregulation of EGF family of growth factors and COX-2 in the uterus during the preattachment and attachment reactions of the blastocyst with the luminal epithelium correlates with implantation failure in LIF-deficient mice. *Mol Endocrinol.* 2000; 14:1147–1161. [PubMed: 10935540]
- Song H, Lim H, Paria BC, Matsumoto H, Swift LL, Morrow J, Bonventre JV, Dey SK. Cytosolic phospholipase A2alpha is crucial [correction of A2alpha deficiency is crucial] for 'on-time' embryo implantation that directs subsequent development. *Development.* 2002; 129:2879–2889. [PubMed: 12050136]
- Soyal SM, Mukherjee A, Lee KY, Li J, Li H, DeMayo FJ, Lydon JP. Cre-mediated recombination in cell lineages that express the progesterone receptor. *Genesis.* 2005; 41:58–66. [PubMed: 15682389]
- Stewart CL, Kaspar P, Brunet LJ, Bhatt H, Gadi I, Kontgen F, Abbondanzo SJ. Blastocyst implantation depends on maternal expression of leukaemia inhibitory factor. *Nature.* 1992; 359:76–79. [PubMed: 1522892]
- Sun X, Zhang L, Xie H, Wan H, Magella B, Whitsett JA, Dey SK. Kruppel-like factor 5 (KLF5) is critical for conferring uterine receptivity to implantation. *Proc Natl Acad Sci U S A.* 2012; 109:1145–1150. [PubMed: 22233806]
- van Amerongen R, Nusse R. Towards an integrated view of Wnt signaling in development. *Development.* 2009; 136:3205–3214. [PubMed: 19736321]
- Veeman MT, Axelrod JD, Moon RT. A second canon. Functions and mechanisms of beta-catenin-independent Wnt signaling. *Dev Cell.* 2003; 5:367–377. [PubMed: 12967557]
- Yamaguchi TP, Bradley A, McMahon AP, Jones S. A Wnt5a pathway underlies outgrowth of multiple structures in the vertebrate embryo. *Development.* 1999; 126:1211–1223. [PubMed: 10021340]
- Ye X, Hama K, Contos JJ, Anliker B, Inoue A, Skinner MK, Suzuki H, Amano T, Kennedy G, Arai H, et al. LPA3-mediated lysophosphatidic acid signalling in embryo implantation and spacing. *Nature.* 2005; 435:104–108. [PubMed: 15875025]

### Highlights

- Normal embryo implantation requires regulated uterine Wnt5a-ROR signaling
- Wnt5a-ROR signaling aids in orienting villi-like epithelial projections
- Aberrant Wnt5a-ROR signaling is reflected in altered Bmp2-Smad signaling
- Defective implantation compromises decidualization, placentation and fertility




**Figure 1. *Wnt5a* is expressed in the periimplantation uterus and its dysregulation results in defective implantation**

(A) In situ hybridization of *Wnt5a* expression patterns on days 1–5 and 8 of pregnancy. Arrowheads, embryo location. Scale bar: days 1–5, 500  $\mu$ m; scale bar: day 8, 1 mm. M, mesometrial pole; AM, anti-mesometrial pole. (B) Litter sizes in pregnant *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females and their *Wnt5a*<sup>fl/fl</sup> littermates (mean  $\pm$  SEM). Percentages and numbers within parentheses indicate number of litters born compared to total number of plug-positive females in each genotype (n = number of mice). (C) In situ hybridization of uterine *Wnt5a* expression in *Wnt5a*<sup>d/d</sup>, *Wnt5a*<sup>GOF</sup> and floxed littermates on days 4 and 5. Arrowheads, embryo location. Scale bars, 250  $\mu$ m. (D) Western blotting of uterine *Wnt5a* levels of *Wnt5a*<sup>d/d</sup>, *Wnt5a*<sup>GOF</sup> and IS of floxed littermates on day 5. Bar diagram, quantification of *Wnt5a* protein levels relative to actin (mean  $\pm$  SEM). (E) Representative TEM images of the apical surface of uterine LE of *Wnt5a*<sup>d/d</sup>, *Wnt5a*<sup>GOF</sup> and floxed littermates on day 5 morning. mv, microvilli. Scale bar, 500 nm. (F) Day 5 IS (blue bands) in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females. Unfilled arrows indicate differential blue band intensities in adjacent IS. Brackets indicate clustering of IS. Unattached blastocysts retrieved from uterine luminal


flushings from representative mice are shown. le, luminal epithelium; s, stroma; myo, myometrium.






**Figure 2. Defective implantation under dysregulated *Wnt5a* levels leads to compromised decidualization and resorption**  
**(A)** Histology of day 6 IS in *Wnt5a<sup>fl/fl</sup>*, *Wnt5a<sup>d/d</sup>*, and *Wnt5a<sup>GOF</sup>* females. Brackets indicate the distance from the implanting embryo to the main LE at the M pole; asterisks, aberrantly short distance. Scale bar, 500  $\mu$ m. **(B)** IF of ZO-1 shows epithelioid cells in the primary decidual zone (PDZ, bracket) on day 6. E-cadherin IF marks the embryo and LE. Scale bar, 200  $\mu$ m. **(C)** Day 8 *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* IS after intravenous blue dye injection. Note differential sizes (arrowheads) and crowding (brackets, asterisks) of IS. **(D)** Immunohistochemistry shows reduced Cox2 expression at the M pole of day 8 *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* IS. Bar, 200  $\mu$ m. **(E)** Histology of day 10 *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* IS show resorptions. Scale bar, 1mm. **(F)** Day 12 *Wnt5a<sup>fl/fl</sup>*, *Wnt5a<sup>d/d</sup>* and *Wnt5a<sup>GOF</sup>* uterine horns.


(G) Resorption rate in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females on day 12. Percentages and numbers within parentheses indicate number of resorptions over total number of IS. n, numbers of mice. M, mesometrial pole; AM, anti-mesometrial pole. Arrowheads, location of embryos or resorption sites.



**Figure 3.** *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females show poor crypt formation with undetectable *Hbegf* expression  
 (A) Histology of longitudinal sections of *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> uteri approaching attachment reaction (day 4, 1900h) show aberrant luminal architecture and villi-like epithelial projections in both M and AM directions compared to regularly spaced projections towards the AM pole in floxed littermates (asterisks). Note normal placement of blastocyst within a crypt (nidus) at the AM pole in floxed mice (top, first panel). In *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females, blastocysts were often situated at the M pole (second and third panels), within the primary lumen (fourth panel), and/or adjoined together within one site (bottom, fifth panel). Left panels, scale bar, 500  $\mu$ m; right panels, scale bar, 250  $\mu$ m. (B) In situ hybridization of *Wnt5a* expression in WT females on day 3 (1000h) and day 4 (1600h and 1900h). Asterisks, locations of epithelial projections towards the AM pole. Scale bar, 500  $\mu$ m. (C) In situ hybridization of *Hbegf* expression in *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> females. Arrowheads, locations of blastocysts. le, luminal epithelium; ge, glandular epithelium; s, stroma; myo, myometrium. Left panels, scale bar, 500  $\mu$ m; right panels, scale bar, 250  $\mu$ m.



**Figure 4. Uterine Wnt5a levels influence ROR levels, and uterine inactivation of *Ror1* and *Ror2* (*Ror1*<sup>d/d</sup>*Ror2*<sup>d/d</sup>) show fertility defects**  
 (A) Western blotting results of ROR1 and ROR2 in day 5 *Wnt5a*<sup>d/d</sup> and *Wnt5a*<sup>GOF</sup> IS. Bar diagram, quantification of ROR levels (\*p<0.05, mean ± SEM). (B) IP-kinase assay for ROR2 tyrosine phosphorylation in a uterine epithelial cell line (Ishikawa) exposed to Wnt5a. (C) IP results show increased tyrosine phosphorylation of ROR2 in day 5 IS. (D) Gel-shift immunoblotting shows upward shift of ROR1 and ROR2 in *Wnt5a*<sup>GOF</sup> sites and downward shift in *Wnt5a*<sup>d/d</sup> IS on day 5. (E) Western blotting results from *Ror2*<sup>d/d</sup> day 5 IS show compensation by ROR1 for ROR2 deficiency.



**Figure 5. Implantation phenotype in *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females resembles that of *Wnt5a<sup>d/d</sup>* females (A–B) Fertility of mice with uterine inactivation of *Ror2* (*Ror2<sup>d/d</sup>*) (A) or both *Ror1* and *Ror2* (*Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>*) (B) and their floxed littermates (litter size, mean ± SEM). Percentages and numbers within parentheses indicate the number of litters over total number of plug-positive females within each genotype. (C) Day 12 *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* uteri. Arrowheads, resorption sites. Bracket, large area of resorption. (D) Resorption rate on day 12 in *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females. Percentages and numbers within parentheses indicate number of resorptions over total number of IS. n, numbers of females. (E) *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* IS on day 5 after blue dye injection. Arrowheads, locations of very weak blue bands in *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females. (F) Western blotting results of day 5 *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* IS show efficient deletion of**

both *Ror1* and *Ror2*. **(G)** Histology of the irregular LE projections of *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* females on day 4 (1900h). Note the location of a blastocyst within a highly branched lumen. Scale bars: Left panel, 500  $\mu\text{m}$ ; right panel, 200  $\mu\text{m}$ . **(H)** In situ hybridization of *Ptgs2* and *Bmp2* in *Wnt5a<sup>d/d</sup>*, *Wnt5a<sup>GOF</sup>*, and *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* IS on day 5. Scale bars, 500  $\mu\text{m}$ . **(I)** IF results for phospho-Smad1/5/8 (pSmad1/5/8) in *Wnt5a<sup>d/d</sup>*, *Wnt5a<sup>GOF</sup>*, and *Ror1<sup>d/d</sup>Ror2<sup>d/d</sup>* day 5 IS. Bars, 100  $\mu\text{m}$ . Arrowheads, locations of blastocysts.