

## **AUTHOR CORRECTION**

## Correction for Anand et al., TprC/D (Tp0117/131), a Trimeric, Pore-Forming Rare Outer Membrane Protein of *Treponema pallidum*, Has a Bipartite Domain Structure

Arvind Anand,<sup>a</sup> Amit Luthra,<sup>a</sup> Star Dunham-Ems,<sup>a</sup> Melissa J. Caimano,<sup>a,e</sup> Carson Karanian,<sup>e</sup> Morgan LeDoyt,<sup>a</sup> Adriana R. Cruz,<sup>d</sup> Juan C. Salazar,<sup>c,e</sup> and Justin D. Radolf<sup>a,b,c,e</sup>

Departments of Medicine, a Genetics and Developmental Biology, and Immunology, Cuniversity of Connecticut Health Center, Farmington, Connecticut, USA; Centro Internacional de Entrenamiento e Investigaciones Médicas (CIDEIM), Cali, Colombiad; and Department of Pediatrics, Connecticut Children's Medical Center, Division of Pediatric Infectious Diseases, Hartford, Connecticut, USAe

Volume 194, no. 9, pages 2321–2333, 2012. Page 2330: The legend to Fig. 6E should read as follows. "Immunoblot of unboiled, recombinant TprC<sup>N</sup> and TprC<sup>C</sup> separated by SDS-PAGE and then probed with rabbit anti-TprC<sup>Fl</sup> polyclonal antiserum (asterisk indicates TprC<sup>C</sup> trimer)."

Page 2331: The legend to Fig. 7C should read as follows. "Reactivities of the anti-Tpr $C^{FI}$  antiserum and IRS used for the opsonophagocytosis assays in Fig. 5 against Tpr $C^N$  and folded, unboiled Tpr $C^C$ . The immunoblot shown in Fig. 6E was stripped and reprobed with IRS. Bottom and top panels show the reactivity of Tpr $C^N$  and Tpr $C^C$  monomers with anti-Tpr $C^{FI}$  antiserum (reproduced from Fig. 6E) and IRS, respectively."

Copyright © 2014, American Society for Microbiology. All Rights Reserved. doi:10.1128/JB.02012-14