



Sources of DNA Double-Strand Breaks and Models of Recombinational DNA Repair

Anuja Mehta and James E. Haber

Rosenstiel Basic Medical Sciences Research Center, MS029 Rosenstiel Center, Brandeis University, Waltham, Massachusetts 02454-9110

Correspondence: haber@brandeis.edu

DNA is subject to many endogenous and exogenous insults that impair DNA replication and proper chromosome segregation. DNA double-strand breaks (DSBs) are one of the most toxic of these lesions and must be repaired to preserve chromosomal integrity. Eukaryotes are equipped with several different, but related, repair mechanisms involving homologous recombination, including single-strand annealing, gene conversion, and break-induced replication. In this review, we highlight the chief sources of DSBs and crucial requirements for each of these repair processes, as well as the methods to identify and study intermediate steps in DSB repair by homologous recombination.

EXOGENOUS AND ENDOGENOUS SOURCES OF DNA DOUBLE-STRAND BREAKS

DNA damage can occur as a result of endogenous metabolic reactions and replication stress or from exogenous sources like radiation and chemotherapeutics. Damage comes in several different varieties: base lesions, intra- and interstrand cross-links, DNA-protein cross-links, and both single- and double-strand breaks (DSBs) (Lindahl 1993). Some types of damage, such as oxidative damage to DNA bases, arise, and are repaired, as often as 10^5 lesions per cell each day (Hoeijmakers 2009). Much less frequent are DNA DSBs, in which the phosphate backbones of the two complementary DNA strands are broken simultaneously, and these are one of the most cytotoxic forms of lesion.

Some well-known exogenous DNA damaging agents (clastogens) are anticancer chemotherapeutic drugs and ionizing radiation (IR). Chemotherapeutic drugs include DNA-alkylating agents such as methyl methanesulfonate and temozolomide, cross-linking agents such as mitomycin C and cisplatin, and radiomimetic compounds such as bleomycin or phleomycin (Chen and Stubbe 2005; Wyrobek et al. 2005). Another class are topoisomerase inhibitors such as camptothecin and etoposide, which induce the formation of single-strand breaks (SSBs) and DSBs, respectively, by trapping covalently linked topoisomerase-DNA cleavage complexes (Koster et al. 2007). Other drugs, such as hydroxyurea and aphidicolin, impair the progression of replication by depleting deoxyribonucleotide pools or inhibiting DNA polymerase.

Editors: Stephen Kowalczykowski, Neil Hunter, and Wolf-Dietrich Heyer
Additional Perspectives on DNA Recombination available at www.cshperspectives.org

Copyright © 2014 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a016428
Cite this article as *Cold Spring Harb Perspect Biol* 2014;6:a016428

Ionizing radiation leads to extensive base damage and, additionally, creates DNA SSBs by producing radiolysis radicals that attack the sugar-phosphate backbone (Ward 1994; Thompson 2012). Frequently, at high doses of irradiation, two such nicks are present in complementary DNA strands within one helical turn leading to DSBs (Milligan et al. 1995). There are about 10 SSBs for each DSB created by IR (Ma et al. 2012). IR breakage frequently leaves “dirty ends,” consisting of phosphoglycolates and terminal nucleotides, that cannot be ligated to “clean” ends consisting of a 5′ phosphate and 3′-OH group, such as those created by endonucleases (Weinfeld and Soderlind 1991).

Even in the absence of exogenously inflicted stress during an unperturbed cell cycle, DNA is vulnerable to suffer damage during replication, which, if unrepaired, can promote genomic instability. There are numerous natural impediments that lead to pausing or blocking of a replication fork, such as unusual DNA and chromatin structures or collisions with transcription machinery (Prado and Aguilera 2005; Aguilera and Gaillard 2014) or DNA-binding proteins (Mirkin and Mirkin 2007; Merrikh et al. 2012). The types of damage produced by normal cellular processes are very similar to those caused by some environmental agents (De Bont and van Larebeke 2004).

One way to estimate the frequency of spontaneous DSBs is to count them in cells in which DSB repair is prevented. In budding yeast, one can examine the fate of a single G₁ cell lacking the *RAD52* gene that is required for DSB repair by homologous recombination (HR). Approximately one cell in eight gives rise to a pair of daughter cells, one of which is inviable (J Haber, unpubl., cited in Coic et al. 2008). This finding implies that there is a DSB that arises during DNA replication that would normally be repaired by sister chromatid recombination in a recombination-proficient cell. Given a genome size of $\sim 1.2 \times 10^7$ bp, this result, hence, suggests that there is about one spontaneous DSB per 10^8 bp. Another study estimates that, in normal human cells, $\sim 1\%$ of single-strand lesions are converted to ~ 50 DSBs per cell per cell cycle, that is, about one DSB per 10^8 bp (Vilen-

chik and Knudson 2003). In vertebrate cells such as chicken DT40, depleted for yet another key recombination protein, Rad51, the estimated rate of breakage is of the same magnitude (Sonoda et al. 2001).

An alternate way to count DSBs in a cell is to monitor the formation of damage-induced foci, either by indirect immunofluorescent staining or the use of fluorescent proteins fused to proteins that are recruited to the sites of DNA damage as part of the DNA damage response. In vertebrate cells, phosphorylation of the minor histone H2A variant, H2AX, to produce so-called γ -H2AX, is often used as an indicator of the incidence of DSBs; however, it is now becoming evident that γ -H2AX can be associated with DNA damage other than DSBs (Soutoglou and Misteli 2008; Löbrich et al. 2010; Valdiguélias et al. 2013) and thus may overestimate their incidence. Binding of other key DNA repair proteins, such as 53BP1, also serves as a surrogate for monitoring DSBs, as do the appearance of RPA (replication protein A) and Rad51 foci (Haaf et al. 1995; Raderschall et al. 1999; Noon and Goodarzi 2011). In budding yeast, the most frequently used live-cell marker of DSB damage is the recruitment of Rad52-YFP (or other colors) into damage-induced foci. The fact that even multiple DSBs result in a single Rad52 focus has been interpreted as evidence that DSBs aggregate into a repair center (Lisby et al. 2003); however, recent studies have suggested that Rad52-fluorescent protein fusion proteins have a remarkable ability to aggregate, so that they mark strongly only one of several independent DSBs (M Brown, I Fitzgerald, B Glick, and D Bishop, unpubl.; C-S Lee and J Haber, unpubl.).

The fact that the majority of spontaneous DSBs appear in the context of DNA replication (Syeda et al. 2014) could suggest that DNA may be nicked before the passage of the replication fork, thus creating a broken chromatid, and, with the arrival of another replication fork in the opposite direction, an intact sister chromatid. However, such breaks can also arise by the processing of stalled replication forks. Stalled forks have been studied by impairing replication, for example, by adding hydroxyurea or introducing replication fork-blocking sequenc-

es. Stalled forks can regress by the unwinding and annealing of the newly synthesized strands to form a “chicken foot” structure, which is known as a Holliday junction (Fig. 1). These Holliday junctions (HJs) can be cleaved by structure-selective nucleases or HJ resolvases, such as Mus81-Mms4 or Yen1, to create a broken end and intact chromatid (Wyatt and West 2014). The broken end can then be processed by 5' to 3' resection (Symington 2014) and, after binding Rad51, engage in recombination-dependent replication restart, that is, break-induced replication (BIR) (Hanada et al. 2007; Petermann and Helleday 2010).

A growing body of evidence implicates transcription as one of the leading causes of DSBs and associated genome instability. Apart from hindering replication fork progression (Prado and Aguilera 2005), R-loops, three-strand nucleic acid structures formed by an RNA:DNA hybrid, plus a displaced single-stranded DNA (ssDNA) identical to the RNA molecule, have been linked to mutagenesis, recombination, and chromosome rearrangements (Huertas

and Aguilera 2003; Kim and Jinks-Robertson 2009; Lin et al. 2010). R-loops are mainly formed by defects in RNA metabolism, but also seem to arise naturally in wild-type cells (Helmrich et al. 2011; Wahba et al. 2011). It was recently reported that formation of these RNA:DNA hybrids, in certain yeast transcription repression and RNA degradation mutants, requires the involvement of the HR machinery, including Rad51 and Rad52 (Wahba et al. 2013). Several high-throughput screens in budding yeast and mammalian cells have identified multiple RNA biogenesis factors whose depletion leads to R-loops-mediated DSB formation and the activation of the DNA damage response (Li and Manley 2005; Paulsen et al. 2009; Stirling et al. 2012; Gavalda et al. 2013). The evolutionarily conserved RNase H (in prokaryotes) or RNase H1 (in eukaryotes), which specifically degrades the RNA portion of hybrids, plays a major role in suppression and removal of such structures and, hence, helps maintain genome stability (Lin et al. 2010; Helmrich et al. 2011; Wahba et al. 2011). Thus, a complex network




Figure 1. Replication fork stalling and restart. DNA replication can be stalled at UV-induced thymidine dimers (TT), as well as DNA secondary structures. A stalled replication fork (A) can undergo regression and pairing of the newly synthesized strands to form a HJ “chicken foot” intermediate (B). The HJ can be cleaved by HJ resolvases (C) to lead to a collapsed fork, effectively a one-ended chromosome break (D). The free end can initiate HR by strand invasion (E) to bypass the lesion and resume replication (F). Newly synthesized DNA is depicted as dashed lines in the same color as the template; arrowheads indicate 3' ends.

is emerging that links DNA damage and RNA metabolism.

PROGRAMMED DSBs

Although DSBs are among the most deleterious lesions, several physiologically and developmentally important processes require the generation of programmed site-specific DSBs and their subsequent repair by various pathways. In mammals, DSBs are produced by the RAG protein complex to initiate V(D)J recombination for assembling immunoglobulin antigen receptor genes, as well as T-cell receptor genes. DSBs also arise during immunoglobulin class switching (Soulas-Sprauel et al. 2007). A recent claim that there are DSBs in stimulated mouse neurons (Suberbielle et al. 2013), possibly dependent on Spo11 endonuclease, will require further investigation. DSB-induced recombination and subsequent crossover formation are crucial for faithful segregation of homologous chromosomes of different parental origins by the meiotic spindle during sexual reproduction (Lam and Keeney 2014). This is catalyzed by the generation of hundreds of programmed DSBs at nonrandom hotspots by the evolutionarily conserved meiosis-specific topoisomerase-II-like Spo11 endonuclease (Keeney 2008).

Ironically, missegregated chromosomes, although undergoing cytokinesis, can also promote the direct and indirect acquisition of DNA damage that may further lead to unbalanced translocations in the daughter cells (Janssen et al. 2011; Ganem and Pellman 2012). Segregation defects formed by merotelic kinetochore attachments or other causes of chromosome nondisjunction may foster the formation of ultrafine chromosome bridges during anaphase. Any delay in clearing the chromosomes from the central spindle and out of the path of the incoming actin-myosin cytokinetic ring can lead to cleavage of the chromatin and formation of DSBs (Hoffelder et al. 2004; Samoshkin et al. 2009; Janssen et al. 2011; Quevedo et al. 2012). Similar problems arise in the mitosis of dicentric chromosomes (Haber and Thorburn 1984; Kramer et al. 1994) or when intertwined sister chromatids are not disentangled by topoisom-

erase II (Spell and Holm 1994; Baxter and Diffley 2008). These broken fragments are recombinogenic and can generate chromosome translocations in the next cell cycle (Janssen et al. 2011; Quevedo et al. 2012). Suppression of cytokinetic furrowing can rescue the formation of DSBs at such chromosome bridges, confirming that they are formed as a result of cytokinesis (Baxter and Diffley 2008; Janssen et al. 2011). Often, anaphase-lagging chromosomes are left so severely behind that on telophase they form their own independent nuclear envelope, creating a micronucleus. It has recently been shown that, because of inefficient nuclear import, micronuclei fail to attain all of the required replication and repair components and, hence, show higher replication stress and DNA fragmentation (Crasta et al. 2012).

In many organisms, the excision of transposable elements creates DSBs that are usually repaired by gene conversion (GC) with a sister chromatid from which excision has not occurred. Recombinational repair can also occur with ectopic homologous sequences (Gloor et al. 1991). In the ciliate *Paramecium*, during the development of the somatic nucleus, programmed DSBs initiate the extensive genome rearrangements that take place at each sexual cycle. In particular, thousands of short non-coding germline sequences, called internal eliminated sequences, are spliced out (Betermier 2004). These developmentally programmed DNA DSBs depend on the domesticated transposase PiggyMac (Baudry et al. 2009) and are repaired using components of the nonhomologous end-joining (NHEJ) pathway (Kapusta et al. 2011).

In budding and fission yeasts, programmed switching of mating-type genes begins with a DSB within the mating-type locus that is repaired by ectopic recombination with donor sequences on the same chromosome, encoding opposite mating-type alleles. In *Schizosaccharomyces pombe*, *mat1* switching depends on a programmed ssDNA nick that, during replication, is converted into a DSB (Arcangioli 2000; Klar 2007). *MAT* switching in *Saccharomyces cerevisiae* uses a site-specific homothallic (HO) endonuclease, which cleaves a degenerate 24-bp

sequence to generate a DSB with 4-bp, 3'-OH overhanging ends (Haber 2012). Curiously, in another budding yeast, *Kluyveromyces lactis*, a very similar *MAT* switching event involving evolutionarily related mating-type sequences occurs without HO endonuclease; instead, a transposon-like sequence apparently excises from the *MAT* locus and leaves hairpin-closed ends that are opened to initiate DSB-mediated switching (Barsoum et al. 2010). Another well-characterized, specialized endonuclease is the mitochondrial enzyme I-*SceI*, which is encoded by the optional intron of the 21S ribosomal RNA gene and responsible for intron mobility (Colleaux et al. 1988). Much of the detailed description of DSB repair in budding yeast has relied on the inducible expression of HO endonuclease or a codon-optimized I-*SceI*, which can cut cleavage sites introduced in different chromosomal contexts (Krogh and Symington 2004; Haber 2006; Wyman and Kanaar 2006; Kass and Jasin 2010).

Thus, DNA DSB formation and repair can be both deleterious and beneficial, and together have played a major role in the evolutionary development and survival of all living organisms.

REPAIR OF DSBs CAN OCCUR IN SEVERAL WAYS

Two mechanistically distinct sets of pathways have evolved to repair DSBs: NHEJ and HR (Fig. 2). This review focuses on HR in somatic cells, but a brief summary of NHEJ is provided.

NHEJ involves modification and ligation of the broken DNA ends with very little or no homology, often creating small deletions or insertions (Fig. 2A). Although NHEJ can occur throughout the cell cycle, it is especially important in the G₁ stage, when a key initial step in HR, the 5' to 3' resection of DSB ends, is blocked (Aylon et al. 2004; Ira et al. 2004). In the special case of haploid cells, such as in the well-studied budding or fission yeasts, G₁ cells lack a homologous chromosome and, hence, can only use NHEJ until chromosomal replication creates a sister chromatid that can be used as a template to repair a DSB (Moore and Haber 1996). NHEJ

essentially consists of several related but distinct mechanisms that have different genetic requirements and lengths of microhomology at the junctions. Precise end joining of 3'-overhanging ends (e.g., religating the ends of a restriction endonuclease cleavage) requires the Ku70 and Ku80 proteins, as well as DNA ligase IV; whereas, more extensive deletions with longer microhomologies at the junction prove to be Ku independent and, in mammals, DNA ligase IV independent. This alternative NHEJ or microhomology-mediated end-joining (MMEJ) pathway is evident even in wild-type cells. The reader is encouraged to read several excellent reviews (Bennardo et al. 2008; McVey and Lee 2008; Mladenov and Iliakis 2011; Chiruvella et al. 2013; Decottignies 2013).

HR

DNA DSBs can be repaired by several different HR pathways (Fig. 2). Single-strand annealing (SSA) is the simplest mechanism, allowing the formation of a deletion between homologous sequences flanking a DSB. Other types of HR all depend on the recognition and pairing of broken DNA ends with intact homologous sequences present on a sister chromatid, an allelic locus, or at some ectopic location in the genome (Pâques and Haber 1999; Krogh and Symington 2004). We will focus on two major mechanisms of HR: GC and BIR. An additional HR mechanism, gene targeting, can account for the integration of foreign DNA into a homologous chromosomal locus (Rothstein 1983). We further distinguish two distinct pathways of GC: one involving the formation of a pair of HJs that often leads to crossovers (COs) accompanying repair and one in which COs are rare.

All HR repair mechanisms require proteins belonging to the evolutionarily conserved *RAD52* epistasis group. Astonishingly, the *Rad52* protein itself, which is the most recombination-essential gene in budding and fission yeasts, appears to have an inconsiderable role in HR in metazoans. *Rad52* is absent in flies and worms and, although it is present in mammals, it plays a minor role in repair, apparently limited to strand annealing. Instead *Rad52*'s central role




Figure 2. Pathways of DNA DSB repair. DSBs are efficiently repaired by HR and NHEJ. DSBs are processed by 5' to 3' end resection producing 3' single-stranded tails. (A) NHEJ involves ligation of broken ends, with little or no base pairing, to produce small deletions or insertions. (B) Single-strand annealing (SSA) takes place when resection reveals flanking homologous repeats that can anneal, leading to deletion of the intervening sequences. (C,D) Repair by two different mechanisms of GC results in a short patch of new DNA synthesis to repair the DSB. (C) In synthesis-dependent strand-annealing (SDSA), the newly synthesized strand dissociates from the D-loop and results in a noncrossover (NCO) outcome with no change to the template DNA. (D) The double Holliday Junction (dHJ) pathway involves second end capture to stabilize the D-loop. The dHJ structure can be resolved either by helicase and topoisomerase-mediated dissolution to give NCO or cleaved by HJ resolvases to produce both crossover (CO) or NCO outcomes. (E) BIR involves both leading and lagging strand synthesis and results in loss of heterozygosity or, if the template is located ectopically, a nonreciprocal translocation. Newly synthesized DNA is depicted as dashed lines in the same color as the template; arrowheads indicate 3' ends.



is replaced by orthologs of the mammalian BRCA2 protein (Jensen et al. 2010; Liu et al. 2010; Holloman 2011). It seems that budding and fission yeasts may have lost their BRCA2 orthologs because another yeast, *Ustilago maydis*, has retained it and is dependent on it for HR (Yang et al. 2005). GC and most BIR depend on Rad51 recombinase, the homolog of bacterial RecA. Rad51 is a DNA-dependent ATPase that forms a filament on ssDNA and promotes strand invasion with a homologous double-stranded partner.

The series of biochemical steps in HR have been best studied in budding yeast, in which

several early steps can be monitored in real time by a variety of techniques analyzing DNA intermediates and the proteins bound to these intermediates after inducing a DSB by the site-specific HO endonuclease (White and Haber 1990; Hicks et al. 2011) or other nucleases such as I-SceI (Fig. 3) (Plessis et al. 1992). Efficiency of DSB formation in a population can be gauged by a Southern blot (Fig. 3D) or monitoring the loss of a polymerase chain reaction (PCR) product with primers placed on opposite sides of the DSB site.

All HR repair mechanisms are reliant on a common critical initial step: extensive 5' to 3'




Figure 3. Key intermediate steps of HR and methods to study them. (A) Key proteins are depicted in sequential early steps in GC in budding yeast. DSB formation (i) is immediately followed by 5' to 3' resection (ii). The 3' tails are stabilized by RPA (iii), which is then replaced with Rad51 recombinase with assistance from accessory proteins like Rad52 (iv). Once the homologous donor is found, strand invasion occurs, resulting in formation of a D-loop (v) by displacement of the identical strand and base pairing with the complementary strand at the donor. (vi) Various components of the replication machinery assemble to start copying from the donor template. (vii) The break is sealed. Small arrows denote positions of PCR primer pairs used to analyze intermediate steps, shown on the right for *MAT* switching in *S. cerevisiae*. (B) Recruitment of Rad51 at DSB site (the *MATa* locus) by chromatin immunoprecipitation (ChIP) followed by quantitative PCR (qPCR) using primers p1 and p2 (solid line). Rad51 binding to the donor template (budding yeast *HMLα* locus) using primers p3 and p4 for qPCR (dotted line). Error bars indicate standard error of the mean. (C) The initiation of new DNA synthesis by primer extension is detected by using PCR primers p5 and p6, which amplifies a unique fragment once new DNA synthesis has been initiated (solid line). A dotted line shows quantitative densitometric analysis of Southern blot (D) that follows GC progression in budding yeast as *MATa* switches to *MATα*. Data from the investigators (A Mehta and J Haber, unpubl.).

processing of the broken DNA ends (Fig. 3Aii). Whereas NHEJ and MMEJ involve limited processing of DNA ends for ligation, a prerequisite for HR is that broken ends are considerably resected to generate 3'-ended ssDNA tails that, once bound by Rad51 recombinase, act as the functionally active agents in searching for homologous template sequences to repair the DSB. 5' to 3' resection is a surprisingly complex

process. An initial incision, removing about 100 nt, is performed by the Mre11-Rad50-Xrs2 complex along with the Sae2 protein. Subsequently, two other excision machines remove DNA at a rate of about 4 kb/hr. Exo1 chews off mononucleotides, whereas Sgs1-Top3-Rmi1-Dna2 acts as a helicase/endonuclease to clip off short oligonucleotides from the 5' strand (Huertas et al. 2008; Mimitou and Symington

2008; Zhu et al. 2008; Cejka et al. 2010; Niu et al. 2010). Resection of a single HO endonuclease-induced DSB is highly limited in G₁ phase of the cell cycle, when cyclin-dependent kinase Cdk1 (Cdc28) activity is low. The rate of resection increases once cells progress through the cell cycle and initiate DNA synthesis followed by mitosis in the G₂/M phase, when Cdk1 activity is high (Aylon et al. 2004; Ira et al. 2004; Barlow et al. 2008). Additionally, Exo1 is impaired by binding of the Ku70-Ku80 proteins that facilitate NHEJ (Balestrini et al. 2013). Analogous machinery is found in mammalian cells, in which Sae2's ortholog, CtIP, plays a central role and Sgs1's ortholog, BLM helicase, appears to be coupled with both EXO1 and DNA2 (Sartori et al. 2007; Nimonkar et al. 2011; Sun et al. 2012).

Initially, the multimeric replication protein A (RPA) binds to ssDNA overhangs presumably to take out kinks and melt DNA secondary structures (internal base pairing) (Fig. 3Aiii). RPA is then replaced with the Rad51 recombinase, which forms a filament along the ssDNA (Fig. 3Aiv). As with the best-studied case of *Escherichia coli* RecA protein, the ssDNA is stretched within the Rad51 nucleoprotein filament to about 1.5 times the length of B-form DNA (Chen et al. 2008). However, this stretching is not uniform. The three bases bound by the RecA recombinase, and presumably by Rad51, are in almost a B-DNA configuration; the stretching happens predominantly between the 3-bp units. This arrangement may facilitate pairing through canonical Watson-Crick hydrogen bonds with complementary triplets in the donor duplex DNA; when RecA or Rad51 binds to dsDNA, it also becomes stretched to 150% of its normal length (Chen et al. 2008). The creation of the Rad51 filament is a complex process. After the binding of RPA, with the aid of the recombination mediator proteins (in yeast Rad52; in vertebrates BRCA2) plus several Rad51 paralog proteins (in yeast Rad55 and Rad57; in vertebrates RAD51B, RAD51C, RAD51D, XRCC2, and XRCC3) (Daley et al. 2014; Morrical 2014), RPA is displaced and the Rad51 nucleoprotein filament is formed. These steps can be seen in budding yeast in real time

through the use of ChIP, using anti-RPA and anti-Rad51 antibodies (Fig. 3B) (Sugawara et al. 2003; Wang and Haber 2004). RPA binding is detected ~10 min before Rad51 binding; no Rad51 binding occurs without Rad52.

ChIP can also be used to visualize the key next step in DSB repair: Rad51 nucleoprotein filament-mediated search for a distant homologous sequence and subsequent strand invasion (synapsis) between the resected end of the DSB and its duplex homologous donor sequence (Fig. 3Av). As shown in Figure 3B, these are remarkably slow steps. In the absence of special pairing sequences (i.e., the recombination enhancer that facilitates *MAT* switching between *MATa* and *HMLα*) (Li et al. 2012), Rad51 and its invading ssDNA strand becomes associated with the donor duplex DNA only after 30–45 min (Hicks et al. 2011; A Mehta and J Haber, unpubl.). In addition to Rad52, the Swi2/Snf2 homolog Rad54 also facilitates functional strand invasion (Sugawara et al. 2003; Kiianitsa et al. 2006; Hicks et al. 2011).

Following strand invasion, new DNA synthesis using either DNA polymerase (Pol)δ or Pole occurs using the 3' invading end as a primer. Initiation of new DNA synthesis can be monitored by a PCR-based primer extension assay using two primers: one complementary to sequences distal to the DSB break-site and one within the donor (Fig. 3Avi,C) (White and Haber 1990). Several pathways can then occur (discussed below), ensuing in completion of repair. Southern blot analysis is used to accurately reveal the rate of repair product formation (Fig. 3C,D). DSB repair by HR can occur with less than 100 bp of homology with the DSB ends, but efficient repair is achieved with a few hundred bp (Sugawara and Haber 1992).

SSA

SSA is restricted to repair of DNA breaks that are flanked by direct repeats that can be as short as 30 nt (Sugawara et al. 2000; Villarreal et al. 2012). Resection exposes the complementary strands of homologous sequences, which recombine resulting in a deletion containing a single copy of the repeated sequence (Fig. 2B).

SSA is therefore considered to be highly mutagenic. The nonhomologous single-stranded tails are removed by the Rad1-Rad10 endonuclease (XPF-ERCC1 in mammals) in a complex that includes both the Msh2-Msh3 mismatch repair proteins and “scaffold” proteins Slx4 and Saw1 (Sugawara et al. 1997; Li et al. 2008; Toh et al. 2010). After tail clipping, remaining gaps must be filled in by DNA synthesis and sealed by ligation. SSA requires the strand-annealing activity of Rad52 and is aided by the Rad52 homolog Rad59 (Sugawara et al. 2000); however, SSA does not involve DNA-strand invasion and thus is independent of the Rad51 recombinase (Ivanov et al. 1996).

GC

GC was initially defined as a nonreciprocal transfer of genetic information from one chromosome to its homolog during meiosis, but its meaning has been broadened to include DSB repair events in which a short patch of new DNA synthesis is copied from the homologous template. In budding yeast, about 10%–20% of interchromosomal allelic mitotic GCs are associated with crossing over, whereas COs are less frequent when the DSB uses an ectopic template that shares only a few kb with the regions around the break (Pâques and Haber 1999; Ira et al. 2003).

GC does not require many components of the normal DNA replication machinery, including DNA Pol α -primase or Cdc45-GINS (complex consisting of four proteins: Sld5, Psf1, Psf2, and Psf3)-Mcm helicase complex (Wang et al. 2004; Lydeard et al. 2010). The efficiency of GC is also not significantly reduced by three mutations that severely impair the alternative HR repair mechanism of BIR, which can involve the synthesis of hundreds of kilobases of DNA: the nonessential DNA Pol δ subunit *pol32* Δ (Lydeard et al. 2007; Jain et al. 2009), the 5' to 3' helicase *pif1* Δ (Chung et al. 2010; Wilson et al. 2013), and the proliferating cell nuclear antigen (PCNA) mutation *pol30-FF248,249AA* (Lydeard et al. 2010). The fact that GC does not require many of the normal DNA replication processivity factors might explain why it is

highly susceptible to mutagenesis. Most mutations have the signatures of template switching during repair DNA synthesis (Strathern et al. 1995; Hicks et al. 2010).

Two different mechanisms, both of which are supported by substantial experimental data, can explain GCs and the different outcomes.

The Double Holliday Junction Mechanism

The double-strand break repair model, now known as the double Holliday junction or dHJ model, was first suggested by Resnick (1976) and later extended and elaborated by Szostak et al. (1983). Following resection and invasion into an intact homologous template, the 3' Rad51-coated nucleoprotein filament base pairs with the complementary strand creating a displacement loop (D-loop) structure consisting of a region of heteroduplex DNA and displaced single strand of DNA. The D-loop can be extended by the initiation of new DNA synthesis from the 3' end of the invading strand or the action of helicases, so that the ssDNA of the opposite side of the DSB can anneal, thus forming a dHJ intermediate (Fig. 2C). Alternatively, two independent strand invasions from both DSB ends, followed by simultaneous DNA synthesis and annealing could also result in a dHJ intermediate. No experimental evidence has shown whether recombination depends on one end or whether both undergo strand invasion. These HJs can be cleaved by one of several HJ resolvases, and, depending on which pair of strands is cut, can yield a noncrossover (NCO) or CO outcome. As an alternative to cleavage, dHJs can be “dissolved” to yield exclusively NCO outcomes (Wu and Hickson 2003).

The formation of dHJs has been extensively studied in budding yeast meiosis, but they have also been visualized in mitotic cells after a DSB is initiated by expression of a site-specific I-SceI endonuclease (Bzymek et al. 2010). Evidence for the presence of dHJ intermediates in budding yeast comes from ectopic recombination experiments in which the components of the dHJ “dissolvase” (Sgs1-Top3-Rmi1, or STR) are deleted. The proportion of GC events accompanied by a CO—in this case, a reciprocal

translocation—increases from 4%–~12% (Ira et al. 2003) when Sgs1 or other cofactors are absent. Assuming that *sgs1Δ* does not alter the proportion of events that form the dHJ intermediate, these results suggest 2/3 of the dHJs are normally dissolved. The proportion of COs increases even more when both Sgs1 and another 3' to 5' helicase, Mph1, are deleted. In this instance, it appears that deleting *MPH1* shifts more of the repair events into the dHJ pathway (Prakash et al. 2009). Another estimate of the fraction of dHJs processed by the STR complex was obtained by Mitchel et al. (2013) by studying the position of heteroduplex DNA relative to the initiating gap in a plasmid-based gap-repair system. Their studies suggested that the fraction of dHJs processed by “resolution” is roughly equal to that processed by “dissolution.”

The Synthesis-Dependent Strand-Annealing Mechanism

Synthesis-dependent strand-annealing (SDSA) begins with the common step of strand invasion and formation of a D-loop (Fig. 2D). This is followed by new DNA synthesis (as monitored by primer extension assay) initiated by DNA Pol δ or Pol ϵ from the invading 3' end. There is mounting evidence that the principal repair polymerase is Pol δ (Li et al. 2009a; Sebesta et al. 2011; Prindle and Loeb 2012), but repair is not eliminated either in a temperature-sensitive mutation of Pol δ or Pol ϵ , suggesting both can be used (Lydeard et al. 2007). Pol α primase is, however, not required (Wang et al. 2004). A carboxy-terminal truncation mutation of Pol δ , *pol3-ct*, which does not affect replication, results in short DNA repair tracts in both mitosis and meiosis (Maloiel et al. 2004, 2008). If one examines mutations created during DSB repair (Hicks et al. 2010), defects of the 3' to 5' proofreading exonuclease activity of Pol ϵ resulted in the appearance of +1 frameshift mutations that were much less evident in wild-type repair. However, a much more dramatic effect was seen in the absence of proofreading activity of Pol δ ; several classes of mutations characterized as template switching defects (–1 frameshifts, quasi-palindrome mutations, and interchromosomal tem-

plate switches) are surprisingly eliminated, leading to the suggestion that the wild-type Pol δ enzyme is responsible for most of these events.

In *Saccharomyces*, there is little effect of ablating either of two translesion DNA polymerases, Pol ζ or Pol η , although the error-prone Pol ζ polymerase has been shown to be the major contributor to mutations that arise in ssDNA regions, including those adjacent to the region in which both strands are newly copied during SDSA (Ratray et al. 2002; Yang et al. 2008). In vertebrates, evidence suggests that the translesion DNA polymerase, Pol η , is a more important player in recombinational repair (Kawamoto et al. 2005). This conclusion is supported by demonstrations that Pol η can perform repair synthesis in vitro (McIlwraith et al. 2005; Sneeden et al. 2013).

What makes SDSA synthesis unusual is that it is not semiconservative, in which the newly copied strand remains base paired to the donor template. Instead, the newly synthesized strand appears to be displaced from a migrating replication “bubble” and eventually pairs with the resected 3' ssDNA on the other side of the DSB, resulting in an NCO outcome (Fig. 2D). Evidence in favor of SDSA includes the fact that the great majority of mitotic DSB repair events occur without an associated CO. Moreover, a number of studies in which GC is associated with the appearance of repair-associated mutations have found that all of the alterations are found in the recipient locus, whereas the donor remains unchanged—a result more consistent with SDSA than the dHJ mechanism (although dissolving dHJs would also result in this outcome) (Pâques and Haber 1999). Direct evidence for this mechanism was obtained by “heavy-light” density transfer experiments that showed that all of the newly copied DNA was located at the recipient locus, whereas the donor sequences were unaltered (Ira et al. 2006), indicating a conservative mode of strand inheritance. How the nascent DNA strand is unwound from the template is not known in detail, but both Mph1 and Srs2 3' to 5' helicases appear to be involved. As noted above, *mph1Δ* strains appear to direct a larger proportion of outcomes to the dHJ pathway, as would occur if Mph1

promoted the strand displacement. In contrast, *srs2Δ* strains show a 70% drop in viability, which results from a specific deficit in recovering NCO outcomes (Ira et al. 2003). This result suggests that Srs2 is required for completing SDSA, but not the dHJ pathway.

Although SDSA is presumed to yield exclusively NCO outcomes, one should not discount models that include the possibility of crossing over wherein the migrating D-loop associated with synthesizing the first strand can be captured by the second end, leading to the formation of a dHJ intermediate that can be resolved into a CO associated with GC (Ferguson and Holloman 1996; Pâques et al. 1998).

BIR

Under some circumstances, a broken chromosome may present only one end for repair, for instance, at collapsed forks during S-phase and in lengthening of telomeres (Doksani and de Lange 2014) in telomerase-deficient cells in pathways known as “alternative lengthening of telomeres” (McEachern and Haber 2006). When only one end of a chromosomal DSB shares homology with a template, cells rely on recombination-dependent DNA replication (i.e., BIR), a process in which strand invasion sets up a unidirectional replication fork capable of copying all of the sequences distal to the site of homology up to the telomere (Fig. 2E) (Malkova et al. 1996, 2005; Davis and Symington 2004). Repair by this mode results in a nonreciprocal translocation and, if the template is a homologous chromosome, extensive loss of heterozygosity. BIR has been observed with as little as 70 bp of shared homology between the DSB end and an ectopic template (Bosco and Haber 1998), but is much more efficient with longer regions of homology.

BIR has been analyzed in budding yeast both by transformation of a linear plasmid that has telomere-forming sequences at one end and homology with a chromosomal template on the other (Morrow et al. 1997; Davis and Symington 2004; Marrero and Symington 2010), and creating broken chromosomes using HO or I-SceI endonucleases (Bosco and Haber

1998; Malkova et al. 2005; Lydeard et al. 2007; Ruiz et al. 2009). The initial steps of BIR, up to homology search and strand invasion appear to be identical to those of GC and require all of the same proteins (Davis and Symington 2004; Jain et al. 2009). However, the subsequent repair DNA synthesis occurs very differently. BIR requires all of the replication factors and the three major DNA polymerases required for leading and lagging strand synthesis, except some proteins that are needed exclusively for the assembly of the prereplication complex at origins of replication (Lydeard et al. 2007, 2010).

Nevertheless, in several respects, BIR is quite different from normal leading- and lagging-strand S-phase replication. First, BIR is remarkably mutagenic compared to normal replication (Deem et al. 2011). Second, at least for the first several kb that are copied, BIR shows a very high frequency of template switching, from one homologous chromosomal template to the other (Smith et al. 2007). This instability may be a reflection of the finding that BIR can be initiated without DNA Pole, but requires this normally leading-strand polymerase for its completion. As noted above, unlike GC, BIR is significantly reduced by the deletion of *POL32* (Lydeard et al. 2007) or *PIF1* (Chung et al. 2010; Wilson et al. 2013), or the PCNA mutation *pol30-FF248,249AA* (Lydeard et al. 2010). This PCNA mutation was found as a suppressor of the cold sensitivity of *pol32Δ*, however, it is by itself as defective for BIR as *pol32Δ* and does not suppress *pol32Δ*'s BIR defect. The mutant has no effect on normal replication; moreover, the location of the mutated sites is unrelated to PCNA mutations involved in recruiting translesion DNA polymerases during replication fork stalling. Analogous mutations created in the *Xenopus* PCNA protein show a defect in an in vitro replication restart assay (Hashimoto et al. 2012). These differences point toward unique features of BIR, either in the establishment of the repair replication fork or its efficiency.

Recently, an explanation for the differences between BIR and normal replication was provided by the work from the Malkova and Lobachev laboratories (Saini et al. 2013), which analyzed intermediates of BIR by DNA combing

and two-dimensional gel electrophoresis. These analyses produced several critical revelations. First, all of the newly copied DNA is found on the extension of the originally broken molecule, a conclusion also reached by Donnianni and Symington (2013). This “conservative” mode of DNA synthesis suggests either that semiconservatively replicated DNA is unwound by branch migration of a HJ formed behind the replication fork or the second new strand is copied from the first (Fig. 2E). Indeed, the latter possibility seems to be the case as the migration of replication intermediates is consistent with that expected for a migrating D-loop (as first proposed by Formosa and Alberts 1986) with a long ssDNA extension. In this respect, BIR would share certain common steps with SDSA but, as noted above, none of the BIR-specific factors (Pol32, Pif1, or the PCNA mutant) have much effect on repair by GC. One can rationalize the lack of effect of mutating these factors by the fact that most GC assays require copying of no more than ~1 kb of new DNA. Indeed, when cells are required to repair a DSB in which the homologous ends of the DSB invade a template separated by ≥ 5 kb, the events become dependent on *POL32* (Jain et al. 2009) and are impaired by the PCNA mutation, *pol30-FF248,249AA* (A Mehta, T Ryu, and J Haber, unpubl.).

Another interesting distinction between GC and BIR is that the latter shows a dramatic delay in the initiation of new DNA synthesis. This delay reflects a replication execution checkpoint (REC) that somehow monitors whether only one or both ends of a DSB can successfully pair with homologous template sequences on the same template and in the proper orientation and distance to complete GC (Jain et al. 2009). The kinetics of strand invasion, as monitored by the appearance of Rad51 at the donor locus by ChIP, are the same for GC and BIR, so the REC acts before the initiation of new DNA synthesis and is seen even when both ends of the DSB can pair with donor sequences, when they are separated by ≥ 5 kb. The strong kinetic barrier to promoting BIR also helps to explain the hierarchy of DSB repair events; in a situation in which both GC and BIR are possible, GC is by far the predominant outcome, although when

homology is removed from one end, BIR proves to be quite efficient (Malkova et al. 2005; Jain et al. 2009).

PERSPECTIVES

In this review, we have summarized the various sources of DSBs, pathways to repair them, and ways to monitor the intermediate steps of repair. Faithful and efficient repair of all DNA lesions is essential to maintain genome integrity and cell viability. Deficiencies and mutations in repair pathways are associated with multiple human diseases and with aging. It is curious to note that DSBs can be detrimental as well as beneficial to organisms. The molecular details connecting DSB repair to other cellular processes are emerging. One interesting aspect that is yet to be sufficiently explored is the role of chromatin structure in HR.

Most of our understanding of HR comes from work in bacteria, yeast, and other model systems. Comparable studies in mammalian systems are now beginning to elucidate the intricacies of DNA repair. The exceedingly more complex mammalian system is likely to require additional hitherto unknown components. Indeed, several proteins, including ssDNA-binding protein (SSB) (Li et al. 2009b) and RAD51-associated protein 1 (RAD51AP1) (Modesti et al. 2007; Wiese et al. 2007) have been identified that have no homologs in lower eukaryotes.

Any deficiency in HR directs the cells along more error-prone pathways like NHEJ. Ironically, recent studies have shown that HR itself is highly error-prone and could be one of the leading causes of mutagenesis (Malkova and Haber 2012). Thus, HR is a double-edged sword whose precise regulation is critical. HR components are targets for emerging therapies, especially in cancer therapeutics. Hence, further elucidation of the mechanisms of HR repair, especially in mammals, is vital.

ACKNOWLEDGMENTS

Work from the Haber laboratory is supported by NIH Grants GM20056, GM61766, and GM76020 to JEH.

REFERENCES

* Reference is also in this collection.

- * Aguilera A, Gaillard H. 2014. Transcription and recombination: When RNA meets DNA. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016543.
- Arcangioli B. 2000. Fate of *mat1* DNA strands during mating-type switching in fission yeast. *EMBO Rep* **1**: 145–150.
- Aylon Y, Liefshitz B, Kupiec M. 2004. The CDK regulates repair of double-strand breaks by homologous recombination during the cell cycle. *EMBO J* **23**: 4868–4875.
- Balestrini A, Ristic D, Dionne I, Liu XZ, Wyman C, Wellinger RJ, Petrini JH. 2013. The Ku heterodimer and the metabolism of single-ended DNA double-strand breaks. *Cell Rep* **3**: 2033–2045.
- Barlow JH, Lisby M, Rothstein R. 2008. Differential regulation of the cellular response to DNA double-strand breaks in G₁. *Mol Cell* **30**: 73–85.
- Barsoum E, Martinez P, Astrom SU. 2010. $\alpha 3$, a transposable element that promotes host sexual reproduction. *Genes Dev* **24**: 33–44.
- Baudry C, Malinsky S, Restituto M, Kapusta A, Rosa S, Meyer E, Betermier M. 2009. PiggyMac, a domesticated *piggyBac* transposase involved in programmed genome rearrangements in the ciliate *Paramecium tetraurelia*. *Genes Dev* **23**: 2478–2483.
- Baxter J, Diffley JE. 2008. Topoisomerase II inactivation prevents the completion of DNA replication in budding yeast. *Mol Cell* **30**: 790–802.
- Bennardo N, Cheng A, Huang N, Stark JM. 2008. Alternative-NHEJ is a mechanistically distinct pathway of mammalian chromosome break repair. *PLoS Genet* **4**: e1000110.
- Betermier M. 2004. Large-scale genome remodelling by the developmentally programmed elimination of germ line sequences in the ciliate *Paramecium*. *Res Microbiol* **155**: 399–408.
- Bosco G, Haber JE. 1998. Chromosome break-induced DNA replication leads to nonreciprocal translocations and telomere capture. *Genetics* **150**: 1037–1047.
- Bzymek M, Thayer NH, Oh SD, Kleckner N, Hunter N. 2010. Double Holliday junctions are intermediates of DNA break repair. *Nature* **464**: 937–941.
- Cejka P, Cannavo E, Polaczek P, Masuda-Sasa T, Pokharel S, Campbell JL, Kowalczykowski SC. 2010. DNA end resection by Dna2-Sgs1-RPA and its stimulation by Top3-Rmi1 and Mre11-Rad50-Xrs2. *Nature* **467**: 112–116.
- Chen J, Stubbe J. 2005. Bleomycins: Towards better therapeutics. *Nat Rev Cancer* **5**: 102–112.
- Chen Z, Yang H, Pavletich NP. 2008. Mechanism of homologous recombination from the RecA-ssDNA/dsDNA structures. *Nature* **453**: 484–489.
- Chiruvella KK, Liang Z, Wilson TE. 2013. Repair of double-strand breaks by end joining. *Cold Spring Harbor Perspect Biol* **5**: a012757.
- Chung WH, Zhu Z, Papusha A, Malkova A, Ira G. 2010. Defective resection at DNA double-strand breaks leads to de novo telomere formation and enhances gene targeting. *PLoS Genet* **6**: e1000948.
- Coïc E, Feldman T, Landman AS, Haber JE. 2008. Mechanisms of Rad52-independent spontaneous and UV-induced mitotic recombination in *Saccharomyces cerevisiae*. *Genetics* **179**: 199–211.
- Colleaux L, D'Auriol L, Galibert F, Dujon B. 1988. Recognition and cleavage site of the intron-encoded omega transposase. *Proc Natl Acad Sci* **85**: 6022–6026.
- Crasta K, Ganem NJ, Dagher R, Lantermann AB, Ivanova EV, Pan Y, Nezi L, Protopopov A, Chowdhury D, Pellman D. 2012. DNA breaks and chromosome pulverization from errors in mitosis. *Nature* **482**: 53–58.
- * Daley JM, Gaines WA, Kwon YH, Sung P. 2014. Regulation of DNA pairing in homologous recombination. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a017954.
- Davis AP, Symington LS. 2004. RAD51-dependent break-induced replication in yeast. *Mol Cell Biol* **24**: 2344–2351.
- De Bont R, van Larebeke N. 2004. Endogenous DNA damage in humans: A review of quantitative data. *Mutagenesis* **19**: 169–185.
- Decottignies A. 2013. Alternative end-joining mechanisms: A historical perspective. *Front Genet* **4**: 48.
- Deem A, Keszthelyi A, Blackgrove T, Vayl A, Coffey B, Mathur R, Chabes A, Malkova A. 2011. Break-induced replication is highly inaccurate. *PLoS Biol* **9**: e1000594.
- * Doksani Y, de Lange T. 2014. The role of DSB repair pathways at functional and dysfunctional telomeres. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016576.
- Donnianni RA, Symington LS. 2013. Break-induced replication occurs by conservative DNA synthesis. *Proc Natl Acad Sci* **110**: 13475–13480.
- Ferguson DO, Holloman WK. 1996. Recombinational repair of gaps in DNA is asymmetric in *Ustilago maydis* and can be explained by a migrating D-loop model. *Proc Natl Acad Sci* **93**: 5419–5424.
- Formosa T, Alberts BM. 1986. DNA synthesis dependent on genetic recombination: Characterization of a reaction catalyzed by purified bacteriophage T4 proteins. *Cell* **47**: 793–806.
- Ganem NJ, Pellman D. 2012. Linking abnormal mitosis to the acquisition of DNA damage. *J Cell Biol* **199**: 871–881.
- Gavalda S, Gallardo M, Luna R, Aguilera A. 2013. R-loop mediated transcription-associated recombination in *trf4Δ* mutants reveals new links between RNA surveillance and genome integrity. *PLoS ONE* **8**: e65541.
- Gloor GB, Nassif NA, Johnson-Schlitz DM, Preston CR, Engels WR. 1991. Targeted gene replacement in *Drosophila* via P element-induced gap repair. *Science* **253**: 1110–1117.
- Haaf T, Golub EI, Reddy G, Radding CM, Ward DC. 1995. Nuclear foci of mammalian Rad51 recombination protein in somatic cells after DNA damage and its localization in synaptonemal complexes. *Proc Natl Acad Sci* **92**: 2298–2302.
- Haber JE. 2006. Transpositions and translocations induced by site-specific double-strand breaks in budding yeast. *DNA Repair* **5**: 998–1009.
- Haber JE. 2012. Mating-type genes and MAT switching in *Saccharomyces cerevisiae*. *Genetics* **191**: 33–64.
- Haber JE, Thorburn PC. 1984. Healing of broken linear dicentric chromosomes in yeast. *Genetics* **106**: 207–226.



- Hanada K, Budzowska M, Davies SL, van Drunen E, Onizawa H, Beverloo HB, Maas A, Essers J, Hickson ID, Kanaar R. 2007. The structure-specific endonuclease Mus81 contributes to replication restart by generating double-strand DNA breaks. *Nat Struct Mol Biol* **14**: 1096–1104.
- Hashimoto Y, Puddu F, Costanzo V. 2012. RAD51- and MRE11-dependent reassembly of uncoupled CMG helicase complex at collapsed replication forks. *Nat Struct Mol Biol* **19**: 17–24.
- Helmrich A, Ballarino M, Tora L. 2011. Collisions between replication and transcription complexes cause common fragile site instability at the longest human genes. *Mol Cell* **44**: 966–977.
- Hicks WM, Kim M, Haber JE. 2010. Increased mutagenesis and unique mutation signature associated with mitotic gene conversion. *Science* **329**: 82–85.
- Hicks WM, Yamaguchi M, Haber JE. 2011. Real-time analysis of double-strand DNA break repair by homologous recombination. *Proc Natl Acad Sci* **108**: 3108–3115.
- Hoeijmakers JHJ. 2009. DNA damage, aging, and cancer. *N Engl J Med* **361**: 1475–1485.
- Hoffelder DR, Luo L, Burke NA, Watkins SC, Gollin SM, Saunders WS. 2004. Resolution of anaphase bridges in cancer cells. *Chromosoma* **112**: 389–397.
- Holloman WK. 2011. Unraveling the mechanism of BRCA2 in homologous recombination. *Nat Struct Mol Biol* **18**: 748–754.
- Huertas P, Aguilera A. 2003. Cotranscriptionally formed DNA:RNA hybrids mediate transcription elongation impairment and transcription-associated recombination. *Mol Cell* **12**: 711–721.
- Huertas P, Cortes-Ledesma F, Sartori AA, Aguilera A, Jackson SP. 2008. CDK targets Sae2 to control DNA-end resection and homologous recombination. *Nature* **455**: 689–692.
- Ira G, Malkova A, Liberi G, Foiani M, Haber JE. 2003. Srs2 and Sgs1-Top3 suppress crossovers during double-strand break repair in yeast. *Cell* **115**: 401–411.
- Ira G, Pelliccioli A, Balijja A, Wang X, Fiorani S, Carotenuto W, Liberi G, Bressan D, Wan L, Hollingsworth NM, et al. 2004. DNA end resection, homologous recombination and DNA damage checkpoint activation require CDK1. *Nature* **431**: 1011–1017.
- Ira G, Satory D, Haber JE. 2006. Conservative inheritance of newly synthesized DNA in double-strand break-induced gene conversion. *Mol Cell Biol* **26**: 9424–9429.
- Ivanov EL, Sugawara N, Fishman-Lobell J, Haber JE. 1996. Genetic requirements for the single-strand annealing pathway of double-strand break repair in *Saccharomyces cerevisiae*. *Genetics* **142**: 693–704.
- Jain S, Sugawara N, Lydeard J, Vaze M, Tanguy Le Gac N, Haber JE. 2009. A recombination execution checkpoint regulates the choice of homologous recombination pathway during DNA double-strand break repair. *Genes Dev* **23**: 291–303.
- Janssen A, van der Burg M, Szuhai K, Kops GJ, Medema RH. 2011. Chromosome segregation errors as a cause of DNA damage and structural chromosome aberrations. *Science* **333**: 1895–1898.
- Jensen RB, Carreira A, Kowalczykowski SC. 2010. Purified human BRCA2 stimulates RAD51-mediated recombination. *Nature* **467**: 678–683.
- Kapusta A, Matsuda A, Marmignon A, Ku M, Silve A, Meyer E, Forney JD, Malinsky S, Betermier M. 2011. Highly precise and developmentally programmed genome assembly in *Paramecium* requires ligase IV-dependent end joining. *PLoS Genet* **7**: e1002049.
- Kass EM, Jasin M. 2010. Collaboration and competition between DNA double-strand break repair pathways. *FEBS Lett* **584**: 3703–3708.
- Kawamoto T, Araki K, Sonoda E, Yamashita YM, Harada K, Kikuchi K, Masutani C, Hanaoka F, Nozaki K, Hashimoto N, et al. 2005. Dual roles for DNA polymerase η in homologous DNA recombination and translesion DNA synthesis. *Mol Cell* **20**: 793–799.
- Keeney S. 2008. Spo11 and the formation of DNA double-strand breaks in meiosis. *Genome Dyn Stab* **2**: 81–123.
- Kiiianitsa K, Solinger JA, Heyer WD. 2006. Terminal association of Rad54 protein with the Rad51-dsDNA filament. *Proc Natl Acad Sci* **103**: 9767–9772.
- Kim N, Jinks-Robertson S. 2009. dUTP incorporation into genomic DNA is linked to transcription in yeast. *Nature* **459**: 1150–1153.
- Klar AJ. 2007. Lessons learned from studies of fission yeast mating-type switching and silencing. *Annu Rev Genet* **41**: 213–236.
- Koster DA, Palle K, Bot ES, Bjornsti MA, Dekker NH. 2007. Antitumour drugs impede DNA uncoiling by topoisomerase I. *Nature* **448**: 213–217.
- Kramer KM, Brock JA, Bloom K, Moore JK, Haber JE. 1994. Two different types of double-strand breaks in *Saccharomyces cerevisiae* are repaired by similar RAD52-independent, nonhomologous recombination events. *Mol Cell Biol* **14**: 1293–1301.
- Krogh BO, Symington LS. 2004. Recombination proteins in yeast. *Annu Rev Genet* **38**: 233–271.
- * Lam I, Keeney S. 2014. Mechanism and regulation of meiotic recombination initiation. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016634.
- Li X, Manley JL. 2005. Inactivation of the SR protein splicing factor ASF/SF2 results in genomic instability. *Cell* **122**: 365–378.
- Li F, Dong J, Pan X, Oum JH, Boeke JD, Lee SE. 2008. Microarray-based genetic screen defines SAW1, a gene required for Rad1/Rad10-dependent processing of recombination intermediates. *Mol Cell* **30**: 325–335.
- Li X, Stith CM, Burgers PM, Heyer WD. 2009a. PCNA is required for initiation of recombination-associated DNA synthesis by DNA polymerase δ . *Mol Cell* **36**: 704–713.
- Li Y, Bolderson E, Kumar R, Muniandy PA, Xue Y, Richard DJ, Seidman M, Pandita TK, Khanna KK, Wang W. 2009b. hSSB1 and hSSB2 form similar multiprotein complexes that participate in DNA damage response. *J Biol Chem* **284**: 23525–23531.
- Li J, Coic E, Lee K, Lee CS, Kim JA, Wu Q, Haber JE. 2012. Regulation of budding yeast mating-type switching donor preference by the FHA domain of Fkh1. *PLoS Genet* **8**: e1002630.



- Lin Y, Dent SY, Wilson JH, Wells RD, Napierala M. 2010. R loops stimulate genetic instability of CTG.CAG repeats. *Proc Natl Acad Sci* **107**: 692–697.
- Lindahl T. 1993. Instability and decay of the primary structure of DNA. *Nature* **362**: 709–715.
- Lisby M, Mortensen UH, Rothstein R. 2003. Colocalization of multiple DNA double-strand breaks at a single Rad52 repair centre. *Nat Cell Biol* **5**: 572–577.
- Liu J, Doty T, Gibson B, Heyer WD. 2010. Human BRCA2 protein promotes RAD51 filament formation on RPA-covered single-stranded DNA. *Nat Struct Mol Biol* **17**: 1260–1262.
- Löbrich M, Shibata A, Beucher A, Fisher A, Ensminger M, Goodarzi AA, Barton O, Jeggo PA. 2010. γ H2AX foci analysis for monitoring DNA double-strand break repair: Strengths, limitations and optimization. *Cell Cycle* **9**: 662–669.
- Lydeard JR, Jain S, Yamaguchi M, Haber JE. 2007. Break-induced replication and telomerase-independent telomere maintenance require Pol32. *Nature* **448**: 820–823.
- Lydeard JR, Lipkin-Moore Z, Sheu YJ, Stillman B, Burgers PM, Haber JE. 2010. Break-induced replication requires all essential DNA replication factors except those specific for pre-RC assembly. *Genes Dev* **24**: 1133–1144.
- Ma W, Halweg CJ, Menendez D, Resnick MA. 2012. Differential effects of poly(ADP-ribose) polymerase inhibition on DNA break repair in human cells are revealed with Epstein-Barr virus. *Proc Natl Acad Sci* **109**: 6590–6595.
- Malkova A, Haber JE. 2012. Mutations arising during repair of chromosome breaks. *Annu Rev Genet* **46**: 455–473.
- Malkova A, Ivanov EL, Haber JE. 1996. Double-strand break repair in the absence of RAD51 in yeast: A possible role for break-induced DNA replication. *Proc Natl Acad Sci* **93**: 7131–7136.
- Malkova A, Naylor ML, Yamaguchi M, Ira G, Haber JE. 2005. RAD51-dependent break-induced replication differs in kinetics and checkpoint responses from RAD51-mediated gene conversion. *Mol Cell Biol* **25**: 933–944.
- Maloisel L, Bhargava J, Roeder GS. 2004. A role for DNA polymerase δ in gene conversion and crossing over during meiosis in *Saccharomyces cerevisiae*. *Genetics* **167**: 1133–1142.
- Maloisel L, Fabre F, Gangloff S. 2008. DNA polymerase δ is preferentially recruited during homologous recombination to promote heteroduplex DNA extension. *Mol Cell Biol* **28**: 1373–1382.
- Marrero VA, Symington LS. 2010. Extensive DNA end processing by exo1 and sgs1 inhibits break-induced replication. *PLoS Genet* **6**: e1001007.
- McEachern MJ, Haber JE. 2006. Break-induced replication and recombinational telomere elongation in yeast. *Annu Rev Biochem* **75**: 111–135.
- McIlwraith MJ, Vaisman A, Liu Y, Fanning E, Woodgate R, West SC. 2005. Human DNA polymerase η promotes DNA synthesis from strand invasion intermediates of homologous recombination. *Mol Cell* **20**: 783–792.
- McVey M, Lee SE. 2008. MMEJ repair of double-strand breaks (director's cut): Deleted sequences and alternative endings. *Trend Genet* **24**: 529–538.
- Merrikh H, Zhang Y, Grossman AD, Wang JD. 2012. Replication-transcription conflicts in bacteria. *Nat Rev Microbiol* **10**: 449–458.
- Milligan JR, Ng JY, Wu CC, Aguilera JA, Fahey RC, Ward JF. 1995. DNA repair by thiols in air shows two radicals make a double-strand break. *Radiat Res* **143**: 273–280.
- Mimitou EP, Symington LS. 2008. Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. *Nature* **455**: 770–774.
- Mirkin EV, Mirkin SM. 2007. Replication fork stalling at natural impediments. *Microbiol Mol Biol Rev* **71**: 13–35.
- Mitchel K, Lehner K, Jinks-Robertson S. 2013. Heteroduplex DNA position defines the roles of the Sgs1, Srs2, and Mph1 helicases in promoting distinct recombination outcomes. *PLoS Genet* **9**: e1003340.
- Mladenov E, Iliakis G. 2011. Induction and repair of DNA double strand breaks: The increasing spectrum of non-homologous end joining pathways. *Mutat Res* **711**: 61–72.
- Modesti M, Budzowska M, Baldeyron C, Demmers JA, Ghirlando R, Kanaar R. 2007. RAD51AP1 is a structure-specific DNA binding protein that stimulates joint molecule formation during RAD51-mediated homologous recombination. *Mol Cell* **28**: 468–481.
- Moore JK, Haber JE. 1996. Cell cycle and genetic requirements of two pathways of nonhomologous end-joining repair of double-strand breaks in *Saccharomyces cerevisiae*. *Mol Cell Biol* **16**: 2164–2173.
- * Morrical SW. 2014. DNA pairing and annealing processes in homologous recombination and homology-directed repair. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016444.
- Morrow DM, Connelly C, Hieter P. 1997. “Break copy” duplication: A model for chromosome fragment formation in *Saccharomyces cerevisiae*. *Genetics* **147**: 371–382.
- Nimonkar AV, Genschel J, Kinoshita E, Polaczek P, Campbell JL, Wyman C, Modrich P, Kowalczykowski SC. 2011. BLM-DNA2-RPA-MRN and EXO1-BLM-RPA-MRN constitute two DNA end resection machineries for human DNA break repair. *Genes Dev* **25**: 350–362.
- Niu H, Chung WH, Zhu Z, Kwon Y, Zhao W, Chi P, Prakash R, Seong C, Liu D, Lu L, et al. 2010. Mechanism of the ATP-dependent DNA end-resection machinery from *Saccharomyces cerevisiae*. *Nature* **467**: 108–111.
- Noon AT, Goodarzi AA. 2011. 53BP1-mediated DNA double strand break repair: Insert bad pun here. *DNA Repair* **10**: 1071–1076.
- Pâques F, Haber JE. 1999. Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* **63**: 349–404.
- Pâques F, Leung WY, Haber JE. 1998. Expansions and contractions in a tandem repeat induced by double-strand break repair. *Mol Cell Biol* **18**: 2045–2054.
- Paulsen RD, Soni DV, Wollman R, Hahn AT, Yee MC, Guan A, Hesley JA, Miller SC, Cromwell EF, Solow-Cordero DE, et al. 2009. A genome-wide siRNA screen reveals diverse cellular processes and pathways that mediate genome stability. *Mol Cell* **35**: 228–239.
- Petermann E, Helleday T. 2010. Pathways of mammalian replication fork restart. *Nat Rev Mol Cell Biol* **11**: 683–687.



- Plessis A, Perrin A, Haber JE, Dujon B. 1992. Site-specific recombination determined by I-SceI, a mitochondrial group I intron-encoded endonuclease expressed in the yeast nucleus. *Genetics* **130**: 451–460.
- Prado F, Aguilera A. 2005. Impairment of replication fork progression mediates RNA polII transcription-associated recombination. *EMBO J* **24**: 1267–1276.
- Prakash R, Satory D, Dray E, Papusha A, Scheller J, Kramer W, Krejci L, Klein H, Haber JE, Sung P, et al. 2009. Yeast Mph1 helicase dissociates Rad51-made D-loops: Implications for crossover control in mitotic recombination. *Genes Dev* **23**: 67–79.
- Prindle MJ, Loeb LA. 2012. DNA polymerase δ in DNA replication and genome maintenance. *Environ Mol Mutagen* **53**: 666–682.
- Quevedo O, Garcia-Luis J, Matos-Perdomo E, Aragon L, Machin F. 2012. Nondisjunction of a single chromosome leads to breakage and activation of DNA damage checkpoint in G₂. *PLoS Genet* **8**: e1002509.
- Raderschall E, Golub EI, Haaf T. 1999. Nuclear foci of mammalian recombination proteins are located at single-stranded DNA regions formed after DNA damage. *Proc Natl Acad Sci* **96**: 1921–1926.
- Rattray AJ, Shafer BK, McGill CB, Strathern JN. 2002. The roles of REV3 and RAD57 in double-strand-break-repair-induced mutagenesis of *Saccharomyces cerevisiae*. *Genetics* **162**: 1063–1077.
- Resnick MA. 1976. The repair of double-strand breaks in DNA; a model involving recombination. *J Theor Biol* **59**: 97–106.
- Rothstein RJ. 1983. One-step gene disruption in yeast. *Methods Enzymol* **101**: 202–211.
- Ruiz JF, Gomez-Gonzalez B, Aguilera A. 2009. Chromosomal translocations caused by either pol32-dependent or pol32-independent triparental break-induced replication. *Mol Cell Biol* **29**: 5441–5454.
- Saini N, Ramakrishnan S, Elango R, Ayyar S, Zhang Y, Deem A, Ira G, Haber JE, Lobachev KS, Malkova A. 2013. Migrating bubble during break-induced replication drives conservative DNA synthesis. *Nature* **502**: 389–392.
- Samoshkin A, Arnaoutov A, Jansen LE, Ouspenski I, Dye L, Karpova T, McNally J, Dasso M, Cleveland DW, Strunnikov A. 2009. Human condensin function is essential for centromeric chromatin assembly and proper sister kinetochore orientation. *PLoS ONE* **4**: e6831.
- Sartori AA, Lukas C, Coates J, Mistrik M, Fu S, Bartek J, Baer R, Lukas J, Jackson SP. 2007. Human CtIP promotes DNA end resection. *Nature* **450**: 509–514.
- Sebesta M, Burkovics P, Haracska L, Krejci L. 2011. Reconstitution of DNA repair synthesis in vitro and the role of polymerase and helicase activities. *DNA Repair* **10**: 567–576.
- Smith CE, Llorente B, Symington LS. 2007. Template switching during break-induced replication. *Nature* **447**: 102–105.
- Sneeden JL, Grossi SM, Tappin I, Hurwitz J, Heyer WD. 2013. Reconstitution of recombination-associated DNA synthesis with human proteins. *Nucleic Acids Res* **41**: 4913–4925.
- Sonoda E, Morrison C, Yamashita YM, Takata M, Takeda S. 2001. Reverse genetic studies of homologous DNA recombination using the chicken B-lymphocyte line, DT40. *Philos Trans R Soc Lond B Biol Sci* **356**: 111–117.
- Soulas-Sprauel P, Rivera-Munoz P, Malivert L, Le Guyader G, Abramowski V, Revy P, de Villartay JP. 2007. V(D)J and immunoglobulin class switch recombinations: A paradigm to study the regulation of DNA end-joining. *Oncogene* **26**: 7780–7791.
- Soutoglou E, Misteli T. 2008. Activation of the cellular DNA damage response in the absence of DNA lesions. *Science* **320**: 1507–1510.
- Spell RM, Holm C. 1994. Nature and distribution of chromosomal intertwinings in *Saccharomyces cerevisiae*. *Mol Cell Biol* **14**: 1465–1476.
- Stirling PC, Chan YA, Minaker SW, Aristizabal MJ, Barrett I, Sipahimalani P, Kobor MS, Hieter P. 2012. R-loop-mediated genome instability in mRNA cleavage and polyadenylation mutants. *Genes Dev* **26**: 163–175.
- Strathern JN, Shafer BK, McGill CB. 1995. DNA synthesis errors associated with double-strand-break repair. *Genetics* **140**: 965–972.
- Suberbielle E, Sanchez PE, Kravitz AV, Wang X, Ho K, Eilertson K, Devidze N, Kreitzer AC, Mucke L. 2013. Physiologic brain activity causes DNA double-strand breaks in neurons, with exacerbation by amyloid- β . *Nat Neurosci* **16**: 613–621.
- Sugawara N, Haber JE. 1992. Characterization of double-strand break-induced recombination: Homology requirements and single-stranded DNA formation. *Mol Cell Biol* **12**: 563–575.
- Sugawara N, Paques F, Colaiacovo M, Haber JE. 1997. Role of *Saccharomyces cerevisiae* Msh2 and Msh3 repair proteins in double-strand break-induced recombination. *Proc Natl Acad Sci* **94**: 9214–9219.
- Sugawara N, Ira G, Haber JE. 2000. DNA length dependence of the single-strand annealing pathway and the role of *Saccharomyces cerevisiae* RAD59 in double-strand break repair. *Mol Cell Biol* **20**: 5300–5309.
- Sugawara N, Wang X, Haber JE. 2003. In vivo roles of Rad52, Rad54, and Rad55 proteins in Rad51-mediated recombination. *Mol Cell Biol* **23**: 209–219.
- Sun J, Lee KJ, Davis AJ, Chen DJ. 2012. Human Ku70/80 protein blocks exonuclease 1-mediated DNA resection in the presence of human Mre11 or Mre11/Rad50 protein complex. *J Biol Chem* **287**: 4936–4945.
- * Syeda AH, Hawkins M, McGlynn P. 2014. Recombination and replication. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016550.
- * Symington LS. 2014. Processing of DNA breaks: Mechanism and regulation. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016536.
- Szostak JW, Orr-Weaver TL, Rothstein RJ, Stahl FW. 1983. The double-strand-break repair model for recombination. *Cell* **33**: 25–35.
- Thompson LH. 2012. Recognition, signaling, and repair of DNA double-strand breaks produced by ionizing radiation in mammalian cells: The molecular choreography. *Mutat Res* **751**: 158–246.
- Toh GW, Sugawara N, Dong J, Toth R, Lee SE, Haber JE, Rouse J. 2010. Mec1/Tel1-dependent phosphorylation of Slx4 stimulates Rad1-Rad10-dependent cleavage of non-homologous DNA tails. *DNA Repair* **9**: 718–726.



- Valdiglesias V, Giunta S, Fenech M, Neri M, Bonassi S. 2013. γ H2AX as a marker of DNA double strand breaks and genomic instability in human population studies. *Mutat Res* **753**: 24–40.
- Vilenchik MM, Knudson AG. 2003. Endogenous DNA double-strand breaks: Production, fidelity of repair, and induction of cancer. *Proc Natl Acad Sci* **100**: 12871–12876.
- Villarreal DD, Lee K, Deem A, Shim EY, Malkova A, Lee SE. 2012. Microhomology directs diverse DNA break repair pathways and chromosomal translocations. *PLoS Genet* **8**: e1003026.
- Wahba L, Amon JD, Koshland D, Vuica-Ross M. 2011. RNase H and multiple RNA biogenesis factors cooperate to prevent RNA:DNA hybrids from generating genome instability. *Mol Cell* **44**: 978–988.
- Wahba L, Gore SK, Koshland D. 2013. The homologous recombination machinery modulates the formation of RNA-DNA hybrids and associated chromosome instability. *eLife* **2**: e00505.
- Wang X, Haber JE. 2004. Role of *Saccharomyces* single-stranded DNA-binding protein RPA in the strand invasion step of double-strand break repair. *PLoS Biol* **2**: E21.
- Wang X, Ira G, Tercero JA, Holmes AM, Diffley JF, Haber JE. 2004. Role of DNA replication proteins in double-strand break-induced recombination in *Saccharomyces cerevisiae*. *Mol Cell Biol* **24**: 6891–6899.
- Ward JF. 1994. The complexity of DNA damage: Relevance to biological consequences. *Int J Radiat Biol* **66**: 427–432.
- Weinfeld M, Soderlind KJ. 1991. 32P-postlabeling detection of radiation-induced DNA damage: Identification and estimation of thymine glycols and phosphoglycolate termini. *Biochemistry* **30**: 1091–1097.
- White CI, Haber JE. 1990. Intermediates of recombination during mating type switching in *Saccharomyces cerevisiae*. *EMBO J* **9**: 663–673.
- Wiese C, Dray E, Groesser T, San Filippo J, Shi I, Collins DW, Tsai MS, Williams GJ, Rydberg B, Sung P, et al. 2007. Promotion of homologous recombination and genomic stability by RAD51AP1 via RAD51 recombinase enhancement. *Mol Cell* **28**: 482–490.
- Wilson MA, Kwon Y, Xu Y, Chung WH, Chi P, Niu H, Mayle R, Chen X, Malkova A, Sung P, et al. 2013. Pif1 helicase and Pol δ promote recombination-coupled DNA synthesis via bubble migration. *Nature* **502**: 393–396.
- Wu L, Hickson ID. 2003. The Bloom's syndrome helicase suppresses crossing over during homologous recombination. *Nature* **426**: 870–874.
- * Wyatt HDM, West SC. 2014. Holliday junction resolvases. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a016485.
- Wyman C, Kanaar R. 2006. DNA double-strand break repair: All's well that ends well. *Annu Rev Genet* **40**: 363–383.
- Wyrobek AJ, Schmid TE, Marchetti F. 2005. Relative susceptibilities of male germ cells to genetic defects induced by cancer chemotherapies. *J Natl Cancer Inst Monogr* **34**: 31–35.
- Yang H, Li Q, Fan J, Holloman WK, Pavletich NP. 2005. The BRCA2 homologue Brh2 nucleates RAD51 filament formation at a dsDNA–ssDNA junction. *Nature* **433**: 653–657.
- Yang Y, Sterling J, Storic F, Resnick MA, Gordenin DA. 2008. Hypermutability of damaged single-strand DNA formed at double-strand breaks and uncapped telomeres in yeast *Saccharomyces cerevisiae*. *PLoS Genet* **4**: e1000264.
- Zhu Z, Chung WH, Shim EY, Lee SE, Ira G. 2008. Sgs1 helicase and two nucleases Dna2 and Exo1 resect DNA double-strand break ends. *Cell* **134**: 981–994.