



Published in final edited form as:

*Am J Geriatr Psychiatry*. 2014 December ; 22(12): 1633–1641. doi:10.1016/j.jagp.2014.02.006.

## Association of Age of Depression Onset with Cognitive Functioning in Individuals with Late Life Depression and Executive Dysfunction

R. Scott Mackin<sup>1,2</sup>, J. Craig Nelson<sup>1</sup>, Kevin L Delucchi<sup>1</sup>, Patrick J Raue<sup>3</sup>, Derek D Satre<sup>1,4</sup>, Dimitris N Kiosses<sup>3</sup>, George S Alexopoulos<sup>3</sup>, and Patricia A Arean<sup>1</sup>

<sup>1</sup>University of California, San Francisco, Department of Psychiatry

<sup>2</sup>Center for Imaging of Neurodegenerative Disease, Veterans Administration Medical Center, San Francisco, CA, USA

<sup>3</sup>Weill Cornell Medical College, Department of Psychiatry

<sup>4</sup>Kaiser Permanente Division of Research

### Abstract

**Objective**—To compare patterns of cognitive performance in older adults with late-onset depression (LOD; ≥ 65 years of age) to that of older adults with early-onset depression (EOD; < 65 years).

**Method**—Participants were 171 adults aged 60 years or older with major depression and executive dysfunction who were participating in a randomized psychotherapy trial. Participants included 72 LOD and 99 EOD individuals. Cognitive performance on measures of verbal learning, memory, and executive functioning were evaluated. Demographic and clinical characteristics, severity of cerebrovascular risk factors, and disability ratings were also compared between groups.

**Results**—The LOD group was older, and had fewer previous episodes of depression, and lower severity of depression compared to EOD participants. The LOD group demonstrated poorer performance on measures of verbal learning,  $F(1,161) = 4.28, p = .04$  and memory  $F(1,160) = 4.65, p = .03$ , than the EOD group. Linear regression analysis demonstrated that LOD and fewer years of education were significant predictors of poorer verbal learning,  $F(7,114) = 6.25, p < 0.001$  and memory,  $F(7,113) = 7.24, p < .001$ . Performance on measures of executive functioning, severity of vascular risk factors, and disability ratings did not differ between the two groups.

**Conclusions**—In older adults with depression and executive dysfunction, LOD was associated with poorer performance on measures of verbal learning and memory. Aging related brain changes associated with LOD may play a more important role leading to dysfunction in these cognitive

---

Corresponding Author: R. Scott Mackin, PhD, UCSF Department of Psychiatry, 401 Parnassus Avenue, San Francisco, CA 94143, scottm@lppi.ucsf.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosures: No other authors report competing interests.

domains than a history of recurrent depressive episodes in older adults with a dysexecutive syndrome.

### Keywords

cognitive functioning; disability; late life depression; age of onset; late onset; memory; executive dysfunction; learning; vascular risk factors

## INTRODUCTION

Major depression in older adults, or Late Life Depression (LLD), is a prevalent and debilitating psychiatric disorder that is commonly linked to a wide range of cognitive dysfunctions (1). LLD is most often associated with deficits of executive functioning, i.e. the depression dysexecutive syndrome of late life (2, 3), which is often conceptualized as representing individuals with concurrent cerebrovascular disease (4). However, cognitive impairments in the domains of verbal learning and memory are also frequently reported (1, 5). Further, there is considerable heterogeneity in the types, severity, and overall number of cognitive impairments reported in LLD samples (1, 6–8). While depression severity contributes to cognitive dysfunction in LLD (9), it is insufficient to explain the range of cognitive dysfunction reported. As cognitive dysfunction in LLD contributes to poor treatment adherence (10), increased cost of mental health treatment (11), and increased burden of disability (12), elucidating the impact of other LLD characteristics on cognitive functioning may represent a significant avenue to improve treatment and outcomes in these patients.

Age of depression onset is one clinical characteristic that has the potential to clarify the etiology of cognitive dysfunction in LLD. Two alternative, but not mutually exclusive, hypotheses have been suggested to explain cognitive dysfunction associated with LLD with respect of age of depression onset. Late onset depression (LOD), typically defined as onset of a first depressive episode after 65 years of age, occurs in approximately 30% of LLD individuals (13) and is often thought to be associated with aging related brain changes (14). This association is supported by consistent findings of cerebrovascular disease, specifically frontal striatal white matter abnormalities and executive dysfunction in LLD (15, 16). LOD has also been associated with temporal lobe atrophy and hippocampal volume loss, (17, 18) which are characteristics of incipient Alzheimer's disease and are associated with memory impairment (19, 20). Together these findings suggest that LOD is associated with aging related brain changes that could adversely impact cognition. The alternate hypothesis suggests that cognitive dysfunction could be the result of damage suffered from repeated lifetime episodes of depression in those with EOD. Specifically, it has been suggested that repeated episodes of depression and associated hypercortisolemia (21, 22) and other factors could result in hippocampal damage and cortical abnormalities (23, 24) that contribute to learning and memory deficits in LLD. Complicating these hypotheses is the potential for cognition in individuals with EOD to be further impacted by neurodegenerative disease in later life. Despite the importance of clarifying the contributing factors to cognitive dysfunction in LLD relatively few studies have been conducted to evaluate these alternative hypotheses.

While early studies initially reported similar patterns of cognitive functioning in EOD and LOD (8), four recent studies reported that LOD was associated with poorer memory performance when compared to EOD (25–28). However, Rapp et al (29) did not find this association. Of note, there have been no previous studies which have evaluated verbal learning, a cognitive domain associated with memory performance, in LOD in contrast to EOD. Poorer performance on measures of executive functioning and information processing speed in LOD relative to EOD have also been reported in some studies (27, 30), but not others (5). These inconsistent findings may result from several factors. Many of these studies utilized small samples of LLD participants. Additionally, few studies controlled for the impact of depression severity on cognitive functioning and relied on dichotomous classifications of age of onset without accounting for the number of prior depressive episodes. Lastly, several previous studies have not controlled for the impact of education history on cognitive performance in LOD and EOD individuals (25, 26, 31). Given the potential for level of previous education, or “cognitive reserve”, to impact cognitive performance in older adults (32), such investigations are particularly important.

This analysis evaluated cognitive functioning in LOD relative to EOD in a sample of older adults participating in a clinical trial of the efficacy of psychotherapy for patients with late life major depression and executive dysfunction. Evaluating the impact of age of depression onset on cognition in this sample, presumably at higher risk for aging related brain changes, offers a significant opportunity to clarify the role of age of depression onset on cognition. Based on the existing literature, we hypothesized that LOD participants would exhibit poorer performance on measures of learning and memory than EOD participants, and that LOD would be a stronger predictor of poor performance on measures of learning and memory than number of prior episodes after taking education level and current depression severity into account. We also examined if LOD was associated with greater burden of cerebrovascular risk factors when compared to EOD and if LOD patients differed from EOD patients on measures of disability status and executive functioning. To our knowledge, this is the first study to examine the effects of age of onset on cognitive functioning in older adults with depression and executive dysfunction.

## METHOD

### Participants

Data for this study were obtained for secondary analysis from a two-site, randomized controlled trial conducted to compare the efficacy of a modified version of Problem Solving Therapy (PST) to Supportive Therapy (ST) in patients ages 60 and over with major depression and executive dysfunction (33). Participants were recruited for participation in this study by community advertisements and referrals from psychiatry clinics at each study site. All participants provided informed consent to participate in the study and all study procedures were approved by a committee for human research institutional review board at each site. Psychiatric diagnoses were made by licensed psychologists utilizing DSM-IV criteria and the Structured Clinical Interview for the Diagnosis of DSM-IV Disorders, and were reviewed at a consensus conference comprised of psychologists, psychiatrists, and social workers. Depression severity at intake was evaluated with the 24-item Hamilton

Depression Rating Scale (HDRS) (34) and participants with moderate to severe depression severity (HDRS score of  $\geq 20$ ) were included in the study. Inclusion criteria with regard to executive functioning were based on screening performance on the Initiation/Perseveration index of the Mattis Dementia Rating Scale (DRS-IP; score  $\geq 33$ ) and the Stroop Word Color Test (SCWT; score  $\geq 25$ ) (33). Exclusion criteria consisted of current involvement in psychotherapy, currently on antidepressant medications or wish to be treated with antidepressants, presence of psychotic depression (SCID-R), high suicide risk (i.e., intent or plan to attempt suicide in near future), any Axis I psychiatric disorder or substance abuse other than unipolar major depression or generalized anxiety disorder, antisocial personality (DSM-IV), history of head trauma, dementia (MMSE $<24$  or diagnosis of dementia by DSM-IV), acute or severe medical illness (i.e., delirium, metastatic cancer, decompensated cardiac, liver or kidney failure, major surgery, stroke or myocardial infarction during the three months prior to entry), drugs known to cause depression (e.g., reserpine, alpha-methyl-dopa, steroids), and inability to perform any activities of daily living even with assistance.

Of the 653 participants referred to the study, 29 either refused to be screened after explanation of the study or did not complete the screen. The remaining 624 underwent a structured interview, which led to exclusion of 345 individuals: 26% did not meet criteria for major depression; 12% did not have a HDRS score  $\geq 20$ ; 40% did not exhibit evidence of executive dysfunction, 1% had low MMSE scores, and 13% had another psychiatric diagnosis. Of the 279 who were eligible for the study, 58 (21%) were not included because they failed to complete baseline evaluation and did not return for further assessment or treatment. Of the 221 participants enrolled in the treatment study, age of depression onset was available for 171 participants.

## Measures

All measures of depression severity, vascular risk factors, disability, and cognitive functioning were obtained for each participant during a single assessment. Late onset of depression (LOD) was diagnosed for individuals who reported a first episode of major depression meeting DSM IV criteria at 65 years of age or older. The specific measures utilized and outcome variables for each of these measures are described below. Additionally, demographic information (age, years of education, gender) and clinical history (age of depression onset, number of lifetime depressive episodes) was obtained for each participant.

## VERBAL LEARNING AND MEMORY

### Hopkins Verbal Learning Test - Revised (HVLT-R) (35)

The HVLT-R is a measure of verbal learning and memory for lists of verbally presented information. The outcome variable utilized for verbal learning performance was the total number of correct responses on the three learning trials of this test (HVLT-L). The total possible score on this measure was 36 points. The outcome variable utilized to evaluate memory (HVLT-M) was the total number of correct responses on the delayed free recall trial of this test and the total possible score was 12.

## EXECUTIVE FUNCTIONING

### **Mattis Dementia Rating Scale-2 Initiation/Perseveration Scale (DRS IP) (36)**

The DRS-IP is a measure of executive functioning for older adults with an information processing speed component; the total number of correct responses was utilized as the primary outcome variable. The total possible score on this measure was 37.

### **Wisconsin Card Sorting Test-64 Computer Version 2 (WCST-64) (37)**

The WCST is a non-speeded measure of problem solving ability, cognitive flexibility, and ability to maintain a cognitive set; the outcome variable was the total number of correct responses. The total possible score on this measure was 64.

### **The Stroop Color and Word Test (SCWT) (38)**

The SCWT is a timed measure of response inhibition, ability to maintain cognitive set, and information processing speed; the total number of correct responses on the color word trial was utilized as the outcome variable.

### **Trail Making Test Part B (TMT) (39)**

Part B of the TMT is a timed measure of sequencing ability, visuospatial speed, and response inhibition. The time required to complete both trials was utilized as the outcome variables for this test.

## DEPRESSION, VASCULAR RISK, DISABILITY

### **Hamilton Depression Rating Scale (34)**

The 24-item HDRS assesses severity of depressive symptoms; high scores indicate greater severity of depression and the total possible score was 75.

Vascular Risk Scale (VRS): the VRS consists of 8 questions evaluating the presence of antihypertensive treatment, history of diabetes, smoking, congestive heart failure/myocardial infarction, atrial fibrillation, ventricular hypertrophy, stroke, and transient ischemic attacks. Scores ranged from 0–8 with high scores representing greater vascular risk.

### **World Health Organization Disability Assessment Scale (WHODAS II) (40)**

The WHODAS II is a 36 item measure of disability and scores range from 0 to 100; higher scores represent greater disability.

### **Data Analysis**

To evaluate clinical characteristics and cognitive functioning between the two groups analysis of variance models were estimated and tested. Evaluation of the assumption of normalcy for dependent measures was conducted and we did not find violations of this assumption. For low frequency count variables we retrospectively utilized non-parametric and our results did not change. Covariates analysis of variance models included age and education. Linear regression models were used to model factors associated with cognitive performance on measures differentiating the two groups. These factors included the effect of

age, education, depression severity, number of previous depressive episodes, vascular risk factors, and age of depression onset (EOD vs LOD). Given correlations among cognitive functioning variables colinearity thresholds were also evaluated for each model. Analyses were conducted using SPSS (version 20).

## RESULTS

The mean age for the sample was 73.0 years ( $SD = 7.7$ ), the mean years of education was 15.2 ( $SD = 2.8$ ), the mean MMSE score was 27.8 ( $SD = 1.7$ ), and 65% of the sample was female. The mean HDRS for the sample was 24.0 ( $SD=4.4$ ), the average age of onset of depression was 55.9 years ( $SD=22.4$ ), the mean number of previous episodes of depression was 2.3 ( $SD = 2.2$ , median = 2.0). The sample demonstrated relatively low ratings of vascular risk (mean = 0.90,  $SD = 1.1$ , median = 1.0) and moderate ratings of disability (WHODAS II mean =26.6,  $SD = 7.3$ ). A high degree of correlation was found between measures of learning and memory and moderate correlations with these measures was seen with measures of executive functioning (Table 1). Forty-two percent of the sample reported late onset of depression (LOD; age of major depression onset  $\geq 65$  years) and age of onset groups did not differ with respect to gender ( $\chi^2=0.32$ ,  $p=0.57$ ).

With respect to group comparisons on clinical characteristics, the LOD group was significantly older,  $F(1,170) = 41.0$ ,  $p < .001$  and had lower depression severity ratings than the EOD group,  $F(1,170) = 6.4$ ,  $p = .01$  (Table 2). The LOD group also reported fewer previous episodes of depression,  $F(1,170) = 94.4$ ,  $p < .001$  but the two groups did not differ with respect to ratings of vascular risk or disability. With respect to cognitive performance (Table 3), after controlling for the effects of age and education, the LOD group performed significantly worse on measures of verbal learning  $F(1,162) = 4.3$ ,  $p = .04$  and memory  $F(1,162) = 4.7$ ,  $p = .03$ . The two groups did not differ with respect to performance on measures of executive functioning. Linear regression analyses demonstrated that LOD and education were the only significant predictors of verbal learning,  $F(7,114) = 6.25$ ,  $p < 0.001$ ; and memory,  $F(7,113)=7.24$ ,  $p < .001$  in the sample (Tables 4,5) accounting for 4% and 5% of the total variance of these models respectively. In separate analyses EOD/LOD was replaced with age of depression onset and age of onset was not a significant predictor of either learning or memory performance.

## DISCUSSION

Our study has two primary findings: 1) in patients with LLD and executive dysfunction, LOD was associated with poorer performance on measures of both verbal learning and memory when compared to EOD participants, 2) number of previous depressive episodes, current depression severity, and age of depression onset as a continuous measure were not associated with learning and memory performance. We also found that education was significantly associated with learning and memory performance and that the LOD and EOD depression groups did not differ with respect to ratings of vascular risk, disability, or performance on measures of executive functioning.



Our finding that LOD was associated with poorer performance on measures of memory is consistent with previous studies (25–28). However, our findings are important as our data suggest that LOD, i.e. onset on or after 65 years of age, is more strongly associated with memory performance than the specific age of first depressive episode, the number of lifetime depressive episodes, or depression severity in this sample. While we acknowledge a modest effect size in our statistical models, these results suggest that late onset of depression may play a more significant role in the development of memory impairment than the repeated insults associated with recurrent depressive episodes. Further, as memory disturbance has been shown to impact medical treatment adherence (10), our findings suggest that memory dysfunction in LOD may also impact both pharmacologic and psychotherapy treatment outcomes in LLD.

Our findings suggest that in addition to memory performance, LOD is also associated with poorer performance on measures of verbal learning. These findings represent an important extension of the prior literature which has primarily focused on memory performance (25–28). Specifically, verbal learning impairments may have important clinical implications particularly for psychotherapy interventions given the inherent learning requirements of these treatments for LLD. Given consistent findings that education level or “cognitive reserve” can impact cognitive performance in older adults (32), we statistically controlled for the impact of both age and education when performing our group comparisons. Our results, therefore, represent a more conservative statistical approach to evaluate differences in cognitive performance between the two depression onset groups across several cognitive domains than has been previously reported. Similarly, in contrast to previous studies (9), in our sample depression severity was not associated with measures of memory and learning. These results suggest that age of depression onset may be an important factor associated with verbal learning and memory dysfunction in older adults with LLD and concurrent executive dysfunction.

This is the first report of the influence of age of onset on cognition among older depressed patients presenting with executive dysfunction. In the current sample all participants were required to exhibit evidence of executive dysfunction, thus we likely limited our ability to detect group differences in this cognitive domain. Previous studies of patients with major depression not selected for executive dysfunction have demonstrated poorer performance on measures of executive functioning in LOD relative to EOD (27) (30). Further, the two depression groups in our sample did not differ with respect to burden of vascular risk factors which is consistent with a number of previous studies (13) (41–43). These findings are, however, relatively inconsistent with previous studies documenting a greater burden of MRI abnormalities associated with cerebrovascular disease in LOD (15, 44). Our findings provide further evidence to suggest that vascular risk measures similar to those utilized in our study may not be good indicators of MRI abnormalities associated with cerebrovascular disease in LLD. Also of interest, ratings of disability severity did not differ between depression groups suggesting that despite poorer cognitive performance, increased disability was not characteristic of LOD with executive dysfunction.

The degree to which verbal learning and memory functioning may represent a phenotypic marker of specific underlying neurodegenerative disease(s) in LOD is not well understood.

There is relatively consistent evidence supporting the impact of cerebrovascular disease on verbal learning and memory performance in LOD and non-depressed adults (45, 46). As such, cerebrovascular disease represents one possible etiology for the cognitive patterns we observed in LOD with executive dysfunction despite a lack of difference between the two groups on ratings of vascular risk factors. Additionally, reports linking LOD to hippocampal volume loss (17), and studies linking hippocampal atrophy to memory performance in LOD (42), suggest that our findings could be also interpreted as suggesting that LOD may be associated with incipient Alzheimer's disease. Our results did not support the role of repeated episodes of major depression as being a primary factor in cognitive dysfunction in this sample. Additional research will be necessary to clarify the etiology of memory and learning dysfunction in LOD.

Our study has several significant strengths including a relatively large sample of older adults with major depression and measures of cognitive functioning in executive functioning, verbal learning, and memory. Further, our statistical approach to control for the effects of education on cognitive performance and our evaluation of depression characteristics as predictors of cognitive functioning are relatively unique. Our study also has limitations. Our measure of vascular risk was not comprehensive, and as a result, the lack of difference on this measure between the two depression groups could reflect poor sensitivity of the measure in identifying cerebrovascular disease. Similarly, our measure of previous depressive episodes did not specify duration of lifetime depression which could be an important consideration for future studies. Additionally, the reliability and validity of self-reported history of age of depression onset, while commonly utilized (8), should be clarified in future studies. Further, our sample was highly selected and therefore our findings may not generalize to the larger population. Finally, nearly all of the cognitive tests we utilized were inter-correlated with each other. While these associations are not unexpected, we do acknowledge that our delineation of cognitive domains based on individual cognitive tests may have limited our ability to recognize broader patterns of differential performance between the two depression onset groups using multivariate analyses.

## Conclusions

Among older patients with major depression and executive dysfunction, late onset of depression was associated with poorer memory and verbal learning performance when compared to individuals with early onset of depression, while number of prior depressive episodes and current depression severity was not associated with cognition. Further, in our sample education level was significantly associated with cognitive performance in these domains. The implications of poor memory and learning performance in LOD on treatment outcomes are yet to be demonstrated. However these cognitive characteristics of LOD may be important factors influencing psychotherapy outcomes as psychotherapy interventions require patients to learn and remember information. Similarly these cognitive characteristics may also influence pharmacological interventions given studies documenting the impact of cognitive dysfunction on treatment adherence.



## Acknowledgments

Grant support: NIMH grants K08 MH081065, R01 MH063982, R01 0977669, R01 MH064099, P30 MH068638; UCSF Leon J. Epstein Endowment and the Sanchez Foundation.

Dr. Alexopoulos received grant support from Forest and has been a member of speakers' bureaus of Forest, Lilly, Merck, Sunovion, and Novartis.

## REFERENCES

1. Butters MA, Whyte EM, Nebes RD, et al. The nature and determinants of neuropsychological functioning in late-life depression. *Arch Gen Psychiatry*. 2004; 61:587–595. [PubMed: 15184238]
2. Alexopoulos GS, Raue PJ, Kanellopoulos D, et al. Problem solving therapy for the depression-executive dysfunction syndrome of late life. *Int J Geriatr Psychiatry*. 2008; 23:782–788. [PubMed: 18213605]
3. Alexopoulos GS, Kiosses DN, Klimstra S, et al. Clinical presentation of the "depression-executive dysfunction syndrome" of late life. *Am J Geriatr Psychiatry*. 2002; 10:98–106. [PubMed: 11790640]
4. Alexopoulos GS, Meyers BS, Young RC, et al. 'Vascular depression' hypothesis. *Arch Gen Psychiatry*. 1997; 54:915–922. [PubMed: 9337771]
5. Sheline YI, Barch DM, Garcia K, et al. Cognitive function in late life depression: relationships to depression severity, cerebrovascular risk factors and processing speed. *Biol Psychiatry*. 2006; 60:58–65. [PubMed: 16414031]
6. Fischer C, Schweizer TA, Atkins JH, et al. Neurocognitive profiles in older adults with and without major depression. *Int J Geriatr Psychiatry*. 2008; 23:851–856. [PubMed: 18311855]
7. Thomas AJ, Gallagher P, Robinson LJ, et al. A comparison of neurocognitive impairment in younger and older adults with major depression. *Psychol Med*. 2009; 39:725–733. [PubMed: 18667097]
8. Sheline YI, Pieper CF, Barch DM, et al. Support for the vascular depression hypothesis in late-life depression: results of a 2-site, prospective, antidepressant treatment trial. *Arch Gen Psychiatry*. 2010; 67:277–285. [PubMed: 20194828]
9. Culang ME, Sneed JR, Keilp JG, et al. Change in cognitive functioning following acute antidepressant treatment in late-life depression. *Am J Geriatr Psychiatry*. 2009; 17:881–888. [PubMed: 19916207]
10. Mackin RS, Arean PA. Cognitive and psychiatric predictors of medical treatment adherence among older adults in primary care clinics. *Int J Geriatr Psychiatry*. 2007; 22:55–60. [PubMed: 17006873]
11. Mackin RS, Delucchi KL, Bennett RW, et al. The Effect of Cognitive Impairment on Mental Healthcare Costs for Individuals With Severe Psychiatric Illness. *Am J Geriatr Psychiatry*. 2011; 19:176–184. [PubMed: 20808129]
12. Kiosses DN, Klimstra S, Murphy C, et al. Executive dysfunction and disability in elderly patients with major depression. *Am J Geriatr Psychiatry*. 2001; 9:269–274. [PubMed: 11481135]
13. Janssen J, Beekman AT, Comijs HC, et al. Late-life depression: the differences between early- and late-onset illness in a community-based sample. *Int J Geriatr Psychiatry*. 2006; 21:86–93. [PubMed: 16323255]
14. Butters MA, Young JB, Lopez O, et al. Pathways linking late-life depression to persistent cognitive impairment and dementia. *Dialogues Clin Neurosci*. 2008; 10:345–357. [PubMed: 18979948]
15. Janssen J, Hulshoff Pol HE, de Leeuw FE, et al. Hippocampal volume and subcortical white matter lesions in late life depression: comparison of early and late onset depression. *J Neurol Neurosurg Psychiatry*. 2007; 78:638–640. [PubMed: 17210630]
16. Alexopoulos GS, Kiosses DN, Choi SJ, et al. Frontal white matter microstructure and treatment response of late-life depression: a preliminary study. *Am J Psychiatry*. 2002; 159:1929–1932. [PubMed: 12411231]
17. Ballmaier M, Narr KL, Toga AW, et al. Hippocampal morphology and distinguishing late-onset from early-onset elderly depression. *Am J Psychiatry*. 2008; 165:229–237. [PubMed: 17986679]

18. Taylor WD, McQuoid DR, Krishnan KR. Medical comorbidity in late-life depression. *Int J Geriatr Psychiatry*. 2004; 19:935–943. [PubMed: 15449369]
19. Teipel SJ, Pruessner JC, Faltraco F, et al. Comprehensive dissection of the medial temporal lobe in AD: measurement of hippocampus, amygdala, entorhinal, perirhinal and parahippocampal cortices using MRI. *J Neurol*. 2006; 253:794–800. [PubMed: 16511646]
20. Mungas D, Reed BR, Farias ST, et al. Age and education effects on relationships of cognitive test scores with brain structure in demographically diverse older persons. *Psychol Aging*. 2009; 24:116–128. [PubMed: 19290743]
21. Sapolsky RM. The possibility of neurotoxicity in the hippocampus in major depression: a primer on neuron death. *Biol Psychiatry*. 2000; 48:755–765. [PubMed: 11063972]
22. Bos EH, Bouhuys AL, Geerts E, et al. Cognitive, physiological, and personality correlates of recurrence of depression. *J Affect Disord*. 2005; 87:221–229. [PubMed: 15979155]
23. Steffens DC, McQuoid DR, Payne ME, et al. Change in hippocampal volume on magnetic resonance imaging and cognitive decline among older depressed and nondepressed subjects in the neurocognitive outcomes of depression in the elderly study. *Am J Geriatr Psychiatry*. 2011; 9:4–12. [PubMed: 20808107]
24. Sheline YI, Gado MH, Kraemer HC. Untreated depression and hippocampal volume loss. *Am J Psychiatry*. 2003; 160:1516–1518. [PubMed: 12900317]
25. Dillon C, Allegri RF, Serrano CM, et al. Late- versus early-onset geriatric depression in a memory research center. *Neuropsychiatr Dis Treat*. 2009; 5:517–526. [PubMed: 19851519]
26. Kohler S, Thomas AJ, Barnett NA, et al. The pattern and course of cognitive impairment in late-life depression. *Psychol Med*. 2010; 40:591–602. [PubMed: 19656429]
27. Salloway S, Malloy P, Kohn R, et al. MRI and neuropsychological differences in early- and late-life-onset geriatric depression. *Neurology*. 1996; 46:1567–1574. [PubMed: 8649550]
28. Mossaheb N, Weissgram S, Zehetmayer S, et al. Late-onset depression in elderly subjects from the Vienna Transdanube Aging (VITA) study. *J Clin Psychiatry*. 2009; 70:500–508. [PubMed: 19358786]
29. Rapp MA, Schnaider Beeri M, Schmeidler J, et al. Relationship of neuropsychological performance to functional status in nursing home residents and community-dwelling older adults. *Am J Geriatr Psychiatry*. 2005; 13:450–459. [PubMed: 15956264]
30. Lesser IM, Boone KB, Mehringer CM, et al. Cognition and white matter hyperintensities in older depressed patients. *Am J Psychiatry*. 1996; 153:1280–1287. [PubMed: 8831435]
31. Rapp MA, Dahlman K, Sano M, et al. Neuropsychological differences between late-onset and recurrent geriatric major depression. *Am J Psychiatry*. 2005; 162:691–698. [PubMed: 15800140]
32. Hall CB, Derby C, LeValley A, et al. Education delays accelerated decline on a memory test in persons who develop dementia. *Neurology*. 2007; 69:1657–1664. [PubMed: 17954781]
33. Arean PA, Raue P, Mackin RS, et al. Problem-solving therapy and supportive therapy in older adults with major depression and executive dysfunction. *Am J Psychiatry*. 2010; 167:1391–1398. [PubMed: 20516155]
34. Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry*. 1960; 23:56–62. [PubMed: 14399272]
35. Brandt, J.; Benedict, RHB. *Hopkins Verbal Learning Test - Revised*. Lutz, FL: Psychological Assessment Resources, Inc.; 2001.
36. Mattis, S. *Dementia Rating Scale-2: Professional Manual*. Odessa, FL: Psychological Assessment Resources; 2004.
37. Heaton, R. *Wisconsin Card Sorting Test: Computer Version 4*. Odessa: Psychological Assessment Resources; 2000.
38. Golden, C.; Freshwater, SM. *Stroop Color and Word Test*. Wood Dale: Stoelting; 2002.
39. Reitan, RM.; Wolfson, D. *The Halstead-Reitan Neuropsychological Test Battery: Theory and Clinical Interpretation*. 2nd Tucson: Neuropsychology Press; 1993.
40. World Health Organization; 2001. *World Health Organization Disability Assessment Schedule II (WHODAS II)*.

41. Baldwin R, Jeffries S, Jackson A, et al. Neurological findings in late-onset depressive disorder: comparison of individuals with and without depression. *Br J Psychiatry*. 2005; 186:308–313. [PubMed: 15802687]
42. Hickie I, Naismith S, Ward PB, et al. Reduced hippocampal volumes and memory loss in patients with early- and late-onset depression. *Br J Psychiatry*. 2005; 186:197–202. [PubMed: 15738499]
43. Holroyd S, Duryee JJ. Differences in geriatric psychiatry outpatients with early- vs late-onset depression. *Int J Geriatr Psychiatry*. 1997; 12:1100–1106. [PubMed: 9427094]
44. Nebes RD, Reynolds CF 3rd, Boada F, et al. Longitudinal increase in the volume of white matter hyperintensities in late-onset depression. *Int J Geriatr Psychiatry*. 2002; 17:526–530. [PubMed: 12112176]
45. Kohler S, Thomas AJ, Lloyd A, et al. White matter hyperintensities, cortisol levels, brain atrophy and continuing cognitive deficits in late-life depression. *Br J Psychiatry*. 196:143–149. [PubMed: 20118461]
46. Duzel S, Schutze H, Stallforth S, et al. A close relationship between verbal memory and SN/VTA integrity in young and older adults. *Neuropsychologia*. 2008; 46:3042–3052. [PubMed: 18601938]

**Table 1**

Pearson Correlations Coefficients Between Cognitive Tests (n=171)

	HVLT-L	HVLT-M	DRS-IP	TMT B	WCST	SCWT
HVLT-L	1.00					
HVLT-M	.734**	1.00				
DRS-IP	.404**	.277**	1.00			
TMT B	-.337**	-.283**	-.245**	1.00		
WCST	.295**	.314**	.108	-.243*	1.00	
SCWT	.270**	.202**	.007	-.189*	.228*	1.00

SCWT = Stroop Color Word Test, DRS-IP = Dementia Rating Scale – Initiation/Perseveration, WCST = Wisconsin Card Sort Test, TMT B= Trail Making Test Part B, HVLT-L = Hopkins Verbal Learning Test-Learning, HVLT-M = Hopkins Verbal Learning Test - Memory

\*  $p < .05$

\*\*  $p < .01$

**Table 2**  
Demographic and Clinical Characteristics of Early and Late Onset Depression groups (n=171)

	Early Onset		Late Onset		Test	p
	n	M (SD)	n	M (SD)		
Age	99	70.2 (8.0)	72	77.1 (5.0)	F(1,170) = 41.0	< 0.001
Education	96	15.7 (2.8)	71	14.9 (2.7)	F(1,166) = 3.0	0.09
HDRS	99	24.7 (4.8)	72	23.0 (3.3)	F(1,170) = 6.4	<b>0.01</b>
Age at first MDD Episode	85	39.7 (18.1)	72	75.0 (5.4)	F(1,156) = 254.0	< <b>0.001</b>
Number of MDD Episodes	89	2.6 (1.4)	71	1.3 (0.8)	F(1,159) = 94.4	< <b>0.001</b>
CMCI	43	38.7 (3.6)	31	38.2 (2.8)	F(1,111) = 0.35	0.55
Vascular risk	68	0.88 (1.2)	56	0.91 (.98)	F(1,124) = 0.02	0.88
WHODAS	94	26.2 (7.3)	66	27.7 (7.4)	F(1,159) = 1.4	0.23

HDRS = Hamilton Depression Rating Scale, CMCI = Charlson Medical Comorbidity Index, WHODAS = World Health Organization Disability Assessment Scale, MDD=Major depression

**Table 3**  
Cognitive Performance for Early and Late Onset Depression Groups (n=171)

	Early Onset (n=99)		Late Onset (n=72)		Test	p
	N	M (SD)	N	M (SD)		
MMSE	95	27.7 (1.6)	70	27.8 (1.7)	F(1,164) = 0.02	0.89
SCWT	95	22.7 (7.4)	71	21.1 (9.1)	F(1,165) = 1.32	0.25
DRS IP	96	32.6 (3.5)	70	32.0 (3.8)	F(1,165) = 1.04	0.30
WCST	94	26.2 (11.1)	66	27.7 (10.9)	F(1,159) = 1.44	0.23
TMT B	87	127.6 (54.7)	57	148.5 (68.0)	F(1,143) = 3.36	0.07
HVLT L	94	21.9 (5.3)	68	19.9 (5.7)	F(1,161) = 4.28	<b>0.04</b>
HVLT M	94	7.5 (3.6)	67	6.2 (3.0)	F(1,160) = 4.65	<b>0.03</b>

MMSE = Mini Mental State Exam, SCWT = Stroop Color Word Test, DRS-IP = Dementia Rating Scale – Initiation/Perseveration, WCST = Wisconsin Card Sort Test, TMT B= Trail Making Test Part B, HVLT-L = Hopkins Verbal Learning Test-Learning, HVLT-M = Hopkins Verbal Learning Test – Memory.

Note: F Statistic corresponds with the group effect of age of depression onset



**Table 4**  
 Summary of Multivariate Regression Analysis for Prediction of Memory Performance in LLD Participants (n=160)

Variable	B	Std. Error	Std. Beta	Test	p
Age	-.010	.037	-.026	t(159) = -0.279	.781
Education	.533	.100	.450	t(158) = 5.317	<.001
HDRS	-.013	.072	-.016	t(159) = -0.185	.854
Number of previous depressive episodes	-.064	.219	-.027	t(154) = -0.293	.770
Vascular risk factors	-.227	.230	-.082	t(124) = -0.984	.327
Early vs late depression onset	-1.437	.660	-.229	t(159) = -2.176	.032

HDRS=Hamilton Depression Rating Scale

**Table 5**  
 Summary of Multivariate Regression Analysis for Prediction of Verbal Learning Performance in LLD Participants (n=161)

Variable	B	Std. Error	Std. Beta	Test	p
Age	-.086	.070	-.118	t(160) = -1.226	.223
Education	.785	.188	.359	t(159) = 4.174	<.001
HDRS	.072	.135	.046	t(160) = 0.534	.594
Number of previous depressive episodes	-.295	.408	-.069	t(155) = -0.722	.472
Vascular risk factors	.016	.431	.003	t(125) = 0.036	.971
Early vs late depression onset	-3.031	1.227	-.265	t(160) = -2.470	.015

HDRS=Hamilton Depression Rating Scale