

DELLAs Function as Coactivators of GAI-ASSOCIATED FACTOR1 in Regulation of Gibberellin Homeostasis and Signaling in *Arabidopsis*^{WOPEN}

Jutarou Fukazawa,^{a,b,c,1} Hiroshi Teramura,^c Satoru Murakoshi,^c Kei Nasuno,^c Naotaka Nishida,^c Takeshi Ito,^a Michiteru Yoshida,^c Yuji Kamiya,^b Shinjiro Yamaguchi,^b and Yohsuke Takahashi^a

^aDepartment of Biological Science, Graduate School of Science, Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan

^bRIKEN Plant Science Center, Tsurumi, Yokohama, Kanagawa 230-0045, Japan

^cDepartment of Biological Science and Technology, Tokyo University of Science, Noda, Chiba 278-8510, Japan

Gibberellins (GAs) are essential regulators of plant development, and DELLAs are negative regulators of GA signaling. The mechanism of GA-dependent transcription has been explained by DELLA-mediated titration of transcriptional activators and their release through the degradation of DELLAs in response to GA. However, the effect of GA on genome-wide expression is predominantly repression, suggesting the existence of unknown mechanisms of GA function. In this study, we identified an *Arabidopsis thaliana* DELLA binding transcription factor, GAI-ASSOCIATED FACTOR1 (GAF1). GAF1 shows high homology to INDETERMINATE DOMAIN1 (IDD1)/ENHYDROUS. GA responsiveness was decreased in the double mutant *gaf1 idd1*, whereas it was enhanced in a GAF1 overexpressor. GAF1 binds to genes that are subject to GA feedback regulation. Furthermore, we found that GAF1 interacts with the corepressor TOPLESS RELATED (TPR) and that DELLAs and TPR act as coactivators and a corepressor of GAF1, respectively. GA converts the GAF1 complex from transcriptional activator to repressor via the degradation of DELLAs. These results indicate that DELLAs turn on or off two sets of GA-regulated genes via dual functions, namely titration and coactivation, providing a mechanism for the integrative regulation of plant growth and GA homeostasis.

INTRODUCTION

Hormonal regulation of transcription is a key control point for plant growth and development. Gibberellins (GAs), which are tetracyclic diterpenoid growth factors, are essential regulators of many aspects of plant development, including seed germination, stem elongation, flower induction, and anther development. The endogenous levels of GAs are fine-tuned by feedback control at several steps in their metabolic pathway, including GA 20-oxidase and GA 3-oxidase (Yamaguchi, 2008). GA feedback regulation has been shown to depend on GA signaling components (Sun, 2011), but its molecular mechanisms are still largely unknown.

DELLA proteins are major plant growth repressors. Upon binding to a soluble GA receptor, GIBBERELLIN INSENSITIVE DWARF1 (GID1), GA triggers the degradation of DELLAs through the ubiquitin-proteasome pathway, thereby promoting plant growth (Ueguchi-Tanaka et al., 2007b; Sun, 2011). *Arabidopsis thaliana* contains five DELLAs, GIBBERELLIN-INSENSITIVE (GAI), REPRESSOR OF *ga1-3* (RGA), RGA-LIKE1

(RGL1), RGL2, and RGL3, which display partially overlapping but also distinct functions in repressing GA responses (Sun and Gubler, 2004). Among the DELLAs, GAI and RGA are the major GA repressors during vegetative growth and floral induction (Mutasa-Göttgens and Hedden, 2009). Because DELLAs localize in the nucleus and show structural similarities to mammalian STAT (for signal transducers and activators of transcription) proteins (Richards et al., 2000), they are thought to be involved in transcription. Several DELLA binding transcription factors have been identified to date. For example, DELLAs regulate hypocotyl elongation by interacting with PHYTOCHROME INTERACTING FACTORS (PIFs) (de Lucas et al., 2008; Feng et al., 2008) and BRASSINAZOLE RESISTANT1 (BZR1) (Bai et al., 2012; Gallego-Bartolomé et al., 2012) and also play a role in plant defense by interacting with JASMONATE ZIM-DOMAIN (JAZ) proteins (Hou et al., 2010). Through these interactions, DELLAs inhibit the activity of these proteins (Hauvermale et al., 2012). Thus, DELLAs function as signaling nodes that mediate the crosstalk of endogenous programs and various environmental stimuli.

Among these transcription factors, PIFs are the most studied. PIFs promote hypocotyl elongation and are negatively regulated by the photoreceptor PHYTOCHROME B. The interaction between DELLAs and PIFs inhibits PIF-induced hypocotyl elongation by blocking the DNA binding activities of PIFs (de Lucas et al., 2008; Feng et al., 2008). GA triggers the degradation of DELLAs, which release PIFs to activate the target genes, including *LIPID TRANSFER PROTEIN3*, β -*EXPANSIN*, and *PACLOBUTRAZOL*

¹ Address correspondence to jutarouf@hiroshima-u.ac.jp.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Jutarou Fukazawa (jutarouf@hiroshima-u.ac.jp).

^{WOPEN} Online version contains Web-only data.

^{OPEN} Articles can be viewed online without a subscription.

www.plantcell.org/cgi/doi/10.1105/tpc.114.125690




Figure 1. Identification of a DELLA Interactor Using a Modified Y2H System.

(A) Schematic representation of DELLA proteins. The fusion of the repression domain of Tup1 repressed the strong transcriptional activity of GAI.
(B) Y2H assay. Tup1-GAI interacts with GAF1 in Y2H assays. Transformed yeast cells were streaked on a plate with His (+His) or without His but with 30 mM aminotriazole (3AT).
(C) β -Galactosidase activity for the Tup1 two-hybrid system. Data are means \pm SD; $n = 3$.

RESISTANCE, and thus promotes hypocotyl elongation (de Lucas et al., 2008). The titration of transcriptional activators by DELLAs partly explains GA-dependent transcriptional activation and how plants integrate environmental stimuli and GA signals to optimize growth and development.

However, genes encoding GA 20-oxidase and GA 3-oxidase are downregulated by GA via feedback regulation. Genome-wide analysis revealed that the effect of GA on gene expression is predominantly through repression, whereas that of DELLAs is through activation (Zentella et al., 2007; Hirano et al., 2012). Furthermore, a chromatin immunoprecipitation (ChIP) assay showed an *in vivo* association of DELLAs with promoters of several genes, although DELLAs lack known DNA binding motifs (Zentella et al., 2007; Lim et al., 2013; Park et al., 2013). All of the DELLA-regulated genes, including GA biosynthetic enzyme genes and GA receptor genes, are repressed by GA and activated by DELLAs (Zentella et al., 2007). These observations cannot be explained by the conventional titration model. Thus, other molecular mechanisms underlying GA-dependent transcriptional regulation must exist.

Here, we describe a previously unknown DELLA binding transcription factor, designated GAI-ASSOCIATED FACTOR1 (GAF1). DELLAs and TOPLESS-RELATED (TPR) act as coactivators and a corepressor of GAF1, respectively, in GA-mediated transcriptional regulation. GA converts the GAF1 complex from transcriptional activator to repressor. DELLAs simultaneously turn on or off two sets of GA-regulated genes via two mechanisms, namely titration and coactivation. Our results thus provide insight into the mechanism of regulation of GA homeostasis and plant growth by DELLA proteins.

RESULTS

Isolation of GAF1, a DELLA-Interacting Protein

Because DELLAs exhibit strong transcriptional activity in yeast, it is difficult to screen for DELLA-interacting proteins using the conventional yeast two-hybrid (Y2H) system with a full-length DELLA as bait (Figures 1A to 1C). To overcome this problem, we developed a modified Y2H system in which a bait protein, the *Arabidopsis* DELLA GAI, was fused to Tup1, a general repressor from yeast. The N-terminal domain of Tup1 (1 to 200 bp), which was sufficient for repression (Jabet et al., 2000; Hirst et al., 2001), reduced the transcriptional activity of GAI in the Tup1-GAI fusion protein (Figures 1B and 1C). We performed a Y2H screen

with Tup1-GAI as bait using an *Arabidopsis* cDNA library. The GAI-interacting protein GAF1 was isolated from 1.6×10^6 transformants. Y2H assays showed that GAF1 interacted with all *Arabidopsis* DELLAs, namely, GAI, RGA, and RGL1 to RGL3 (Figure 1D), and pull-down assays showed direct interaction between GAI and GAF1 (Figure 1E).

To investigate protein-protein interaction between GAI and GAF1 in plant cells, we performed bimolecular fluorescence complementation (BiFC) analysis using *Arabidopsis* T87 cultured cells. The reconstituted yellow fluorescent protein (YFP) signal, caused by interaction between YFP^N-GAI and GAF1-YFP^C, was observed in the nucleus of the protoplasts of T87 cells; no YFP signal was observed when YFP^N-GAI was cotransfected with YFP^C (Figures 1F and 1G). These results suggest that GAF1 binds to GAI in the nucleus of plant cells.

GAF1 Belongs to the IDD Transcription Factor Family

GAF1 encodes a transcription factor with zinc finger motifs that shows similarity to maize (*Zea mays*) INDETERMINATE1 (ID1) (Colasanti et al., 1998, 2006). Mutations in maize *ID1* have severe effects on floral transition (Singleton, 1946; Colasanti et al., 1998), resulting in late flowering, demonstrating that ID1 is essential for normal floral transition in monocots. ID1 appears to be specific to monocots, and functional orthologs are not found in *Arabidopsis* (Colasanti et al., 2006). Although the *Arabidopsis* genome contains 16 ID1-related proteins (IDD, for ID1 domain protein), their amino acid sequence similarity to ID1 is limited to the zinc finger motif that is important for DNA binding (Figure 2A). Maize ID1 binds to the consensus sequence TTTTGTCG (Kozaki et al., 2004). To determine whether GAF1 binds to this sequence, we performed a gel retardation assay. Recombinant GAF1 protein specifically bound to the ID1 binding sequence (Figure 2B), suggesting that GAF1 is a sequence-specific DNA binding protein.

To investigate the expression pattern of *GAF1* in plants, we monitored the activity of β -glucuronidase (GUS) driven by the *GAF1* promoter (Figure 2C). Histochemical analysis indicated that the *GAF1* promoter is mainly expressed in hypocotyls, petioles, shoot apices, root tips, and trichomes (Figure 2C). This observation is consistent with the microarray data derived from different stages of *Arabidopsis* development (AtGenExpress). Expression of *GAF1* mRNAs in various organs was confirmed by quantitative real-time PCR (Figure 2D). We further investigated the expression of GAF1 protein by immunoblot analysis using antibody raised against recombinant GAF1. Our analysis

Figure 1. (continued).

(D) GAF1 and IDD1 interact with five DELLA proteins in yeast β -galactosidase assays. Data are means \pm sd; $n = 3$. vec indicates empty vector used as a negative control.

(E) *In vitro* pull-down assays with GST-GAI protein. GST and GST-GAI proteins were incubated with recombinant 6 \times His-GAF1 protein bound to Glutathione Sepharose 4B and then eluted and analyzed by immunoblotting (IB) with anti-GAF1 antibody (top) and anti-GST antibody (bottom).

(F) BiFC analysis showing interaction between GAF1 and GAI. *CaMV35S:GAF1-YFP^C* and *CaMV35S:YFP^N-GAI* plasmids were introduced and transiently expressed in protoplasts of T87 *Arabidopsis* cultured cells (left). *CaMV35S:YFP^C* and *CaMV35S:YFP^N-GAI* plasmids were introduced into protoplasts of T87 *Arabidopsis* cultured cells as a negative control (right). DIC, differential interference contrast.




Figure 2. Characterization of GAF1.

detected GAF1 in transgenic plants expressing GAF1 under the control of the cauliflower mosaic virus (CaMV) 35S promoter but not in wild-type plants (Figure 2E). These results indicate that GAF1 is mainly expressed in vegetative tissues at low levels.

The SAW DOMAIN of GAI Is Important for GAF1 Binding

DELLAs belong to a subclass of the plant-specific GRAS family, a family of transcriptional regulators. Like all GRAS proteins, DELLAs share a conserved C-terminal domain that is involved in transcriptional regulation and is characterized by two Leu heptad repeats (LHRI and LHRII) and three conserved motifs, VHIIID, PFYRE, and SAW (Itoh et al., 2002; Ueguchi-Tanaka et al., 2007a) (Figure 1A). DELLAs are distinguished from the rest of the GRAS family by a specific N-terminal sequence containing two conserved domains: the DELLA domain (which gives them their name) and the TVHYNP domain. To determine which region of GAI is important for interaction with GAF1, we generated various deletion mutants of GAI (Figure 2F) and performed Y2H assays. As shown in Figure 2F, deletion of the N-terminal region of GAI did not affect GAF1 binding. By contrast, deletion of the C-terminal eight amino acids containing the core sequence of the SAW domain abolished GAF1 binding (Δ SAW). Because the mutation in the SAW domain of SLENDER RICE1 (SLR1), a rice (*Oryza sativa*) DELLA, caused slender phenotypes in rice (Ikeda et al., 2001), the SAW domain of DELLA is thought to be necessary for the repression of GA responses (Itoh et al., 2002; Ueguchi-Tanaka et al., 2007a). These results suggest that GAF1 might be involved in GA responses through binding to the SAW domain of DELLA.

The *gaf1 idd1* Double Mutant and GAF1 Overexpressor Lines Show GA-Related Phenotypes

The most closely related protein to GAF1 among the *Arabidopsis* IDD family is IDD1. The two proteins share 68.5% amino acid sequence similarity. IDD1 is also known as ENHYDROUS (ENY) and promotes the transition of seeds to germination by regulating light and hormonal signaling during their maturation (Feurtado et al., 2011). IDD1/ENY also interacts with DELLAs (Feurtado et al., 2011; Figure 1D); however, the functional

significance of this remains unclear. *GAF1* and *IDD1* show distinct but overlapping expression patterns (i.e., *GAF1* is expressed mainly in vegetative tissues) (Figure 2C), while *IDD1/ENY* is expressed mainly in seeds and at lower levels in vegetative tissues (Feurtado et al., 2011). These observations suggest that *GAF1* and *IDD1* could play partially redundant roles in plants.

To investigate the function of GAF1, we compared the *gaf1* and *idd1* mutants with GAF1 overexpressors (Supplemental Figure 1). A T-DNA insertional mutant line for *GAF1* (SALK_070916) was obtained and confirmed as a transcriptional knockout (Figure 3A; Supplemental Figure 2A). However, the only available mutant line for *IDD1* (SALK_022425) has a T-DNA insertion in the promoter region, leading to reduced expression of *IDD1* as compared with that of the Columbia-0 (Col-0) wild type (Figure 3A; Supplemental Figure 2A).

Because *gaf1* showed slightly late flowering under short-day (SD) conditions, which is a GA-related phenotype, we generated and characterized a *gaf1 idd1* double mutant (Figure 3). GAs affect flowering: GA-deficient mutants are late flowering and a DELLA quadruple mutant was shown to be early flowering under SD conditions (Cheng et al., 2004). We examined the effects of mutations and the overexpression of *GAF1* on flowering. The *gaf1 idd1* double mutant flowered later than the wild type, especially under SD conditions (Figures 3B to 3E), while the *GAF1* overexpressor flowered earlier (Figures 3D to 3F). In addition to this, the *gaf1 idd1* double mutant showed a semidwarf phenotype (Figures 3G and 3H). To confirm that the late flowering of *gaf1 idd1* under SD conditions is due to the mutations in *GAF1* and *IDD1*, we performed a complementation experiment. The late-flowering phenotype of *gaf1 idd1* was rescued by the expression of a *GAF1-GFP* fusion gene under the control of the *GAF1* promoter (Supplemental Figure 3). The *GAF1* overexpressor plants were also taller (reaching 100 cm) than the wild type under SD conditions (Figure 3F). Furthermore, treatment with GA_4 , which restores the growth of *ga1-3*, a GA-deficient mutant, did not affect the semidwarf phenotype of *gaf1 idd1* (Figure 3I), suggesting that GA responsiveness is reduced in *gaf1 idd1*.

As GAs promote the elongation of the hypocotyl, we examined the effects of bioactive GA application on the hypocotyl

Figure 2. (continued).

(A) Alignments of the zinc finger domains of GAF1, IDD1, and maize ID1 proteins. Identical residues are highlighted in black. The solid circles indicate the Cys and His residues in the C_2H_2 -type zinc finger motifs. The solid triangles indicate the nuclear localization signal.

(B) Gel retardation assay using recombinant GAF1 protein. Oligonucleotides containing *ID1-cis* (WT; lanes 1 to 4) or *mtID1-cis* (mt; lane 5) were used as probes. The mutated bases are highlighted. WT and mt, competition with a 1000-fold excess of unlabeled wild-type and mutated probe, respectively. The specific GAF1-DNA complexes are indicated by the arrowhead. +, addition to the reaction mixtures; -, omission from the reaction mixtures.

(C) GUS expression pattern in transgenic plants expressing *GAF1 promoter:GUS*. Analysis was performed in 3-, 6-, and 10-d-old seedlings. Bars = 5 mm.

(D) Quantitative real-time PCR analysis of *GAF1* expression in different tissues. *UBQ11* was used as an internal control. Data are means \pm SD; $n = 3$.

(E) Immunoblot analysis of GAF1 protein expression in different tissues of Col-0 and 35S:*GAF1*-overexpressing transgenic *Arabidopsis*. Coomassie blue (CBB) staining was used as a loading control.

(F) Identification by Y2H assay of the domain of GAI that binds to GAF1. Schematic representations of GAI-truncated proteins used for Y2H assays. Yeast was transformed with a combination of the indicated plasmids, and subsequently, β -galactosidase activity was determined. Data are means \pm SD; $n = 3$.




Figure 3. Phenotypes of *GAF1* Overexpressor and the *gaf1 idd1* Mutant.

(A) Positions of the T-DNA insertions within At3g50700 in line SALK_070916 (*gaf1*) and At5g66730 in line SALK_022425 (*idd1*). Bar = 200 bp.

(B) A 25-d-old of *gaf1 idd1* double mutant plant exhibiting a semidwarf phenotype under long-day (LD) conditions.

length of *gaf1 idd1* and *GAF1* overexpressor. The hypocotyl of the *GAF1* overexpressor was longer than that of the wild type either in the presence or absence of exogenous GA, whereas that of *gaf1 idd1* was shorter even in the presence of GA (Figure 3J). Another mutant allele of *IDD1* (*TL-7*) exhibited similar phenotypes in the *gaf1* background (Supplemental Figure 4). These results suggest that the phenotype of *gaf1 idd1* results from reduced responsiveness to GA and that both *GAF1* and *IDD1* are involved in GA signaling.

Binding Domain of GAF1 to GAI

To identify the domain of *GAF1* that interacts with *GAI*, we generated various deletion mutants of *GAF1* (Figure 4A). *GAI* bound to the C-terminal half of *GAF1* (189 to 452) but not to the zinc finger domain of *GAF1* (34 to 196) in the Y2H assay (Figure 4A). Internal deletion of 16 amino acids from 325 to 340 of *GAF1*, which we refer to hereafter as the PAM domain, abolished *GAI* binding (Figure 4A; Supplemental Figure 5A); however, this deletion did not affect binding to another protein, *TPR* (see below). These results suggest that the PAM domain of *GAF1* is necessary for *GAI* binding.

Suppression of GA-Deficient and GA-Insensitive Mutants by GAF1 and ΔPAM

GA-deficient mutants display characteristic phenotypes, including reduced germination, dark green leaves, and a dwarf growth habit attributable to reduced stem elongation. A GA-insensitive semidominant mutant, *gai-1*, in which *GAI* is resistant to GA-dependent proteolysis because of a lack of the N-terminal DELLA domain, shows phenotypes similar to those of GA-deficient mutants except for reduced germination. To investigate the biological role of *GAI* binding to *GAF1*, we generated transgenic plants expressing *GAF1* or the mutant version of *GAF1* that cannot bind *GAI* (Δ PAM) under the control of the *CaMV* 35S promoter in the *ga1-3* or *gai-1* background (Supplemental Figures 2B to 2D). The expression of Δ PAM rescued the reduced germination phenotype of *ga1-3* more effectively than did that of wild-type *GAF1* (Figure 4B). Although both wild-type *GAF1* and the *GAF1* mutant Δ PAM could partially rescue the dwarf phenotypes of *ga1-3* and *gai-1*, Δ PAM was found to be more effective (Figures 4B to 4F). These results suggest that the *GAF1* mutant Δ PAM is more effective in suppressing the GA-deficient

mutant *ga1-3* and the GA-insensitive mutant *gai-1*. As accumulation and reduction of DELLAs inhibits and promotes plant growth, respectively, we examined whether the overexpression of *GAF1* promotes plant growth by repressing the expression of *GAI*. Immunoblot analysis showed that *GAI* protein levels were not reduced by the expression of *GAF1* or the *GAF1* mutant Δ PAM (Figures 4G and 4H), suggesting that overexpression of *GAF1*s can rescue plant growth in *ga1-3* and *gai-1* without affecting the expression of *GAI*.

TOPLESS Is a GAF1 Binding Protein

The fact that the mutant version of *GAF1* (Δ PAM), which cannot bind to *GAI*, suppresses the dwarf phenotypes of *ga1-3* or *gai-1* more effectively than *GAF1* (Figure 4) raised the possibility that *GAF1* plays a role in promoting plant growth after DELLAs are degraded in response to GAs. Transcription factors often form multiprotein complexes with other proteins, including general transcription factors, cofactors, or enzymes. To investigate the non-DELLA-dependent function of *GAF1*, we searched for *GAF1* binding proteins other than DELLAs using a conventional Y2H system. From 1.5×10^5 transformants, we identified *TPR1* and *TPR4* as *GAF1* binding proteins in addition to DELLAs. In *Arabidopsis*, *TOPLESS* (*TPL*) and *TPRs* form a family of transcriptional corepressors (Long et al., 2006; Szemenyei et al., 2008). The *TPL/TPR* family belongs to a larger Groucho/Tup1 family, first identified in *Drosophila* and yeasts, respectively. This family represents an ancient class of corepressors recruited by a range of DNA binding transcription factors to elicit a repressed chromosomal state (Liu and Kamarkar, 2008; Krogan et al., 2012; Wang et al., 2013). Recent studies showed that *TPL/TPR* corepressors interact with transcription complexes involved in auxin and jasmonate signal transduction, meristem maintenance, and defense responses (Long et al., 2006; Szemenyei et al., 2008; Pauwels et al., 2010).

Using BiFC analysis, we confirmed the *in vivo* interaction between *GAF1* and *TPR4*, and *GAF1* and *GAI* in the nucleus of *Arabidopsis* T87 cultured protoplast cells (Figure 5A). GAs release the growth inhibition imposed by DELLAs via the degradation of DELLAs through the ubiquitin-26S proteasome pathway (Sun, 2010). We examined whether the interactions between *GAF1* and *GAI* and between *GAF1* and *TPR4* are regulated by GAs. Treatment with GA induced the disappearance of the interaction between *GAF1* and *GAI* within 60 min, while it did

Figure 3. (continued).

(C) A 40-d-old of *gaf1 idd1* double mutant plant exhibiting a semidwarf phenotype under SD conditions.

(D) and **(E)** Flowering time analysis (rosette leaf number) under LD and SD conditions. The *GAF1* overexpressor exhibits early flowering under LD and SD conditions. The *gaf1 idd1* mutant exhibits slightly delayed flowering in LD conditions and extremely delayed flowering under SD conditions ($n > 8$). Asterisks represent significance by Student's *t* test compared with Col-0 (** $P < 0.01$).

(F) One hundred-day-old *GAF1*-overexpressing plants exhibiting early flowering and taller phenotypes. Bar = 10 cm.

(G) Phenotypes of Col-0, *gaf1*, *idd1*, *gaf1 idd1*, and *GAF1* overexpressor plants grown for 90 d under SD conditions.

(H) Extremely delayed flowering phenotype of the *gaf1 idd1* mutant (8 months old) under SD conditions.

(I) The *gaf1 idd1* mutant is GA insensitive. Plants were treated or not with bioactive GA_4 (1 μ M). GA_4 was applied once per week for 2 months.

(J) Hypocotyl lengths of 11-d-old *gaf1 idd1*, Col-0, and *GAF1* overexpressor plants grown on GA_3 -containing medium. Data are means \pm sd; $n = 10$. The photographs in the top panel show hypocotyl lengths of 11-d-old *gaf1 idd1*, Col-0, and *GAF1* overexpressor plants grown on medium containing 0, 1, or 10 μ M GA_3 . Asterisks represent significance by Student's *t* test compared with Col-0 (** $P < 0.01$ and * $P < 0.05$).




Figure 4. Effect of Overexpression of *GAF1* or Δ *PAM* in the *gai-3* or *gai-1* Mutant.

(A) Schematic representations of *GAF1* proteins used for Y2H assays. Levels of DELLA binding activity are indicated by + or -. The + and - signs indicate interaction and no interaction between truncated *GAF1* and *GAI*, respectively.

not affect the interaction between GAF1 and TPR4 (Figures 5B and 5C). However, treatment with MG132, a proteasome inhibitor, blocked the GA-induced disappearance of BiFC fluorescence. The subcellular localization and stability of GAF1-GFP were not affected by treatment with GAs (Supplemental Figures 5B to 5D). These observations were confirmed by coimmunoprecipitation experiments using transgenic plants expressing myc-tagged GAF1 under the control of the CaMV 35S promoter. The myc-tagged GAF1 protein was immunoprecipitated with anti-myc antibodies, and GAF1-bound materials were blotted and probed with anti-GAI antibodies or anti-TPR4 antibodies. As shown in Figure 5D, GAI and TPR4 coimmunoprecipitated with myc-tagged GAF1. GAs promoted the loss of interaction between GAI and GAF1 but did not affect the interaction between TPR4 and GAF1 (Figure 5B). These results suggest that the binding partners of GAF1 in plant cells are regulated by GA levels.

The Domain of GAF1 That Binds TPR4

To identify the interaction domain of GAF1 for TPR4, we used various deletion mutants of GAF1 (Supplemental Figure 6A). We found that internal deletion of 18 amino acids from residues 367 to 384 of GAF1 diminished TPR4 binding in the Y2H assay, but the deletion did not affect GAI binding (Figures 6A and 6B; Supplemental Figure 5A). This region includes the ethylene-responsive element binding factor-associated amphiphilic repression (EAR) motif, which is a target of TPLs (Tiware et al., 2004; Szemenyei et al., 2008). The EAR motif was found only in GAF1 and IDD1 among *Arabidopsis* IDD proteins. These results suggest that GAF1 binds to GAI and TPR4 through independent binding motifs.

GAI and TPR4 Act as Transcriptional Cofactors of GAF1 in Plant Cells

Previous studies showed that DELLAs negatively regulate transcriptional activators, including PIFs, MYC2, and BZR1, by inhibiting DNA binding (Bai et al., 2012; Hong et al., 2012). GAs relieve the activities of transcriptional activators by promoting the degradation of DELLAs via the ubiquitin-26S proteasome pathway. Nevertheless, this model does not explain the feedback regulation of GA biosynthetic genes, which are upregulated in the presence of accumulated DELLA protein under

GA-deficient conditions. However, the fact that TPR4 is a corepressor and GAI exhibits transcriptional activity in yeast (Figure 1C) suggests that GAI and TPR4 may function as coactivator and corepressor of GAF1, respectively. This hypothesis would provide a molecular mechanism for GA feedback regulation, namely transcriptional activation of GA biosynthetic genes by GAF1 and DELLAs under GA-deficient conditions.

To test this, we performed transient assays using *Arabidopsis* T87 protoplasts. The promoter of *AtGA20ox2* encoding GA 20-oxidase fused with *GUS* was used as a reporter because *AtGA20ox2* is under GA feedback regulation and activated by the induction of DELLAs within 2 h (Zentella et al., 2007) and contains the putative GAF1 target sequences in its promoter. Although neither GAF1 nor GAI alone affected the expression of *AtGA20ox2:GUS*, cotransfection of GAI with GAF1 greatly enhanced the expression of the reporter gene (Figure 6C; Supplemental Figure 6B). Cotransfection of GAI with the mutant version of GAF1 (Δ PAM) that cannot bind to GAI failed to activate the reporter gene, indicating that GAI binding to GAF1 is indispensable for transcriptional activation. Contrary to GAI, TPR4 showed corepressor activity with GAF1 in the transient assay. Cotransfection of TPR4 with the mutant version of GAF1 (Δ EAR) that cannot bind TPR4 did not repress the expression of the reporter gene, indicating that TPR4 binding to GAF1 is necessary for transcriptional repression. IDD1 also exhibited similar activity (Figure 6D).

With respect to feedback regulation of GA, the expression of *AtGA20ox2* is negatively regulated by GAs. Thus, we examined if transcriptional regulation of *AtGA20ox2* by the GAF1 transcription complex is modulated by GAs. As expected, the transcriptional activation of *AtGA20ox2* by GAF1 and GAI was completely inhibited by treatment with gibberellic acid (GA_3) in the presence of TPR4 (Figure 6E).

The GAF1 Complex Is Involved in GA Feedback Regulation of *AtGA20ox2*

To evaluate the contribution of the GAF1 complex to GA feedback regulation of *AtGA20ox2* in plants, we investigated *AtGA20ox2* mRNA levels in the wild type (Col-0), the GA-deficient mutant *gai-3*, and a transgenic line expressing Δ PAM in the *gai-3* background. The upregulation of *AtGA20ox2* in the GA-deficient mutant *gai-3* was repressed by the expression of Δ PAM, showing that the GAF1-DELLA complex is important for

Figure 4. (continued).

(B) Germination percentage of seeds of the *gai-3* mutant compared with that of *gai-3/35S:GAF1* or *gai-3/35S: Δ PAM*. Two days after stratification, seeds were transferred to medium without GA. Germination was monitored every 12 h between 24 and 96 h after imbibition.

(C) Representative plants of *gai-3*, *gai-3/35S:GAF1*, and *gai-3/35S: Δ PAM* grown in soil for 40 d.

(D) Representative plants of *gai-1* and *gai-1/35S:GAF1* grown in soil for 70 d.

(E) Representative plants of *gai-1* and *gai-1/35S: Δ PAM* grown in soil for 70 d.

(F) Final height of *gai-1*, *gai-1/35S:GAF1*, and *gai-1/35S: Δ PAM* transgenic plants. Error bars represent sd .

(G) and **(H)** Immunoblot analysis of *gai-1*, *gai-1/35S:GAF1*, and *gai-1/35S: Δ PAM* transgenic plants with anti-GAI antibody. Seven-day-old Col-0 seedlings were transferred to half-strength MS plates with or without 1 μ M PAC or 1 μ M PAC and 10 μ M GA_3 and incubated for 1 week. *gai-1*, *gai-1/35S:GAF1*, and *gai-1/35S: Δ PAM* transgenic plants were grown without PAC and GA_3 for 2 weeks. Total proteins were extracted from whole plants and analyzed by immunoblotting with affinity-purified anti-GAI antibody. Coomassie blue staining was used to confirm equal loading. Arrowheads indicate the positions of the GAI protein (top) and the *gai-1* protein (bottom). Immunoblot analysis used overexpression of GAF1 in *gai-1* **(G)** and overexpression of Δ PAM in *gai-1* **(H)**.




Figure 5. In Vivo Interaction between GAF1 and GAI or TPR4 with or without GA.

(A) BiFC analysis of the interaction between GAF1 and TPR4 in T87 *Arabidopsis* cultured cells. CaMV35S:GAF1-YFP^C and CaMV35S:YFP^N-TPR4 plasmids were introduced and transiently expressed in protoplasts of T87 *Arabidopsis* cultured cells. DIC, differential interference contrast.

(B) BiFC analysis of the interaction between GAF1 and GAI or TPR4 in T87 *Arabidopsis* cultured cells. GA treatment disrupts the interaction between GAF1 and GAI. The top three rows show the interaction between GAF1 and GAI, while the bottom row shows the interaction between GAF1 and TPR4. The cells in the second and bottom rows were treated with 10 μM GA₃. The cells in third row were treated with 10 μM GA₃ and 10 μM MG132. Each cell was observed at 0, 5, 10, 20, 40, 60, 90, and 120 min after treatment with or without GA₃ or GA₃ + MG132. Bars = 10 μm.

(C) Measure of the interaction between GAF1 and GAI or TPR4 in the presence or absence (Normal) of GA and GA + MG132 using BiFC. Error bars represent SD (*n* = 3).

GA feedback regulation in plants (Figure 7A). Furthermore, we confirmed that GAF1 associates with DELLAs and TPR on the *AtGA20ox2* promoter in plants. ChIP assay using anti-GAF1, anti-GAI, or anti-TPR4 antibody showed that GAF1, GAI, or TPR4 binds to the *AtGA20ox2* promoter (–1030 to –1011) in plants, indicating that the coactivator GAI and the corepressor TPR4 bind to the target gene through the DNA binding transcription factor GAF1 (Figures 7B and 7C). These results suggest that GAF1 directly regulates *AtGA20ox2* either positively or negatively with the coactivator GAI or the corepressor TPR4, respectively, in response to GA levels.

Target Genes of GAF1

To identify other target genes of GAF1, we focused on DELLA-regulated genes (Zentella et al., 2007) because DELLAs are thought to bind to target genes indirectly through GAF1 or through an unknown DNA binding protein. We isolated the promoter regions of several DELLA-regulated genes and fused them to *GUS*. In addition to *AtGA20ox2*, GAF1 and GAI activated the promoters of *AtGA3ox1* encoding a GA 3-oxidase and *GID1b* encoding a GA receptor among the DELLA-regulated genes tested (Figure 8A). Putative GAF1 target sequences were found in the *AtGA20ox2* and *GID1b* promoters. A gel retardation assay indicated that recombinant GAF1 directly binds to these sequences (Figure 8B). Furthermore, ChIP assays using transgenic plants expressing myc-GAF1 under the control of the CaMV 35S promoter showed that GAF1 binds to the *AtGA20ox2* and *GID1b* promoters in vivo (Figures 8C and 8D). These results suggest that GAF1 directly regulates *GID1b* as well as *AtGA20ox2* and is involved in the feedback regulation of GA biosynthesis.

DISCUSSION

Defining the precise molecular mechanisms that determine patterns of transcription in response to specific signals is essential for understanding development and homeostasis. Transcriptional cofactors, in addition to DNA binding transcription factors, play key roles in such biological responses. Transcriptional complexes can coordinately turn target genes on or off depending on the functions of coactivators and corepressors.

To date, the molecular mechanism of GA-dependent transcriptional regulation has been explained by the titration of transcriptional activators, including PIFs, with DELLAs and their release through the degradation of DELLAs in response to GA (de Lucas et al., 2008; Feng et al., 2008). However, the effect of GA on gene expression is predominantly repression, and DELLAs associate with the promoters of DELLA-induced genes

even though DELLA proteins lack DNA binding motifs (Zentella et al., 2007; Hirano et al., 2012; Lim et al., 2013; Park et al., 2013). These observations suggest the existence of hitherto unknown mechanisms of GA-dependent transcriptional regulation. In this study, we found that DELLAs promote the transcription of genes encoding GA biosynthetic enzymes and GA receptors as coactivators of a DNA binding transcription factor, GAF1. Upon GA perception, these genes are actively repressed by a GAF1 corepressor-TPR complex that appears upon the degradation of DELLAs via the ubiquitin-26S proteasome pathway. Thus, the transcriptional state of GAF1 target genes is determined by the balance between coactivator and corepressor. This mechanism clearly accounts for GA feedback regulation and explains why the effect of GA on gene expression is predominantly repression, neither of which can be explained by the conventional titration model (Figure 9).

Mammalian nuclear receptors are the best investigated examples of proteins with dual transcriptional regulation properties (Rosenfeld et al., 2006). Nuclear receptors for steroids, retinoic acid, and thyroid hormone are potent repressors in the absence of ligand, while they function as activators of transcription when bound to their cognate ligands. Upon ligand binding, the conformational change in the ligand binding domain of these receptors induces corepressors to dislodge and coactivators to bind, allowing transactivation (Rosenfeld et al., 2006). The exchange of coactivators/corepressors is an interesting common molecular framework of the hormonal regulation of transcription in two evolutionally distant organisms (i.e., plants and mammals). However, there is a significant difference: the GA-induced functional conversion of the GAF1 complex depends on the degradation of coactivator DELLAs through the ubiquitin-proteasome pathway, whereas ligand-induced activation of the transcription of nuclear receptors in mammals depends on a conformational change.

Many of the cofactors do not operate in isolation but, rather, are part of large multiprotein complexes. Transcriptional complexes exhibit a diversity of enzymatic activities that can be divided into two generic classes: enzymes capable of remodeling the structure of the nucleosome in an ATP-dependent manner and enzymes capable of covalently modifying histone tails. The latter group includes those with acetylating and deacetylating activities or methylating and demethylating activities, as well as kinases and phosphatases, poly(ADP) ribosylases, and ubiquitin and SUMO ligases (reviewed in Rosenfeld et al., 2006). Serial posttranslational modifications of histones and transcription factors are involved in the specific activation or repression of the genes. DELLAs could have activities for chromatin remodeling or covalent modifications; however, they do not have conserved catalytic motifs. Alternatively, DELLA proteins could recruit these enzymes to the promoters of target genes. In this context, it is worth noting that animal STAT, which shows structural

Figure 5. (continued).

(D) In vivo interaction between GAF1 and GAI or TPR4 in *Arabidopsis*. Extracts from 35S:4×myc-GAF1 seedlings were immunoprecipitated (IP) using anti-myc antibody. The coimmunoprecipitated proteins were detected by either anti-GAI or anti-TPR4 antibody. Each plant was grown with or without GA and PAC. N, control; P, 1 μ M PAC; P+G, 1 μ M PAC and 10 μ M GA₃.




Figure 6. A DELLA and TPR Act as a Coactivator and a Corepressor of GAF1, Respectively.

(A) Interaction domain of GAF1 for GAI or TPR4. Positive and no interactions are indicated by + and -, respectively. The PAM domain, which is necessary to bind to GAI, and the EAR-like motif, which is necessary to bind to TPR4, are indicated.

(B) β -Galactosidase activity of Y2H assay showing that Δ PAM or Δ EAR cannot interact with GAI or TPR but can interact with TPR or DELLAs, respectively. Data are means \pm SD; $n = 3$.

(C) Transactivation assay of GAF1, GAI, and TPR4. The reporter, effector, and internal control constructs used in the assay are shown in the left panel. The reporter, effector, and internal control plasmids were cotransfected into *Arabidopsis* T87 cell protoplasts. The transfected protoplasts were incubated for 20 h, and then the GUS and LUC activities were measured. The results are shown as GUS/LUC activity. Error bars indicate SD ($n = 3$).

(D) The functions of IDD1 are similar to that of GAF1. The results are shown as GUS/LUC activity. Error bars indicate SD ($n = 3$).

(E) Transactivation assay showing that GAF1-GAI activates while GAF1-TPR represses the activity of the GA20ox2 promoter. GA represses the transcriptional activation of the GA20ox2 promoter via the GAF1-GAI-TPR complex. Transfected protoplasts were incubated for 20 h with or without 10 μ M GA, and then GUS and LUC activities were measured. Error bars indicate SD ($n = 3$).

similarity to DELLA proteins, associates with various cofactors, including histone acetyltransferase p300/CBP (CREB binding protein) and PCAF (p300/CBP-associated factor), histone deacetylase HDAC, and chromatin remodeler BRG1 (*brm/SWI2-related gene 1*) to regulate transcription (Rosenfeld et al., 2006).

In addition to this coactivator function, DELLAs inhibit transcriptional activators by preventing DNA binding. Thus, DELLAs can have opposite transcriptional effects on two sets of GA-regulated genes by acting as a coactivator of GAF1 and by titrating transcriptional activators, including PIFs and BZR1 (Bai




Figure 7. The GAF1 Complex Is Involved in GA Feedback Regulation of *AtGA20ox2*.

(A) The overexpression of ΔPAM in *ga1-3* represses the upregulation of *GA20ox2* by GA feedback regulation. Relative expression levels of *GA20ox2* in Col-0, *ga1-3*, and 35S:ΔPAM in *ga1-3* are shown. Error bars indicate the SD of three biological replicates ($n = 3$).

(B) ChIP analysis of GAF1, GAI, and TPR4 binding to the GAF1 binding regions of *GA20ox2* promoters (−1030 to −1011). ChIP was performed with Col-0 using anti-GAF1, anti-GAI, and anti-TPR4 antibodies (+) or preimmune serum (−). The *GA20ox2* promoter was detected by PCR and DNA gel blot hybridization. The values at the bottom of each panel indicate the relative signal strength. The value for preimmune serum was set to 1.0.

(C) ChIP assays performed with preimmune serum, anti-GAF1, anti-GAI, or anti-TPR4 antibody in Col-0. The coprecipitated level of each DNA fragment was quantified by real-time PCR and normalized with respect to the input DNA. The relative coprecipitated level of each DNA fragment using preimmune serum was set to 1, and the relative enrichment of each DNA fragment using anti-GAF1, anti-GAI, or anti-TPR4 is shown. Error bars indicate the SD of three biological replicates ($n = 3$).




Figure 8. Identification of GAF1 Target Genes.

(A) Transient transactivation assay showing that the GAF1/GAI complex also activates the *GA3ox1* and *GID1b* promoters. The reporter plasmids consist of a 3-kb promoter region of *GA3ox1*, *GID1a*, *GID1b*, *RGL1*, and *MYB* fused with the *GUS* reporter gene. The results are shown as GUS/LUC activity. Error bars indicate *SD* (*n* = 3).




Figure 9. A Coactivator and Corepressor Model for GA Signaling.

Under GA-deficient conditions, DELLA proteins are stable and localized in nuclei. DELLAs titrate PIF and BZR transcription factors by inhibiting DNA binding activity while exhibiting high transcriptional activity with GAF1. In the presence of GA, DELLAs are degraded via the 26S proteasome pathway. On the one hand, PIF and BZR activate target genes such as *PACLOBUTRAZOL RESISTANCE (PRE)* and *EXPANSIN*. On the other hand, the GAF1-TPR complex exhibits transcriptional repression activity. GA-induced functional conversion of the GAF1 complex in plants depends on the degradation of coactivator DELLA proteins.

et al., 2012; Hong et al., 2012). This mechanism remarkably suits GA signaling. A decrease in GA levels will turn off the expression of genes involved in the promotion of plant growth by titration of transcriptional activators with DELLA proteins and simultaneously turn on the expression of genes subject to GA feedback regulation by a GAF1-DELLA complex. Conversely, an increase in GA will have opposite effects on the two sets of genes. Such dual properties of a cofactor introduce a new paradigm for hormone action through which variations in the levels of GA can coordinately turn on or off two sets of genes, thus enabling the integrative regulation of plant growth and GA homeostasis. Furthermore, because DELLAs mediate the crosstalk of internal and external stimuli through the interaction

with key regulators of signaling, including PIFs of light, BZR1 of brassinosteroids, and JAZ of jasmonates (Bai et al., 2012; Hong et al., 2012), they serve as central regulators of plant development.

Our understanding of GA signaling has advanced considerably in recent years (Hauvermale et al., 2012); however, the mechanisms by which DELLAs repress plant growth remain to be elucidated. Sequence-specific transcription factors collectively function as the key interface between genetic information encoded in the DNA sequence and the signal transduction systems in response to internal and external stimuli. Our results suggest that the DNA binding transcription factor GAF1 regulates genes encoding GA biosynthetic enzymes and GA

Figure 8. (continued).

(B) Gel retardation assay showing that recombinant GAF1 binds to promoters of *GA20ox2* and *GID1b*. Oligonucleotides (20 or 21 bp) were used as probes. The numbers adjacent to the gene names indicate base pairs upstream of the initiation ATG of each gene. The specific GAF1-DNA complexes are indicated by the arrowhead. +, addition to the reaction mixtures; -, omission from the reaction mixtures.

(C) ChIP analysis of 4×myc-GAF1 binding to the GAF1 binding regions of *GA20ox2* and *GID1b* promoters described in **(B)**. ChIP was performed with a 35S:4×myc-GAF1 transgenic plant using anti-myc or anti-GST antibody. The *GA20ox2* and *GID1b* promoters and the *GA20ox2* coding region were detected by PCR and DNA gel blot hybridization. The values at the bottom of each panel indicate the relative signal strength. The value for anti-GST antibody was set to 1.0.

(D) ChIP assays performed with anti-GST or anti-myc antibody in a 35S:4×myc-GAF1 transgenic plant. The coprecipitated level of each DNA fragment was quantified by real-time PCR and normalized with respect to the input DNA. The relative coprecipitated level of each DNA fragment using anti-GST antibody was set to 1, and the relative enrichment of each DNA fragment using anti-myc antibody compared with anti-GST antibody is shown. Error bars indicate the SD of three biological replicates ($n = 3$).

receptors either positively or negatively depending on GA levels. The mutant version of GAF1 that cannot bind GAI (Δ PAM) rescued vegetative growth more effectively than did wild-type GAF1 in DELLA-accumulating plants (Figure 4), suggesting that the GAF1-TPR repressor complex rather than the GAF1-DELLA activator complex plays a role in promoting plant growth. Conversely, we found that GAF1 binds to the DELLA SAW domain (Figure 2F), which is necessary for the repression of GA responses by DELLAs, suggesting that GAF1 is involved in DELLA-mediated growth repression. One possible interpretation of these seemingly conflicting observations is that the GAF1-DELLA complex regulates the transcription of a gene encoding a growth repressor as well as that of genes encoding GA biosynthetic enzymes and a GA receptor. A decrease in GA levels could promote the expression of a growth repressor gene through a GAF1-DELLA complex, resulting in growth arrest, while an increase in GA levels could actively repress the growth repressor gene through the GAF1-TPR complex, promoting plant growth. Expression of SLR1, the rice DELLA protein, fused to the activation domain of the herpes simplex virus protein VP16 has been shown to more severely inhibit plant growth than that of wild-type SLR1 (Hirano et al., 2012), suggesting that DELLAs suppress plant growth through transcriptional activity. This supports the hypothesis that an unknown growth repressor gene is one of the target genes of the GAF1-DELLA complex. Identification of such a gene will provide an important clue for understanding the mechanisms downstream of DELLAs in GA signaling, which influences many aspects of plant growth and development.

METHODS

Plant Material and Growth Conditions

All mutant and transgenic lines in this study were derived from *Arabidopsis thaliana* ecotype Col-0 (wild type), with the exception of the *gai-1* mutant, which was isolated from the Landsberg *erecta* ecotype. The *gai-3* mutant was introgressed into Col-0 by backcrossing six times (Tyler et al., 2004). The *gaf1-1* (SALK_070916), *idd1* (SALK_022425), and *gai-1* (CS63) mutants were obtained from the ABRC (Ohio State University). *gaf1 idd1* double mutants were generated by crossing the *gaf1* and *idd1* mutants. To generate transgenic plants overexpressing AtGAF1 or Δ PAM, GAF1 or Δ PAM cDNA was cloned into the *NotI-XhoI* site of the binary pBIJ4 vector, which contains a CaMV 35S promoter with an Ω sequence and a Kozak sequence to enhance translation activity, as described previously (Fukazawa et al., 2010). To generate transgenic plants overexpressing myc-tagged GAF1, the coding sequence for the 4 \times myc tag was amplified and cloned into the *NotI* site of pBIJ4-GAF1. To generate GAF1 promoter-driven GAF1-GFP transgenic plants in the *gaf1 idd1* background, the GAF1 promoter was cloned into the *HindIII-PstI* site of the binary vector pBI101, and GAF1-GFP was cloned into *XbaI-SacI* site of GAF1 promoter-GUS plasmid. *Agrobacterium tumefaciens*-mediated *Arabidopsis* transformation was performed according to the floral dip method (Clough and Bent, 1998). The primer sets for cloning are listed in Supplemental Table 1. Plants were grown in a controlled growth chamber at 22°C under long days (16 h of light/8 h of dark) or short days (8 h of light/16 h of dark) with white light illumination.

Y2H Assay

The coding region of GAI was cloned into the *BamHI* site of pGBT9 vector (Clontech). The Tup1 repression domain was cloned into the *EcoRI* site of the pGBT9-GAI plasmid construct. The coding region of GAF1 was also

cloned into the *BamHI* site of pGBDU vector (Clontech). The truncated GAI was cloned into the *BamHI-PstI* site of pGBU-C1 vector (Clontech). The truncated GAF1 was generated by PCR using this plasmid as template and cloned into the *BamHI-PstI* site of pGBDU vector (Clontech). The coding regions of *IDD1* were cloned into the *BamHI-PstI* site of pGBDU vector (Clontech). The coding regions of GAI and RGA were cloned into the *BamHI* site of pGAD vector (Clontech). The coding regions of *RGL1*, *RGL2*, *RGL3*, and the N-terminal region of *TPR4* were cloned into the *SalI-BglII* site of pGAD vector (Clontech). The primers used for all cloning are listed in Supplemental Table 1. *Saccharomyces cerevisiae* PJ69-4A was cotransformed with a bait and a prey plasmid. An *Arabidopsis* seedling cDNA library (Clontech) was screened on plates containing SD/-Leu/-Trp/-His with 3-aminotriazole (50 mM). About 2000 positive colonies were restreaked on plates containing SD/-Leu/-Trp/-His with 3-aminotriazole (50 mM). The -His plate assay and β -galactosidase assay were performed according to the manufacturer's protocol for the Matchmaker Two Hybrid system (Clontech).

In Vitro Pull-Down Assay

The coding region of GAI was cloned into the *BamHI* site of pGEX4T-3 vector (GE Healthcare). The fusion proteins from *Escherichia coli* BL21 cells harboring pGEX4T-3-GAI and pGEX4T-3 plasmids were purified with Glutathione Sepharose 4B beads (GE Healthcare). The beads bound with GST-GAI and GST were washed with PBS buffer. The coding region of GAF1 was cloned into the *NotI-XhoI* site of pET30b vector. The recombinant protein from *E. coli* BL21 cells harboring pET30b-GAF1 plasmid was purified with chelating Sepharose Fast Flow (GE Healthcare). After being added to Glutathione Sepharose 4B beads carrying a GST fusion protein, purified GAF1 protein mixtures were incubated for 2 h at 4°C. After washing seven times with PBS, proteins were eluted with 10 mM reduced glutathione. Samples were separated by 10% SDS-PAGE and immunodetection with anti-GST antibody at 1:5000 dilution and anti-GAF1 antibody at 1:1000 dilution.

Application of GA₄ to Col-0, *gai-3*, and *gaf1 idd1*

Seven-day-old seedlings of Col-0, *gai-3*, and *gaf1 idd1* grown on half-strength Murashige and Skoog (MS) agar medium were transferred to soil and grown for an additional 2 months. To investigate GA sensitivity, plants were treated or not with 1 μ M GA₄. GA₄ was applied by spraying once per week for 2 months.

Transactivation Assay

The 3-kb promoters of *GA20ox2*, *RGL*, *GA3ox1*, and *GID1b* from positions -1 to -3000 (where +1 is the initiation codon) were cloned into the *SphI-SalI* site of p-less GUS vector, which is a pUC18-based plasmid containing the GUS gene cassette of pBI101 (Takahashi et al., 1995). The 3-kb promoters of *GID1a* and *MYB* from positions -1 to -3000 (where +1 is the initiation codon) were cloned into the *PstI-BamHI* site of the p-less GUS vector using the In-Fusion cloning kit (Clontech). All primers used for transient assay analysis are shown in Supplemental Table 1. GAF1, truncated GAF1, *IDD1*, GAI, and *TPR* were cloned into the *NotI-XhoI* site of the pJ4 vector carrying the CaMV 35S promoter with a viral translation enhancer, the Ω sequence (Fukazawa et al., 2000), to be used as effectors. Protoplasts were prepared from T87 *Arabidopsis* cultured cells, and transfection of protoplasts was performed as described previously (Maliga et al., 1976; Satoh et al., 2004). Relative GUS activity was calculated by the normalization of LUC activity, and the data presented are averages of three independent biological replicates.

Gel Retardation Assay

The gel retardation assays were performed following the procedure described previously (Fukazawa et al., 2000, 2010). GAF1 was cloned into the

NotI-XhoI site of pET30b vector (Novagen). Recombinant protein 6xHis-GAF1 was expressed and affinity-purified from *E. coli* BL21 (DE3) pLysE using Ni²⁺ resin (Novagen). Nucleotide sequences of the double-stranded oligonucleotides used for the gel mobility shift assays are described in Supplemental Table 1. The oligonucleotides were annealed and then labeled using [α -³²P]dCTP and the Klenow fragment of DNA polymerase I. Binding mixtures contained 50 fmol of labeled probe, 1 μ g of purified recombinant GAF1 or 1 μ g of control extract of *E. coli*, and 2 μ g of poly(dI/dC). DNA competitor was used at 100-fold molar excess. The binding buffer consisted of 20 mM Tris-HCl, pH 7.5, 3 mM MgCl₂, 50 mM KCl, 1 mM EDTA, 10% (v/v) glycerol, and 2 μ M ZnCl₂. Reactions were incubated at 4°C for 30 min and loaded onto 4% polyacrylamide gels containing 6.7 mM Tris-HCl, pH 7.5, 1 mM EDTA, and 3.3 mM sodium acetate.

Histochemical Staining

The promoter of *GAF1* (−1500 to −1) was amplified by PCR from *Arabidopsis* genomic DNA and cloned into the *HindIII-PstI* site of binary vector pBI101 to generate a *GUS* fusion gene. The primers are listed in Supplemental Table 1. Kanamycin-resistant transgenic *Arabidopsis* plants were histochemically stained to detect *GUS* activity by immersing seedlings in a staining solution (100 mM sodium phosphate buffer, pH 7.0, with 50 mM NaCl, 1 mM potassium ferricyanide, 0.1% [v/v] Triton X-100 and 1 mM 5-bromo-4-chloro-3-indolyl- β -D-glucuronide) overnight at 37°C. After staining, samples were immersed in a fixing solution (5% [v/v] formaldehyde, 5% [v/v] acetic acid, and 20% [v/v] ethanol), followed by dechlorophyllation in 70% (v/v) ethanol.

Gene Expression Analysis

Total RNA from shoot, leaves, stems, flower, and root of Col-0 were extracted using the RNeasy Mini Kit (Qiagen) according to the manufacturer's protocol. For the *ga1-3*, *ga1-3/35S:GAF1*, *ga1-3/35S: Δ PAM*, *gai-1*, *gai-1/35S:GAF1*, *gai-1/35S: Δ PAM*, and Col-0 lines, total RNA was isolated from 14-d-old plants. For the *gaf1* and *idd1* mutants and Col-0, total RNA was isolated from shoots of 10-d-old seedlings. A 1- μ g aliquot of total RNA was used for cDNA synthesis using the QuantiTect Reverse Transcription Kit (Qiagen). Quantitative RT-PCR was performed on the ABI PRISM7000 sequence detection system using the THUNDERBIRD SYBER qPCR kit (Toyobo) according to the manufacturer's instructions. *UBIQUITIN11* expression was used as an internal standard.

BiFC

The *YFP^N* and *YFP^C* sequences were generated by PCR using *EYFP* DNA (Clontech) as a template and the primers listed in Supplemental Table 1. Each PCR product was cloned into pJ4 vector (Fukazawa et al., 2000). The full-length *GAF1* cDNA was cloned into pJ4:*YFP^C*, whereas *GAI* and *TPR4* were cloned into pJ4:*YFP^N* vector. Each plasmid was introduced into T87 *Arabidopsis* protoplast cells as described above. The transfected *Arabidopsis* protoplast cells were cultured for 24 h, and the YFP fluorescence was observed using a Zeiss LSM Pascal confocal microscope.

Preparation of Antibodies

The anti-GAF1 and anti-GAI antibodies were generated against the recombinant full-length GAF1 and GAI, respectively, in rabbits. The anti-TPR4 antibody was generated against the recombinant N-terminal 242 amino acids of TPR4 in rabbits. Recombinant 6xHis-GAF1, 6xHis-GAI, and 6xHis-N-terminal-TPR4 proteins as antigens were produced using *E. coli* BL21 (DE3) pLysE using pET30 (Novagen). The coding regions for *GAI* and the N-terminal region of *TPR4* were cloned into the

BamHI and *Sall-NotI* sites of pET30a vector, respectively (Novagen). The coding region of *GAF1* was cloned into the *NotI-XhoI* site of pET30b vector (Novagen). The primer sets for cloning are listed in Supplemental Table 1.

Immunoblot Analysis

To investigate protein levels in 7- and 14-d-old seedlings of CaMV35S:GAF1 overexpressor transgenic plants, young leaves, mature leaves, flower buds, flowers, and roots of light-grown plants were harvested in liquid nitrogen. Total protein was extracted by SDS sample buffer and boiled for 10 min. For the *gai-1*, *gai-1/35S:GAF1*, *gai-1/35S: Δ PAM*, and Col-0 lines, total protein was isolated from 14-d-old plants. For the GA₃- or paclobutrazol (PAC)-treated Col-0 plants, 7-d-old Col-0 seedlings were transferred to half-strength MS plates with or without 1 μ M PAC or 1 μ M PAC and 10 μ M GA₃, and these plant were incubated for 1 week. Proteins were separated by SDS-PAGE and immunoblotting with anti-GAF1 antibody at 1:1000 dilution.

Coimmunoprecipitation Assay

Coimmunoprecipitation assays of GAF1, GAI, and TPR4 were performed with 21-d-old 35S:4 \times myc-GAF1 seedlings. Plant materials were cross-linked with 1% formaldehyde, ground in liquid nitrogen, and then extracted with NEB buffer (20 mM HEPES-KOH, pH 7.5, 40 mM KCl, 1 mM EDTA, 0.5% Triton X-100, and 1 \times protease inhibitor cocktail [Roche]) at a ratio of 2 mL buffer/g tissue. After centrifugation, the supernatant was incubated with anti-myc antibody (MBL) and Protein G Sepharose beads (GE Healthcare). Proteins bound to the beads were resolved by SDS-PAGE and then detected by anti-myc HRP conjugate (MBL), anti-GAI (homemade), or anti-TPR4 (homemade) antibody. Anti-myc HRP conjugate was used at 1:5000. Anti-GAI and anti-TPR4 antibodies were used at 1:1000 dilution.

ChIP Assay

The ChIP experiment was performed following the procedure described previously (Fukazawa et al., 2010) with some modifications. In brief, 2-week-old 4 \times myc-GAF1 transgenic or Col-0 plants were cross-linked for 10 min in 1% (v/v) formaldehyde by vacuum filtration and incubated at 4°C for 1 h. Aliquots of each protein sample were immunoprecipitated with anti-GST (Santa Cruz Biotechnology), anti-myc (MBL International), anti-GAF1, anti-GAI, and anti-TPR (homemade) antibodies for 12 h at 4°C. The chromatin-antibody complexes were precipitated with salmon sperm DNA/protein G agarose beads for 2 h at 4°C. The amount of immunoprecipitated chromatin was determined by PCR. PCR cycles were 30 cycles for the *GA20ox2* and *GID1b* promoters and 35 cycles for the *GA20ox2* coding region. The PCR products were separated on 1% (w/v) agarose gels, blotted onto Hybond XL membranes (GE Healthcare), and hybridized with gene-specific DNA probes that were labeled using the AlkPhos Direct Labeling and Detection System (GE Healthcare). After hybridization, the signals were detected using the Image Reader LAS-3000 (Fujifilm). Primers used for ChIP analysis are listed in Supplemental Table 1. The coprecipitated level of each DNA fragment was quantified by real-time PCR using specific primer sets and normalized with input DNA level. The relative coprecipitated levels using preimmune serum or anti-GST antibody (immunoprecipitated DNA/input DNA) were set to 1.

Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases under the following accession numbers: GAF1/IDD2 (At3g50700), IDD1/ENY (At5g66730), GAI

(At1g14920), RGA (At2g01570), RGL1 (At1g66350), RGL2 (At3g03450), RGL3 (At5g63970), TPR1 (At1g80490), and TPR4 (At3g15880).

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure 1. Phenotypes of *GAF1* Overexpressor and the *gaf1* Mutant.

Supplemental Figure 2. Relative Expression Levels of *GAF1* and *IDD1* in Several Plants.

Supplemental Figure 3. Complementation of *gaf1 idd1* by *GAF1pro:GAF1-GFP*.

Supplemental Figure 4. *gaf1 idd1-2* Mutant Exhibits Similar Phenotypes to the *gaf1 idd1* Mutant.

Supplemental Figure 5. Binding Domain of GAF1 for GAI and Subcellular Localization of GAF1-GFP Protein.

Supplemental Figure 6. Domain of GAF1 Binding to TPR4 and Transactivation Assay Showing That GAI and TPR4 Do Not Exhibit Transcriptional Activity without GAF1.

Supplemental Table 1. Primers Used in This Work.

ACKNOWLEDGMENTS

We thank Kiminori Toyooka (RIKEN) for helpful discussions and support with confocal microscopy analysis. We thank Belay T. Ayele (University of Manitoba) for helpful comments on the article. We also thank Rie Yoshikawa for technical assistance. This study was supported by the Japan Society for the Promotion of Science (Grant 23770058 to J.F. and Grant 21370022 to Y.T.) and the Ministry of Education, Culture, Sports, Science, and Technology of Japan (Grant 24116525 to J.F. and Grant 24118004 to Y.T.).

AUTHOR CONTRIBUTIONS

J.F., Y.K., S.Y., and Y.T. designed the research. J.F., H.T., S.M., N.N., and K.N. performed research. J.F., M.Y., T.L., S.Y., and Y.T. analyzed data. J.F. and Y.T. wrote the article.

Received March 26, 2014; revised June 11, 2014; accepted June 25, 2014; published July 17, 2014.

REFERENCES

- Bai, M.Y., Shang, J.X., Oh, E., Fan, M., Bai, Y., Zentella, R., Sun, T.P., and Wang, Z.Y. (2012). Brassinosteroid, gibberellin and phytochrome impinge on a common transcription module in *Arabidopsis*. *Nat. Cell Biol.* **14**: 810–817.
- Cheng, H., Qin, L., Lee, S., Fu, X., Richards, D.E., Cao, D., Luo, D., Harberd, N.P., and Peng, J. (2004). Gibberellin regulates *Arabidopsis* floral development via suppression of DELLA protein function. *Development* **131**: 1055–1064.
- Clough, S.J., and Bent, A.F. (1998). Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**: 735–743.
- Colasanti, J., Tremblay, R., Wong, A.Y., Coneva, V., Kozaki, A., and Mable, B.K. (2006). The maize INDETERMINATE1 flowering time regulator defines a highly conserved zinc finger protein family in higher plants. *BMC Genomics* **7**: 158.
- Colasanti, J., Yuan, Z., and Sundaresan, V. (1998). The indeterminate gene encodes a zinc finger protein and regulates a leaf-generated signal required for the transition to flowering in maize. *Cell* **93**: 593–603.
- de Lucas, M., Davière, J.M., Rodríguez-Falcón, M., Pontin, M., Iglesias-Pedraz, J.M., Lorrain, S., Fankhauser, C., Blázquez, M.A., Titarenko, E., and Prat, S. (2008). A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480–484.
- Feng, S., et al. (2008). Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**: 475–479.
- Feurtado, J.A., Huang, D., Wicki-Stordeur, L., Hemstock, L.E., Potentier, M.S., Tsang, E.W., and Cutler, A.J. (2011). The *Arabidopsis* C2H2 zinc finger INDETERMINATE DOMAIN1/ENHYDROUS promotes the transition to germination by regulating light and hormonal signaling during seed maturation. *Plant Cell* **23**: 1772–1794.
- Fukazawa, J., Nakata, M., Ito, T., Yamaguchi, S., and Takahashi, Y. (2010). The transcription factor RSG regulates negative feedback of NtGA20ox1 encoding GA 20-oxidase. *Plant J.* **62**: 1035–1045.
- Fukazawa, J., Sakai, T., Ishida, S., Yamaguchi, I., Kamiya, Y., and Takahashi, Y. (2000). Repression of shoot growth, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. *Plant Cell* **12**: 901–915.
- Gallego-Bartolomé, J., Minguet, E.G., Grau-Enguix, F., Abbas, M., Locascio, A., Thomas, S.G., Alabadi, D., and Blázquez, M.A. (2012). Molecular mechanism for the interaction between gibberellin and brassinosteroid signaling pathways in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **109**: 13446–13451.
- Hauvermale, A.L., Ariizumi, T., and Steber, C.M. (2012). Gibberellin signaling: A theme and variations on DELLA repression. *Plant Physiol.* **160**: 83–92.
- Hirano, K., Kouketu, E., Katoh, H., Aya, K., Ueguchi-Tanaka, M., and Matsuoka, M. (2012). The suppressive function of the rice DELLA protein SLR1 is dependent on its transcriptional activation activity. *Plant J.* **71**: 443–453.
- Hirst, M., Ho, C., Sabourin, L., Rudnicki, M., Penn, L., and Sadowski, I. (2001). A two-hybrid system for transactivator bait proteins. *Proc. Natl. Acad. Sci. USA* **98**: 8726–8731.
- Hong, G.J., Xue, X.Y., Mao, Y.B., Wang, L.J., and Chen, X.Y. (2012). *Arabidopsis* MYC2 interacts with DELLA proteins in regulating sesquiterpene synthase gene expression. *Plant Cell* **24**: 2635–2648.
- Hou, X., Lee, L.Y., Xia, K., Yan, Y., and Yu, H. (2010). DELLAs modulate jasmonate signaling via competitive binding to JAZs. *Dev. Cell* **19**: 884–894.
- Ikeda, A., Ueguchi-Tanaka, M., Sonoda, Y., Kitano, H., Koshioka, M., Futsuhara, Y., Matsuoka, M., and Yamaguchi, J. (2001). *slender rice*, a constitutive gibberellin response mutant, is caused by a null mutation of the *SLR1* gene, an ortholog of the height-regulating gene *GAI/RGA/RHT/D8*. *Plant Cell* **13**: 999–1010.
- Itoh, H., Ueguchi-Tanaka, M., Sato, Y., Ashikari, M., and Matsuoka, M. (2002). The gibberellin signaling pathway is regulated by the appearance and disappearance of SLENDER RICE1 in nuclei. *Plant Cell* **14**: 57–70.
- Jabet, C., Sprague, E.R., VanDemark, A.P., and Wolberger, C. (2000). Characterization of the N-terminal domain of the yeast transcriptional repressor Tup1. Proposal for an association model of the repressor complex Tup1 x Ssn6. *J. Biol. Chem.* **275**: 9011–9018.
- Kozaki, A., Hake, S., and Colasanti, J. (2004). The maize ID1 flowering time regulator is a zinc finger protein with novel DNA binding properties. *Nucleic Acids Res.* **32**: 1710–1720.
- Krogan, N.T., Hogan, K., and Long, J.A. (2012). APETALA2 negatively regulates multiple floral organ identity genes in *Arabidopsis* by recruiting

- the co-repressor TOPLESS and the histone deacetylase HDA19. *Development* **139**: 4180–4190.
- Lim, S., Park, J., Lee, N., Jeong, J., Toh, S., Watanabe, A., Kim, J., Kang, H., Kim, D.H., Kawakami, N., and Choi, G.** (2013). ABA-INSENSITIVE3, ABA-INSENSITIVE5, and DELLAs interact to activate the expression of *SOMNUS* and other high-temperature-inducible genes in imbibed seeds in *Arabidopsis*. *Plant Cell* **25**: 4863–4878.
- Liu, Z., and Karmarkar, V.** (2008). Groucho/Top1 family co-repressors in plant development. *Trends Plant Sci.* **13**: 137–144.
- Long, J.A., Ohno, C., Smith, Z.R., and Meyerowitz, E.M.** (2006). TOPLESS regulates apical embryonic fate in *Arabidopsis*. *Science* **312**: 1520–1523.
- Maliga, P., Lázár, G., Sváb, Z., and Nagy, F.** (1976). Transient cycloheximide resistance in a tobacco cell line. *Mol. Gen. Genet.* **149**: 267–271.
- Mutasa-Göttgens, E., and Hedden, P.** (2009). Gibberellin as a factor in floral regulatory networks. *J. Exp. Bot.* **60**: 1979–1989.
- Park, J., Nguyen, K.T., Park, E., Jeon, J.S., and Choi, G.** (2013). DELLA proteins and their interacting RING finger proteins repress gibberellin responses by binding to the promoters of a subset of gibberellin-responsive genes in *Arabidopsis*. *Plant Cell* **25**: 927–943.
- Pauwels, L., et al.** (2010). NINJA connects the co-repressor TOPLESS to jasmonate signalling. *Nature* **464**: 788–791.
- Richards, D.E., Peng, J., and Harberd, N.P.** (2000). Plant GRAS and metazoan STATs: One family? *BioEssays* **22**: 573–577.
- Rosenfeld, M.G., Lunnyak, V.V., and Glass, C.K.** (2006). Sensors and signals: A coactivator/corepressor/epigenetic code for integrating signal-dependent programs of transcriptional response. *Genes Dev.* **20**: 1405–1428.
- Satoh, R., Fujita, Y., Nakashima, K., Shinozaki, K., and Yamaguchi-Shinozaki, K.** (2004). A novel subgroup of bZIP proteins functions as transcriptional activators in hypoosmolarity-responsive expression of the ProDH gene in *Arabidopsis*. *Plant Cell Physiol.* **45**: 309–317.
- Singleton, W.R.** (1946). Inheritance of indeterminate growth in maize. *J. Hered.* **37**: 61–64.
- Sun, T.P.** (2010). Gibberellin-GID1-DELLA: A pivotal regulatory module for plant growth and development. *Plant Physiol.* **154**: 567–570.
- Sun, T.P.** (2011). The molecular mechanism and evolution of the GA-GID1-DELLA signaling module in plants. *Curr. Biol.* **21**: R338–R345.
- Sun, T.P., and Gubler, F.** (2004). Molecular mechanism of gibberellin signaling in plants. *Annu. Rev. Plant Biol.* **55**: 197–223.
- Szemenyei, H., Hannon, M., and Long, J.A.** (2008). TOPLESS mediates auxin-dependent transcriptional repression during *Arabidopsis* embryogenesis. *Science* **319**: 1384–1386.
- Takahashi, Y., Sakai, T., Ishida, S., and Nagata, T.** (1995). Identification of auxin-responsive elements of parB and their expression in apices of shoot and root. *Proc. Natl. Acad. Sci. USA* **92**: 6359–6363.
- Tiwari, S.B., Hagen, G., and Guilfoyle, T.J.** (2004). Aux/IAA proteins contain a potent transcriptional repression domain. *Plant Cell* **16**: 533–543.
- Tyler, L., Thomas, S.G., Hu, J., Dill, A., Alonso, J.M., Ecker, J.R., and Sun, T.P.** (2004). DELLA proteins and gibberellin-regulated seed germination and floral development in *Arabidopsis*. *Plant Physiol.* **135**: 1008–1019.
- Ueguchi-Tanaka, M., Nakajima, M., Katoh, E., Ohmiya, H., Asano, K., Saji, S., Hongyu, X., Ashikari, M., Kitano, H., Yamaguchi, I., and Matsuoka, M.** (2007a). Molecular interactions of a soluble gibberellin receptor, GID1, with a rice DELLA protein, SLR1, and gibberellin. *Plant Cell* **19**: 2140–2155.
- Ueguchi-Tanaka, M., Nakajima, M., Motoyuki, A., and Matsuoka, M.** (2007b). Gibberellin receptor and its role in gibberellin signaling in plants. *Annu. Rev. Plant Biol.* **58**: 183–198.
- Wang, L., Kim, J., and Somers, D.E.** (2013). Transcriptional corepressor TOPLESS complexes with pseudoresponse regulator proteins and histone deacetylases to regulate circadian transcription. *Proc. Natl. Acad. Sci. USA* **110**: 761–766.
- Yamaguchi, S.** (2008). Gibberellin metabolism and its regulation. *Annu. Rev. Plant Biol.* **59**: 225–251.
- Zentella, R., Zhang, Z.L., Park, M., Thomas, S.G., Endo, A., Murase, K., Fleet, C.M., Jikumaru, Y., Nambara, E., Kamiya, Y., and Sun, T.P.** (2007). Global analysis of DELLA direct targets in early gibberellin signaling in *Arabidopsis*. *Plant Cell* **19**: 3037–3057.