

WJG 20<sup>th</sup> Anniversary Special Issues (14): Pancreatic cancer

## Therapeutic options for the management of pancreatic cancer

Maria L Rossi, Azeem A Rehman, Christopher S Gondi

Maria L Rossi, Azeem A Rehman, Christopher S Gondi, Department of Medicine, University of Illinois College of Medicine-Peoria, Peoria, IL 61656-1649, United States

Author contributions: Rossi ML and Gondi CS provided the conception and design the article; Rossi ML, Rehman AA and Gondi CS drafted the article; Rossi ML and Gondi CS revised the critically important content, and provided the final approval of the version to be published.

Supported by Grant from the William E McElroy Foundation, Springfield, IL 62794-9264, United States to Gondi CS; and the Carrie M. Craig Medical Student Summer Research Fellowship Program sponsored by the University of Illinois College of Medicine at Peoria, Box 1649, Peoria, IL 61656-1649, United States to Rossi ML

Correspondence to: Christopher S Gondi, PhD, Assistant Professor, Department of Medicine, University of Illinois College of Medicine-Peoria, Box 1649, Peoria, IL 61656-1649, United States. [gondi@uic.edu](mailto:gondi@uic.edu)

Telephone: +1-309-4958167 Fax: +1-309-6557732

Received: October 25, 2013 Revised: January 11, 2014

Accepted: May 28, 2014

Published online: August 28, 2014

### Abstract

Since its initial characterization, pancreatic ductal adenocarcinoma has remained one of the most devastating and difficult cancers to treat. Pancreatic cancer is the fourth leading cause of death in the United States, resulting in an estimated 38460 deaths annually. With few screening tools available to detect this disease at an early stage, 94% of patients will die within five years of diagnosis. Despite decades of research that have led to a better understanding of the molecular and cellular signaling pathways in pancreatic cancer cells, few effective therapies have been developed to target these pathways. Other treatment options have included more sophisticated pancreatic cancer surgeries and combination therapies. While outcomes have improved modestly for these patients, more effective treatments

are desperately needed. One of the greatest challenges in the future of treating this malignancy will be to develop therapies that target the tumor microenvironment and surrounding pancreatic cancer stem cells in addition to pancreatic cancer cells. Recent advances in targeting pancreatic stellate cells and the stroma have encouraged researchers to shift their focus to the role of desmoplasia in pancreatic cancer pathobiology in the hopes of developing newer-generation therapies. By combining novel agents with current cytotoxic chemotherapies and radiation therapy and personalizing them to each patient based on specific biomarkers, the goal of prolonging a patient's life could be achieved. Here we review the most effective therapies that have been used for the treatment of pancreatic cancer and discuss the future potential of therapeutic options.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

**Key words:** Pancreatic cancer; Pancreatic cancer stem cells; Microenvironment; Surgical resection; Neoadjuvant therapy; Adjuvant therapy; Chemotherapy; Radiation; Personalized therapy

**Core tip:** Pancreatic ductal adenocarcinoma has challenged researchers for decades. It remains one of the most deadly cancers due to the complex molecular and genetic makeup of its cancer cells and their surrounding microenvironment. In addition, there are no valid screening tests available to detect pancreatic cancer in its early stages. Yet, as knowledge of this cancer has evolved over time, so have novel methods for treating it. Researchers have a deeper understanding of pancreatic cancer now than ever before. The future holds much promise for new breakthroughs that will significantly improve patient outcomes.

Rossi ML, Rehman AA, Gondi CS. Therapeutic options for the management of pancreatic cancer. *World J Gastroenterol* 2014;

20(32): 11142-11159 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v20/i32/11142.htm> DOI: <http://dx.doi.org/10.3748/wjg.v20.i32.11142>

## INTRODUCTION

Despite the improved survival rates noted in numerous cancers, including breast<sup>[1-3]</sup>, prostate<sup>[4]</sup> and colon cancer<sup>[5]</sup>, the overall survival rates for patients diagnosed with pancreatic cancer have shown little improvement over the past thirty years<sup>[6-8]</sup>. Pancreatic ductal adenocarcinoma (PDA) remains one of the most rapidly progressive and deadly malignancies worldwide<sup>[4]</sup>. The prevention of pancreatic cancer is difficult to assess, due to limited studies identifying potential risk factors compounded with the multifactorial, heterogeneous nature of the disease. Cigarette smoking has been noted to double the risk of pancreatic cancer, yet only accounts for 20%-25% of the cases<sup>[9,10]</sup>. Additionally, family history may also contribute a significant role as 5%-10% of individuals with pancreatic cancer report an incidence of pancreatic cancer in a close family member<sup>[11]</sup>. This risk is further substantiated when there is a larger number of family members with pancreatic cancer and a decrease in age of onset in kindred<sup>[12]</sup>. Other noted risk factors include alcohol abuse<sup>[13]</sup>, a high-fat diet<sup>[14,15]</sup>, and certain trace elements<sup>[16]</sup>.

The challenge of diagnosing PDA at an early stage further contributes to the dismal five-year survival rate that is projected for patients. Located in the retroperitoneum of patients who present with non-specific symptoms, PDA is not diagnosed until it has reached an advanced clinical stage in 80% of patients<sup>[17]</sup>. In addition, lack of effective screening and early biomarker detection methods have prevented clinicians from identifying this cancer in a pre-malignant stage. Ideally, visual evaluation *via* computerized tomography (CT) and magnetic resonance imaging (MRI) should be incorporated upon suspicion of pancreatic cancer for detection and resectability assessment<sup>[18]</sup>. Although CT scan has often been utilized to detect pancreatic cancer<sup>[19-21]</sup>, reliance on MRI, particularly in regard to assessing local invasion and metastasis, has increased<sup>[22]</sup>. Other imaging may also provide certain benefits, such as endoscopic ultrasound for investigating vascular invasion<sup>[23]</sup>, fludeoxyglucose-positron emission tomography scanning for recurrent tumors<sup>[24]</sup>, and laparoscopy for more accurate staging<sup>[25]</sup>. While the use of these techniques remains helpful to determine prognosis and treatment regimen for patients diagnosed with pancreatic cancer, none have been validated as effective screening tests for general or high risk populations.

Once diagnosed, a number of chemotherapy, radiation and combination therapy regimens have been used to treat patients with ductal pancreatic tumors. Unfortunately, the dynamic molecular and cellular makeup of individual pancreatic tumors, renders many of them resistant to the majority of these therapies. Although surgical resection has been shown to increase patient survival

by 10 mo<sup>[26]</sup>, the majority of patients who undergo these procedures experience comorbidities and recurrence. Current research has identified additional sources of therapeutic resistance in the microenvironment of these tumors. Characterized by stromal proliferation, reduced angiogenesis and a unique subset of cells known as cancer stem cells (CSCs), the tumor microenvironment has become a target of new therapeutic agents.

While improved understanding of pancreatic cancer biology has lead to several therapeutic breakthroughs in the treatment of PDA, major progress toward improving survival rates in patients has been extremely slow. However, as our understanding of this tumor's therapeutic resistant nature improves, so will future progress in treating pancreatic cancer.

## CLINICAL PRESENTATION AND DIAGNOSIS

One of the greatest challenges in treating pancreatic ductal adenocarcinoma (PDA) is discovering it in the pre-malignant stage. The average patient diagnosed with pancreatic cancer is in their seventh decade of life and presents to their primary care physician with general symptoms of abdominal pain and weight loss<sup>[27]</sup>. Not only is the pancreas difficult to palpate due to its retroperitoneal location, but there are also no specific blood tests to confirm suspicion of malignancy. More specific symptoms, such as unexplained jaundice<sup>[28]</sup>, onset of diabetes<sup>[29]</sup> and development of thromboembolic disease<sup>[30]</sup> are more diagnostic of pancreatic cancer, but do not present until later stages of the disease. The primary comorbidities associated with PDA include biliary obstruction, infection, ascites, pancreatic insufficiency and in advanced stages of the disease, cachexia<sup>[31]</sup>. Unfortunately, once a patient presents with these symptoms, the disease has often already reached its malignant stage and the patient may never be able to receive treatment.

Effective screening tests to provide early diagnosis of pancreatic cancer could potentially prevent these symptoms. The ones that do exist are not validated. For example, although endoscopic ultrasounds provide a higher yield of detecting pancreatic cancer in its early stages, the comorbidities associated with this procedure render it an unsuitable screening test in the general population. As a result, studies are currently underway to identify high risk individuals who may benefit from this invasive procedure<sup>[32-34]</sup>. Other techniques, such as cross-sectional imaging could be used to identify asymptomatic pancreatic neoplasms for surgical resection<sup>[34]</sup> as long as they are confirmed by CT or MRI techniques which provide better resolution between normal and neoplastic pancreatic tissue<sup>[35]</sup>.

Although a greater understanding of the molecular events in pancreatic cancer tumorigenesis has lead to the discovery of biomarkers that help to predict the tumor's response to treatment, there has been no use of these markers in cancer drug development<sup>[36]</sup>. The only biomarker

**Table 1 Cellular mechanisms of therapeutic resistance in pancreatic cancer**

Cellular pathways	Mutated gene	Ref.
Cell-cycle control	CDK2NA (90%); APC2	Almoguera <i>et al</i> <sup>[54]</sup> ; Schutte <i>et al</i> <sup>[71]</sup> ; Hahn <i>et al</i> <sup>[72]</sup>
RAS	KRAS (90%); MAP2K4	Almoguera <i>et al</i> <sup>[54]</sup> ; Hruban <i>et al</i> <sup>[56]</sup> ; Pellegata <i>et al</i> <sup>[57]</sup> ; Hezel <i>et al</i> <sup>[58]</sup> ; Maitra <i>et al</i> <sup>[59]</sup>
DNA damage repair	TP53 (75%-90%)	Almoguera <i>et al</i> <sup>[54]</sup> ; Redston <i>et al</i> <sup>[67]</sup> ; Olive <i>et al</i> <sup>[68]</sup>
TGF-β	DPC4 (50%), SMAD4	Almoguera <i>et al</i> <sup>[54]</sup> ; Yachida <i>et al</i> <sup>[73]</sup>
Apoptosis	CASP10; CAD	Jones <i>et al</i> <sup>[77]</sup>
Cell adhesion	FAT; PCDH9	Jones <i>et al</i> <sup>[77]</sup>
Hedgehog	GLI1; GLI3	Jones <i>et al</i> <sup>[77]</sup>
Integrin	ILK; LAMA1	Jones <i>et al</i> <sup>[77]</sup>
JNK	MAP4K3; TNF	Jones <i>et al</i> <sup>[77]</sup>
Small GTPases	PLCB3; RP1	Jones <i>et al</i> <sup>[77]</sup>
Wnt-β-catenin	MYC; TSC2	Jones <i>et al</i> <sup>[77]</sup>

APC: Adenomatous polyposis coli; CDK2NA: Cyclin-dependent kinase inhibitor 2 A; CAD: Carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase; CASP10: Caspase 10; DPC4: deleted in pancreatic cancer, locus 4; FAT: Fat tumor suppressor; GLI 1: Glioma-associated oncogene; GLI3: Glioma-associated oncogene 3; GTPases: Guanosine triphosphate; Wnt-B-catenin: Wingless type B-catenin; ILK: Integrin-linked kinase; JNK: c-Jun N-terminal kinases; KRAS: Kristen rat sarcoma; LAMA1: Laminin A-1 chain; MAP2K4: Mitogen-activated protein kinase kinase 4; MAP4K3: Mitogen-activated protein-3 kinase-3; TNF: Tumor necrosis factor; MYC: Myelocytomatosis oncogene; PCDH9: Protocadherin 9; PLCB3: Phospholipase C, beta 3; RP1: Retinitis pigmentosa 1; SMAD4: Mothers against decapentaplegic homolog 4; TGF-β: Transforming growth factor β; TP53: Tumor protein 53; TSC2: Tuberous sclerosis 2; RAS: Rat sarcoma.

that has shown a great deal of promise in therapeutic monitoring and in identifying the recurrence of pancreatic cancer after treatment is the carbohydrate antigen 19-9 (CA 19-9), a sialylated Lewis blood group A antigen secreted by many pancreatic lesions<sup>[37]</sup>. Yet, CA 19-9 is not specific for pancreatic cancer and therefore cannot be used to screen for this tumor. Several other conditions, including hepatobiliary diseases, pancreatic diseases and gastrointestinal malignancies, bronchitis, congestive heart failure, cystic fibrosis, diverticulitis, lung cancer, ovarian cysts, pleural effusions, renal cysts and rheumatoid arthritis may present with elevated levels of CA 19-9<sup>[38]</sup>. In addition, approximately 10% of patients with pancreatic cancer are negative for Lewis antigen a or b. As a result, these patients are unable to synthesize CA 19-9 and will have no detectable levels of this biomarker, even in advanced stages of pancreatic cancer<sup>[39]</sup>. Although measurement of serum CA 19-9 levels has clinical significance in determining treatment and prognosis for patients with known pancreatic cancer, its usefulness as a diagnostic screening tool of the disease is not substantiated<sup>[40]</sup>.

Upon diagnosis of pancreatic cancer, treatment and management of patients should utilize a multidisciplinary team including, primary care physicians, medical oncolo-

gists, radiation oncologists, surgeons, endocrinologists, radiologists and pathologists<sup>[41]</sup>. Cancer staging subsequently follows, with the American Joint Committee on Cancer providing the standard model, based upon the tumor-node-metastasis system<sup>[42]</sup>. However, not all criteria regarding tumor staging can be measured prior to surgical intervention. As a result, the majority of staging for pancreatic cancer incorporates imaging results and liver function tests. From these results, patients with pancreatic cancer can often be classified into three major cohorts: (1) patients with a resectable tumor or borderline resectable tumor; (2) patients with a locally invasive tumor without metastasis; and (3) patients with a systemically disseminated tumor.

Appropriately designating cases into the proper subgroup is vital to ensure appropriate treatment and management of patients presenting with pancreatic cancer. Often, fine needle aspiration guided by endoscopic ultrasonography is necessary to obtain histological confirmation<sup>[43,44]</sup>, especially prior to the initiation of chemotherapy and radiation. Throughout the treatment process, CA 19-9 should be continuously measured<sup>[45,46]</sup>. Nonetheless, previous studies still support the usefulness of CA 19-9 in predicting patient response to chemotherapy<sup>[47,48]</sup>, preoperative prognosis<sup>[49]</sup>, as well as assessing treatment response<sup>[50,51]</sup>, overall survival<sup>[51-53]</sup>, and recurrence<sup>[51]</sup>.

## THERAPEUTIC RESISTANCE IN PANCREATIC CANCER

### Cellular mechanisms of therapeutic resistance

In an effort to understand the therapeutic resistant (Table 1) nature of PDA, researchers have attempted to characterize the molecular and cellular components of the pancreatic cancer cells as well as the desmoplasia that surrounds them. Although pancreatic cancer displays pathologic and clinical heterogeneity, data suggests the majority of PDA express a successive accumulation of highly penetrant genetic alterations that occur at four genetic loci: K-ras, p53, cdkn2a and smad4/DPC4<sup>[54]</sup>. Originating in the ductal epithelium, pancreatic cancer evolves from a premalignant lesion to a highly invasive metastatic disease<sup>[55]</sup>.

Ninety percent of tumors have point mutations that are specific for the KRAS2 oncogene, resulting in the constitutive production of the Ras protein<sup>[56-59]</sup>. Occurring early in tumorigenesis, these point mutations are essential for maintaining the malignant phenotype because once activated, Ras initiates a signal transduction cascade that activates proliferative and cell survival pathways and increases cell invasion<sup>[60,61]</sup>. The majority of the point mutations occur on codon 12 of the ras protein and give rise to pancreatic tumor-specific neo-antigens. Several studies demonstrated these antigens are recognized by helper T-cells and cytotoxic T-cells<sup>[62,63]</sup>. Using this knowledge, scientists developed personalized peptide vaccines corresponding to the K-ras mutations present in the tumors of patients enrolled in one clinical trial<sup>[64]</sup>. The vaccine was

**Table 2** Extracellular mechanisms of therapeutic resistance in pancreatic cancer

Potential therapeutic targets	Extracellular response	Ref.
K-ras mutant oncogene	Proliferation of desmoplastic reaction (leukocytes, fibroblasts, endothelial cells, neuronal cells, collagen, hyaluron): upregulation of GM-CSF	Chu <i>et al</i> <sup>[78]</sup> ; Neesse <i>et al</i> <sup>[79]</sup> ; Ying <i>et al</i> <sup>[81]</sup> ; Nolan-Stevaux <i>et al</i> <sup>[82]</sup> ; Bayne <i>et al</i> <sup>[87]</sup>
Sonic hedgehog (SHH)	Growth and differentiation of stromal fibroblasts	Bailey <i>et al</i> <sup>[83]</sup> ; Tian <i>et al</i> <sup>[84]</sup> ; Olive <i>et al</i> <sup>[85]</sup>
Tumor associated macrophages (TAMs); cancer associated fibroblasts (CAFs); regulatory T-cells (Treg); myeloid derived suppressor cells	Evasion of the immune system	Bayne <i>et al</i> <sup>[87]</sup> ; Pylayeva-Gupta <i>et al</i> <sup>[88]</sup>
Desmoplastic reaction	Anti-angiogenesis; hypoxic tumor environment	Komar <i>et al</i> <sup>[86]</sup>

K-ras: Kinase-rat sarcoma; GM-CSF: Granulocyte macrophage colony-stimulating factor.

proven safe and tolerable and resulted in a more efficient immunologic attack on the tumor<sup>[65]</sup>. As a result, patients given the vaccine demonstrated a more favorable clinical outcome than those not given the vaccine. Combined with surgical resection, a long-term immune response initiated by the K-ras vaccine has resulted in a 10-year survival in some patients. These results may implicate a role for the K-ras vaccine as an adjuvant treatment option in the future<sup>[66,67]</sup>.

Rather than being activated like the mutated KRAS2 oncogene, the p53 tumor suppressor gene is inactivated in 75%-90% of pancreatic tumors<sup>[68,69]</sup>. As a result, there is an impaired response to DNA damage in pancreatic epithelial cells, impaired apoptosis and impaired cell cycle control<sup>[70,71]</sup>. Two other tumor suppressor genes, *p16Ink4a* and *p15ARF* are encoded by the *cdkn2a* locus. Inactivation mutations in these genes are present in about 90% of human PDA<sup>[72,73]</sup>.

A fourth common mutation seen in more than half of pancreatic cancers causes an alteration in DPC4<sup>[74]</sup>. These mutations confer a metastatic phenotype. The genetic makeup of the patient determines the number and combination of these mutations that will be present in their PDA. Patients with three or four mutated genes will have a much worse prognosis than those with one or two. The variable expressivity of these tumors presents a challenge in effectively treating them<sup>[75]</sup>.

In addition to these four primary genetic alterations in PDA many other less-frequent mutations occur as well<sup>[76-78]</sup>. According to a comprehensive genetic analysis of 24 pancreatic cancers, an average of more than 60 genetic abnormalities, primarily point mutations, per tu-

mor were noted in the PDA phenotype. While these mutations have been organized into 12 functional cancer-relevant pathways, not all tumors have alterations in each of these pathways. In addition, key mutations in select pathways appear to differ from one tumor to another<sup>[78]</sup>. These pathways may confer therapeutic resistance in the pancreatic tumor. Significant genomic instability in pancreatic cancer may also reduce the effectiveness of therapeutic agents by contributing to acquired chemoresistance.

### Extracellular mechanisms of therapeutic resistance

Paracrine signals from pancreatic cancer cells stimulate the extracellular proliferation of leukocytes, fibroblasts, endothelial cells, neuronal cells, collagen and hyaluron (Table 2). This extracellular proliferation of cells is known as a desmoplastic reaction. It forms a thick stromal environment around the pancreatic cancer cells<sup>[79,80]</sup>. Studies have demonstrated that the signals that influence the proliferation of the desmoplastic reaction originate from the K-ras mutant oncogene in the epithelium of the pancreatic cancer cells<sup>[81,82]</sup>.

In addition to the K-ras mutant signaling pathway, there has been an effort by researchers to understand the roles of other signaling pathways between the pancreatic cancer cells and their microenvironment. Sonic hedgehog (SHH) functions similarly to the K-ras mutant. Although it is over expressed in pancreatic cancer cells during the early stages of their development, SHH does not act on the SHH pathway in these cells<sup>[83]</sup>. Instead, it acts in a paracrine fashion in the extracellular fibroblasts, resulting in their growth and differentiation<sup>[84,85]</sup>.

The desmoplastic reaction not only provides a mechanical barrier to the pancreatic cancer cells, but it is also thought to contribute to the anti-angiogenic environment that is characteristic of pancreatic ductal adenocarcinoma. Both properties directly affect therapeutic efficacy. Inadequate drug delivery to the site of the tumor is directly correlated with a negative patient outcome<sup>[86]</sup>.

Researchers have also suggested a role for the tumor stroma in the T-lymphocyte depleted microenvironment of the PDA. Several cell types found in the desmoplastic reaction have been associated with tumor associated macrophages, cancer associated fibroblasts, regulatory T-cells and myeloid derived suppressor cells. In addition, a role for a K-ras dependent signaling molecule has been shown to up-regulate granulocyte-macrophage colony stimulating factor when activated. This cytokine promotes the maturation of immature myeloid progenitor cells into myeloid derived suppressor cells<sup>[87,88]</sup>.

## TREATMENT OF PANCREATIC CANCER

### Surgical resection

Although surgical resection offers hope for curative therapy, only 20% of patients present with potentially resectable tumors<sup>[89,90]</sup>. It is important to note that surgical resection is only considered in patients with completely



**Table 3 Therapies for the management of pancreatic cancer**

Therapeutic option	Subset	Ref.
Surgical resection	Cephalic pancreatoduodenectomy Distal pancreatectomy Total pancreatectomy	Hidalgo <sup>[41]</sup>
Chemotherapy	Neoadjuvant	Lemmens <i>et al</i> <sup>[101]</sup> ; Gillen <i>et al</i> <sup>[102]</sup> ;
	Gemcitabine	Neoptolemos <i>et al</i> <sup>[108]</sup> ; Burris <i>et al</i> <sup>[118]</sup> ;
	Adjuvant	Heinemann <i>et al</i> <sup>[119]</sup> ; Reni <i>et al</i> <sup>[120]</sup> ;
	Gemcitabine	Moore <i>et al</i> <sup>[122]</sup> ; Neesse <i>et al</i> <sup>[79]</sup>
	5-Fluorouracil	
	Advanced disease	
	Gemcitabine	
	Gemcitabine + fluoropyrimidines	
	Gemcitabine + platinum analogs	
	Gemcitabine + erlotinib	
Radiation therapy	FOLFIRINOX	
	Nab-paclitaxel	
	Neoadjuvant	Pisters <i>et al</i> <sup>[131]</sup> ; Hong <i>et al</i> <sup>[133]</sup> ;
	Radiation + 5-fluorouracil	Yeo <i>et al</i> <sup>[140]</sup> ;
	Radiation + paclitaxel	Regine <i>et al</i> <sup>[138]</sup> ;
	Proton beam radiation + capecitabine	Neoptolemos <i>et al</i> <sup>[137]</sup> ; Moertel <i>et al</i> <sup>[144]</sup> ;
	Adjuvant	Schellenberg <i>et al</i> <sup>[147]</sup>
	Radiation + 5-Fluorouracil	
	Radiation + gemcitabine	
	Radiation + chemotherapy	
Personalized therapy	Advanced	
	Radiation + 5-fluorouracil	
	Radiation + chemotherapy	
	Stereotactic body radiotherapy	
Personalized therapy	Target specific point mutations	Jones <i>et al</i> <sup>[77]</sup> ;
	Mitomycin C	Villarroel <i>et al</i> <sup>[153]</sup> ;
	Immune system stimulation	Yanagimoto <i>et al</i> <sup>[154]</sup>

resectable or borderline-resectable tumors (Table 3). Depending on the size and location of the tumor, three operative procedures are potentially utilized, as noted by Hidalgo<sup>[41]</sup>, with additional removal of adjacent lymph nodes: (1) cephalic pancreatoduodenectomy (whipple procedure); (2) distal pancreatectomy; and (3) total pancreatectomy.

Although additional palliative care is often utilized, controversy surrounds the potential benefits. For example, almost 80% of patients presenting with tumors in the pancreatic head exhibit jaundice due to biliary obstruction<sup>[91,92]</sup>. However, previous investigations have conflicting results regarding preoperative biliary drainage with certain studies reporting a decrease in perioperative morbidity and mortality<sup>[93]</sup> while others concluding recognizable benefit<sup>[94,95]</sup>.

Preoperative biliary stenting doubled between 1992 and 2007 due to evidence demonstrating a higher risk of postoperative complications in patients presenting with a tumor in the head of the pancreas. Biliary drainage was further supported by evidence demonstrating its ability to improve liver function, nutritional status and cell-mediated immune function<sup>[93]</sup>. Despite intentions to reduce post-operative morbidity and mortality by improving liver function, extensive clinical studies have demonstrated preoperative biliary stenting prolongs time to operation, increases preoperative infection and is associated with overall increased complication rates after surgical proce-

dures<sup>[94,95]</sup>. As a result, most studies have advised against routinely performing preoperative biliary drainage and have recommended that patients presenting with jaundice due to a resectable and non-metastatic tumor in the head of the pancreas should undergo early surgery without preoperative biliary resection<sup>[95]</sup>. Currently, the only indications for preoperative biliary decompression are for patients who present with severe jaundice, are undergoing neoadjuvant therapy, or have had their surgery postponed due to logistics<sup>[94,95]</sup>.

Several poor predictors for successful resection have been identified, including lymph node involvement<sup>[96]</sup>, high tumor grade<sup>[97]</sup>, large tumor size<sup>[98]</sup>, elevated CA 19-9 levels<sup>[46]</sup>, and positive margins of tumor following resection<sup>[41]</sup>. These same factors also contribute to recurrence of pancreatic tumors. Even with surgical resection, 5-year survival rates remain dismal, at approximately 20% following surgery<sup>[90]</sup>. However, perioperative complications and mortality have significantly decreased over the past decade, likely due to greater hospital clinical volume through centralization<sup>[99,100]</sup>.

**Chemotherapy**

**Neoadjuvant therapy:** Certain patients might receive neoadjuvant therapy, especially if the tumor presents with borderline resectability. In a study utilizing gemcitabine-based chemotherapy, improved tumor resection with increased survival rates was noted in border-line resect-

able cases<sup>[101,102]</sup>. However, these effects may only occur in select tumors, with influences by both the genetic composition and microenvironment of the pancreatic cancer<sup>[103,104]</sup>. For example, White *et al.*<sup>[105]</sup> noted p53 mutations were more common in patients with large residual tumors following treatment with chemoradiation. Moreover, outcomes for neoadjuvant therapy prior to surgically-resectable tumors did not differ when chemotherapy was provided post-operatively<sup>[106]</sup>. Chemotherapy with radiation has also been shown to improve survival, but not stage, of cases presenting with locally invasive tumors without metastasis<sup>[107]</sup>. However, previous studies do note that surgical interventions are more challenging and increased postoperative stay is associated with patients undergoing resection after neoadjuvant chemoradiation therapy for locally invasive cancer<sup>[106]</sup>. Since metastatic pancreatic cancer cannot be completely resected, surgical options are unavailable and hence no neoadjuvant therapy can be provided. Lastly, it is important to note that prior to initiating neoadjuvant therapy, histological confirmation of pancreatic adenocarcinoma is required, unlike surgical resection, which often relies solely on imaging.

**Adjuvant chemotherapy:** Even following complete, successful resection of pancreatic tumors, overall survival and prognosis remains discouraging. Hence, postoperative chemotherapy or chemoradiation is almost always incorporated in the therapeutic regimen. Postoperative chemotherapy often utilizes gemcitabine or 5-fluorouracil (with concurrent leucovorin as a rescue agent). Both drugs have demonstrated significant increases in patient survival, regardless of initial case presentation. These drugs may also be given simultaneously, however, significant toxicity (especially gastrointestinal) has been reported. Although gemcitabine has often been considered the standard, previous studies do differ on which agents are associated with the most optimal benefits. In a phase III, randomized control trial, Neoptolemos *et al.*<sup>[108]</sup> noted no significance difference in survivorship between gemcitabine and 5-fluorouracil (with folinic acid) in patients with resected tumors. In a separate study published in JAMA, the authors concluded that gemcitabine alone should be favored over 5-fluorouracil with leucovorin due to its decreased toxicity<sup>[108]</sup>.

Developments of other complementary agents to enhance chemotherapeutic effects are currently under review. For example, possible inhibition of Hedgehog signaling<sup>[84]</sup> or concurrent use of Smac mimetics<sup>[109]</sup>, microRNAs<sup>[110]</sup>, Resveratrol<sup>[111]</sup>, capecitabine<sup>[112]</sup>, thymoquinone<sup>[113]</sup> or heat-shock protein complements<sup>[114]</sup> may promote tumor uptake and damage of administered drugs, such as Gemcitabine administered with concurrent curcumin may also be a potential option, especially in tumors exhibiting gemcitabine-resistance<sup>[115]</sup>. In addition to utilizing CA 19-9 and imaging to monitor patient response to chemotherapy, other markers, such as human equilibrative nucleoside transporter 1 levels have also shown to be useful<sup>[116]</sup>. Other gene expression levels, as noted in Fujita

*et al.*<sup>[117]</sup>, may also be predictive of treatment efficacy, particularly with gemcitabine. Further investigation is required as to whether adjuvant chemotherapy should be administered if prior neoadjuvant therapy before surgery had already been provided.

**Chemotherapy for advanced disease:** Due to its poor detection rate, 60%-70% of patients present with metastatic pancreatic cancer upon initial diagnosis. In the advanced stage of disease, pancreatic cancer causes imminent mortality for the majority of affected patients and median survival rate is typically 6-8 mo. Therefore, treatment of patients with metastatic pancreatic adenocarcinoma incorporates chemotherapy, targeted-therapy, comorbid conditions, intensive supportive treatment and psychosocial support.

Gemcitabine is currently considered the chemotherapeutic standard of care in treatment of advanced pancreatic cancer<sup>[118]</sup>. It has been shown to prolong the average survival rate by 4 mo. In an attempt to improve survival rates, several phase II and phase III trials combined Gemcitabine with fluoropyrimidines and platinum analogs. Most of these combinations failed to show statistically significant survival benefit, however compared to Gemcitabine alone<sup>[119]</sup>. In another attempt to prolong patient survival, scientists have developed several Gemcitabine-based polychemotherapy regimens involving 3-4 cytotoxic agents. When Reni *et al.*<sup>[120]</sup> performed a randomized trial to test the cisplatin, epirubicin, fluorouracil and gemcitabine (PEFG-regimen) against gemcitabine alone, those patients treated with the PEFG saw a significant decrease in cancer progression, when compared to those treated with gemcitabine alone<sup>[120]</sup>. Yet in regards to survival, the infusional 5-FU/folinic acid, irinotecan, and oxaliplatin (FOLFIRINOX) regimen has been shown to be superior to the PEFG-regimen.

According to Conroy *et al.*<sup>[121]</sup>, FOLFIRINOX is the new standard in the treatment of advanced stage pancreatic cancer. Compared to gemcitabine alone, FOLFIRINOX demonstrated a better objective response rate, progression-free survival, overall survival and one-year survival. While the toxicity levels associated with FOLFIRINOX are greater than those caused by gemcitabine, the effects did not seem to have a significant impact on quality of life. In addition, few toxic deaths have been reported.

Inhibitors of epidermal growth factor receptor (EGFR) have also been tested for treatment of metastatic pancreatic cancer. Cetuximab, an anti-EGFR directed antibody and erlotinib, an oral EGFR tyrosine kinase inhibitor have been tested in several randomized trials. However, Moore *et al.*<sup>[122]</sup> demonstrated that combining gemcitabine with erlotinib is the only targeted-therapeutic agent that has clinical efficacy. Compared with gemcitabine alone, gemcitabine plus erlotinib showed significant decrease in tumor progression and concurrently increased overall survival rates.

Other targeted therapies have focused on targeting

the desmoplastic stroma, one of the key components of pancreatic cancer that may contribute to impaired drug delivery and thus chemotherapy resistance<sup>[79]</sup>. Nab-paclitaxel is one therapy that has been developed to diminish this stromal tissue network. Studies have demonstrated that albumin interacts with secreted protein acidic and rich in cysteine (SPARC), a matrix glycoprotein that has a role in tumor invasion, facilitating the uptake of paclitaxel by the tumor<sup>[123]</sup>. Infante *et al.*<sup>[124]</sup> have demonstrated that overexpression of SPARC in peritumoral fibroblasts was a negative prognostic indicator in patients with advanced pancreatic cancer<sup>[124]</sup>. In a phase I / II trial, Von Hoff *et al.*<sup>[125]</sup> demonstrated the ability of nab-paclitaxel to increase median survival rate in patients with metastatic pancreatic cancer.

Similar to the mechanism of action of nab-paclitaxel, new therapies are being developed that target the peritumoral stroma in order to increase tumor perfusion. One such preclinical strategy inhibits the hedgehog signaling pathway, depleting the stroma and increasing angiogenesis to improve delivery of chemotherapeutic agents to the tumor<sup>[126]</sup>.

Phase II clinical trials have demonstrated a benefit of second-line treatment for patients who are resistant to gemcitabine treatment<sup>[127]</sup>. Second-line treatments typically consist of fluoropyrimidines in combination with oxaliplatin<sup>[128]</sup>. Limited data exists on how to treat patients who do not tolerate FOLFIRINOX as a first-line therapy. However, Conroy *et al.*<sup>[121]</sup> have demonstrated the benefit of using gemcitabine-based therapies in these instances.

The primary prognostic indicators for patient survival are both patient and tumor related. Based on the genetic and morphologic heterogeneity that exists within each individual pancreatic tumor, therapy, dose and length of therapy administration will need to be customized for each Individual patient to ensure optimal treatment.

## Radiation

**Neoadjuvant radiation therapy:** Many studies have demonstrated the important roles for chemotherapy and radiation therapy in preventing the recurrence and improving the resectability of pancreatic tumors. While surgery is currently the only potential curative treatment modality for pancreatic cancer, more than 80% of patients who undergo surgical resection will experience tumor recurrence within 12 mo of their procedure<sup>[129]</sup>. Therefore, a great deal of focus has not only been placed on developing effective neoadjuvant and adjuvant therapies, but also on effective preoperative staging techniques to determine candidates who will benefit most from surgical resection<sup>[130]</sup>. Since surgery is associated with high rates of morbidity and mortality, many patients do not begin adjuvant therapy until after they have recovered. As a result, there is a long delay before they receive adjuvant therapy. In order to begin more potent treatments earlier and to potentially shrink the tumor before surgery, researchers developed neoadjuvant therapeutic regimens.

Multiple trials of 5-fluorouracil-based chemoradia-

tion have been done to date. At the conclusion of these studies, researchers determined that a combined treatment modality with preoperative rapid-fractionation chemoradiation, Whipple procedure, and intra-operative radiation therapy resulted in minimal toxicity and a small recurrence rate<sup>[131]</sup>. In a follow-up study, paclitaxel replaced 5-fluorouracil and was used to treat 35 patients who presented with resectable pancreatic tumors<sup>[132]</sup>. Based on the results of this study, researchers concluded that preoperative paclitaxel-based chemotherapy with rapid-fractionation chemoradiation, Whipple procedure and intraoperative radiation therapy resulted in similar outcomes as the previous study, but toxicity levels were greater than those from 5-fluorouracil. In another study, researchers treated patients who presented with tumors in the head of the pancreas with a neoadjuvant chemoradiation regimen of capecitabine with proton beam radiation<sup>[133]</sup>. No dose limiting toxicities were observed and the authors concluded that this form of neoadjuvant therapy was feasible. In several other prospective neoadjuvant chemoradiation trials in patients with resectable pancreatic cancers, the rate of resection was high in all studies, ranging from 87%-100%<sup>[107,134,135]</sup>.

**Adjuvant radiation therapy:** The median survival rate of patients who undergo surgical resection of a pancreatic tumor is 15-22 mo. Only 20% of patients survive for five years following surgery<sup>[136]</sup>. The most common site for pancreatic cancer recurrence is the retroperitoneum. Therefore, adjuvant therapy is needed to improve patient prognosis. In the United States, adjuvant therapy is currently delivered in the form of chemotherapy, chemoradiotherapy or chemotherapy followed by chemoradiotherapy. Standard adjuvant treatment in Europe is chemotherapy alone. These guidelines were based on previous randomized trials that showed improved survival in patients given adjuvant therapy following surgical resection.

The first prospective trial for adjuvant chemoradiotherapy was conducted by the Gastrointestinal Tumor Study Group in 1985. The trial enrolled patients with resectable pancreatic cancer. The protocol called for external beam radiation delivered with 5-fluorouracil. The patients were then given a maintenance dose of 5-fluorouracil for an additional two years following initial treatment. Patients treated with adjuvant chemoradiation achieved a longer median and 2-year survival rate than those not treated with adjuvant therapy. As a result, adjuvant chemoradiation became the most frequently used adjuvant treatment for resectable pancreatic cancer in the United States.

To further assess adjuvant radiation therapy for resectable pancreatic cancer, the European Study Group for Pancreatic Cancer (ESPAC-1) conducted the largest randomized trial to date in 2004<sup>[137]</sup>. In order to evaluate the effects of chemoradiotherapy and chemotherapy on patient survival following surgical resection, patients with resectable pancreatic ductal adenocarcinoma were divided into one of four groups: chemotherapy alone;

chemoradiotherapy alone; chemoradiotherapy followed by chemotherapy; or no further treatment. Patients who received chemotherapy followed by chemoradiotherapy had a 5-year survival rate that was 10% less than those who received chemotherapy alone. In addition, patients who received chemotherapy treatment showed a 5-year survival benefit when compared to those who received no chemotherapy. As a result of these findings, the standard of adjuvant treatment in Europe shifted towards chemotherapy only, abandoning postoperative chemoradiation.

A phase III trial was conducted by the Radiation Oncology Group and GI Intergroup around the same time as the ESPAC-1 trial<sup>[138]</sup>. This trial compared the 5-fluorouracil-based chemoradiation to gemcitabine-based chemoradiation. Patients receiving gemcitabine-based chemoradiation had a median survival of 20.6 mo, 3.5 mo more than those given 5-fluorouracil-based chemotherapy. The Charite Onkologie Clinical Studies in GI Cancer 001 (CONKO-001) trial in Germany and Austria showed similar median survival in patients given gemcitabine-based chemotherapy alone<sup>[139]</sup>.

Additionally, reports from several institutions, including the Mayo Clinic, Johns Hopkins Medical Center and Virginia Mason University have all reported the benefit of adjuvant chemoradiation therapy following resectable pancreatic cancer compared to those who received surgery alone<sup>[140-142]</sup>.

#### Management of locally advanced pancreatic cancer:

Patients with locally advanced pancreatic cancer achieve little benefit from surgical resection because their cancer meets the criteria for unresectable cancer: (1) distant metastasis and/or pancreatic lymph node involvement; (2) encasement or occlusion of the superior mesenteric vein or superior mesenteric vein/portal vein confluence; and/or (3) direct involvement of the celiac axis, aorta, inferior vena cava, or superior mesenteric artery<sup>[143]</sup>. As a result, chemoradiation is recommended for these patients based on data from several studies.

A 1981 trial conducted by the Gastrointestinal Tumor Study Group compared the effects of high-dose radiation therapy alone; moderate dose radiation combined with 5-fluorouracil and high dose radiation combined with 5-fluorouracil in 194 patients with locally advanced pancreatic cancer. Researchers found that patients administered 5-fluorouracil in combination with low or high dose radiation showed a greater median survival than those treated with radiation alone<sup>[144]</sup>. In a follow-up study, the same group demonstrated that chemotherapy, when combined with radiotherapy afforded patients with locally advanced pancreatic cancer a greater median survival when compared to combination chemotherapy alone<sup>[145]</sup>. These results were verified by the ECOG trial as well, which demonstrated an increased median survival rate in patients treated with gemcitabine and radiotherapy together as opposed to those treated with gemcitabine alone<sup>[146]</sup>.

Although chemoradiation has been shown to provide

an increased median survival in patients with locally advanced pancreatic cancer by 9-13 mo, many of these patients progress to the metastatic stage of disease shortly after therapy. Perhaps a better approach to these patients would be to begin a chemotherapy regimen, restage their cancer after completion of initial treatment, and follow up with chemoradiation in patients who do not demonstrate metastatic disease progression. Radiation in these patients could relieve pain associated with disease by slowing local progression.

**Stereotactic body radiotherapy:** An evolving radiation therapy for treatment of locally advanced pancreatic cancer is stereotactic body radiation therapy. This newer technique uses image guidance to deliver toxic radiation doses directly to tumors. As a result, there is less systemic involvement, and patient outcomes are improved without having to undergo daily treatments. However, the major challenge of this novel therapy is accurately characterizing the tumor in terms of size, number and location. In order to do so, precise diagnostic tests and real-time imaging techniques are used. In addition, each treatment regimen is tailored to each individual patient. To date, scientific literature suggests stereotactic body radiotherapy does slow local progression in patients with locally advanced pancreatic cancer<sup>[147-149]</sup>. However, it does not increase overall survival rate because patient mortality is due to distant metastases.

**Advances in radiation therapy techniques:** Over the past decade, major advances in radiation therapy have been in treatment planning and more precise delivery methodologies. One technique, intensity-modulated radiotherapy, decreases systemic toxicities in patients by modifying radiation dose delivery specifically to the tumor sites, sparing surrounding normal tissue<sup>[150,151]</sup>. Another technique, image-guided radiotherapy, has provided more accurate visualization and real-time tracking of viscera-located tumors and thus has enabled more precise delivery of high-dose therapeutic beams of radiation to these tumors and prevented adverse effects in normal tissue<sup>[152]</sup>.

In order to improve patient outcome and prolong median survival rate, additional studies are needed to define the optimal role of adjuvant and neoadjuvant treatment in patients with resectable pancreatic cancer. As radiation therapies become more precise and customized to individual patients, it will be necessary to continue to investigate their future role in the treatment of pancreatic adenocarcinoma, especially as a greater understanding of the molecular pathways involved in the carcinogenesis and progression of this disease are understood.

**Personalized therapy:** In an article published in *Science*, Jones *et al.*<sup>[77]</sup> performed a comprehensive genome assessment on 24 different pancreatic cancers. Results revealed an average of 63 genetic mutations per cancer, spanning 12 separate signal transduction pathways. This study sup-



ports the notion of pancreatic cancer being a genetically heterogeneous malignancy, partially accounting for its notable resistance to therapy as well as varied responses to treatment. Moreover, this finding likely explains why no candidate gene has been identified in the treatment of pancreatic cancer. This heterogeneity will likely dictate an individualized, unique approach for each particular case, which has already shown to be effective against even advanced pancreatic cancer stages. In one such case report, Villarreal *et al.*<sup>[153]</sup> identified Mitomycin C, a DNA-damaging agent, as a highly effective agent by utilizing a xenograft derived from the patient's tumor. Upon administration of this drug, the patient exhibited notable clinical benefits for over three years, despite the tumor previously being gemcitabine resistant. Personalized immune system stimulation may also be a viable option in treatment of unresectable disease. For example, Yanagimoto *et al.*<sup>[154]</sup> incorporated a vaccine containing individualized, reactive peptides with concurrent gemcitabine treatment, noting a significant correlation between immune boosting and survivorship.

Due to the ongoing advances in DNA sequencing, personalized genomic therapy appears more plausible. Moreover, as scientists continue to identify regions of the genome with high potential for tumor-pathogenesis, this method will only become more efficient. Upon identification, cases can be distributed into cohorts based upon their tumor's genetic composition and administered treatment previously demonstrated to be effective in that particular subgroup. This method would not only identify pertinent biomolecules in pancreatic pathogenesis, but also lead to tumor-specific treatment, which is likely necessary if we are to see any significant improvement in the prognosis of pancreatic cancer.

## FUTURE PERSPECTIVES

Despite decades of effort by the scientific community to design sophisticated chemotherapeutic and radiation techniques to combat pancreatic ductal adenocarcinoma, less than 5% of patients with this disease have a 5-year survival rate. The majority of patients have a median survival period of 4-6 mo<sup>[155,156]</sup>. A combination of factors including few early symptoms, few accurate biomarkers for early detection, rapid metastasis to the lymphatic system and distant organs, and few effective treatment options, makes this disease one of the most deadly cancers today<sup>[157,158]</sup>. Although current therapeutic agents have had limited effects on patient care, there has been substantial advancement in the understanding of the molecular and biological makeup of pancreatic adenocarcinoma. This knowledge has the potential to lead to the development of novel therapies that could significantly improve the lifespan of individuals suffering with this disease.

Such advances in understanding this complex disease have been achieved with genetically-engineered mouse models and patient-derived xenografts. These studies have demonstrated the genetic diversity of pancreatic

ductal adenocarcinoma results from successive accumulation of mutations in several primary oncogenes and tumor suppressor genes, leading to its heterogeneity, instability and early tumor metastasis<sup>[77]</sup>. Pancreatic ductal adenocarcinoma is composed of several compartments. In addition to a mature cancer cell population, some researchers have characterized cancer cells that display stem cell properties and are resistant to chemotherapy and radiation therapy, potentiating their ability to metastasize<sup>[56]</sup>. Another area of interest is the dense tumor microenvironment that surrounds the pancreatic cancer cells. Composed of collagen I, activated fibroblasts, and inflammatory cells, it has been shown to interact with pancreatic cancer cells in order to foster tumor development, act as a barrier to optimal drug delivery and aid the tumor in invasion and metastasis<sup>[59]</sup>. Furthermore, this dense stroma creates a hypoxic microenvironment that pancreatic cancer cells thrive in<sup>[159]</sup>. However, the mechanisms by which these cancer cells adapt to these conditions are currently being identified and may serve as additional therapeutic targets in the near future.

### Pancreatic cancer cells

Pancreatic ductal adenocarcinoma most likely originates in the ductal epithelium of pancreatic cells<sup>[160]</sup>. Neoplastic cells contain one or more of four primary genetic mutations that will ultimately give rise to the invasive form of this disease. Ninety percent of these tumors have mutations in the KRAS2 oncogene, resulting in the activation of proliferative survival signaling pathways. Ninety-five percent have a mutation in the CDKN2A tumor suppressor gene, resulting in the loss of the p16 protein and thus loss of regulation of the G1-S transition of the cell cycle. An abnormal TP53 gene has been identified in 50%-75% of characterized cancer cells, allowing cells to avoid DNA damage control checkpoints and subsequently, apoptotic signals. Another 50% have a deleted SMAD4/MADH4 gene, resulting in aberrant signaling by the TGF- $\beta$  cell surface receptor<sup>[153]</sup>.

One study performed genetic analysis on 24 pancreatic ductal adenocarcinomas and reported that each tumor has an average of 63 clinically relevant genetic abnormalities. While these abnormalities differ from one cancer to another, they all seem to play a role in 12 functional cancer-related pathways<sup>[161]</sup>. Recently, two studies compared the genetic makeup of distant metastases to their primary metastatic lesions. They found that over time, the distant metastases accumulated additional mutations to those present in the clonal cells from which they arose, adding to the complexity this disease<sup>[162]</sup>. Such genetic diversity not only results in different prognoses for patients, but also causes individual tumors to respond differently to common therapeutic agents used in treating pancreatic ductal adenocarcinoma<sup>[163]</sup>.

The varying degrees of genetic instability that exist between individual pancreatic ductal adenocarcinomas present a greater need for genomic sequencing of individual tumors, followed by personalized therapies to target spe-

cific genes and pathways that have been altered<sup>[164]</sup>. Several clinical trials have begun exploring this idea<sup>[165]</sup>. In order to incorporate this treatment modality into the clinical setting, several criteria must be met: (1) a high quality tumor tissue sample must be attained at the time of diagnosis; (2) sophisticated bioinformatic analysis of the data must be performed to identify the most relevant mutations in each tissue sample; and (3) model systems must be designed to experimentally test varying treatment options to determine the most effective one for the patient. Perhaps the greatest challenge lies in developing a drug once specific genetic abnormalities have been identified.

### **Pancreatic cancer stem cells**

Recently, investigators have characterized pancreatic cancer cells with stem cell properties<sup>[162]</sup>. Known as pancreatic cancer stem cells, these cells have the ability to regrow new tumors when placed into naïve mouse models and are able to maintain long-term tumorigenic potential<sup>[163]</sup>. Studies have shown that pancreatic CSCs are not only capable of self renewal, but may also confer therapeutic resistance, and play a role in tumor formation and disease progression<sup>[164,165]</sup>. In addition, different cancer stem cell populations perform different biological functions. One of the most recent findings has demonstrated that these cells may transition between epithelial and mesenchymal states, contributing to their highly metastatic potential<sup>[165]</sup>. Therefore, eliminating or inhibiting these CSCs with new therapeutic designs could significantly improve patient outcomes. Current therapies have already been designed to target cancer stem cell-specific antigens in order to inhibit their roles in cell survival, adhesion, self renewal and differentiation. A greater understanding of individual CSC populations and how they interact with one another will enable further progress in the treatment of pancreatic cancer<sup>[164,165]</sup>.

Therapeutic targets of pancreatic ductal adenocarcinoma cancer stem cells include genes located in developmental pathways such as hedgehog, Wnt, Notch, CXCR4 and Met. In addition, targeting apoptotic pathways such as DR5 and nodal-activin could have a significant therapeutic implications. Several preclinical trials have been conducted to target these pathways in models of human pancreatic ductal adenocarcinoma cancer stem cells. By inhibiting these pathways, investigators were able to confer longer-term tumor control when compared to current standard chemotherapeutic regimens, in which tumor regression was significantly shorter-lived. In one recent trial, salinomycin was shown to induce cell death in epithelial-mesenchymal transition-induced cancer stem cells<sup>[124]</sup>.

Due to the heterogeneity of the cancer stem cell population, future drugs designed to target pancreatic ductal adenocarcinoma stem cells may require clinical trials in which therapies are designed specifically for pancreatic tumors in each individual patient. These customized therapies could potentially serve as adjuvant treatment options for patients following pancreatic tumor resection. Similar to previous clinical trial designs, adjuvant thera-

pies targeting cancer stem cells could be given to patients with or without current conventional chemotherapy and/or chemoradiation to determine which option confers the greatest overall survival rate in patients following surgical resection.

### **Tumor microenvironment**

One of the primary characteristics of pancreatic ductal adenocarcinoma is the dense stroma surrounding the pancreatic cancer cells. Composed of fibroblasts, collagen I and other fibrillar elements, this desmoplastic reaction has become a primary target of current drug therapies<sup>[125]</sup>. The key players in the formation and turnover of this dense stroma are pancreatic stellate cells. Certain growth factors [TGF- $\beta$ 1, platelet-derived growth factor (PDGF) and fibroblast growth factor] activate these cells to myofibroblasts. Not only do these activated myofibroblasts secrete components of the extracellular matrix, but they are also responsible for the poor vascularization of the pancreatic tumor<sup>[166,167]</sup>. In addition to forming a mechanical barrier around the pancreatic cancer cells, the stroma has an important role in tumor formation, progression, invasion and metastasis<sup>[168]</sup>. Many proteins expressed by stromal cells have been directly correlated with poor prognosis and resistance to current therapies [Cox-2, PDGF receptor, VEGF, stromal-derived factor, chemokines, integrins, secreted protein acidic and rich in cysteine (SPARC), and hedgehog elements].

Pre-clinical models have demonstrated that targeting these receptors and enzymes is associated with antitumor effects. Perhaps one of the most promising targets to date is the hedgehog signaling pathway. Some studies have demonstrated that targeting smoothed resulted in a depletion of the stroma and thereby increased delivery of gemcitabine to the tumor cells<sup>[169]</sup>. Another target for therapeutic trials has been SPARC (osteonectin) and hyaluronic acid. SPARC is an extracellular matrix protein that plays a role in collagen turnover in the dense stroma. It is associated with invasion and metastasis in pancreatic ductal adenocarcinoma, and thus poor prognosis in patients with elevated levels<sup>[170]</sup>. As mentioned previously, SPARC is the target of the albumin-bound chemotherapy agent, nab-paclitaxel. Phase I / II clinical trials have shown that administration of this drug breaks down the stroma and improves delivery of the chemotherapeutic agent to the site of the tumor<sup>[171]</sup>. In addition, in a mouse model of pancreatic ductal adenocarcinoma, investigators demonstrated that administration of pegylated hyaluronidase eliminated hyaluronic acid content, thereby relieving pressure on the blood vessels surrounding the tumor and allowed for increased perfusion of the chemotherapeutic agent to the site of the tumor<sup>[172]</sup>.

The immunosuppressive nature of the tumor microenvironment has been another stromal characteristic targeted by recent therapeutic development. Using a CD40 antibody combined with gemcitabine chemotherapy, researchers have attempted to reverse immune suppression and drive antitumor T-cell responses in patients with non-resectable pancreatic ductal adenocarcinoma. Studies

have shown that this agent results in tumor regression by stimulating tumor macrophages to attack and deplete the pancreatic cancer stroma<sup>[173]</sup>.

To date, targeting pancreatic ductal adenocarcinoma has proved most effective when treating patients with locally advanced disease, especially patients with tumors characterized by wild-type DPC4. These tumors are known to be less prone to metastasis and possess higher stromal content. Other tumors, especially those in late stages of the disease, characterized by distant metastases, have not been effectively treated with current stromal-targeting therapeutic agents. This is due to the fact that although pancreatic ductal adenocarcinoma has a rich and hypovascularized stroma, metastases arising from this cancer do not and are not different from other tumors. Therefore, patients who may benefit most from treatment with agents targeting the dense stroma microenvironment would be those with resectable tumors that have not progressed to the advanced stages of disease<sup>[174]</sup>.

### Metabolic pathways

Another conventional way to target pancreatic ductal adenocarcinoma would be to inhibit its major metabolic pathways. In order to do so, researchers would need to prevent its supply of glucose and glutamine; interrupt the pathways that enable it to exist in a hypoxic environment<sup>[175]</sup>; and prevent its ability to digest intracellular organelles for energy<sup>[176]</sup>.

Investigators have identified several key metabolic enzymes to target [hexokinase, pyruvate kinase, lactate dehydrogenase A (LDHA) and ampicillin-activated protein kinase (AMPK)]. Several preclinical trials have demonstrated the anti-tumor effects of agents directed against these enzymes. One study demonstrated a potential clinical application for the LDHA inhibitor, FX11. By blocking the conversion of lactate to pyruvate in cells with p53 mutations, FX11 has antitumor potential. However, to date, there are only two therapies that have shown potential for targeting pancreatic ductal adenocarcinoma metabolism. One of these medications, metformin, is an activator of AMPK. It has been shown to decrease the potential for patients with diabetes to develop pancreatic cancer and to increase survival in diabetic patients with this disease<sup>[177,178]</sup>. The other drug used to target the metabolic pathways of is rapamycin. An inhibitor of mTOR, rapamycin has been shown to decrease glucose uptake by reducing levels of Glut1 in pancreatic cancer<sup>[179,180]</sup>.

In order to inhibit autophagy, a significant mechanism for pancreatic cancer cell survival, investigators have used chloroquine, the antimalarial drug<sup>[181]</sup>. In preclinical trials with both allografts and xenografts, chloroquine has been shown to decrease tumorigenesis in a transgenic model and is currently being tested in clinical trials.

### CONCLUSION

A greater understanding of the molecular and cellular makeup of pancreatic cancer over the past four decades

has resulted in innovative therapeutic designs to target this aggressive malignancy. We now know that pancreatic cancer is a dynamic, heterogenous and genetically unstable tumor that results from successive mutations early in disease and gives rise to metastases that continue to garner mutations as they travel to distant locations. An equally important understanding of the role the peritumor microenvironment, composed of a dense desmoplastic stroma, plays in tumor development, metastases and as a barrier to chemotherapy delivery has been elucidated over the years. More recently, a role for pancreatic cancer stem cells in resistance to chemotherapy and radiation therapy was discovered. In addition, a deeper understanding of the metabolic pathways responsible for adaptation of pancreatic cancer cells to hypoxic environments has significant implications for future therapeutic development.

While some of these discoveries have resulted in novel therapeutic targets and treatment strategies and others are currently being tested in preclinical trials, efficient and effective drug development to combat pancreatic ductal adenocarcinoma is a necessity for the future. The majority of clinical trials have been conducted in patients with advanced stage disease. In the future, it will be necessary to design clinical trials to enroll patients with earlier stages of pancreatic cancer in an attempt to cure their cancer before it can metastasize.

### REFERENCES

- 1 **American Cancer Society.** Cancer Facts & Figures 2013. Atlanta: American Cancer Society, 2013: 25-32
- 2 **Kalager M, Zelen M, Langmark F, Adami HO.** Effect of screening mammography on breast-cancer mortality in Norway. *N Engl J Med* 2010; **363**: 1203-1210 [PMID: 20860502 DOI: 10.1056/NEJMoa1000727]
- 3 **DeSantis C, Ma J, Bryan L, Jemal A.** Breast cancer statistics, 2013. *CA Cancer J Clin* 2013; **64**: 52-62 [PMID: 24114568 DOI: 10.3322/caac.21203]
- 4 **Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D.** Global cancer statistics. *CA Cancer J Clin* 2011; **61**: 69-90 [PMID: 21296855 DOI: 10.3322/caac.20107]
- 5 **Center MM, Jemal A, Smith RA, Ward E.** Worldwide variations in colorectal cancer. *CA Cancer J Clin* 2009; **59**: 366-378 [PMID: 19897840 DOI: 10.3322/caac.20038]
- 6 **Sener SF, Fremgen A, Menck HR, Winchester DP.** Pancreatic cancer: a report of treatment and survival trends for 100,313 patients diagnosed from 1985-1995, using the National Cancer Database. *J Am Coll Surg* 1999; **189**: 1-7 [PMID: 10401733 DOI: 10.1016/S1072-7515(99)00075-7]
- 7 **Simard EP, Ward EM, Siegel R, Jemal A.** Cancers with increasing incidence trends in the United States: 1999 through 2008. *CA Cancer J Clin* 2012; Epub ahead of print [PMID: 22281605 DOI: 10.3322/caac.20141]
- 8 **Siegel R, Naishadham D, Jemal A.** Cancer statistics, 2012. *CA Cancer J Clin* 2012; **62**: 10-29 [PMID: 22237781 DOI: 10.3322/caac.20138]
- 9 **Petersen GM, de Andrade M, Goggins M, Hruban RH, Bondy M, Korczak JF, Gallinger S, Lynch HT, Syngal S, Rabe KG, Seminara D, Klein AP.** Pancreatic cancer genetic epidemiology consortium. *Cancer Epidemiol Biomarkers Prev* 2006; **15**: 704-710 [PMID: 16614112 DOI: 10.1158/1055-9965.EPI-05-0734]
- 10 **Blackford A, Parmigiani G, Kensler TW, Wolfgang C, Jones**



- S, Zhang X, Parsons DW, Lin JC, Leary RJ, Eshleman JR, Goggins M, Jaffee EM, Iacobuzio-Donahue CA, Maitra A, Klein A, Cameron JL, Olin K, Schulick R, Winter J, Vogelstein B, Velculescu VE, Kinzler KW, Hruban RH. Genetic mutations associated with cigarette smoking in pancreatic cancer. *Cancer Res* 2009; **69**: 3681-3688 [PMID: 19351817 DOI: 10.1158/0008-5472.CAN-09-0015]
- 11 **Shi C**, Hruban RH, Klein AP. Familial pancreatic cancer. *Arch Pathol Lab Med* 2009; **133**: 365-374 [PMID: 19260742]
  - 12 **Brune KA**, Lau B, Palmisano E, Canto M, Goggins MG, Hruban RH, Klein AP. Importance of age of onset in pancreatic cancer kindreds. *J Natl Cancer Inst* 2010; **102**: 119-126 [PMID: 20068195 DOI: 10.1093/jnci/djp466]
  - 13 **Genkinger JM**, Spiegelman D, Anderson KE, Bergkvist L, Bernstein L, van den Brandt PA, English DR, Freudenheim JL, Fuchs CS, Giles GG, Giovannucci E, Hankinson SE, Horn-Ross PL, Leitzmann M, Männistö S, Marshall JR, McCullough ML, Miller AB, Reding DJ, Robien K, Rohan TE, Schatzkin A, Stevens VL, Stolzenberg-Solomon RZ, Verhage BA, Wolk A, Ziegler RG, Smith-Warner SA. Alcohol intake and pancreatic cancer risk: a pooled analysis of fourteen cohort studies. *Cancer Epidemiol Biomarkers Prev* 2009; **18**: 765-776 [PMID: 19258474 DOI: 10.1158/1055-9965.EPI-08-0880]
  - 14 **Zhang J**, Dhakal IB, Gross MD, Lang NP, Kadlubar FF, Harnack LJ, Anderson KE. Physical activity, diet, and pancreatic cancer: a population-based, case-control study in Minnesota. *Nutr Cancer* 2009; **61**: 457-465 [PMID: 19838917 DOI: 10.1080/01635580902718941]
  - 15 **Philip B**, Roland CL, Daniluk J, Liu Y, Chatterjee D, Gomez SB, Ji B, Huang H, Wang H, Fleming JB, Logsdon CD, Cruz-Monserrate Z. A high-fat diet activates oncogenic Kras and COX2 to induce development of pancreatic ductal adenocarcinoma in mice. *Gastroenterology* 2013; **145**: 1449-1458 [PMID: 23958541 DOI: 10.1053/j.gastro.2013.08.018]
  - 16 **Amaral AF**, Porta M, Silverman DT, Milne RL, Kogevinas M, Rothman N, Cantor KP, Jackson BP, Pumarega JA, López T, Carrato A, Guarner L, Real FX, Malats N. Pancreatic cancer risk and levels of trace elements. *Gut* 2012; **61**: 1583-1588 [PMID: 22184070 DOI: 10.1136/gutjnl-2011-301086]
  - 17 **Stathis A**, Moore MJ. Advanced pancreatic carcinoma: current treatment and future challenges. *Nat Rev Clin Oncol* 2010; **7**: 163-172 [PMID: 20101258 DOI: 10.1038/nrclinonc.2009.236]
  - 18 **Sheridan MB**, Ward J, Guthrie JA, Spencer JA, Craven CM, Wilson D, Guillou PJ, Robinson PJ. Dynamic contrast-enhanced MR imaging and dual-phase helical CT in the preoperative assessment of suspected pancreatic cancer: a comparative study with receiver operating characteristic analysis. *AJR Am J Roentgenol* 1999; **173**: 583-590 [PMID: 10470884 DOI: 10.2214/ajr.173.3.10470884]
  - 19 **Horton KM**, Fishman EK. Adenocarcinoma of the pancreas: CT imaging. *Radiol Clin North Am* 2002; **40**: 1263-1272 [PMID: 12479710 DOI: 10.1016/S0033-8389(02)00041-6]
  - 20 **Faria SC**, Tamm EP, Loyer EM, Szklaruk J, Choi H, Charnsangavej C. Diagnosis and staging of pancreatic tumors. *Semin Roentgenol* 2004; **39**: 397-411 [PMID: 15372753 DOI: 10.1016/j.ro.2004.06.012]
  - 21 **House MG**, Yeo CJ, Cameron JL, Campbell KA, Schulick RD, Leach SD, Hruban RH, Horton KM, Fishman EK, Lillemoe KD. Predicting resectability of periaampullary cancer with three-dimensional computed tomography. *J Gastrointest Surg* 2004; **8**: 280-288 [PMID: 15019924 DOI: 10.1016/j.gassur.2003.12.011]
  - 22 **Vachiranubhap B**, Kim YH, Balci NC, Semelka RC. Magnetic resonance imaging of adenocarcinoma of the pancreas. *Top Magn Reson Imaging* 2009; **20**: 3-9 [PMID: 19687720 DOI: 10.1097/RMR.0b013e3181b48392]
  - 23 **Tellez-Avila FI**, Chavez-Tapia NC, López-Arce G, Franco-Guzmán AM, Sosa-Lozano LA, Alfaro-Lara R, Chan-Núñez C, Giovannini M, Elizondo-Rivera J, Ramírez-Luna MA. Vascular invasion in pancreatic cancer: predictive values for endoscopic ultrasound and computed tomography imaging. *Pancreas* 2012; **41**: 636-638 [PMID: 22460727 DOI: 10.1097/MPA.0b013e31823e3632]
  - 24 **Sperti C**, Pasquali C, Bissoli S, Chierichetti F, Liessi G, Pedrazzoli S. Tumor relapse after pancreatic cancer resection is detected earlier by 18-FDG PET than by CT. *J Gastrointest Surg* 2010; **14**: 131-140 [PMID: 19777315 DOI: 10.1007/s11605-009-1010-8]
  - 25 **Warshaw AL**, Tepper JE, Shipley WU. Laparoscopy in the staging and planning of therapy for pancreatic cancer. *Am J Surg* 1986; **151**: 76-80 [PMID: 2936262 DOI: 10.1016/0002-9610(86)90015-2]
  - 26 **Bilimoria KY**, Bentrem DJ, Ko CY, Ritchey J, Stewart AK, Winchester DP, Talamonti MS. Validation of the 6th edition AJCC Pancreatic Cancer Staging System: report from the National Cancer Database. *Cancer* 2007; **110**: 738-744 [PMID: 17580363]
  - 27 **Bakkevold KE**, Arnesjø B, Kambestad B. Carcinoma of the pancreas and papilla of Vater--assessment of resectability and factors influencing resectability in stage I carcinomas. A prospective multicentre trial in 472 patients. *Eur J Surg Oncol* 1992; **18**: 494-507 [PMID: 1426302]
  - 28 **Porta M**, Fabregat X, Malats N, Guarner L, Carrato A, de Miguel A, Ruiz L, Jariod M, Costafreda S, Coll S, Alguacil J, Corominas JM, Solà R, Salas A, Real FX. Exocrine pancreatic cancer: symptoms at presentation and their relation to tumour site and stage. *Clin Transl Oncol* 2005; **7**: 189-197 [PMID: 15960930 DOI: 10.1007/BF02712816]
  - 29 **Chari ST**, Leibson CL, Rabe KG, Ransom J, de Andrade M, Petersen GM. Probability of pancreatic cancer following diabetes: a population-based study. *Gastroenterology* 2005; **129**: 504-511 [PMID: 16083707 DOI: 10.1016/j.gastro.2005.05.007]
  - 30 **Khorana AA**, Fine RL. Pancreatic cancer and thromboembolic disease. *Lancet Oncol* 2004; **5**: 655-663 [PMID: 15522652 DOI: 10.1016/S1470-2045(04)01606-7]
  - 31 **Pausch T**, Hartwig W, Hinz U, Swolana T, Bundy BD, Hackert T, Grenacher L, Büchler MW, Werner J. Cachexia but not obesity worsens the postoperative outcome after pancreatoduodenectomy in pancreatic cancer. *Surgery* 2012; **152**: S81-S88 [PMID: 22770957 DOI: 10.1016/j.surg.2012.05.028]
  - 32 **Langer P**, Kann PH, Fendrich V, Gabbe N, Schneider M, Sina M, Slater EP, Heverhagen JT, Gress TM, Rothmund M, Bartsch DK. Five years of prospective screening of high-risk individuals from families with familial pancreatic cancer. *Gut* 2009; **58**: 1410-1418 [PMID: 19470496 DOI: 10.1136/gut.2008.171611]
  - 33 **Verna EC**, Hwang C, Stevens PD, Rotterdam H, Stavropoulos SN, Sy CD, Prince MA, Chung WK, Fine RL, Chabot JA, Frucht H. Pancreatic cancer screening in a prospective cohort of high-risk patients: a comprehensive strategy of imaging and genetics. *Clin Cancer Res* 2010; **16**: 5028-5037 [PMID: 20876795 DOI: 10.1158/1078-0432.CCR-09-3209]
  - 34 **Canto MI**, Hruban RH, Fishman EK, Kamel IR, Schulick R, Zhang Z, Topazian M, Takahashi N, Fletcher J, Petersen G, Klein AP, Axilbund J, Griffin C, Syngal S, Saltzman JR, Mortelet KJ, Lee J, Tamm E, Vikram R, Bhosale P, Margolis D, Farrell J, Goggins M. Frequent detection of pancreatic lesions in asymptomatic high-risk individuals. *Gastroenterology* 2012; **142**: 796-804; quiz e14-5 [PMID: 22245846]
  - 35 **Erkan M**, Hausmann S, Michalski CW, Fingerle AA, Dobritz M, Kleeff J, Friess H. The role of stroma in pancreatic cancer: diagnostic and therapeutic implications. *Nat Rev Gastroenterol Hepatol* 2012; **9**: 454-467 [PMID: 22710569 DOI: 10.1038/nrgastro.2012.115]
  - 36 **Goggins M**. Molecular markers of early pancreatic cancer. *J Clin Oncol* 2005; **23**: 4524-4531 [PMID: 16002843 DOI: 10.1200/JCO.2005.19.711]
  - 37 **Steinberg W**. The clinical utility of the CA 19-9 tumor-asso-



- ciated antigen. *Am J Gastroenterol* 1990; **85**: 350-355 [PMID: 2183589]
- 38 **Ballehaninna UK**, Chamberlain RS. The clinical utility of serum CA 19-9 in the diagnosis, prognosis and management of pancreatic adenocarcinoma: An evidence based appraisal. *J Gastrointest Oncol* 2012; **3**: 105-119 [PMID: 22811878 DOI: 10.3978/j.issn.2078-6891.2011.021]
- 39 **Frebourg T**, Bercoff E, Manchon N, Senant J, Basuyau JP, Breton P, Janvresse A, Brunelle P, Bourreille J. The evaluation of CA 19-9 antigen level in the early detection of pancreatic cancer. A prospective study of 866 patients. *Cancer* 1988; **62**: 2287-2290 [PMID: 3179943]
- 40 **Nakai Y**, Isayama H, Sasaki T, Takahara N, Hamada T, Uchino R, Mizuno S, Miyabayashi K, Yamamoto K, Mohri D, Kogure H, Yamamoto N, Hirano K, Ijichi H, Tateishi K, Tada M, Koike K. A retrospective analysis of early CA19-9 change in salvage chemotherapy for refractory pancreatic cancer. *Cancer Chemother Pharmacol* 2013; **72**: 1291-1297 [PMID: 24121456 DOI: 10.1007/s00280-013-2313-8]
- 41 **Hidalgo M**. Pancreatic cancer. *N Engl J Med* 2010; **362**: 1605-1617 [PMID: 20427809 DOI: 10.1056/NEJMra0901557]
- 42 **Edge SB**, Compton CC. The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging manual and the future of TNM. *Ann Surg Oncol* 2010; **17**: 1471-1474 [PMID: 20180029]
- 43 **Chang KJ**, Nguyen P, Erickson RA, Durbin TE, Katz KD. The clinical utility of endoscopic ultrasound-guided fine-needle aspiration in the diagnosis and staging of pancreatic carcinoma. *Gastrointest Endosc* 1997; **45**: 387-393 [PMID: 9165320 DOI: 10.1016/S0016-5107(97)70149-4]
- 44 **Gress F**, Gottlieb K, Sherman S, Lehman G. Endoscopic ultrasonography-guided fine-needle aspiration biopsy of suspected pancreatic cancer. *Ann Intern Med* 2001; **134**: 459-464 [PMID: 11255521 DOI: 10.7326/0003-4819-134-6-200103200-0010]
- 45 **Perkins GL**, Slater ED, Sanders GK, Prichard JG. Serum tumor markers. *Am Fam Physician* 2003; **68**: 1075-1082 [PMID: 14524394]
- 46 **Goonetilleke KS**, Siriwardena AK. Systematic review of carbohydrate antigen (CA 19-9) as a biochemical marker in the diagnosis of pancreatic cancer. *Eur J Surg Oncol* 2007; **33**: 266-270 [PMID: 17097848 DOI: 10.1016/j.ejso.2006.10.004]
- 47 **Tsutsumi K**, Kawamoto H, Hirao K, Sakakihara I, Yamamoto N, Noma Y, Fujii M, Kato H, Ogawa T, Ishida E, Kuwaki K, Nouse K, Okada H, Yamamoto K. Monitoring of CA19-9 and SPan-1 can facilitate the earlier confirmation of progressing pancreatic cancer during chemotherapy. *Pancreatology* 2012; **12**: 409-416 [PMID: 23127528 DOI: 10.1016/j.pan.2012.07.009]
- 48 **Chung MH**, Gupta RK, Bilchik AJ, Ye W, Yee R, Morton DL. Preoperative serum TA90-IC as an adjunct to serum CA 19-9 in the diagnosis of pancreatic malignancy: a pilot study. *Curr Surg* 2002; **59**: 194-198 [PMID: 16093131 DOI: 10.1016/S0149-7944(01)00594-3]
- 49 **Ishii H**, Okada S, Sato T, Wakasugi H, Saisho H, Furuse J, Ishikawa O, Matsuno S, Yokoyama S. CA 19-9 in evaluating the response to chemotherapy in advanced pancreatic cancer. *Hepatogastroenterology* 1997; **44**: 279-283 [PMID: 9058159]
- 50 **Wong D**, Ko AH, Hwang J, Venook AP, Bergsland EK, Tempero MA. Serum CA19-9 decline compared to radiographic response as a surrogate for clinical outcomes in patients with metastatic pancreatic cancer receiving chemotherapy. *Pancreas* 2008; **37**: 269-274 [PMID: 18815548 DOI: 10.1097/MPA.0b013e31816d8185]
- 51 **Montgomery RC**, Hoffman JP, Riley LB, Rogatko A, Ridge JA, Eisenberg BL. Prediction of recurrence and survival by post-resection CA 19-9 values in patients with adenocarcinoma of the pancreas. *Ann Surg Oncol* 1997; **4**: 551-556 [PMID: 9367020 DOI: 10.1007/BF02305535]
- 52 **Halm U**, Schumann T, Schiefke I, Witzigmann H, Mössner J, Keim V. Decrease of CA 19-9 during chemotherapy with gemcitabine predicts survival time in patients with advanced pancreatic cancer. *Br J Cancer* 2000; **82**: 1013-1016 [PMID: 10737382]
- 53 **Berger AC**, Garcia M, Hoffman JP, Regine WF, Abrams RA, Safran H, Konski A, Benson AB, MacDonald J, Willett CG. Postresection CA 19-9 predicts overall survival in patients with pancreatic cancer treated with adjuvant chemotherapy: a prospective validation by RTOG 9704. *J Clin Oncol* 2008; **26**: 5918-5922 [PMID: 19029412 DOI: 10.1200/JCO.2008.18.6288]
- 54 **Almoguera C**, Shibata D, Forrester K, Martin J, Arnheim N, Perucho M. Most human carcinomas of the exocrine pancreas contain mutant c-K-ras genes. *Cell* 1988; **53**: 549-554 [PMID: 2453289 DOI: 10.1016/0092-8674(88)90571-5]
- 55 **Vogelstein B**, Kinzler KW. Cancer genes and the pathways they control. *Nat Med* 2004; **10**: 789-799 [PMID: 15286780 DOI: 10.1038/nm1087]
- 56 **Hruban RH**, Maitra A, Goggins M. Update on pancreatic intraepithelial neoplasia. *Int J Clin Exp Pathol* 2008; **1**: 306-316 [PMID: 18787611]
- 57 **Pellegata NS**, Sessa F, Renault B, Bonato M, Leone BE, Solcia E, Ranzani GN. K-ras and p53 gene mutations in pancreatic cancer: ductal and nonductal tumors progress through different genetic lesions. *Cancer Res* 1994; **54**: 1556-1560 [PMID: 8137263]
- 58 **Hezel AF**, Kimmelman AC, Stanger BZ, Bardeesy N, Depinho RA. Genetics and biology of pancreatic ductal adenocarcinoma. *Genes Dev* 2006; **20**: 1218-1249 [PMID: 16702400 DOI: 10.1101/gad.1415606]
- 59 **Maitra A**, Hruban RH. Pancreatic cancer. *Annu Rev Pathol* 2008; **3**: 157-188 [PMID: 18039136 DOI: 10.1146/annurev.pathmechdis.3.121806.154305]
- 60 **Bardeesy N**, Cheng KH, Berger JH, Chu GC, Pahler J, Olson P, Hezel AF, Horner J, Lauwers GY, Hanahan D, DePinho RA. Smad4 is dispensable for normal pancreas development yet critical in progression and tumor biology of pancreas cancer. *Genes Dev* 2006; **20**: 3130-3146 [PMID: 17114584 DOI: 10.1101/gad.1478706]
- 61 **Takaori K**, Hruban RH, Maitra A, Tanigawa N. Pancreatic intraepithelial neoplasia. *Pancreas* 2004; **28**: 257-262 [PMID: 15084967 DOI: 10.1097/00006676-200404000-00008]
- 62 **Gedde-Dahl T**, Eriksen JA, Thorsby E, Gaudernack G. T-cell responses against products of oncogenes: generation and characterization of human T-cell clones specific for p21 ras-derived synthetic peptides. *Hum Immunol* 1992; **33**: 266-274 [PMID: 1639630 DOI: 10.1016/0198-8859(92)90334-J]
- 63 **Fossium B**, Gedde-Dahl T, Hansen T, Eriksen JA, Thorsby E, Gaudernack G. Overlapping epitopes encompassing a point mutation (12 Gly-> Arg) in p21 ras can be recognized by HLA-DR, -DP and -DQ restricted T cells. *Eur J Immunol* 1993; **23**: 2687-2691 [PMID: 7691613 DOI: 10.1002/eji.1830231045]
- 64 **Gjertsen MK**, Bakka A, Breivik J, Saeterdal I, Solheim BG, Søreide O, Thorsby E, Gaudernack G. Vaccination with mutant ras peptides and induction of T-cell responsiveness in pancreatic carcinoma patients carrying the corresponding RAS mutation. *Lancet* 1995; **346**: 1399-1400 [PMID: 7475823 DOI: 10.1016/S0140-6736(95)92408-6]
- 65 **Abou-Alfa GK**, Chapman PB, Feilchenfeldt J, Brennan MF, Capanu M, Gansukh B, Jacobs G, Levin A, Neville D, Kelsen DP, O'Reilly EM. Targeting mutated K-ras in pancreatic adenocarcinoma using an adjuvant vaccine. *Am J Clin Oncol* 2011; **34**: 321-325 [PMID: 20686403 DOI: 10.1097/COC.0b013e3181e84b1f]
- 66 **Wedén S**, Klemp M, Gladhaug IP, Møller M, Eriksen JA, Gaudernack G, Buanes T. Long-term follow-up of patients with resected pancreatic cancer following vaccination against mutant K-ras. *Int J Cancer* 2011; **128**: 1120-1128 [PMID: 20473937 DOI: 10.1002/ijc.25449]
- 67 **Redston MS**, Caldas C, Seymour AB, Hruban RH, da Costa L, Yeo CJ, Kern SE. p53 mutations in pancreatic carcinoma

- and evidence of common involvement of homocopolymer tracts in DNA microdeletions. *Cancer Res* 1994; **54**: 3025-3033 [PMID: 8187092]
- 68 **Olive KP**, Tuveson DA, Ruhe ZC, Yin B, Willis NA, Bronson RT, Crowley D, Jacks T. Mutant p53 gain of function in two mouse models of Li-Fraumeni syndrome. *Cell* 2004; **119**: 847-860 [PMID: 15607980 DOI: 10.1016/j.cell.2004.11.004]
- 69 **Morton JP**, Timpson P, Karim SA, Ridgway RA, Athineos D, Doyle B, Jamieson NB, Oien KA, Lowy AM, Brunton VG, Frame MC, Evans TR, Sansom OJ. Mutant p53 drives metastasis and overcomes growth arrest/senescence in pancreatic cancer. *Proc Natl Acad Sci USA* 2010; **107**: 246-251 [PMID: 20018721 DOI: 10.1073/pnas.0908428107]
- 70 **Caldas C**, Hahn SA, da Costa LT, Redston MS, Schutte M, Seymour AB, Weinstein CL, Hruban RH, Yeo CJ, Kern SE. Frequent somatic mutations and homozygous deletions of the p16 (MTS1) gene in pancreatic adenocarcinoma. *Nat Genet* 1994; **8**: 27-32 [PMID: 7726912 DOI: 10.1038/ng0994-27]
- 71 **Schutte M**, Hruban RH, Geradts J, Maynard R, Hilgers W, Rabindran SK, Moskaluk CA, Hahn SA, Schwarte-Waldhoff I, Schmiegel W, Baylin SB, Kern SE, Herman JG. Abrogation of the Rb/p16 tumor-suppressive pathway in virtually all pancreatic carcinomas. *Cancer Res* 1997; **57**: 3126-3130 [PMID: 9242437]
- 72 **Hahn SA**, Hoque AT, Moskaluk CA, da Costa LT, Schutte M, Rozenblum E, Seymour AB, Weinstein CL, Yeo CJ, Hruban RH, Kern SE. Homozygous deletion map at 18q21.1 in pancreatic cancer. *Cancer Res* 1996; **56**: 490-494 [PMID: 8564959]
- 73 **Yachida S**, White CM, Naito Y, Zhong Y, Brosnan JA, Macgregor-Das AM, Morgan RA, Saunders T, Laheru DA, Herman JM, Hruban RH, Klein AP, Jones S, Velculescu V, Wolfgang CL, Iacobuzio-Donahue CA. Clinical significance of the genetic landscape of pancreatic cancer and implications for identification of potential long-term survivors. *Clin Cancer Res* 2012; **18**: 6339-6347 [PMID: 22991414 DOI: 10.1158/1078-0432.CCR-12-1215]
- 74 **Hansel DE**, Kern SE, Hruban RH. Molecular pathogenesis of pancreatic cancer. *Annu Rev Genomics Hum Genet* 2003; **4**: 237-256 [PMID: 14527303 DOI: 10.1146/annurev.genom.4.070802.110341]
- 75 **Biankin AV**, Waddell N, Kassahn KS, Gingras MC, Muthuswamy LB, Johns AL, Miller DK, Wilson PJ, Patch AM, Wu J, Chang DK, Cowley MJ, Gardiner BB, Song S, Harliwong I, Idrisoglu S, Nourse C, Nourbakhsh E, Manning S, Wani S, Gongora M, Pajic M, Scarlett CJ, Gill AJ, Pinho AV, Rooman I, Anderson M, Holmes O, Leonard C, Taylor D, Wood S, Xu Q, Nones K, Fink JL, Christ A, Bruxner T, Cloonan N, Kolle G, Newell F, Pinese M, Mead RS, Humphris JL, Kaplan W, Jones MD, Colvin EK, Nagrial AM, Humphrey ES, Chou A, Chin VT, Chantrill LA, Mawson A, Samra JS, Kench JG, Lovell JA, Daly RJ, Merrett ND, Toon C, Epari K, Nguyen NQ, Barbour A, Zeps N, Kakkar N, Zhao F, Wu YQ, Wang M, Muzny DM, Fisher WE, Brunicardi FC, Hodges SE, Reid JG, Drummond J, Chang K, Han Y, Lewis LR, Dinh H, Buhay CJ, Beck T, Timms L, Sam M, Begley K, Brown A, Pai D, Panchal A, Buchner N, De Borja R, Denroche RE, Yung CK, Serra S, Onetto N, Mukhopadhyay D, Tsao MS, Shaw PA, Petersen GM, Gallinger S, Hruban RH, Maitra A, Iacobuzio-Donahue CA, Schulick RD, Wolfgang CL, Morgan RA, Lawlor RT, Cappelli P, Corbo V, Scardoni M, Tortora G, Tempero MA, Mann KM, Jenkins NA, Perez-Mancera PA, Adams DJ, Largaespada DA, Wessels LF, Rust AG, Stein LD, Tuveson DA, Copeland NG, Musgrove EA, Scarpa A, Eshleman JR, Hudson TJ, Sutherland RL, Wheeler DA, Pearson JV, McPherson JD, Gibbs RA, Grimmond SM. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nature* 2012; **491**: 399-405 [PMID: 23103869 DOI: 10.1038/nature11547]
- 76 **Pérez-Mancera PA**, Rust AG, van der Weyden L, Kristiansen G, Li A, Sarver AL, Silverstein KA, Grützmann R, Aust D, Rümmele P, Knösel T, Herd C, Stemple DL, Kettleborough R, Brosnan JA, Li A, Morgan R, Knight S, Yu J, Stegeman S, Collier LS, ten Hoeve JJ, de Ridder J, Klein AP, Goggins M, Hruban RH, Chang DK, Biankin AV, Grimmond SM, Wessels LF, Wood SA, Iacobuzio-Donahue CA, Pilarsky C, Largaespada DA, Adams DJ, Tuveson DA. The deubiquitinase USP9X suppresses pancreatic ductal adenocarcinoma. *Nature* 2012; **486**: 266-270 [PMID: 22699621]
- 77 **Jones S**, Zhang X, Parsons DW, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Kamiyama H, Jimeno A, Hong SM, Fu B, Lin MT, Calhoun ES, Kamiyama M, Walter K, Nikolskaya T, Nikolsky Y, Hartigan J, Smith DR, Hidalgo M, Leach SD, Klein AP, Jaffee EM, Goggins M, Maitra A, Iacobuzio-Donahue C, Eshleman JR, Kern SE, Hruban RH, Karchin R, Papadopoulos N, Parmigiani G, Vogelstein B, Velculescu VE, Kinzler KW. Core signaling pathways in human pancreatic cancers revealed by global genomic analyses. *Science* 2008; **321**: 1801-1806 [PMID: 18772397 DOI: 10.1126/science.1164368]
- 78 **Chu GC**, Kimmelman AC, Hezel AF, DePinho RA. Stromal biology of pancreatic cancer. *J Cell Biochem* 2007; **101**: 887-907 [PMID: 17266048 DOI: 10.1002/jcb.21209]
- 79 **Neesse A**, Michl P, Frese KK, Feig C, Cook N, Jacobetz MA, Lolkema MP, Buchholz M, Olive KP, Gress TM, Tuveson DA. Stromal biology and therapy in pancreatic cancer. *Gut* 2011; **60**: 861-868 [PMID: 20966025 DOI: 10.1136/gut.2010.226092]
- 80 **Collins MA**, Bednar F, Zhang Y, Brisset JC, Galbán S, Galbán CJ, Rakshit S, Flannagan KS, Adsay NV, Pasca di Magliano M. Oncogenic Kras is required for both the initiation and maintenance of pancreatic cancer in mice. *J Clin Invest* 2012; **122**: 639-653 [PMID: 2232209 DOI: 10.1172/JCI59227]
- 81 **Ying H**, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sananikone E, Locasale JW, Son J, Zhang H, Colloff JL, Yan H, Wang W, Chen S, Viale A, Zheng H, Paik JH, Lim C, Guimaraes AR, Martin ES, Chang J, Hezel AF, Perry SR, Hu J, Gan B, Xiao Y, Asara JM, Weissleder R, Wang YA, Chin L, Cantley LC, DePinho RA. Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* 2012; **149**: 656-670 [PMID: 22541435 DOI: 10.1016/j.cell.2012.01.058]
- 82 **Nolan-Stevaux O**, Lau J, Truitt ML, Chu GC, Hebrok M, Fernández-Zapico ME, Hanahan D. GLI1 is regulated through Smoothed-independent mechanisms in neoplastic pancreatic ducts and mediates PDAC cell survival and transformation. *Genes Dev* 2009; **23**: 24-36 [PMID: 19136624 DOI: 10.1101/gad.1753809]
- 83 **Bailey JM**, Swanson BJ, Hamada T, Eggers JP, Singh PK, Caffery T, Ouellette MM, Hollingsworth MA. Sonic hedgehog promotes desmoplasia in pancreatic cancer. *Clin Cancer Res* 2008; **14**: 5995-6004 [PMID: 18829478 DOI: 10.1158/1078-0432.CCR-08-0291]
- 84 **Tian H**, Callahan CA, DuFree KJ, Darbonne WC, Ahn CP, Scales SJ, de Sauvage FJ. Hedgehog signaling is restricted to the stromal compartment during pancreatic carcinogenesis. *Proc Natl Acad Sci USA* 2009; **106**: 4254-4259 [PMID: 19246386 DOI: 10.1073/pnas.0813203106]
- 85 **Olive KP**, Jacobetz MA, Davidson CJ, Gopinathan A, McIntyre D, Honess D, Madhu B, Goldgraben MA, Caldwell ME, Allard D, Frese KK, Denicola G, Feig C, Combs C, Winter SP, Ireland-Zecchini H, Reichelt S, Howat WJ, Chang A, Dhara M, Wang L, Rückert F, Grützmann R, Pilarsky C, Izeradjene K, Hingorani SR, Huang P, Davies SE, Plunkett W, Egorin M, Hruban RH, Whitebread N, McGovern K, Adams J, Iacobuzio-Donahue C, Griffiths J, Tuveson DA. Inhibition of Hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer. *Science* 2009; **324**: 1457-1461 [PMID: 19460966 DOI: 10.1126/science.1171362]
- 86 **Komar G**, Kauhanen S, Liukko K, Seppänen M, Kajander S, Ovaska J, Nuutila P, Minn H. Decreased blood flow with increased metabolic activity: a novel sign of pancreatic tumor

- aggressiveness. *Clin Cancer Res* 2009; **15**: 5511-5517 [PMID: 19706808 DOI: 10.1158/1078-0432.CCR-09-0414]
- 87 **Bayne LJ**, Beatty GL, Jhala N, Clark CE, Rhim AD, Stanger BZ, Vonderheide RH. Tumor-derived granulocyte-macrophage colony-stimulating factor regulates myeloid inflammation and T cell immunity in pancreatic cancer. *Cancer Cell* 2012; **21**: 822-835 [PMID: 22698406 DOI: 10.1016/j.ccr.2012.04.025]
- 88 **Pylayeva-Gupta Y**, Lee KE, Hajdu CH, Miller G, Bar-Sagi D. Oncogenic Kras-induced GM-CSF production promotes the development of pancreatic neoplasia. *Cancer Cell* 2012; **21**: 836-847 [PMID: 22698407 DOI: 10.1016/j.ccr.2012.04.024]
- 89 **Li D**, Xie K, Wolff R, Abbruzzese JL. Pancreatic cancer. *Lancet* 2004; **363**: 1049-1057 [PMID: 15051286 DOI: 10.1016/S0140-6736(04)15841-8]
- 90 **Wagner M**, Redaelli C, Lietz M, Seiler CA, Friess H, Büchler MW. Curative resection is the single most important factor determining outcome in patients with pancreatic adenocarcinoma. *Br J Surg* 2004; **91**: 586-594 [PMID: 15122610 DOI: 10.1002/bjs.4484]
- 91 **Uchida H**, Shibata K, Iwaki K, Kai S, Ohta M, Kitano S. Ampullary cancer and preoperative jaundice: possible indication of the minimal surgery. *Hepatogastroenterology* 2009; **56**: 1194-1198 [PMID: 19760968]
- 92 **van der Gaag NA**, Kloek JJ, de Castro SM, Busch OR, van Gulik TM, Gouma DJ. Preoperative biliary drainage in patients with obstructive jaundice: history and current status. *J Gastrointest Surg* 2009; **13**: 814-820 [PMID: 18726134 DOI: 10.1007/s11605-008-0618-4]
- 93 **Sewnath ME**, Karsten TM, Prins MH, Rauws EJ, Obertop H, Gouma DJ. A meta-analysis on the efficacy of preoperative biliary drainage for tumors causing obstructive jaundice. *Ann Surg* 2002; **236**: 17-27 [PMID: 12131081 DOI: 10.1097/0000658-200207000-00005]
- 94 **van der Gaag NA**, Rauws EA, van Eijck CH, Bruno MJ, van der Harst E, Kubben FJ, Gerritsen JJ, Greve JW, Gerhards MF, de Hingh IH, Klinkenbijl JH, Nio CY, de Castro SM, Busch OR, van Gulik TM, Bossuyt PM, Gouma DJ. Preoperative biliary drainage for cancer of the head of the pancreas. *N Engl J Med* 2010; **362**: 129-137 [PMID: 20071702 DOI: 10.1056/NEJMoa0903230]
- 95 **Jenkins LJ**, Parmar AD, Han Y, Duncan CB, Sheffield KM, Brown KM, Riall TS. Current trends in preoperative biliary stenting in patients with pancreatic cancer. *Surgery* 2013; **154**: 179-189 [PMID: 23889947 DOI: 10.1016/j.surg.2013.03.016]
- 96 **Riediger H**, Keck T, Wellner U, zur Hausen A, Adam U, Hopt UT, Makowiec F. The lymph node ratio is the strongest prognostic factor after resection of pancreatic cancer. *J Gastrointest Surg* 2009; **13**: 1337-1344 [PMID: 19418101 DOI: 10.1007/s11605-009-0919-2]
- 97 **Wasif N**, Ko CY, Farrell J, Wainberg Z, Hines OJ, Reber H, Tomlinson JS. Impact of tumor grade on prognosis in pancreatic cancer: should we include grade in AJCC staging? *Ann Surg Oncol* 2010; **17**: 2312-2320 [PMID: 20422460 DOI: 10.1245/s10434-010-1071-7]
- 98 **Fortner JG**, Klimstra DS, Senie RT, Maclean BJ. Tumor size is the primary prognosticator for pancreatic cancer after regional pancreatectomy. *Ann Surg* 1996; **223**: 147-153 [PMID: 8597508 DOI: 10.1097/0000658-199602000-00006]
- 99 **Skipworth RJ**, Parks RW, Stephens NA, Graham C, Brewster DH, Garden OJ, Paterson-Brown S. The relationship between hospital volume and post-operative mortality rates for upper gastrointestinal cancer resections: Scotland 1982-2003. *Eur J Surg Oncol* 2010; **36**: 141-147 [PMID: 19879717 DOI: 10.1016/j.ejso.2009.10.004]
- 100 **Finks JF**, Osborne NH, Birkmeyer JD. Trends in hospital volume and operative mortality for high-risk surgery. *N Engl J Med* 2011; **364**: 2128-2137 [PMID: 21631325 DOI: 10.1056/NEJMsa1010705]
- 101 **Lemmens VE**, Bosscha K, van der Schelling G, Brennk-meijer S, Coebergh JW, de Hingh IH. Improving outcome for patients with pancreatic cancer through centralization. *Br J Surg* 2011; **98**: 1455-1462 [PMID: 21717423 DOI: 10.1002/bjs.7581]
- 102 **Gillen S**, Schuster T, Meyer Zum Büschenfelde C, Friess H, Kleeff J. Preoperative/neoadjuvant therapy in pancreatic cancer: a systematic review and meta-analysis of response and resection percentages. *PLoS Med* 2010; **7**: e1000267 [PMID: 20422030 DOI: 10.1371/journal.pmed.1000267]
- 103 **McClaine RJ**, Lowy AM, Sussman JJ, Schmulowitz N, Grisell DL, Ahmad SA. Neoadjuvant therapy may lead to successful surgical resection and improved survival in patients with borderline resectable pancreatic cancer. *HPB (Oxford)* 2010; **12**: 73-79 [PMID: 20495649 DOI: 10.1111/j.1477-2574.2009.00136.x]
- 104 **Le Scodan R**, Mornex F, Partensky C, Mercier C, Valette PJ, Ychou M, Roy P, Scazec JY. Histopathological response to preoperative chemoradiation for resectable pancreatic adenocarcinoma: the French Phase II FFCD 9704-SFRO Trial. *Am J Clin Oncol* 2008; **31**: 545-552 [PMID: 19060585 DOI: 10.1097/COC.0b013e318172d5c5]
- 105 **White RR**, Xie HB, Gottfried MR, Czito BG, Hurwitz HI, Morse MA, Blobe GC, Paulson EK, Baillie J, Branch MS, Jowell PS, Clary BM, Pappas TN, Tyler DS. Significance of histological response to preoperative chemoradiotherapy for pancreatic cancer. *Ann Surg Oncol* 2005; **12**: 214-221 [PMID: 15827813 DOI: 10.1245/ASO.2005.03.105]
- 106 **Kim HJ**, Czischke K, Brennan MF, Conlon KC. Does neoadjuvant chemoradiation downstage locally advanced pancreatic cancer? *J Gastrointest Surg* 2002; **6**: 763-769 [PMID: 12399067 DOI: 10.1016/S1091-255X(02)00017-3]
- 107 **Massucco P**, Capussotti L, Magnino A, Sperti E, Gatti M, Muratore A, Sgotto E, Gabriele P, Aglietta M. Pancreatic resections after chemoradiotherapy for locally advanced ductal adenocarcinoma: analysis of perioperative outcome and survival. *Ann Surg Oncol* 2006; **13**: 1201-1208 [PMID: 16955382 DOI: 10.1245/s10434-006-9032-x]
- 108 **Neoptolemos JP**, Stocken DD, Bassi C, Ghaneh P, Cunningham D, Goldstein D, Padbury R, Moore MJ, Gallinger S, Mariette C, Wente MN, Izbicki JR, Friess H, Lerch MM, Dervenis C, Oláh A, Butturini G, Doi R, Lind PA, Smith D, Valle JW, Palmer DH, Buckels JA, Thompson J, McKay CJ, Rawcliffe CL, Büchler MW. Adjuvant chemotherapy with fluorouracil plus folinic acid vs gemcitabine following pancreatic cancer resection: a randomized controlled trial. *JAMA* 2010; **304**: 1073-1081 [PMID: 20823433 DOI: 10.1001/jama.2010.1275]
- 109 **Dineen SP**, Roland CL, Greer R, Carbon JG, Toombs JE, Gupta P, Bardeesy N, Sun H, Williams N, Minna JD, Brekken RA. Smac mimetic increases chemotherapy response and improves survival in mice with pancreatic cancer. *Cancer Res* 2010; **70**: 2852-2861 [PMID: 20332237 DOI: 10.1158/0008-5472.CAN-09-3892]
- 110 **Hwang JH**, Voortman J, Giovannetti E, Steinberg SM, Leon LG, Kim YT, Funel N, Park JK, Kim MA, Kang GH, Kim SW, Del Chiaro M, Peters GJ, Giaccone G. Identification of microRNA-21 as a biomarker for chemoresistance and clinical outcome following adjuvant therapy in resectable pancreatic cancer. *PLoS One* 2010; **5**: e10630 [PMID: 20498843 DOI: 10.1371/journal.pone.0010630]
- 111 **Harikumar KB**, Kunnumakkara AB, Sethi G, Diagaradjane P, Anand P, Pandey MK, Gelovani J, Krishnan S, Guha S, Aggarwal BB. Resveratrol, a multitargeted agent, can enhance antitumor activity of gemcitabine in vitro and in orthotopic mouse model of human pancreatic cancer. *Int J Cancer* 2010; **127**: 257-268 [PMID: 19908231]
- 112 **Cunningham D**, Chau I, Stocken DD, Valle JW, Smith D, Steward W, Harper PG, Dunn J, Tudur-Smith C, West J, Falk S, Crellin A, Adab F, Thompson J, Leonard P, Ostrowski J, Eatock M, Scheithauer W, Herrmann R, Neoptolemos JP.



- Phase III randomized comparison of gemcitabine versus gemcitabine plus capecitabine in patients with advanced pancreatic cancer. *J Clin Oncol* 2009; **27**: 5513-5518 [PMID: 19858379 DOI: 10.1200/JCO.2009.24.2446]
- 113 **Banerjee S**, Kaseb AO, Wang Z, Kong D, Mohammad M, Padhye S, Sarkar FH, Mohammad RM. Antitumor activity of gemcitabine and oxaliplatin is augmented by thymoquinone in pancreatic cancer. *Cancer Res* 2009; **69**: 5575-5583 [PMID: 19549912 DOI: 10.1158/0008-5472.CAN-08-4235]
- 114 **Baylot V**, Andrieu C, Katsogiannou M, Taieb D, Garcia S, Giustiano S, Acunzo J, Iovanna J, Gleave M, Garrido C, Rocchi P. OGX-427 inhibits tumor progression and enhances gemcitabine chemotherapy in pancreatic cancer. *Cell Death Dis* 2011; **2**: e221 [PMID: 22012255 DOI: 10.1038/cddis.2011.104]
- 115 **Kanai M**, Yoshimura K, Asada M, Imaizumi A, Suzuki C, Matsumoto S, Nishimura T, Mori Y, Masui T, Kawaguchi Y, Yanagihara K, Yazumi S, Chiba T, Guha S, Aggarwal BB. A phase I/II study of gemcitabine-based chemotherapy plus curcumin for patients with gemcitabine-resistant pancreatic cancer. *Cancer Chemother Pharmacol* 2011; **68**: 157-164 [PMID: 20859741 DOI: 10.1007/s00280-010-1470-2]
- 116 **Farrell JJ**, Elsaleh H, Garcia M, Lai R, Ammar A, Regine WF, Abrams R, Benson AB, Macdonald J, Cass CE, Dicker AP, Mackey JR. Human equilibrative nucleoside transporter 1 levels predict response to gemcitabine in patients with pancreatic cancer. *Gastroenterology* 2009; **136**: 187-195 [PMID: 18992248 DOI: 10.1053/j.gastro.2008.09.067]
- 117 **Fujita H**, Ohuchida K, Mizumoto K, Itaba S, Ito T, Nakata K, Yu J, Kayashima T, Souzaki R, Tajiri T, Manabe T, Ohtsuka T, Tanaka M. Gene expression levels as predictive markers of outcome in pancreatic cancer after gemcitabine-based adjuvant chemotherapy. *Neoplasia* 2010; **12**: 807-817 [PMID: 20927319]
- 118 **Burris HA**, Moore MJ, Andersen J, Green MR, Rothenberg ML, Modiano MR, Cripps MC, Portenoy RK, Storniolo AM, Tarassoff P, Nelson R, Dorr FA, Stephens CD, Von Hoff DD. Improvements in survival and clinical benefit with gemcitabine as first-line therapy for patients with advanced pancreas cancer: a randomized trial. *J Clin Oncol* 1997; **15**: 2403-2413 [PMID: 9196156]
- 119 **Heinemann V**, Boeck S, Hinke A, Labianca R, Louvet C. Meta-analysis of randomized trials: evaluation of benefit from gemcitabine-based combination chemotherapy applied in advanced pancreatic cancer. *BMC Cancer* 2008; **8**: 82 [PMID: 18373843 DOI: 10.1186/11471-2407-8-82]
- 120 **Renì M**, Cordio S, Milandri C, Passoni P, Bonetto E, Oliani C, Luppi G, Nicoletti R, Galli L, Bordonaro R, Passardi A, Zerbi A, Balzano G, Aldrighetti L, Staudacher C, Villa E, Di Carlo V. Gemcitabine versus cisplatin, epirubicin, fluorouracil, and gemcitabine in advanced pancreatic cancer: a randomised controlled multicentre phase III trial. *Lancet Oncol* 2005; **6**: 369-376 [PMID: 15925814 DOI: 10.1016/S1470-2045(05)70175-3]
- 121 **Conroy T**, Desseigne F, Ychou M, Bouché O, Guimbaud R, Bécouarn Y, Adenis A, Raoul JL, Gourgou-Bourgade S, de la Fouchardière C, Bennouna J, Bachet JB, Khemissa-Akouz F, Péré-Vergé D, Delbaldo C, Assenat E, Chauffert B, Michel P, Montoto-Grillot C, Ducreux M. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med* 2011; **364**: 1817-1825 [PMID: 21561347 DOI: 10.1056/NEJMoa1011923]
- 122 **Moore MJ**, Goldstein D, Hamm J, Figer A, Hecht JR, Gallinger S, Au HJ, Murawa P, Walde D, Wolff RA, Campos D, Lim R, Ding K, Clark G, Voskoglou-Nomikos T, Ptasynski M, Parulekar W. Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* 2007; **25**: 1960-1966 [PMID: 17452677 DOI: 10.1200/JCO.2006.07.9525]
- 123 **Edmonds C**, Cengel KA. Tumor-Stroma interactions in pancreatic cancer: Will this SPARC prove a raging fire? *Cancer Biol Ther* 2008; **7**: 1816-1817 [PMID: 19029794 DOI: 10.4161/cbt.7.11.7285]
- 124 **Infante JR**, Matsubayashi H, Sato N, Tonascia J, Klein AP, Riall TA, Yeo C, Iacobuzio-Donahue C, Goggins M. Peritumoral fibroblast SPARC expression and patient outcome with resectable pancreatic adenocarcinoma. *J Clin Oncol* 2007; **25**: 319-325 [PMID: 17235047 DOI: 10.1200/JCO.2006.07.8824]
- 125 **Von Hoff DD**, Ramanathan RK, Borad MJ, Laheru DA, Smith LS, Wood TE, Korn RL, Desai N, Trieu V, Iglesias JL, Zhang H, Soon-Shiong P, Shi T, Rajeshkumar NV, Maitra A, Hidalgo M. Gemcitabine plus nab-paclitaxel is an active regimen in patients with advanced pancreatic cancer: a phase I/II trial. *J Clin Oncol* 2011; **29**: 4548-4554 [PMID: 21969517 DOI: 10.1200/JCO.2011.36.5742]
- 126 **Caridi G**, Gigante M, Ravani P, Trivelli A, Barbano G, Scolarì F, Dagnino M, Murer L, Murtas C, Edefonti A, Allegri L, Amore A, Coppo R, Emma F, De Palo T, Penza R, Gesualdo L, Ghiggeri GM. Clinical features and long-term outcome of nephrotic syndrome associated with heterozygous NPHS1 and NPHS2 mutations. *Clin J Am Soc Nephrol* 2009; **4**: 1065-1072 [PMID: 19406966 DOI: 10.2215/CJN.03910808]
- 127 **Custodio A**, Puente J, Sastre J, Díaz-Rubio E. Second-line therapy for advanced pancreatic cancer: a review of the literature and future directions. *Cancer Treat Rev* 2009; **35**: 676-684 [PMID: 19758760 DOI: 10.1016/j.ctrv.2009.08.012]
- 128 **Boeck S**, Ankerst DP, Heinemann V. The role of adjuvant chemotherapy for patients with resected pancreatic cancer: systematic review of randomized controlled trials and meta-analysis. *Oncology* 2007; **72**: 314-321 [PMID: 18187951 DOI: 10.1159/000113054]
- 129 **Lim KH**, Chung E, Khan A, Cao D, Linehan D, Ben-Josef E, Wang-Gillam A. Neoadjuvant therapy of pancreatic cancer: the emerging paradigm? *Oncologist* 2012; **17**: 192-200 [PMID: 22250057 DOI: 10.1634/theoncologist.2011-0268]
- 130 **Evans DB**, Rich TA, Byrd DR, Cleary KR, Connelly JH, Levin B, Charnsangavej C, Fenoglio CJ, Ames FC. Preoperative chemoradiation and pancreaticoduodenectomy for adenocarcinoma of the pancreas. *Arch Surg* 1992; **127**: 1335-1339 [PMID: 1359851 DOI: 10.1001/archsurg.1992.01420110083017]
- 131 **Pisters PW**, Abbruzzese JL, Janjan NA, Cleary KR, Charnsangavej C, Goswitz MS, Rich TA, Raijman I, Wolff RA, Lenzi R, Lee JE, Evans DB. Rapid-fractionation preoperative chemoradiation, pancreaticoduodenectomy, and intraoperative radiation therapy for resectable pancreatic adenocarcinoma. *J Clin Oncol* 1998; **16**: 3843-3850 [PMID: 9850029]
- 132 **Krishnan S**, Rana V, Evans DB, Varadhachary G, Das P, Bhatia S, Delclos ME, Janjan NA, Wolff RA, Crane CH, Pisters PW. Role of adjuvant chemoradiation therapy in adenocarcinomas of the ampulla of vater. *Int J Radiat Oncol Biol Phys* 2008; **70**: 735-743 [PMID: 17980502 DOI: 10.1016/j.ijrobp.2007.07.2327]
- 133 **Hong TS**, Ryan DP, Blaszkowsky LS, Mamon HJ, Kwak EL, Mino-Kenudson M, Adams J, Yeap B, Winrich B, DeLaney TF, Fernandez-Del Castillo C. Phase I study of preoperative short-course chemoradiation with proton beam therapy and capecitabine for resectable pancreatic ductal adenocarcinoma of the head. *Int J Radiat Oncol Biol Phys* 2011; **79**: 151-157 [PMID: 20421151 DOI: 10.1016/j.ijrobp.2009.10.061]
- 134 **Hadley DW**, Jenkins JF, Steinberg SM, Liewehr D, Moller S, Martin JC, Calzone KA, Soballe PW, Kirsch IR. Perceptions of cancer risks and predictors of colon and endometrial cancer screening in women undergoing genetic testing for Lynch syndrome. *J Clin Oncol* 2008; **26**: 948-954 [PMID: 18281669 DOI: 10.1200/JCO.2007.13.9014]
- 135 **Patel M**, Hoffe S, Malafa M, Hodul P, Klapman J, Centeno B, Kim J, Helm J, Valone T, Springett G. Neoadjuvant GTX chemotherapy and IMRT-based chemoradiation for borderline resectable pancreatic cancer. *J Surg Oncol* 2011; **104**: 155-161



- [PMID: 21520097 DOI: 10.1002/jso.21954]
- 136 **Yeo CJ**, Cameron JL, Sohn TA, Lillemoe KD, Pitt HA, Talamini MA, Hruban RH, Ord SE, Sauter PK, Coleman J, Zahurak ML, Grochow LB, Abrams RA. Six hundred fifty consecutive pancreaticoduodenectomies in the 1990s: pathology, complications, and outcomes. *Ann Surg* 1997; **226**: 248-57; discussion 257-60 [PMID: 9339931 DOI: 10.1097/0000658-199709000-00004]
  - 137 **Neoptolemos JP**, Stocken DD, Friess H, Bassi C, Dunn JA, Hickey H, Beger H, Fernandez-Cruz L, Dervenis C, Lacaine F, Falconi M, Pederzoli P, Pap A, Spooner D, Kerr DJ, Büchler MW. A randomized trial of chemoradiotherapy and chemotherapy after resection of pancreatic cancer. *N Engl J Med* 2004; **350**: 1200-1210 [PMID: 15028824 DOI: 10.1056/NEJMoa032295]
  - 138 **Regine WF**, Winter KA, Abrams R, Safran H, Hoffman JP, Konski A, Benson AB, Macdonald JS, Rich TA, Willett CG. Fluorouracil-based chemoradiation with either gemcitabine or fluorouracil chemotherapy after resection of pancreatic adenocarcinoma: 5-year analysis of the U.S. Intergroup/ RTOG 9704 phase III trial. *Ann Surg Oncol* 2011; **18**: 1319-1326 [PMID: 21499862 DOI: 10.1245/s10434-011-1630-6]
  - 139 **Oettle H**, Post S, Neuhaus P, Gellert K, Langrehr J, Ridwelski K, Schramm H, Fahlke J, Zuelke C, Burkart C, Gutberlet K, Kettner E, Schmalenberg H, Weigang-Koehler K, Bechthold WO, Niedergethmann M, Schmidt-Wolf I, Roll L, Doerken B, Riess H. Adjuvant chemotherapy with gemcitabine vs observation in patients undergoing curative-intent resection of pancreatic cancer: a randomized controlled trial. *JAMA* 2007; **297**: 267-277 [PMID: 17227978 DOI: 10.1001/jama.297.3.267]
  - 140 **Yeo CJ**, Abrams RA, Grochow LB, Sohn TA, Ord SE, Hruban RH, Zahurak ML, Dooley WC, Coleman J, Sauter PK, Pitt HA, Lillemoe KD, Cameron JL. Pancreaticoduodenectomy for pancreatic adenocarcinoma: postoperative adjuvant chemoradiation improves survival. A prospective, single-institution experience. *Ann Surg* 1997; **225**: 621-33; discussion 633-6 [PMID: 9193189 DOI: 10.1097/0000658-199705000-00018]
  - 141 **Picozzi VJ**, Kozarek RA, Traverso LW. Interferon-based adjuvant chemoradiation therapy after pancreaticoduodenectomy for pancreatic adenocarcinoma. *Am J Surg* 2003; **185**: 476-480 [PMID: 12727570 DOI: 10.1016/S0002-9610(03)00051-5]
  - 142 **Hsu CC**, Herman JM, Corsini MM, Winter JM, Callister MD, Haddock MG, Cameron JL, Pawlik TM, Schulick RD, Wolfgang CL, Laheru DA, Farnell MB, Swartz MJ, Gunderson LL, Miller RC. Adjuvant chemoradiation for pancreatic adenocarcinoma: the Johns Hopkins Hospital-Mayo Clinic collaborative study. *Ann Surg Oncol* 2010; **17**: 981-990 [PMID: 20087786 DOI: 10.1245/s10434-009-0743-7]
  - 143 **Wayne JD**, Abdalla EK, Wolff RA, Crane CH, Pisters PW, Evans DB. Localized adenocarcinoma of the pancreas: the rationale for preoperative chemoradiation. *Oncologist* 2002; **7**: 34-45 [PMID: 11854545 DOI: 10.1634/theoncologist.7-1-34]
  - 144 **Moertel CG**, Frytak S, Hahn RG, O'Connell MJ, Reitemeier RJ, Rubin J, Schutt AJ, Weiland LH, Childs DS, Holbrook MA, Lavin PT, Livstone E, Spiro H, Knowlton A, Kalser M, Barkin J, Lessner H, Mann-Kaplan R, Ramming K, Douglas HO, Thomas P, Nave H, Bateman J, Lokich J, Brooks J, Chaffey J, Corson JM, Zamcheck N, Novak JW. Therapy of locally unresectable pancreatic carcinoma: a randomized comparison of high dose (6000 rads) radiation alone, moderate dose radiation (4000 rads + 5-fluorouracil), and high dose radiation + 5-fluorouracil: The Gastrointestinal Tumor Study Group. *Cancer* 1981; **48**: 1705-1710 [PMID: 7284971]
  - 145 Treatment of locally unresectable carcinoma of the pancreas: comparison of combined-modality therapy (chemotherapy plus radiotherapy) to chemotherapy alone. Gastrointestinal Tumor Study Group. *J Natl Cancer Inst* 1988; **80**: 751-755 [PMID: 2898536]
  - 146 **Loehrer PJ**, Feng Y, Cardenes H, Wagner L, Brell JM, Cella D, Flynn P, Ramanathan RK, Crane CH, Alberts SR, Benson AB. Gemcitabine alone versus gemcitabine plus radiotherapy in patients with locally advanced pancreatic cancer: an Eastern Cooperative Oncology Group trial. *J Clin Oncol* 2011; **29**: 4105-4112 [PMID: 21969502 DOI: 10.1200/JCO.2011.34.8904]
  - 147 **Schellenberg D**, Kim J, Christman-Skieller C, Chun CL, Colombo LA, Ford JM, Fisher GA, Kunz PL, Van Dam J, Quon A, Desser TS, Norton J, Hsu A, Maxim PG, Xing L, Goodman KA, Chang DT, Koong AC. Single-fraction stereotactic body radiation therapy and sequential gemcitabine for the treatment of locally advanced pancreatic cancer. *Int J Radiat Oncol Biol Phys* 2011; **81**: 181-188 [PMID: 21549517 DOI: 10.1016/j.ijrobp.2010.05.006]
  - 148 **Didolkar MS**, Coleman CW, Brenner MJ, Chu KU, Olexa N, Stanwyck E, Yu A, Neerchal N, Rabinowitz S. Image-guided stereotactic radiosurgery for locally advanced pancreatic adenocarcinoma results of first 85 patients. *J Gastrointest Surg* 2010; **14**: 1547-1559 [PMID: 20839073 DOI: 10.1007/s11605-010-1323-7]
  - 149 **Rwigema JC**, Heron DE, Parikh SD, Zeh HJ, Moser JA, Bahary N, Ashby K, Burton SA. Adjuvant stereotactic body radiotherapy for resected pancreatic adenocarcinoma with close or positive margins. *J Gastrointest Cancer* 2012; **43**: 70-76 [PMID: 20809393 DOI: 10.1007/s12029-010-9203-7]
  - 150 **Leibel SA**, Fuks Z, Zelefsky MJ, Hunt M, Burman CM, Mageras GS, Chui CS, Jackson A, Amols HI, Ling CC. Technological advances in external-beam radiation therapy for the treatment of localized prostate cancer. *Semin Oncol* 2003; **30**: 596-615 [PMID: 14571409]
  - 151 **Ling CC**, Burman C, Chui CS, Kutcher GJ, Leibel SA, LoSasso T, Mohan R, Bortfeld T, Reinstein L, Spirou S, Wang XH, Wu Q, Zelefsky M, Fuks Z. Conformal radiation treatment of prostate cancer using inversely-planned intensity-modulated photon beams produced with dynamic multileaf collimation. *Int J Radiat Oncol Biol Phys* 1996; **35**: 721-730 [PMID: 8690638 DOI: 10.1016/0360-3016(96)00174-5]
  - 152 **Varadarajulu S**, Trevino JM, Shen S, Jacob R. The use of endoscopic ultrasound-guided gold markers in image-guided radiation therapy of pancreatic cancers: a case series. *Endoscopy* 2010; **42**: 423-425 [PMID: 20232282 DOI: 10.1055/s-0029-1243989]
  - 153 **Villarreal MC**, Rajeshkumar NV, Garrido-Laguna I, De Jesus-Acosta A, Jones S, Maitra A, Hruban RH, Eshleman JR, Klein A, Laheru D, Donehower R, Hidalgo M. Personalizing cancer treatment in the age of global genomic analyses: PALB2 gene mutations and the response to DNA damaging agents in pancreatic cancer. *Mol Cancer Ther* 2011; **10**: 3-8 [PMID: 21135251 DOI: 10.1158/1535-7163.MCT-10-0893]
  - 154 **Yanagimoto H**, Shiomi H, Satoi S, Mine T, Toyokawa H, Yamamoto T, Tani T, Yamada A, Kwon AH, Komatsu N, Itoh K, Noguchi M. A phase II study of personalized peptide vaccination combined with gemcitabine for non-resectable pancreatic cancer patients. *Oncol Rep* 2010; **24**: 795-801 [PMID: 20664989]
  - 155 **Bapat AA**, Hostetter G, Von Hoff DD, Han H. Perineural invasion and associated pain in pancreatic cancer. *Nat Rev Cancer* 2011; **11**: 695-707 [PMID: 21941281 DOI: 10.1038/nrc3131]
  - 156 **Vincent A**, Herman J, Schulick R, Hruban RH, Goggins M. Pancreatic cancer. *Lancet* 2011; **378**: 607-620 [PMID: 21620466 DOI: 10.1016/S0140-6736(10)62307-0]
  - 157 **Kim EJ**, Simeone DM. Advances in pancreatic cancer. *Curr Opin Gastroenterol* 2011; **27**: 460-466 [PMID: 21778878 DOI: 10.1097/MOG.0b013e328349e31f]
  - 158 **Costello E**, Neoptolemos J. Enhancing the translation of cancer biomarkers. *Br J Surg* 2011; **98**: 1039-1040 [PMID: 21560125 DOI: 10.1002/bjs.7556]
  - 159 **Yachida S**, Jones S, Bozic I, Antal T, Leary R, Fu B, Kamiyama M, Hruban RH, Eshleman JR, Nowak MA, Velculescu VE, Kinzler KW, Vogelstein B, Iacobuzio-Donahue CA.

- Distant metastasis occurs late during the genetic evolution of pancreatic cancer. *Nature* 2010; **467**: 1114-1117 [PMID: 20981102 DOI: 10.1038/nature09515,]
- 160 **Collisson E**, Tempero M. Blinded by the light: molecular imaging in pancreatic adenocarcinoma. *Clin Cancer Res* 2011; **17**: 203-205 [PMID: 21106729 DOI: 10.1158/1078-0432]
- 161 **Liang WS**, Craig DW, Carpten J, Borad MJ, Demeure MJ, Weiss GJ, Izatt T, Sinari S, Christoforides A, Aldrich J, Kurdoglu A, Barrett M, Phillips L, Benson H, Tembe W, Braggio E, Kiefer JA, Legendre C, Posner R, Hostetter GH, Baker A, Egan JB, Han H, Lake D, Stites EC, Ramanathan RK, Fonseca R, Stewart AK, Von Hoff D. Genome-wide characterization of pancreatic adenocarcinoma patients using next generation sequencing. *PLoS One* 2012; **7**: e43192 [PMID: 23071490 DOI: 10.1371/journal.pone.0043192]
- 162 **Hermann PC**, Huber SL, Herrler T, Aicher A, Ellwart JW, Guba M, Bruns CJ, Heeschen C. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell* 2007; **1**: 313-323 [PMID: 18371365 DOI: 10.1016/j.stem.2007.06.002]
- 163 **Li C**, Heidt DG, Dalerba P, Burant CF, Zhang L, Adsay V, Wicha M, Clarke MF, Simeone DM. Identification of pancreatic cancer stem cells. *Cancer Res* 2007; **67**: 1030-1037 [PMID: 17283135 DOI: 10.1158/0008-5472.CAN-06-2030]
- 164 **Penchev VR**, Rasheed ZA, Maitra A, Matsui W. Heterogeneity and targeting of pancreatic cancer stem cells. *Clin Cancer Res* 2012; **18**: 4277-4284 [PMID: 22896694 DOI: 10.1158/1078-0432.CCR-11-3112]
- 165 **Rhim AD**, Mirek ET, Aiello NM, Maitra A, Bailey JM, McAllister F, Reichert M, Beatty GL, Rustgi AK, Vonderheide RH, Leach SD, Stanger BZ. EMT and dissemination precede pancreatic tumor formation. *Cell* 2012; **148**: 349-361 [PMID: 22265420 DOI: 10.1016/j.cell.2011.11.025]
- 166 **Jimeno A**, Feldmann G, Suárez-Gauthier A, Rasheed Z, Solomon A, Zou GM, Rubio-Viqueira B, García-García E, López-Ríos F, Matsui W, Maitra A, Hidalgo M. A direct pancreatic cancer xenograft model as a platform for cancer stem cell therapeutic development. *Mol Cancer Ther* 2009; **8**: 310-314 [PMID: 19174553 DOI: 10.1158/1535-7163.MCT-08-924]
- 167 **Rajeshkumar NV**, Rasheed ZA, García-García E, López-Ríos F, Fujiwara K, Matsui WH, Hidalgo M. A combination of DR5 agonistic monoclonal antibody with gemcitabine targets pancreatic cancer stem cells and results in long-term disease control in human pancreatic cancer model. *Mol Cancer Ther* 2010; **9**: 2582-2592 [PMID: 20660600 DOI: 10.1158/1535-7163.mCT-10-0370]
- 168 **Zhang GN**, Liang Y, Zhou LJ, Chen SP, Chen G, Zhang TP, Kang T, Zhao YP. Combination of salinomycin and gemcitabine eliminates pancreatic cancer cells. *Cancer Lett* 2011; **313**: 137-144 [PMID: 22030254 DOI: 10.1016/j.canlet.2011.05.030]
- 169 **Mahadevan D**, Von Hoff DD. Tumor-stroma interactions in pancreatic ductal adenocarcinoma. *Mol Cancer Ther* 2007; **6**: 1186-1197 [PMID: 17406031 DOI: 10.1158/1535-7163.MCT-06-0686]
- 170 **Masamune A**, Shimosegawa T. Signal transduction in pancreatic stellate cells. *J Gastroenterol* 2009; **44**: 249-260 [PMID: 19271115 DOI: 10.1007/s00535-009-0013-2]
- 171 **Erkan M**, Reiser-Erkan C, Michalski CW, Deucker S, Sauiunaite D, Streit S, Esposito I, Friess H, Kleeff J. Cancer-stellate cell interactions perpetuate the hypoxia-fibrosis cycle in pancreatic ductal adenocarcinoma. *Neoplasia* 2009; **11**: 497-508 [PMID: 19412434]
- 172 **Jacobetz MA**, Chan DS, Neesse A, Bapiro TE, Cook N, Frese KK, Feig C, Nakagawa T, Caldwell ME, Zecchini HI, Lolkema MP, Jiang P, Kultti A, Thompson CB, Maneval DC, Jodrell DJ, Frost GL, Shepard HM, Skepper JN, Tuveson DA. Hyaluronan impairs vascular function and drug delivery in a mouse model of pancreatic cancer. *Gut* 2013; **62**: 112-120 [PMID: 22466618 DOI: 10.1136/gutjnl-2012-302529]
- 173 **Beatty GL**, Chiorean EG, Fishman MP, Saboury B, Teitelbaum UR, Sun W, Huhn RD, Song W, Li D, Sharp LL, Toriggian DA, O'Dwyer PJ, Vonderheide RH. CD40 agonists alter tumor stroma and show efficacy against pancreatic carcinoma in mice and humans. *Science* 2011; **331**: 1612-1616 [PMID: 21436454 DOI: 10.1126/science.1198443]
- 174 **Schrader H**, Wiese M, Ellrichmann M, Belyaev O, Uhl W, Tannapfel A, Schmidt W, Meier J. Diagnostic value of quantitative EUS elastography for malignant pancreatic tumors: relationship with pancreatic fibrosis. *Ultraschall Med* 2012; **33**: E196-E201 [PMID: 21630184 DOI: 10.1055/s-0031-1273256]
- 175 **Von Hoff DD**, Korn R, Mousses S. Pancreatic cancer--could it be that simple? A different context of vulnerability. *Cancer Cell* 2009; **16**: 7-8 [PMID: 19573807 DOI: 10.1016/j.ccr.2009.06.011]
- 176 **Le A**, Rajeshkumar NV, Maitra A, Dang CV. Conceptual framework for cutting the pancreatic cancer fuel supply. *Clin Cancer Res* 2012; **18**: 4285-4290 [PMID: 22896695 DOI: 10.1158/1078-0432.CCR-12-0041]
- 177 **Li D**, Yeung SC, Hassan MM, Konopleva M, Abbruzzese JL. Antidiabetic therapies affect risk of pancreatic cancer. *Gastroenterology* 2009; **137**: 482-488 [PMID: 19375425 DOI: 10.1053/j.gastro.2009.04.013]
- 178 **Kisfalvi K**, Eibl G, Sinnott-Smith J, Rozengurt E. Metformin disrupts crosstalk between G protein-coupled receptor and insulin receptor signaling systems and inhibits pancreatic cancer growth. *Cancer Res* 2009; **69**: 6539-6545 [PMID: 19679549 DOI: 10.1158/0008-5472.CAN-09-0418]
- 179 **Ma WW**, Jacene H, Song D, Vilardell F, Messersmith WA, Laheru D, Wahl R, Endres C, Jimeno A, Pomper MG, Hidalgo M. [18F]fluorodeoxyglucose positron emission tomography correlates with Akt pathway activity but is not predictive of clinical outcome during mTOR inhibitor therapy. *J Clin Oncol* 2009; **27**: 2697-2704 [PMID: 19380450 DOI: 10.1200/JCO.2008.18.8383]
- 180 **Garrido-Laguna I**, Tan AC, Uson M, Angenendt M, Ma WW, Villaroel MC, Zhao M, Rajeshkumar NV, Jimeno A, Donehower R, Iacobuzio-Donahue C, Barrett M, Rudek MA, Rubio-Viqueira B, Laheru D, Hidalgo M. Integrated preclinical and clinical development of mTOR inhibitors in pancreatic cancer. *Br J Cancer* 2010; **103**: 649-655 [PMID: 20664591 DOI: 10.1038/sj.bjc.6605819]
- 181 **Guo JY**, Chen HY, Mathew R, Fan J, Strohecker AM, Karsli-Uzunbas G, Kamphorst JJ, Chen G, Lemons JM, Karantza V, Collier HA, Dipaola RS, Gelinis C, Rabinowitz JD, White E. Activated Ras requires autophagy to maintain oxidative metabolism and tumorigenesis. *Genes Dev* 2011; **25**: 460-470 [PMID: 21317241 DOI: 10.1101/gad.2016311]

**P- Reviewer:** Liu X, Muscarella P, Kurniali PC **S- Editor:** Ma YJ  
**L- Editor:** A **E- Editor:** Wang CH





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

