



Published in final edited form as:

J Antibiot (Tokyo). 2014 August ; 67(8): 571–575. doi:10.1038/ja.2014.37.

Mullinamides A and B, New Cyclopeptides Produced by the Ruth Mullins Coal Mine Fire Isolate *Streptomyces* sp. RM-27-46

Xiachang Wang^{1,2}, Khaled A. Shaaban^{1,2}, Sherif I. Elshahawi^{1,2}, Larissa V. Ponomareva^{1,2}, Manjula Sunkara³, Gregory C. Copley⁴, James C. Hower⁴, Andrew J. Morris³, Madan K. Kharel^{5,*}, and Jon S. Thorson^{1,2,*}

¹Center for Pharmaceutical Research and Innovation, College of Pharmacy, University of Kentucky, Lexington, Kentucky 40536

²Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, Kentucky 40536

³Division of Cardiovascular Medicine, University of Kentucky, Lexington, KY 40536

⁴Center for Applied Energy Research, University of Kentucky, Lexington, KY, 40511

⁵University of Maryland Eastern Shore, Princess Anne, Maryland 21853, United States

Abstract

Two new cyclopeptides, mullinamides A [cyclo-(*-L*-Gly-*-L*-Glu-*-L*-Val-*-L*-Ile-*-L*-Pro-)] and B [cyclo-(*-L*-Glu-*-L*-Met-*-L*-Pro-)] were isolated from the crude extract of terrestrial *Streptomyces* sp. RM-27-46 along with the three known cyclopeptides surugamide A [cyclo-(*-L*-Ile-*-D*-Ile-*-L*-Lys-*-L*-Ile-*-D*-Phe-*-D*-Leu-*-L*-Ile-*-D*-Ala-)], cyclo-(*-L*-Pro-*-L*-Phe-) and cyclo-(*-L*-Pro-*-L*-Leu-). The structures of the new compounds were elucidated by the cumulative analyses of NMR spectroscopy and high resolution mass spectrometry. While mullinamides A and B displayed no appreciable antimicrobial/fungal activity or cytotoxicity, this study highlights the first reported antibacterial activity of surugamide A.

Keywords

surugamide; cyclopeptide; natural product; pharmacognosy; bioprospecting; antibiotic

INTRODUCTION

Coal fires, initiated either via natural ignition or human intervention, and can persist uncontrolled for decades in underground coal mines, coal waste piles, and unmined coal beds. They occur worldwide in all coal-bearing geographies and produce substantial greenhouse gas and toxic emissions, which, in the case of underground coal fires, are

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*To whom correspondence should be addressed. jsthorson@uky.edu; mkkharel@umes.edu.

Supplementary Information including the full spectroscopic data (NMR and mass) of mullinamides A (1) and B (2) are available on the *Journal of Antibiotics* website (<http://www.nature.com/ja>).

exhausted through thermal vents and seams.¹⁻⁴ As such, these exhaust ports and adjacent environments offer highly unique soil/atmospheric conditions likely to impact upon the metabolic output of terrestrial actinomycetes. Within this context, we have focused upon two underground coal mine fire sites in eastern Kentucky (Ruth Mullins and Truman Shepard)⁵⁻⁸ and this effort has facilitated the discovery of a range of new secondary metabolites from corresponding *Streptomyces* strains including the recently reported unique tetracyclic polyketide ruthmycin.⁹⁻¹³ As an extension of this ongoing effort, herein we describe the isolation, structure elucidation and preliminary biological activity assessment of five cyclic peptides (Figure 1) from the previously unreported Ruth Mullins isolate *Streptomyces* sp. RM-27-46. Notably, two of the cyclic peptides, mullinamides A [cyclo-(*L*-Gly-*L*-Glu-*L*-Val-*L*-Ile-*L*-Pro-)] (**1**) and B [cyclo-(*L*-Glu-*L*-Met-*L*-Pro-)] (**2**), are new and a third, surugamide A, is a cyclic octapeptide recently reported as a metabolite of a marine *Streptomyces* isolated near Japan.¹⁴ While surugamides were previously found to weakly inhibit caspase in vitro, the current study is also the first to divulge the moderate antibacterial activity of surugamide A (*Staphylococcus aureus* MIC of 10 μ M).

RESULTS AND DISCUSSION

Compound **1** was obtained as white powder (Table 1). The molecular formula of **1** was established as C₂₃H₃₇N₅O₇ on the basis of HR-ESI-MS (Table 1) and NMR data, indicating eight degrees of unsaturation. The proton NMR (Table 2) of **1** in DMSO-*d*₆ displayed typical features of a peptide-derived compound, including four amide protons at δ_{H} 8.07 (t, 5.4), 8.00 (d, 6.8), 7.87 (d, 8.7) and 7.84 (d, 8.3), and five α -protons at δ_{H} 4.42, 4.19, 4.12, 4.09, 4.00 and 3.80 ppm. The ¹³C-NMR of **1** (Table 2) was also consistent with a peptidic nature with six amide/acid carbonyls (δ_{C} 181.7, 175.5, 174.6 (two overlapped carbon signals), 174.0, 169.5) and five α -carbon signals (δ_{C} 61.5, 59.4, 59.0, 58.3, 42.7). The 1D (¹H and ¹³C) and 2D (COSY, TOCSY, HSQC and HMBC) NMR data revealed that **1** was composed of five amino acid residues. Specifically, the independent spin system of the type XCHCH₂CH₂X' suggested the presence of a glutamic acid unit while the spin systems of XCHCH(CH₃)₂, XCHCH(CH₃)CH₂CH₃ and XCHCH₂CH₂CH₂X' identified from TOCSY experiments were indicative of valine, isoleucine, and proline residues. One remaining CH₂ group, which lacked correlations with other protons in the COSY and TOCSY spectra, suggested the presence of a glycine moiety. Five carbonyl signals and the proline ring accounted for 7 of the requisite 8 degrees of unsaturation as dictated by the molecular formula, suggesting a cyclic form. Key connections among the five amino acid residues were established via HMBC correlations (Figure 2): 2-NH (δ_{H} 8.07) to C-2 (δ_{C} 42.7) and C-3 (δ_{C} 175.5); 4-NH (δ_{H} 8.00) to C-3 (δ_{C} 175.5) and C-8 (δ_{C} 174.6); 9-NH (δ_{H} 7.84) to C-8 (δ_{C} 174.6) and C-13 (δ_{C} 174.0); 14-NH (δ_{H} 7.87) to C-13 (δ_{C} 174.0) and C-19 (δ_{C} 174.6); and H-23 (δ_{H} 3.43, 3.52) to C-1 (δ_{C} 169.5). Importantly, these connections established the sequence **1** as cyclo-(*L*-Gly-*L*-Glu-*L*-Val-*L*-Pro-). Subsequent Marfey's analysis¹⁵ of the total hydrolysate of **1** using 6 N HCl overnight revealed the building blocks of **1** to be proteinogenic amino acids (*L*-Glu, *L*-Ile, *L*-Pro and *L*-Val). Thus, the structure of **1** was assigned as cyclo-(*L*-Gly-*L*-Glu-*L*-Val-*L*-Ile-*L*-Pro-) and, as a new cyclopeptide, was named mullinamide A in reference to its point of origin.

Compound **2** was obtained as white powder and its molecular formula was deduced as C₁₅H₂₃N₃O₅S from its HR-ESI-MS at *m/z* 358.1434 [M + H]⁺ (Calcd. 358.1437 for C₁₅H₂₄N₃O₅S [M + H]⁺), Table 1. The ¹H NMR spectrum (Table 2) of **2** also revealed peptide features with three α-protons at δ_H 4.64, 4.24 and 4.05 and two amide protons at δ_H 8.28 and 7.80 ppm. The planar structure of **2** was similarly established by 1D and 2D NMR analysis (Figure 2) to be cyclo-(-Glu-Met-Pro-), and the absolute amino acid configurations subsequently identified by the total hydrolysis of **2** using 6 N HCl, followed by Marfey's analysis method as described for **1**. Based upon the cumulative data, compound **2** was identified as cyclo-(-L-Glu-L-Met-L-Pro-), and, as a new cyclopeptide, was named mullinamide B in reference to its point of origin.

Three additional compounds were also isolated from *Streptomyces* sp. RM-27-46 and identified as surugamide A [cyclo-(-L-Ile-D-Ile-L-Lys-L-Ile-D-Phe-D-Leu-L-Ile-D-Ala-)]¹⁴, cyclo-(-L-Pro-L-Phe-)^{16,17} and cyclo-(-L-Pro-L-Leu-)¹⁷ based upon correlation of cumulative NMR and HRMS data to that reported in the literature.

Mullinamides A (**1**) and B (**2**) displayed no antibacterial (*Staphylococcus aureus* ATCC 6538, *Salmonella enterica* ATCC 10708) or antifungal (against *Saccharomyces cerevisiae* ATCC 204508) activities at or below 100 μM and no cytotoxicity against human cancer cell lines (A549 non-small cell lung, PC3 prostate) at or below 10 μM. In contrast, surugamide A (previously only reported to inhibit bovine cathepsin B with an IC₅₀ of 21 μM)¹⁴ displayed moderate growth inhibition against *Staphylococcus aureus* (MIC of 10 μM). While the biosynthetic loci encoding surugamides were previously noted to be widely distributed among marine *Streptomyces* species collected near Japan,¹⁴ the current study suggests surugamide production may be a much broader phenomenon than previously appreciated.

MATERIALS AND METHODS

General experimental procedures

Optical rotation was recorded on a Jasco DIP-370 Digital Polarimeter. All NMR data was recorded at 500 MHz or 400 MHz for ¹H and 100 MHz for ¹³C with Varian Inova NMR spectrometers (Agilent, Santa Clara, CA). LC-MS was conducted with a Waters 2695 LC module (Waters, Milford, MA), equipped with a micromass ZQ and a Symmetry Anal C₁₈ column (4.6 × 250 mm, 5 μm; Waters, Milford, MA). HR-ESI-MS spectra were recorded on an AB SCIEX Triple TOF 5600 System (AB Sciex, Framingham, MA, USA). HPLC analyses were performed on an Agilent 1260 system equipped with a photodiode array detector (PDA) detector and a Phenomenex C₁₈ column (250 × 4.6 mm, 5 μm; Phenomenex, Torrance, CA). Semi-preparative HPLC separation was performed on a Varian Prostar 210 HPLC system equipped with a PDA detector 330 using a Supelco Discovery®Bio wide pore C₁₈ column (250 × 21.2 mm, 10 μm; flow rate, 8 ml/min; Sigma-Aldrich, St. Louis, MO). All solvents used were of ACS grade and purchased from Pharmco-AAPER (Brookfield, CT). Sephadex LH-20 (25 ~ 100 μm) was purchased from GE Healthcare (Little Chalfont, United Kingdom). C₁₈-functionalised silica gel (40 ~ 63 μm) was purchased from Material Harvest Ltd. (Cambridge, United Kingdom). Amberlite®XAD16N resin (20–60 mesh) was purchased from Sigma-Aldrich (Saint Louis, MO, United States). TLC silica gel plates (60 F₂₅₄) were purchased from EMD Chemicals Inc. (Darmstadt, Germany).

Isolation of *Streptomyces* sp. RM-27-46

The soil sample was collected from the Ruth Mullins underground coal mine fire site, Perry County, KY (coordinates: N 37° 18.725' and W 83° 10.3335'). For strain purification, 0.5 g of soil sample was suspended in 1.0 ml of sterile water and the suspension was heated at 75 °C for 10 minutes to eliminate non-sporulating bacteria. Following serial dilution (10^{-1} , 10^{-2} , 10^{-3}) of the suspension with sterile water, a 100 µl aliquot was spread on oatmeal agar (Difco™, Ref: 255210, Becton, Dickinson and Company, Sparks, MD) and on ISP4 agar (Difco™, Ref: 277210, Becton, Dickinson and Company, Sparks, MD) plates supplemented with nalidixic acid (75 µg/ml) and cycloheximide (50 µg/ml). A number of sporulating bacterial colonies were observed after a week of incubation at 28 °C and each colony was subsequently streaked on a M₂-agar plate [glucose (4.0 g), malt extract (10.0 g), yeast extract (4.0 g), CaCO₃ (1.0 g) and agar (15.0 g) were dissolved in 1 liter of demineralized water]. Individual bacterial colonies were isolated from the second generation plates. AntiBase¹⁸ comparison of HPLC–high-resolution mass spectrometry (HPLC-HRMS) profiles of the culture extracts of 46 actinomycete strains isolated from a single soil sample collected near a thermal vent associated with the Ruth Mullins underground coal mine fire indicated that one of these, *Streptomyces* sp. RM-27-46, was capable of unique metabolite production.

Identification of *Streptomyces* sp. RM-27-46

Genomic DNA was isolated from a fully grown colony using the GeneJET Genomic DNA Purification Kit (Thermo Scientific, Rockford, IL, USA). The partial 16S rRNA gene fragment was amplified using universal primers (27F, 5'-AGAGTTTGATCMTGGCTCAG-3'; 1492R, 5'-GGTTACCTTGTTACGACTT-3')¹⁹ and Phusion High Fidelity DNA polymerase (New England Biolabs, Ipswich, MA, USA) and was gel-purified using a QIAquick gel extraction kit (Qiagen, Valencia, CA, USA). The amplified fragment (1,311 bp) displayed 99% identity (BLAST search) to the 16S rRNA gene sequence of *Streptomyces exfoliatus* strain NBRC 13475. The sequence of 16S rRNA has been deposited in the NCBI nucleotide database under accession number KF732715.

Host Fermentation and Metabolite Extraction, Isolation and Purification

Streptomyces sp. RM-27-46 was cultivated on M₂-agar plates at 28 °C for 7 days. Chunks of agar with the fullygrown strain were used to inoculate five 250 ml Erlenmeyer flasks, each containing 50 ml of medium A [glucose (10.0 g), yeast extract (5.0 g), soluble starch (20.0 g), peptone (5.0 g), NaCl (4.0 g), K₂HPO₄ (0.5 g), MgSO₄·7H₂O (0.5), and CaCO₃ (2 g) were dissolved in 1 liter of demineralized water]. Individual cultures were grown at 28 °C with shaking (200 rpm) for 3 days and subsequently used as seed cultures for the scale-up fermentation. The seed cultures were used to inoculate 100 Erlenmeyer flasks (250 ml) each containing 100 ml of medium A. The fermentation (10 L) was continued for seven days at 28 °C with 200 rpm agitation. The obtained culture broth was centrifuged at 5000 rpm for 30 minutes. The biomass (mycelium) was extracted with MeOH (3 × 500 ml) and then the recovered organics were evaporated *in vacuo* at 40 °C to yield 5.1 g of crude extract. The supernatant was mixed with 3% (w/v) XAD-16 resin and stirred overnight, followed by filtration. The resin was washed with water (3 × 600 ml) and then extracted with MeOH

until the eluant was colorless. The methanol extract was subsequently evaporated to afford 15.6 g of crude extract. Both extracts (obtained from the biomass and supernatant) revealed a similar metabolite profile based upon HPLC and TLC analysis and were therefore combined (20.7 g) prior to further isolation. The crude extract was then subjected to a silica gel column chromatography (15 × 8 cm, 250 g) eluted with a gradient of CHCl₃-MeOH (100:0 ~ 0:100) to yield eight fractions, A-H. Fraction B (850 mg) was subjected to Sephadex LH-20 column followed by semi-preparative HPLC (mobile phase: 5%~35% aqueous CH₃CN over 25min, 8 ml/min) to yield 24.0 mg of cyclo-(*-L*-Pro-*-L*-Phe-) (retention time: 15.0 min) and 15.0 mg of cyclo-(*-L*-Pro-*-L*-Leu-) (retention time: 9.1 min). Fraction D (320 mg) was first subjected to a Sephadex LH-20 column (MeOH) to obtain four sub-fractions, D-1 - D-4. Sub-fraction D-2 (102 mg) was resolved by semi-preparative HPLC (mobile phase: 10%~35% aqueous CH₃CN in 25min, flow rate: 8 ml/min) to yield reginamide A (14.0 mg) (retention time: 11.4 min). Compound **2** (3 mg, retention time: 10.2 min) was isolated as white powder from the sub-fraction D-3 (52.0 mg) also via semi-preparative HPLC (mobile phase: 20~60% aqueous CH₃CN over 25min, 8 ml/min). Fraction E (200 mg) was subjected to a Sephadex LH-20 column (50 × 2 cm) using methanol to elute target compounds. The major fraction obtained from the Sephadex LH-20 column was dried and then further purified using semi-preparative HPLC (mobile phase: 10%~35% aqueous CH₃CN in 25min, flow rate: 8 ml/min) to yield compound **1** (25.0 mg, retention time: 11.6 min). Based upon HPLC and TLC, the remaining fractions (A, C, and F-H) and sub-fractions lacked additional metabolites of interest and were therefore excluded from further consideration.

Determination of Amino Acid Configuration

The absolute configuration of each amino acid residue was determined following Marfey's method.¹⁵ Specifically, compounds **1** or **2** (each 1.0 mg) were hydrolyzed in 6 N HCl (1 ml) at 110°C for 14 h. After drying under nitrogen, the corresponding hydrolysate was dissolved in 2 ml of EtOAc-H₂O (1:1). The aqueous layer was dried in vacuo, to which a solution of 1% Marfey's reagent in acetone (200 µl) was subsequently added, followed by 1 M NaHCO₃ (50 µl). The reaction was heated to 40°C for 1 h, cooled to room temperature, and acidified with 2 N HCl (25 µl). The reaction mixture was diluted with MeOH (0.5 ml) and analyzed by HPLC using the following gradient: 0–55 min, linear gradient from 10%–55% CH₃CN in 50 mM TEAP buffer (Phenomenex C₁₈ column, 250 × 4.6 mm, 5 µm; 1ml/min; 430 nm). Derivatized standards were prepared from the authentic *D*- and *L*-glutamic acid, valine, isoleucine, proline and methionine (50 µl of a 50 mM stock) following an identical procedure. The retention times for Marfey's derivatives of authentic *D*-Glu, *L*-Glu, *D*-Val, *L*-Val, *D*-Ile, *L*-Ile, *D*-Pro, *L*-Pro, *D*-Met, *L*-Met were 24.7, 22.6, 36.8, 31.8, 41.2, 36.1, 28.5, 26.3, 35.0 and 30.4 min, respectively. Those for the Marfey's derivatives of Glu, Val, Ile, Pro in the hydrolysate of **1** were 22.5 (*L*-Glu), 31.7 (*L*-Val), 36.1 (*L*-Ile) and 26.2 (*L*-Pro) min, respectively. Those of **2** were 22.6 (*L*-Glu), 30.3 (*L*-Met), 26.2 (*L*-Pro) min, respectively.

Cell Viability Assay

A resazurin-based cytotoxicity assay, also known as AlamarBlue assay, was used to assess the cytotoxicity of agents against the human lung non-small cell carcinoma cell line A549 and human prostate cancer cell lines PC3 where the degree of cytotoxicity was based upon

residual metabolic activity as assessed via reduction of resazurin (7-hydroxy-10-oxido-phenoxazin-10-ium-3-one) to its fluorescent product resorufin. A549 and PC3 cells, purchased from ATCC (Manassas, VA, USA), were grown in DMEM/F-12 Kaighn's modification and MEM/EBSS media, respectively (Thermo scientific HyClone, Logan, UT, USA), with 10% heat-inactivated fetal bovine serum (FBS), 100 U/ml penicillin, 100 µg/ml streptomycin, 2 mM L-glutamine. Cells were seeded at a density 2×10^3 cells per well onto 96-well culture plates with a clear bottom (Corning, NY, USA), incubated 24 hrs at 37 °C in a humidified atmosphere containing 5% CO₂ and were exposed to test agents for 2 days (positive controls: 1.5 mM hydrogen peroxide, 10 µg/ml actinomycin D). Resazurin (150 µM final concentration) was subsequently added to each well and the plates were shaken briefly for 10 seconds and were incubated for another 3 h at 37 °C to allow viable cells to convert resazurin into resorufin. The fluorescence intensity for resorufin was detected on a scanning microplate spectrofluorometer FLUOstar Omega (BMG LABTECH GmbH, Ortenberg, Germany) using an excitation wavelength of 560 nm and an emission wavelength of 590 nm. The assay was repeated in 3 independent experimental replications. In each replication, the emission of fluorescence of resorufin values in treated cells were normalized to, and expressed as a percent of, the mean resorufin emission values of untreated control (metabolically active cells; 100%, all cells are viable).

Antibacterial and Antifungal Activity Tests

Antibacterial assay—The protocol used for the determination of MIC was based upon minor modifications of previously published protocol.²⁰ Two bacterial strains [*Staphylococcus aureus* (ATCC 6538) and *Salmonella enterica* (ATCC 10708)] were used as model strains for the assay. Single colonies from each strain were grown in 5 ml of tryptic soy broth (BD 211825) and nutrient broth (BD 234000) media, respectively, and the cultures were allowed to grow overnight at 37 °C with shaking at 220 rpm. The overnight culture for each strain was diluted to OD₆₀₀ 0.8–1.1 and 150 µl of the diluted culture was added to each well of a sterile clear polystyrene 96-well plate (TPP, Fisher Scientific). Test agents were dissolved in DMSO, serially diluted and aliquots added to the plate-based wells (kanamycin, positive control; DMSO, vehicle control). Plates were incubated at 37 °C for 16 h with shaking (150 rpm). Samples (final concentrations 1–60 µM) and controls were tested in triplicate. The OD₆₀₀ was subsequently measured using a scanning microplate spectrofluorometer FLUOstar Omega (BMG Labtech, Cary, NC, USA). The acquired OD₆₀₀ values were normalized to the negative control wells (as 100% viability). The minimal concentration of the tested sample that caused growth inhibition was recorded as the MIC.

Antifungal assay—The yeast *Saccharomyces cerevisiae* (ATCC 204508) was used in a broth microdilution antifungal assay. The assay was performed in a 96-well plate using the CLSI (formerly NCCLS) guidelines.²¹ Individual colonies from an overnight plate were used to inoculate 5 ml YAPD (ATCC medium number 1069) and incubated overnight at 37 °C. The overnight culture was diluted to OD₆₀₀ of 0.1 using sterile YAPD and 150 µl of the diluted culture was transferred to each well of a sterile clear polystyrene 96-well plate (TPP, Fisher Scientific). Test agents were dissolved in DMSO, serially diluted and aliquots added to the plate-based wells (amphotericin B, positive control; DMSO, vehicle control). The plate was incubated at 37 °C for 16 h with shaking (200 rpm) and the OD₆₀₀ was

subsequently measured using a scanning microplate spectrofluorometer FLUOstar Omega (BMG Labtech, Cary, NC, USA). The acquired OD₆₀₀ values were normalized to the negative control wells (as 100% viability). The minimal concentration of the tested sample that caused growth inhibition was recorded as the MIC. Samples (final concentrations 1–60 μM) and controls were tested in triplicate.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgements

This work was supported in part by the University of Kentucky College of Pharmacy, the University of Kentucky Markey Cancer Center and the National Center for Advancing Translational Sciences (UL1TR000117).


References

1. Hower JC, et al. The tiptop coal-mine fire, Kentucky: preliminary investigation of the measurement of mercury and other hazardous gases from coal-fire gas vents. *Int. J. Coal. Geol.* 2009; 80:63–67.
2. O'Keefe JMK, et al. Old smokey coal fire, Floyd County, Kentucky: estimates of gaseous emission rates. *Int. J. Coal. Geol.* 2011; 87:150–156.
3. Hower JC, O'Keefe JMK, Henke KR, Bagherieh A. Time series analysis of CO emissions from an Eastern Kentucky coal fire. *Int. J. Coal. Geol.* 2011; 88:227–231.
4. Hower JC, O'Keefe JMK, Copley GC, Hower JM. The further adventures of Tin Man: vertical temperature gradients at the Lotts Creek coal mine fire, Perry County, Kentucky. *Int. J. Coal Geol.* 2012; 101:16–20.
5. O'Keefe JMK, et al. CO₂, CO, and Hg emissions from the Truman Shepherd and Ruth Mullins coal fires, Eastern Kentucky, USA. *Sci. Total. Environ.* 2010; 408:1628–1633. [PubMed: 20071005]
6. Silva LFO, et al. Nanominerals and ultrafine particles in sublimates from the Ruth Mullins coal fire, Perry County, Eastern Kentucky, USA. *Int. J. Coal Geol.* 2011; 85:237–245.
7. Silva LFO, et al. Geochemistry of carbon nanotube assemblages in coal fire soot, Ruth Mullins fire, Perry County, Kentucky. *Int. J. Coal Geol.* 2012; 94:206–213.
8. Hower JC, et al. Gaseous emissions and sublimates from the Truman Shepherd coal fire, Floyd County, Kentucky: a re-investigation following attempted mitigation of the fire. *Int. J. Coal Geol.* 2013; 116:63–74.
9. Wang X, et al. Ruthmycin, a new tetracyclic polyketide from *Streptomyces* sp. RM-4-15. *Org. Lett.* 2014; 16:456–459. [PubMed: 24341358]
10. Wang X, et al. Frenolicins C–G, pyranonaphthoquinones from *Streptomyces* sp. RM-4-15. *J. Nat. Prod.* 2013; 76:1441–1447. [PubMed: 23944931]
11. Shaaban KA, et al. Herbimycins D–F, ansamycin analogues from *Streptomyces* sp. RM-7-15. *J. Nat. Prod.* 2013; 76:1619–1626. [PubMed: 23947794]
12. Shaaban KA, et al. The native production of the sesquiterpene isopterocarpolone by *Streptomyces* sp. RM-14-6. *Nat. Prod. Res.* 2013
13. Shaaban KA, et al. Venturicidin C, a new 20-membered macrolide produced by *Streptomyces* sp. TS-2-2. *J. Antibio.* 2013
14. Takada K, et al. Surugamides A–E, cyclic octapeptides with four d-amino acid residues, from a marine *Streptomyces* sp.: LC–MS-aided inspection of partial hydrolysates for the distinction of d- and l-amino acid residues in the sequence. *J. Org. Chem.* 2013; 78:6746–6750. [PubMed: 23745669]
15. Marfey P. Determination of d-amino acids. II. Use of a bifunctional reagent, 1,5-difluoro-2,4-dinitrobenzene. *Carlsberg. Res. Commun.* 1984; 49:591–596.


16. Jayatilake GS, Thornton MP, Leonard AC, Grimwade JE, Baker BJ. Metabolites from an antarctic sponge-associated bacterium, *Pseudomonas aeruginosa*. *J. Nat. Prod.* 1996; 59:293–296. [PubMed: 8882433]
17. Gong J, et al. Cyclic dipeptides in actinomycete *Brevibacterium* sp. associated with sea cucumber *Apostichopus japonicus* Selenka: isolation and identification. *Acad. J. Sec. Mil. Med. Univ.* 2012; 33:1284–1287.
18. Laatsch, H. *AntiBase*. Weinheim, Germany: Wiley-VCH; 2012.
19. Lane, DJ. *Nucleic acid techniques in bacterial systematics*. Stackebrandt, E.; Goodfellow, M., editors. New York: John Wiley and Sons Ltd; 1991. p. 115-147.
20. Wiegand I, Hilpert K, Hancock REW. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat. Protoc.* 2008; 3:163–175. [PubMed: 18274517]
21. NCCLS. Reference method for broth dilution antifungal susceptibility testing of yeasts; Approved standard M27-A2. 15 Vol. 22. National Committee on Clinical Laboratory Standards; 2002.




mullinamide A (1)




surugamide A



mullinamide B (2)



cyclo-(-L-Pro-L-Phe-)



cyclo-(-L-Pro-L-Leu-)

Figure 1.
Structures of compounds isolated from *Streptomyces* sp. RM-27-46.




Figure 2. $^1\text{H}, ^1\text{H}$ -COSY (—) and selected HMBC (→) correlations of mullinamides A (1) and B (2).

Table 1Physico-chemical properties of mullinamides A (**1**) and B (**2**)

	mullinamide A (1)	mullinamide B (2)
Molecular Formula	C ₂₃ H ₃₇ N ₅ O ₇	C ₁₅ H ₂₃ N ₃ O ₅ S
Appearance	White powder	White powder
UV	End absorption	End absorption
[α] _D ²⁵	−36.9° (c 2.5, MeOH)	−10.6° (c 0.3, MeOH)
(+)-ESI-MS: <i>m/z</i>	496 [M + H] ⁺ , 991 [2 M + H] ⁺	358 [M + H] ⁺ , 715 [2 M + H] ⁺
(−)-ESI-MS: <i>m/z</i>	494 [M − H] [−]	-
(+)-HR-ESI-MS: <i>m/z</i> Calcd.	496.2767 [M + H] ⁺ , 518.2590 [M + Na] ⁺ , 496.2771 for C ₂₃ H ₃₈ N ₅ O ₇ [M + H] ⁺ , 518.2591 for C ₂₃ H ₃₇ N ₅ O ₇ Na [M + Na] ⁺	358.1434 [M + H] ⁺ , 358.1437 for C ₁₅ H ₂₄ N ₃ O ₅ S [M + H] ⁺
(−)-HR-ESI-MS: <i>m/z</i> Calcd	494.2627 [M − H] [−] 494.2615 for C ₂₃ H ₃₆ N ₅ O ₇ [M − H] [−]	356.1287 [M − H] [−] 356.1280 for C ₁₅ H ₂₂ N ₃ O ₅ S [M − H] [−]

Table 2

¹H and ¹³C NMR data for mullinamides A (1) and B (2)

		Mullinamide A (1)				Mullinamide B (2)					
Position	δ_{H}^a (J in Hz)	δ_{H}^b (J in Hz)	δ_{C}^c , type	Position	δ_{H}^a (J in Hz)	δ_{H}^b (J in Hz)	δ_{C}^c , type	Position	δ_{H}^a (J in Hz)	δ_{H}^b (J in Hz)	δ_{C}^c , type
Gly	1		169.5, qC	Glu	1		175.5, qC				175.3, qC
	2a	3.97, d (16.8)	3.80, dd (5.8, 17.1)		2	4.22, m	42.7, CH ₂		4.05, m	4.05, m	57.8, CH
	2b	4.16, d (16.8)	4.00, dd (4.9, 17.1)		3	2.29, 2.45, m			2.16, m	2.16, m	26.7, CH ₂
	NH		8.07, t (5.4)		4	2.06, 2.38, m			1.84, m	1.84, m	30.9, CH ₂
Glu	3		175.5, qC		5		175.5, qC				181.7, qC
	4	4.22, m	4.12, m		NH		58.3, CH		7.80, d (4.6)		
	5	2.11, 2.48, m	2.42, m	Met	6		26.8, CH ₂		4.81, m	4.64, m	175.1, qC
	6	2.32, 2.42, m	2.04, 2.12, m		7		30.6, CH ₂		1.80, m	1.80, m	51.7, CH
	7				8		181.7, qC		1.96, 2.08, m	1.80, m	31.9, CH ₂
Val	8		8.00, d (6.8)		9		174.6, qC		2.63, m	2.52, m	30.6, CH ₂
	9	4.32, m	4.09, m		NH		59.0, CH		2.11, s	2.03, s	15.4, CH ₃
	10	2.18, m	2.01, m	Pro	11		31.9, CH		8.28, d (6.7)		172.4, qC
	11	0.96 ^d	0.94 ^d		12		18.4, CH ₃		4.45, m	4.24, m	60.5, CH
	12	0.96 ^d	0.94 ^d		13		19.7, CH ₃		2.02, 2.29, m	1.81, 2.15, m	30.3, CH ₂
	NH		7.84, d (8.3)		14		174.0, qC		2.06, m	1.90, m	26.0, CH ₂
Ile	13				15		59.4, CH		3.75, 3.87, m	3.59, 3.68	48.7, CH ₂
	14	4.25, m	4.19, m				38.0, CH				
	15	1.86, m	1.70, m				25.8, CH ₂				
	16	1.20, 1.60, m	1.06, 1.45, m				11.4, CH ₃				
	17	0.90, t (7.6, 7.2)	0.80, t (7.4)				16.0, CH ₃				
	18	0.96 ^d	0.93 ^d				174.6, qC				
	NH		7.87, d (8.7)				61.5, CH				
Pro	19						174.6, qC				
	20	4.49, m	4.42, m								

Mullinamide A (1)			Mullinamide B (2)			
Position	δ_{H}^a (J in Hz)	δ_{H}^b (J in Hz)	Position	δ_{H}^a (J in Hz)	δ_{H}^b (J in Hz)	δ_{C}^c , type
21	1.94, 2.19, m	1.75, 1.96, m				30.7, CH ₂
22	2.00, m	1.86, m				26.1, CH ₂
23	3.61, m	3.43, 3.52, m				47.9, CH ₂

^a measured in CD₃OD, 400MHz.

^b measured in DMSO-*d*₆, 500MHz.

^c measured in CD₃OD, 100MHz.

^d overlapped.