



Published in final edited form as:

Gerontology. 2014 ; 60(5): 381–385. doi:10.1159/000358875.

Tau Based Therapeutic Approaches for Alzheimer's Disease

Allal Boutajangout^{1,3,4,#} and Thomas Wisniewski^{1,2,3}

¹Department of Neurology, New York University School of Medicine, 450 East 29th Street, New York. NY. 10016

²Department of Pathology, New York University School of Medicine, 450 East 29th Street, New York. NY. 10016

³Department of Psychiatry, New York University School of Medicine, 450 East 29th Street, New York. NY. 10016

⁴Department of Physiology & Neuroscience, New York University School of Medicine, 450 East 29th Street, New York. NY. 10016

Abstract

The accumulation of aggregated, hyperphosphorylated tau as neurofibrillary tangles (NFTs) and neuropil threads (NT) are cardinal features of Alzheimer's disease (AD). The other lesions found in AD include amyloid plaques and congophilic amyloid angiopathy, both associated with the extracellular accumulation of the amyloid β (A β) peptide. AD is the most common cause of dementia globally. Currently there are no effective means to treat AD or even to slow it down. The dominant theory for the causation of AD is the amyloid cascade hypothesis, which suggests that the aggregation of A β as oligomers and amyloid plaques is central to the pathogenesis of AD. Numerous therapies have been developed directed to A β relate pathology, in particular various immunotherapeutic approaches. So far all of these have failed in clinical trials. Recently there has been more focus on therapy directed to tau related pathology, which correlates better with the cognitive status of patients, compared to the amyloid burden. Immunotherapeutic targeting of tau pathology has showed great potential to treat tau pathologies in mouse models of AD. A number of studies have shown the efficacy of both passive and active immunization. This review summarizes recent advances on therapy targeting pathological tau protein, in particular focusing on immunotherapeutic approaches which are showing great promise.

Keywords

tau protein; Alzheimer Disease (AD); immunotherapy; active immunization; passive immunization; aggregation; GSK3 β ; CDK5

Corresponding Author: Thomas Wisniewski, M.D., New York University School of Medicine, Alexandria ERSP, Rm 802, 450 East 29th St., New York, NY 10016. Tel: (212) 263-7993, Fax: (212) 263-7528, thomas.wisniewski@nyumc.org.

[#]King Abdulaziz University, School of Medicine. Jeddah. Saudi Arabia

Introduction

Alzheimer disease (AD) is a progressive, neurodegenerative disorder that is characterized by extracellular amyloid-beta ($A\beta$) deposition, in the form of amyloid plaques and congophilic amyloid angiopathy, as well as intracellular neurofibrillary tangles (NFTs), containing pathological tau aggregates [1]. The neuropathological definition of AD requires the presence of both neuritic amyloid plaques and NFTs [1]. The most toxic species of $A\beta$ and aggregated tau are thought to be oligomeric, with both of these pathologies spreading via extracellular soluble oligomers, which under some conditions have been shown to use a “prion-like” mechanism [2]. Alzheimer’s disease (AD) is the most common cause of dementia globally, affecting approximately 36 million people currently and ~115 million by 2050 [3]. The associated costs are enormous, being estimated in the USA alone to be ~\$200 billion in 2013. Presently available treatments have minimal, or no, effect on the course of disease. Numerous therapies are being developed directed to $A\beta$ related pathology. Among the most advanced approaches for $A\beta$ pathology are various immunotherapeutic approaches; however, two recent phase III clinical trials of passive immunization directed to $A\beta$ failed to show any significant clinical benefit [4]. Partially as a result of these clinical failures with anti- $A\beta$ therapies, there has been more focus on approaches directed to tau related pathology.

Pathogenesis of Alzheimer’s Disease and Anti- $A\beta$ Directed Immunotherapy

The dominant theory for the causation of AD has been the amyloid cascade hypothesis [5]. This theory suggests that the accumulation of $A\beta$ peptides, particularly in a highly toxic oligomeric form, is the primary pathogenic driver that downstream leads to tau hyperphosphorylation, NFT formation and ultimately to synaptic and neuronal loss. Missense mutations in amyloid precursor protein (APP) or in the presenilin genes PRES 1 and 2 cause early onset, familial forms of AD (FAD) affecting <1% of AD patients [6]. The most common form of AD is sporadic and late-onset. Extensive evidence supports the amyloid cascade hypothesis in FAD patients and in models of FAD [6]. However, evidence proving that $A\beta$ is central to the pathogenesis of the common late-onset sporadic form of AD (LOAD) is more limited. Recent genome-wide association studies in LOAD have implicated a number of different genes involved in innate immunity, cholesterol metabolism and endocytosis, suggesting greater etiological heterogeneity [6]. Potentially conflicting evidence to the amyloid cascade hypothesis has also come from clinical and autopsy data from the initial human active vaccination trial directed to reducing amyloid plaque pathology [7]. Post-mortem analysis was available from nine subjects, which showed a considerable degree of plaque removal and reduced $A\beta$ load compared to comparable non-immunized controls [8]. Despite this, there were no differences between placebo and active immunization groups in terms of long-term survival outcome, time to severe dementia and in outcome measures such as ADAS-Cog, MMSE or DAD. Several passive immunization trials are underway directed to $A\beta$ pathology; however, the two most advanced Phase III trials of both Bapineuzumab and Solanezumab were recently reported and both failed to show overall clinical improvement or disease modifying outcomes [4;9]. The lack of clinical efficacy in these trials may have been related to the immunization having begun too late in the disease process, at a point when extensive tau pathology already exists and can progress

independent of the initial trigger of A β related pathology [4]. Alternatively, one can use this data to suggest that the amyloid cascade hypothesis is an oversimplification. A number of investigators have suggested alternative theories, whereby accumulation of aggregated, toxic forms of A β and tau are dual pathways both downstream from a common upstream pathogenic deficit [4]. In either of these scenarios it is essential for therapy to specifically address tau related pathology to be highly effective in clinically symptomatic AD. This is particularly important as numerous studies have shown that the degree of tau related pathology correlates better with the degree of dementia compared to amyloid plaque burden, making tau an attractive target in AD [1].

The Tau Protein and Tau Directed Therapy

Tau is microtubule associated protein that is primarily expressed in the cell bodies and axons of neuronal cells of the central nervous system [10]. In the CNS, human tau is expressed in 6 isoforms arising from alternative mRNA splicing from a single gene on chromosome 17q21, containing 16 Exons [10]. The size range of the six isoforms is between 352 and 441 amino acids, which differ by the absence or presence of 29 (Exon 2) or 58 (Exon 2 + Exon 3) amino acids inserts in the amino-terminal. The carboxy-terminal half of tau contains three or four semi-homologous repeats of 31 or 32 amino acids, encoded by Exon 10. The repeats (3R, 4R) correspond to the microtubule binding region of protein tau. Tau has several phosphorylation sites, with phosphorylated tau binding microtubules with lesser affinity and/or having a greater propensity to aggregate into paired helical filaments, producing NFTs.

Given the importance of tau phosphorylation in AD a number of studies have tried inhibition of kinases involved in the process or the activation of protein phosphatase to dephosphorylate tau (see Table 1) [11;12]. Kinases involved in tau phosphorylation include glycogen synthase kinase 3 (GSK-3 β), cdk5, p38, JNK, CK and DYRK1A [11]. Lithium and valproate have been shown to inhibit GSK and when given to Tg models reduce tau related pathology [13]; however, this effect was not corroborated in small clinical trials for both compounds [14;15]. Several small molecule inhibitors of GSK-3 β such as SB 216763, CHIR-98014 and SRN-003-556 are in preclinical development [12]. SB 216763 was shown to reduce the amount of tau phosphorylation; however, adverse effects were detected in control animals [11]. Tideglusib is another inhibitor of GSK-3 β which was in phase IIb trials for mild to moderate AD that are completed but the results have not been reported (clinicaltrials.gov). In a small pilot study Tideglusib produced positive trends in MMSE, ADAS-cog, GDS, and GCA cognitive scales without statistical significance in the small sample of 30 patients (see Table 2) [16].

Another tau pathology directed approach is inhibition of tau aggregation [17]. The most advanced such agents are methylene blue and its derivatives [17;18]. Methylene blue has a range of effects in addition to tau aggregation inhibition [12]. The first methylene blue derivative was Rember (TauRx Therapeutics Ltd) which showed stabilization of progression of AD over 50 weeks [12]. A next generation derivative, LMTX, is currently in a phase III clinical trial.

A further strategy to reduce NFT pathology is to increase tau degradation. Tau degradation has been shown to be inhibited by Hsp 90, a chaperone protein that is involved in folding of denatured proteins. Curcumin has a wide range of activities, among which is to inhibit Hsp 90. A trial in a Tau Tg model showed reductions in pathology and cognitive benefits [19;20]. A number of Hsp 90 inhibitors are known; however, the targeting of a chaperone protein has the potential for adverse effects [12].

Immunotherapeutic approaches have been very successful at reducing amyloid related pathology in AD models; although this approach has yet to be shown as successful in clinical trials. In part driven by the success of immune therapy in amyloid pathology reduction, our group tested the first active immunization against tau pathology in a P301L tau transgenic (Tg) mouse model in 2007 [21]. This model develops neurofibrillary tangles in several brain regions and the spinal cord. The immunization was performed with a tau fragment peptide (amino acids 379–408, with phosphorylated Ser396 and Ser404), using alum as an adjuvant. Two groups were immunized from 2 to 5 months and from 2 to 8 months. Immunohistochemical analysis using PHF-1 and MC1 antibodies showed a significant reduction in tau related pathology, compared to controls. In addition, an amelioration in the vaccinated groups was seen on a number of sensorimotor tasks. Antibodies generated by this vaccination were found to cross the BBB, bind to phosphorylated tau and reduce pathology without significant adverse effect [21]. Subsequently, we tested the same immunogen (Tau 379–408[P-Ser396,404]) in another active immunization study using a novel mouse tau Tg model (htau/PS1), developed by one of us (AB) [22;23]. The animals were immunized from 3–4 months of age for 4 months. A significant reduction of tau pathology was observed in the brain of vaccinated mice compared to controls, which correlated with a cognitive rescue, tested using three separate tasks [23]. A further active immunization study used a mixture of three peptides containing the phosphorylation epitopes: Tau195–213 [P202/205], Tau207–220[P212/214] and Tau224–238[P231] [24]. This immunization was performed in double-mutant tau [K257T/P301S] Tg model that develops NFT like inclusions at 6 months of age. A significant decrease of tau pathology was observed in the hippocampus, the cortex and the brain stem at 8 months post immunization which was accompanied by an increase in the microglial burden. Recently, another tau immunogen was tested containing phospho-serine 422 in THY Tau22 Tg mice, which develop late stage hippocampal pathology and cognitive defects. The mice that were immunized for 18 weeks showed a reduction in tau pathology which correlated with improved performance on spatial memory [25]. Passive immunization with monoclonal antibodies (mAbs) to phosphorylated tau has also shown benefits in Tg models. The first study, conducted by our group, used the PHF-1 antibody (recognizes phosphorylated serine 396–404 tau) in P301L tau Tg mice by intraperitoneal injections and showed a reduction in tau pathology both biochemically and immunohistochemically, in association with behavioral benefits [26]. These results were confirmed by another group using both the PHF1 and the MC1 antibodies (MC1 recognizes an abnormal tau conformation) in two tau Tg models (P301L and P301S tg mice) [27]. A reduction of tau pathology was observed in both models as well as a reduction of neurofilament positive axonal spheroids in the spinal cord with an improvement in locomotor activity [27]. A further study compared DA31 (a pan-tau antibody), MC1 and PHF1 in P301L Tg tau mice

[28]. MC1 injected mice from 7 to 10 months showed a reduction of tau related pathology both immunohistochemically and biochemically [28]. Anti-tau oligomer antibodies which block tau aggregation have also been shown to be effective in P301S Tg mice; however, this study used cranial intra-ventricular injections, limiting its clinical translatability [29]. An additional recent novel approach is the use of active immunization that produces an immune response to both pathological A β and tau conformers concurrently; this has been shown to be effective in a number of different AD Tg models [30]. These various preclinical tau pathology directed immunotherapy studies have led to the first Phase I vaccine trial, AADvac1, which uses a pathology related tau peptide conjugated to KLH for active immunization; this trial is on-going (www.clinicaltrials.gov). At first examination it is difficult to understand how antibodies to a protein which is accumulating intra-cellularly can have beneficial effects. Recent studies have demonstrated that anti-tau antibodies can cross the BBB, be taken up by neurons via low-affinity Fc receptors and bind to pathological tau within the endosomal/lysosomal system [31]. In addition, evidence has shown that injection of fibrillar tau brain extract into the brains of transgenic wild-type expressing mice can induce the formation of human tau into filaments, as well as the spread of pathology from the site of injection into neighboring brain regions in a prion like manner [32]. Therefore, if the spread of certain pathological forms of tau, that lead to PHF pathology in AD occurs via such mechanism, antibodies to disease associated forms of tau would not necessarily need to enter cells in order to be effective.

Conclusions

There is a critical need to develop better treatments for AD. It is clear from studies in AD animal models that tau directed immunotherapy and related approaches can reduce pathology and produce cognitive benefits. The disappointing results with A β directed therapies and the knowledge that tau related pathology correlates more directly with the cognitive status of patients, makes this area of research particularly important and exciting currently. The importance of A β and tau oligomers as the most toxic species and the growing realization that both pathologies can spread in a prion like mechanism, make these the most critical targets for novel therapies. Many questions remain, in particular related to the uncertainties regarding the etiology or etiologies of LOAD as well as to what are the best models for testing effective AD therapies. We hypothesize that the best hope for effective therapies for symptomatic AD will be approaches that reduced pathological conformers of tau alone or in combination with A β oligomers.

Acknowledgement

This manuscript was supported by NIH grant NS073502, AG20245 and AG08051, as well as the Alzheimer's Association, and the Saudi Cultural Mission (USA).

Reference List

1. Nelson PT, Alafuzoff I, Bigio EH, Bouras C, Braak H, Cairns N, Davies P, Tredici KD, Duyckaerts C, Frosch MP, Hof PR, Hulette C, Hyman BT, Iwatsubo T, Jellinger KA, Jicha GA, Kovari E, Kukull WA, Leverenz JB, Love S, Mackenzie IR, Mann DM, Masliah E, McKee A, Montine TJ, Morris JC, Schneider JA, Sonnen JA, Thal DR, Trojanowski JQ, Troncoso JC, Wisniewski T,

- Woltjer RL, Beach TG. Correlation of Alzheimer's disease neuropathologic changes with cognitive status: a review of the literature. *J Neuropath Exp Neurol.* 2012; 71:362–381. [PubMed: 22487856]
2. Ashe KH, Aguzzi A. Prions, prionoids and pathogenic proteins in Alzheimer disease. *Prion.* 2013; 7:55–59. [PubMed: 23208281]
 3. Gaugler, J.; James, B.; Johnson, T.; Scholz, K.; Weuve, J. Alzheimer's disease facts and figures 2013. Alzheimer's Disease Association; 2013.
 4. Wisniewski T, Goni F. Immunotherapy for Alzheimer's disease. *Biochem Pharmacol.* In press.
 5. Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science.* 2002; 297:353–356. [PubMed: 12130773]
 6. Bettens K, Sleegers K, Van BC. Genetic insights in Alzheimer's disease. *Lancet Neurol.* 2013; 12:92–104. [PubMed: 23237904]
 7. Bayer AJ, Bullock R, Jones RW, Wilkinson D, Paterson KR, Jenkins L, Millais SB, Donoghue S. Evaluation of the safety and immunogenicity of synthetic A β 42 (AN1792) in patients with AD. *Neurol.* 2005; 64:94–101.
 8. Holmes C, Boche D, Wilkinson D, Yadegarfar G, Hopkins V, Bayer A, Jones RW, Bullock R, Love S, Neal JW, Zotova E, Nicoll JAR. Long term effects of A β 42 immunization in Alzheimer's disease: immune response, plaque removal and clinical function. *Lancet.* 2008; 372:216–223. [PubMed: 18640458]
 9. Farlow MR, Brosch JR. Immunotherapy for Alzheimer's disease. *Neurol Clin.* 2013; 31:869–878. [PubMed: 23896510]
 10. Yoshiyama Y, Lee VM, Trojanowski JQ. Therapeutic strategies for tau mediated neurodegeneration. *J Neurol Neurosurg Psychiatry.* 2013; 84:784–795. [PubMed: 23085937]
 11. Tell V, Hilgeroth A. Recent developments of protein kinase inhibitors as potential AD therapeutics. *Front Cell Neurosci.* 2013; 7:189. [PubMed: 24312003]
 12. Anand R, Gill KD, Mahdi AA. Therapeutics of Alzheimer's disease: Past, present and future. *Neuropharmacology.* 2014; 76 Pt A:27–50. [PubMed: 23891641]
 13. Engel T, Goni-Oliver P, Lucas JJ, Avila J, Hernandez F. Chronic lithium administration to FTDP-17 tau and GSK-3beta overexpressing mice prevents tau hyperphosphorylation and neurofibrillary tangle formation, but pre-formed neurofibrillary tangles do not revert. *J Neurochem.* 2006; 99:1445–1455. [PubMed: 17059563]
 14. Hampel H, Ewers M, Burger K, Annas P, Mortberg A, Bogstedt A, Frolich L, Schroder J, Schonknecht P, Riepe MW, Kraft I, Gasser T, Leyhe T, Moller HJ, Kurz A, Basun H. Lithium trial in Alzheimer's disease: a randomized, single-blind, placebo-controlled, multicenter 10-week study. *J Clin Psychiatry.* 2009; 70:922–931. [PubMed: 19573486]
 15. Tariot PN, Aisen PS. Can lithium or valproate untie tangles in Alzheimer's disease? *J Clin Psychiatry.* 2009; 70:919–921. [PubMed: 19573485]
 16. Del ST, Steinwachs KC, Gertz HJ, Andres MV, Gomez-Carrillo B, Medina M, Vericat JA, Redondo P, Fleet D, Leon T. Treatment of Alzheimer's disease with the GSK-3 inhibitor tideglusib: a pilot study. *J Alzheimers Dis.* 2013; 33:205–215. [PubMed: 22936007]
 17. Bulic B, Pickhardt M, Mandelkow E. Progress and developments in tau aggregation inhibitors for Alzheimer disease. *J Med Chem.* 2013; 56:4135–4155. [PubMed: 23484434]
 18. Wischik CM, Edwards PC, Lai RY, Roth M, Harrington CR. Selective inhibition of Alzheimer disease-like tau aggregation by phenothiazines. *Proc Natl Acad Sci U S A.* 1996; 93:11213–11218. [PubMed: 8855335]
 19. Giommarelli C, Zuco V, Favini E, Pisano C, Dal PF, De TN, Zunino F. The enhancement of antiproliferative and proapoptotic activity of HDAC inhibitors by curcumin is mediated by Hsp90 inhibition. *Cell Mol Life Sci.* 2010; 67:995–1004. [PubMed: 20039095]
 20. Ma QL, Zuo X, Yang F, Ubeda OJ, Gant DJ, Alaverdyan M, Teng E, Hu S, Chen PP, Maiti P, Teter B, Cole GM, Frautschy SA. Curcumin suppresses soluble tau dimers and corrects molecular chaperone, synaptic, and behavioral deficits in aged human tau transgenic mice. *J Biol Chem.* 2013; 288:4056–4065. [PubMed: 23264626]
 21. Asuni AA, Boutajangout A, Quartermain D, Sigurdsson EM. Immunotherapy targeting pathological tau conformers in a tangle mouse model reduces brain pathology with associated functional improvements. *J Neurosci.* 2007; 27:9115–9129. [PubMed: 17715348]

22. Boutajangout A, Frangione B, Wisniewski T, Brion JP, Sigurdsson EM. Influence of presenilin mutation on tau pathology in novel Alzheimer's disease mouse model. *Alz Dementia*. 2009; 5:P445–P446.
23. Boutajangout A, Quartermain D, Sigurdsson EM. Immunotherapy targeting pathological tau prevents cognitive decline in a new tangle mouse model. *J Neurosci*. 2010; 30:16559–16566. [PubMed: 21147995]
24. Boimel M, Grigoriadis N, Lourbopoulos A, Haber E, Abramsky O, Rosenmann H. Efficacy and safety of immunization with phosphorylated tau against neurofibrillary tangles in mice. *Exp Neurol*. 2010; 224:472–485. [PubMed: 20546729]
25. Troquier L, Caillierez R, Burnouf S, Fernandez-Gomez FJ, Grosjean ME, Zommer N, Sergeant N, Schraen-Maschke S, Blum D, Buee L. Targeting phospho-Ser422 by active Tau Immunotherapy in the THY Tau22 mouse model: a suitable therapeutic approach. *Curr Alzheimer Res*. 2012; 9:397–405. [PubMed: 22272619]
26. Boutajangout A, Ingadottir J, Davies P, Sigurdsson EM. Passive immunization targeting pathological phospho-tau protein in a mouse model reduces functional decline and clears tau aggregates from the brain. *J Neurochem*. 2011; 118:658–667. [PubMed: 21644996]
27. Chai X, Wu S, Murray TK, Kinley R, Cella CV, Sims H, Buckner N, Hanmer J, Davies P, O'Neill MJ, Hutton ML, Citron M. Passive immunization with anti-Tau antibodies in two transgenic models: reduction of Tau pathology and delay of disease progression. *J Biol Chem*. 2011; 286:34457–34467. [PubMed: 21841002]
28. d'Abramo C, Acker CM, Jimenez HT, Davies P. Tau passive immunotherapy in mutant P301L mice: antibody affinity versus specificity. *PLoS ONE*. 2013; 8:e62402. [PubMed: 23638068]
29. Yanamandra K, Kfoury N, Jiang H, Mahan TE, Ma S, Maloney SE, Wozniak DF, Diamond MI, Holtzman DM. Anti-Tau Antibodies that Block Tau Aggregate Seeding In Vitro Markedly Decrease Pathology and Improve Cognition In Vivo. *Neuron*. 2013; 80:402–414. [PubMed: 24075978]
30. Goni F, Herline K, Peyser D, Wong K, Ji Y, Sun Y, Mehta PD, Wisniewski T. Immunomodulation targeting both A β and tau pathological conformers ameliorates Alzheimer's Disease pathology in TgSwDI and 3 \times Tg mouse models. *Journal of Neuroinflammation*. 2013; 10:150. [PubMed: 24330773]
31. Congdon EE, Gu J, Sait HB, Sigurdsson EM. Antibody Uptake into Neurons Occurs Primarily via Clathrin Dependent Fc γ Receptor Endocytosis, and is a Prerequisite for Acute Tau Clearance. *J Biol Chem*. 2013; 288:35452–35465. [PubMed: 24163366]
32. Clavaguera F, Bolmont T, Crowther RA, Abramowski D, Frank S, Probst A, Fraser G, Stalder AK, Beibel M, Staufenbiel M, Jucker M, Goedert M, Tolnay M. Transmission and spreading of tauopathy in transgenic mouse brain. *Nat Cell Biol*. 2009; 11:909–913. [PubMed: 19503072]

Table 1

Preclinical Approaches

Approach	Mechanism of action	references
Glycogen Synthase Kinase3 Inhibitors		
SB216763, CHIR98014	Reduces tau phosphorylation	[12]
SRN-003-556	Reduce soluble phosphorylated and aggregated tau	[12]
Lithium	Reduces tau pathology	[14]
Valproate	Reduces tau pathology	[15]
Other inhibitors		
HSP90	Inhibition of tau degradation	[12]
Active immunization		
Tau 379-408[P-Ser396-404]	Reduces tau pathology and improves cognitive tasks	[21]
Tau195-231[P202/205] Tau207-220[P212/214] Tau224-238[P231]	Significant decrease of tau pathology with increase of the microglia burden	[24]
Y10A peptide [Ser422]	Reduces tau pathology and	[25]
pBri	Reduces tau related pathology	[30]
Passive immunization		
PHF-1 Antibody	Decreases tau pathology and functional impairments	[26]
PHF-1 and MC1	Decreases tau pathology with improvement of locomotor activity	[27]
DA31,MC1,PHF-1	Reduction of tau pathology	[28]
Anti-tau oligomers	Reduces hyperphosphorylated tau and blocks tau aggregation	[29]

Table II

Clinical Trials

Approach	Mechanism of action	references
Tideglusib	Inhibitor of GSK 3 β (in Phase IIb) Completed but results not yet reported	[16]
TauRx	Inhibits tau aggregation (in Phase II)	[12]
LMTX	Reduces the level of aggregated and misfolded tau (in Phase III)	[12]
AADvac1	Peptide targeting pathological tau (in Phase I)	Clinicaltrials.gov

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript