

Nicotine Pharmacokinetics in Rats Is Altered as a Function of Age, Impacting the Interpretation of Animal Model Data[§]

Evelyn L. Craig, Bin Zhao, Jason Z. Cui, Maria Novalen, Sharon Miksys, and Rachel F. Tyndale

Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Departments of Pharmacology & Toxicology and Psychiatry, University of Toronto, Toronto, Ontario, Canada (E.L.C., B.Z., M.N., S.M., R.F.T.) and Child & Family Research Institute, Department of Anesthesiology, Pharmacology, and Therapeutics, University of British Columbia, Vancouver, British Columbia, Canada (J.Z.C.)

Received April 22, 2014; accepted June 30, 2014

ABSTRACT

Several behavioral studies report that adolescent rats display a preference for nicotine compared with adults. However, age-related pharmacokinetic differences may confound the interpretation of these findings. Thus, differences in pharmacokinetic analyses of nicotine were investigated. Nicotine was administered via acute s.c. (1.0 mg base/kg) or i.v. (0.2 mg base/kg) injection to early adolescent (EA; postnatal day 25) and adult (AD; postnatal day 71) male Wistar rats. Nicotine and its primary metabolite, cotinine, and additional metabolites nornicotine, nicotine-1'-N-oxide, *trans*-3'-hydroxycotinine, and nornicotine were sampled from 10 minutes to 8 hours (plasma) and 2 to 8 hours (brain) post nicotine and analyzed by liquid chromatography–tandem mass spectrometry. Following s.c. nicotine, the EA cohort had *lower* levels of plasma nicotine, cotinine, and nicotine-1'-N-oxide at multiple time points, resulting

in a lower area under the plasma concentration–time curve (AUC) for nicotine ($P < 0.001$), cotinine ($P < 0.01$), and nicotine-1'-N-oxide ($P < 0.001$). Brain levels were also lower for these compounds. In contrast, the EA cohort had *higher* plasma and brain AUCs ($P < 0.001$) for the minor metabolite nornicotine. Brain-to-plasma ratios varied for nicotine and its metabolites, and by age. Following i.v. nicotine administration, similar age-related differences were observed, and this route allowed detection of a 1.6-fold-larger volume of distribution and 2-fold higher plasma clearance in the EA cohort compared with the AD cohort. Thus, unlike in humans, there are substantial age differences in nicotine pharmacokinetics such that for a given nicotine dose, adolescent rats will have lower plasma and brain nicotine compared with adults, suggesting that this should be considered when interpreting animal model data.

Introduction

Tobacco dependence is primarily mediated through the pharmacological actions of nicotine (Benowitz et al., 2009; Caille et al., 2012). The study of the pharmacokinetics of nicotine provides insight into variation in nicotine levels and how this may subsequently influence nicotine-mediated behaviors. Among the factors that influence nicotine pharmacokinetics is age (Benowitz et al., 2009). For example, in humans over the age of 65, nicotine clearance (CL) is reduced (Molander et al., 2001), and in neonates, the nicotine half-life ($t_{1/2}$) is

longer compared with adults (Dempsey et al., 2000). No differences in nicotine pharmacokinetics have been observed between human adolescents and adults (Gourlay and Benowitz, 1996; Al Koudsi et al., 2010). However, differences in nicotine plasma levels between adolescent and adult rats given the same nicotine dose in a behavioral study were noted, suggesting potential age-related variation in nicotine pharmacokinetics (Shram et al., 2008).

Nicotine is inactivated and eliminated from the body by various metabolic pathways, primarily mediated by hepatic enzymes (Nakajima and Yokoi, 2005). Nicotine's primary route of elimination in most mammalian species is via its conversion to cotinine by C-oxidation (Messina et al., 1997; Nakajima et al., 1998). Oxidative nicotine metabolism also produces minor nicotine metabolites including nornicotine (Green et al., 2001) and nicotine-1'-N-oxide (Ochiai et al., 2006). Measuring the pharmacokinetics of nicotine and its metabolites over time, preferably in both the plasma and brain, enhances the interpretation of nicotine's effects (Hukkanen et al., 2005).

The majority of current adult smokers began smoking in adolescence, with 88% of first cigarette use occurring before age 18 (Liang et al., 2003; US Department of Health and Human Services, 2012). Human adolescents have a higher risk of developing nicotine dependence, relative to adults, even with limited tobacco exposure (Rose et al., 2012). Initiation of smoking during adolescence has been associated with a lower likelihood of quitting in adulthood, increased cigarette

Parts of this work were previously presented at the following meeting: Craig E, Khokhar JY, Zhao B, Novalen M, Miksys SM, and Tyndale RF (2013) Altered nicotine pharmacokinetics in adolescent versus adult rats impacts the interpretation of animal model data. *Society for Research on Nicotine and Tobacco Annual Meeting*; 2013 Mar 13–16; Boston, MA.

This study was conducted with the support of an Endowed Chair in Addictions (to R.F.T.) and the Canadian Institute of Health Research [Grant MOP97751], National Institutes of Health [Grant U01-DA020830], Canada Foundation for Innovation [Grants 20289 and 16014], Campbell Family Mental Health Research Institute Centre for Addiction and Mental Health, and Ontario Ministry of Research and Innovation.

R.F.T. has participated in one-day consulting meetings for Novartis and McNeil. dx.doi.org/10.1124/dmd.114.058719.

§This article has supplemental material available at dmd.aspetjournals.org.

ABBREVIATIONS: AD, adult; AUC, area under the concentration–time curve; CL, nicotine clearance; EA, early adolescent; HPLC, high-performance liquid chromatography; K_p , brain-to-plasma partition coefficient; LOQ, limit of quantification; PND, postnatal day; $t_{1/2}$, half-life; VD, volume of distribution.

consumption, and increased difficulty quitting (Khuder et al., 1999). Animal models can be used to investigate aspects of the nicotine dependence cycle that cannot be easily investigated in humans, such as the initiation of use (Casey and Jones, 2010). For example, using the conditioned place preference paradigm, which assesses drug reward in rodents (Pema et al., 2011), acute low doses of s.c. nicotine elicited a preference for the nicotine-paired chamber in adolescent but not in adult rats (Vastola et al., 2002; Belluzzi et al., 2004). When first exposed to nicotine in adolescence in an i.v. self-administration paradigm, both male (Levin et al., 2007) and female (Levin et al., 2003) rats displayed enhanced rates of responding compared with animals first exposed in adulthood. These behavioral differences between adolescents and adults may be attributed to a reduced sensitivity to nicotine's aversive effects (Vastola et al., 2002), differences in reward set points during development (Koob and Le Moal, 2001), and/or pharmacokinetic differences (Levin et al., 2003). Plasma nicotine and cotinine levels, measured at a single time point, were significantly lower in adolescent compared with adult rats (Shram et al., 2008). Adolescent rats also had enhanced upregulation of brain nicotinic acetylcholine receptors in response to nicotine and significantly lower brain nicotine levels than those reported in adults (Doura et al., 2008). These studies suggest that pharmacokinetic differences may contribute to the behavioral differences observed between adolescent and adult rats.

This study was conducted to directly assess these potential age-specific pharmacokinetic differences in nicotine and its metabolites and to assist in understanding the potential confound of extrapolating the results of rat behavioral studies to the interpretation of human nicotine-mediated behaviors, given the apparent species differences in nicotine pharmacokinetics as a function of age. Acute dosing of s.c. and i.v. nicotine was used to assess the impact of age on the plasma and brain levels of nicotine and its metabolites in early adolescent and adult rats.

Materials and Methods

Animals. Male Wistar rats were purchased from Charles River Laboratories (St-Constant, QC, Canada). Rats were aged by postnatal days (PNDs); early adolescent (EA; PND 25) and adult (AD; PND 71) rats were used in each

experiment. These age groups were selected according to the previously established developmental periods of EA (PND 21–34) and AD (PND >60) (Spear, 2000; McCormick and Mathews, 2010). Male rats were chosen, as they had been used in the behavioral studies and to avoid the sex-specific influence of sexually dimorphic growth hormone release on drug metabolism, which occurs in rats and is not relevant in humans (Shapiro et al., 1995); see *Discussion*. All EA (81 ± 8 g) and AD (349 ± 25 g) animals were housed in pairs or groups of three and maintained on a 12 hour light/dark and temperature-controlled cycle. Water and food were available ad libitum. All experimental procedures were conducted in accordance with the guidelines provided by the Canadian Council on Animal Care and approved by the Animal Care Committee of the University of Toronto.

Reagents. (–)-Nicotine hydrogen tartrate salt was purchased from Sigma (St. Louis, MO). Nicotine- d_4 , rac-cotinine, rac-cotinine- d_3 , *trans*-3'-hydroxycotinine, *trans*-3'-hydroxycotinine- d_3 , (*R,S*)-norcotinine, (*R,S*)-nornicotine, (1'*S*,2'*S*)-nicotine-1'-oxide, and *rac-trans*-nicotine-1'-oxide-methyl- d_3 were obtained from Toronto Research Chemicals (North York, ON, Canada). The purity of all compounds used was >98%.

In Vivo Nicotine Pharmacokinetics. Animals received either an acute s.c. (1.0 mg free base/kg body weight, a higher dose allowing better quantification of nicotine and metabolites) or i.v. (0.2 mg free base/kg body weight, a lower dose required to avoid toxicity) administration of (–)-nicotine hydrogen tartrate dissolved in sterile saline (0.9% NaCl). Fresh nicotine solutions were prepared and titrated to a pH of 7.4. Venous blood (saphenous vein, 100–200 μ l) was collected at 10 minutes, 30 minutes, 1 hour, 2 hours, and 4 hours after nicotine injection, as well as trunk blood at sacrifice (2, 4, or 8 hours after nicotine injection). For each animal, two timed blood samples were obtained from the saphenous vein and the third sample from trunk blood at time of sacrifice. The sample timing and animal numbers were based on previous nicotine studies performed in our laboratory in adult rats, e.g., Micu et al. (2003), and pilot s.c. nicotine injection studies performed in adolescent rats. Thus, the total number of animals used was 30 for the EA and 14 for the AD groups for the s.c. study, and 25 for the EA and 23 for the AD groups for the i.v. dosing study; as each animal provided three of the six blood sampling times, this resulted in an average of 10, 9, 6, and 7 samples per time point, respectively. Blood samples were placed on ice, and plasma was collected after centrifugation at 3500g for 10 minutes and stored at -30°C until analyzed. Following sacrifice, brains were immediately dissected on ice, frozen on dry ice, and stored at -80°C .

Plasma Sample Preparation. Sample preparation and extraction was adapted from a previously described method (Vieira-Brock et al., 2011). Briefly,




Fig. 1. Plasma nicotine levels and $AUC_{(10\text{min}-4\text{h})}$ were lower in the EA_(s.c.) group following s.c. nicotine administration. (A) Plasma nicotine levels measured over time. (B) Plasma nicotine $AUC_{(10\text{min}-4\text{h})}$. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus AD_(s.c.) group.

TABLE 1

Pharmacokinetic parameters of plasma nicotine and metabolites in EA and AD rats following s.c. nicotine

Data are presented as mean (S.D.).

Parameter	NIC		COT		NNIC		NNO	
	EA	AD	EA	AD	EA	AD	EA	AD
C_{max} (ng/ml)	217 (37)	253 (65)	206 (73)	361 (112)	13** (2.5)	9 (1.8)	66** (23)	105 (8.4)
$t_{1/2}$ (min)	62	74	446	543	123	181	66	89
$AUC_{(10min-4h)}$ (ng·h/ml)	278*** (29)	479 (56)	658** (52)	960 (55)	39*** (1.7)	26 (1.7)	144*** (9.4)	255 (12)
$AUC_{(2-8h)}$ (ng·h/ml)	87* (29)	249 (47)	988* (193)	1893 (272)	37 (4.8)	34 (6.8)	66** (16)	148 (20)

COT, cotinine; NIC, nicotine; NNIC, normicotine; NNO, nicotine-1'-N-oxide.
 * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus AD_(s.c.) group.

100 μ l of 20 ng/ml deuterium-labeled internal standard in 0.01 M HCl, 900 μ l of water, and 100 μ l of 30% perchloric acid were added to 100 μ l of plasma or 100 μ l of standards, for a final volume of 1.2 ml. After vortex-mixing and centrifugation at 5400g for 5 minutes, the supernatant was added to Oasis MCX

solid-phase extraction cartridges (3 cc, 60 mg) (Waters, Milford, MA) that had been preconditioned with 2 ml of high-performance liquid chromatography (HPLC)-grade methanol followed by 2 ml of 2% formic acid. The cartridges were washed with 1 ml of 2% formic acid and 1 ml of HPLC-grade methanol. Analytes




Fig. 2. Plasma cotinine and nicotine-1'-N-oxide levels and $AUC_{(10min-4h)}$ were lower and plasma normicotine levels and $AUC_{(10min-4h)}$ were higher in the EA_(s.c.) group following s.c. nicotine administration. Plasma levels of cotinine (A), normicotine (B), and nicotine-1'-N-oxide (C) measured over time. (D) Plasma cotinine (COT), nicotine-1'-N-oxide (NNO), and normicotine (NNIC) $AUC_{(10min-4h)}$. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus AD_(s.c.) group.

TABLE 2

Plasma *trans*-3'-hydroxycotinine (3HC) and norcotinine (NCOT) levels in EA and AD rats following s.c. nicotine

Data are presented as mean (S.D.).

Time Point	Plasma 3HC Level		Plasma NCOT Level	
	EA	AD	EA	AD
	<i>ng/ml plasma</i>			
10 min	< LOQ	< LOQ	< LOQ	< LOQ
30 min	< LOQ	1.4 (3.7)	2.4 (0.8)	2.7 (3.8)
1 h	1.4 (0.8)	1.4 (0.1)	3.5 (1.3)	3.3 (0.3)
2 h	2.9 (1.6)	2.8 (0.9)	7.7 (3.1)	5.2 (1.8)
4 h	2.9 (1.2)	4.0 (1.9)	7.6 (2.6)	7.8 (3.1)
8 h	1.7 (0.4)	4.0 (0.5)	3.8 (0.7)	6.3 (1.0)

were eluted with 1.5 ml of 5% (v/v) ammonium hydroxide in methanol and 1.5 ml of dichloromethane/isopropanol/ammonium hydroxide, 78:20:2 (v/v/v). The samples were fortified with 100 μ l of 10% (v/v) hydrochloric acid in methanol to enhance nicotine recovery and then dried under a stream of nitrogen at 37°C. Dry samples were then reconstituted in 100 μ l of 100 mM ammonium acetate in water/methanol/acetic acid, 79:20:1 (v/v/v), and 50 μ l were injected onto a liquid chromatography–tandem mass spectrometer.

Brain Sample Preparation. Brain samples (whole brain) were thawed on ice, weighed, homogenized in three volumes of ice-cold saline (0.9% NaCl), and centrifuged at 3000g for 10 minutes. The supernatant was collected and frozen at -80°C until time of analysis. The brain supernatant preparation was performed as described above using 1.0 ml of brain supernatant (without the addition of water), as previously described (Vieira-Brock et al., 2011). The mean recovery ($n = 5$) was $>80\%$ for all compounds assessed, using labeled standards. The recovery of nicotine was 85.4% in EA (range, 82.6–87.3%) and 85.3% in AD (range, 83.2–88.4%) brain samples; the recovery of cotinine was 85.5% in EA (range, 81.2–90.1%) and 85.8% in AD (range, 81.6–89.9%); the recovery of normnicotine was 89.0% in EA (range, 86.3–93.3%) and 90.3% in AD (range, 88.8–93.3%); the recovery of nicotine-1'-*N*-oxide was 86.7% in EA (range, 83.8–90.2%) and 84.6% in AD (range, 82.9–87.3%); the recovery of *trans*-3'-hydroxycotinine was 85.3% in EA (range, 84.5–86.6%) and 84.3% in AD (range, 82.8–86.5%); and the recovery of norcotinine was 81.1% in EA (range, 77.2–86.1%) and 80.7% in AD (range, 76.4–85.7%).

Liquid Chromatography–Mass Spectrometry Conditions. The chromatographic conditions were adapted from an established method (Jacob et al., 2011) and were performed using an Agilent 1260 HPLC system (Agilent Technologies, Santa Clara, CA) equipped with a 4.6 mm \times 150 mm Synergi Polar-RP column (4 μ m) (Phenomenex, Torrance, CA). The mobile phase consisted of a 10 mM ammonium acetate/0.1% acetic acid in water (solvent A) and 10 mM ammonium acetate/0.1% acetic acid in methanol (solvent B). The flow rate was 0.7 ml/min, and the following gradient was used. The initial composition was 80% solvent A, changing to 100% solvent B over 6.5 minutes. From 6.5 to 8 minutes, 100% solvent B was maintained and was then changed to 80% solvent A at 8.1 minutes and maintained at this composition until the end of the run; total run time was 13 minutes. The spectrometric analysis was performed on an Agilent 6430 triple-quadrupole mass spectrometer equipped with an atmospheric pressure chemical ionization ion source operated in the positive ion mode. The vaporizer temperature was optimized to 450°C, the gas temperature was 350°C, and the corona discharge current was set at 5 μ A. The limit of quantification (LOQ) was 1 ng/ml for all analytes. Intraday coefficients of variation were assessed for all metabolites to determine intrarun assay precision. The coefficient of variation values were 2.3% for nicotine, 2.5% for cotinine, 1.9% for normnicotine, 1.6% for nicotine-1'-*N*-oxide, 1.1% for *trans*-3'-hydroxycotinine, and 3.0% for norcotinine.

Pharmacokinetic Analysis. In vivo pharmacokinetic parameters were determined using noncompartmental analysis and were calculated using PK Functions for Microsoft Excel (J.I. Usansky, A. Desia, D. Tan-Liu, Department of Pharmacokinetics and Drug Metabolism, Allergan, Irvine, CA). The area under the concentration-time curves from 10 minutes to 4 hours ($\text{AUC}_{(10\text{min}-4\text{h})}$) or 2 to 8 hours ($\text{AUC}_{(2-8\text{h})}$) in plasma and the $\text{AUC}_{(2-8\text{h})}$ in brain were determined using the log-linear trapezoidal rule. Half-life was estimated using the equation $t_{1/2} = \ln(2)/k_{\text{el}}$, where k_{el} is the terminal elimination rate constant. The terminal elimination rate constant was estimated by linear regression of the terminal phase of the semilogarithmic concentration-time curve. The brain-to-plasma partition coefficient (K_{p}) was calculated as a ratio of brain $\text{AUC}_{(2-8\text{h})}$ to plasma $\text{AUC}_{(2-8\text{h})}$. Following acute i.v. nicotine administration, the initial volume of distribution (VD) was calculated as Dose/C_0 , where C_0 is the initial plasma nicotine concentration, extrapolated to time zero of the concentration-time curve. The plasma CL was calculated as $\ln(2) \times V_c/t_{1/2}$ where V_c is the initial volume of distribution (Toutain and Bousquet-Mélou, 2004c).

Statistical Analysis. Statistical analyses were performed using GraphPad Prism 6.0 software (GraphPad Software Inc., La Jolla, CA). Variables were described by their mean \pm S.D. and were compared between the EA and AD




Fig. 3. Brain nicotine levels were lower in the EA_(s.c.) group following s.c. nicotine administration. (A) Brain nicotine levels measured over time. (B) There was no significant difference in mean brain nicotine $\text{AUC}_{(2-8\text{h})}$, though a trend was noted ($P = 0.06$). (C) The brain-to-plasma partition coefficient (K_{p}) was >1 in both the EA_(s.c.) and AD_(s.c.) groups. $**P < 0.01$ versus AD_(s.c.) group.

TABLE 3

Pharmacokinetic parameters of brain nicotine and metabolites in EA and AD rats following s.c. nicotine

Data are presented as mean (S.D.).

Parameter	NIC		COT		NNIC		NNO	
	EA	AD	EA	AD	EA	AD	EA	AD
$t_{1/2}$ (min)	159	97	449	347	139	137	172	112
AUC _(2-8h) (ng·h/g)	343 (105)	534 (59)	706** (58)	972 (46)	120* (20)	78 (13)	13 (3.4)	19 (3.0)

COT, cotinine; NIC, nicotine; NNIC, normicotine; NNO, nicotine-1'-N-oxide.

* $P < 0.05$, ** $P < 0.01$ versus AD_(s.c.) group.

groups using a one-tailed, unpaired Student's t -test with Welch's correction. Due to the study design, whereby individual animals were sampled across varying time points, only group means are reported for $t_{1/2}$, VD, and CL.

Results

Plasma Nicotine and Metabolite Levels Differ between Early Adolescent and Adult Rats. Plasma nicotine levels were lower in the EA_(s.c.) group compared with the AD_(s.c.) group at 1 hour ($P < 0.05$), 2 hours ($P < 0.001$), and 4 hours ($P < 0.01$) following acute s.c. nicotine administration (Fig. 1A); pharmacokinetic parameters are reported in Table 1. The mean plasma nicotine AUC_(10min-4h) was also lower in the EA_(s.c.) compared with the AD_(s.c.) group ($P < 0.001$), indicative of lower total nicotine exposure in the EA_(s.c.) animals (Fig. 1B). If the lower plasma nicotine levels in the EA_(s.c.) group were due to an increased rate of conversion to cotinine, the major metabolic route for nicotine, higher plasma cotinine levels might be expected in the younger animals. However, plasma cotinine levels were lower in the EA_(s.c.) group at 2 hours ($P < 0.05$), 4 hours ($P < 0.05$), and 8 hours ($P < 0.01$) compared with those in the AD_(s.c.) group (Fig. 2A). The mean plasma cotinine AUC_(10min-4h) was also lower in the EA_(s.c.) compared with the AD_(s.c.) group ($P < 0.01$) (Fig. 2D). We also investigated whether the formation of the minor metabolites normicotine and nicotine-1'-N-oxide contributed to the observed age differences in nicotine levels. Plasma normicotine levels were low, as expected, and higher at 30 minutes ($P < 0.05$), 1 hour ($P < 0.001$), and 2 hours ($P < 0.01$) in the EA_(s.c.) compared with the AD_(s.c.) group (Fig. 2B), resulting in a higher AUC_(10min-4h) ($P < 0.001$) (Fig. 2D). The mean normicotine C_{max} was also higher in the EA_(s.c.) versus the AD_(s.c.) group ($P < 0.001$) (Table 1). The plasma levels of nicotine-1'-N-oxide

were lower in the EA_(s.c.) group at 10 minutes ($P < 0.05$), 1 hour ($P < 0.01$), 2 hours ($P < 0.001$), 4 hours ($P < 0.001$), and 8 hours ($P < 0.015$) following nicotine administration compared with the AD_(s.c.) group (Fig. 2C), resulting in a lower mean nicotine-1'-N-oxide AUC_(10min-4h) ($P < 0.001$) (Fig. 2D). The mean plasma C_{max} for nicotine-1'-N-oxide was also lower in the EA_(s.c.) compared with the AD_(s.c.) group ($P < 0.01$) (Table 1). We also assessed the plasma levels of two additional minor nicotine metabolites. The mean plasma levels of *trans*-3'-hydroxycotinine and norcotinine were very low, near the LOQ (1 ng/ml), in both the EA_(s.c.) and AD_(s.c.) groups (Table 2).

Brain Levels of Nicotine and its Metabolites Differ between Early Adolescent and Adult Rats. Brain nicotine levels were lower in the EA_(s.c.) compared with the AD_(s.c.) group at 2 hours ($P < 0.01$) following acute s.c. nicotine administration (Fig. 3A; Table 3), consistent with a modestly lower AUC_(2-8h) ($P = 0.06$) (Fig. 3B). The brain nicotine K_p ratio from 2 to 8 hours was 3.9 in the EA_(s.c.) group and 2.2 in the AD_(s.c.) group, indicating higher partitioning of nicotine in the brain versus the plasma (Fig. 3C). Brain cotinine levels were lower in the EA_(s.c.) group at 2 hours ($P < 0.05$), 4 hours ($P < 0.05$), and 8 hours ($P < 0.05$) compared with the AD_(s.c.) group (Fig. 4A), as was the brain cotinine AUC_(2-8h) ($P < 0.01$) (Fig. 4B). The brain cotinine K_p ratio from 2 to 8 hours was < 1 for both the EA_(s.c.) and AD_(s.c.) groups, indicating that there was lower partitioning of cotinine into the brain (Fig. 4C). Brain normicotine levels were higher in the EA_(s.c.) group at 2 hours ($P < 0.05$) and 8 hours ($P < 0.05$) compared with the AD_(s.c.) group (Fig. 4A), as was the brain normicotine AUC_(2-8h) ($P < 0.05$) (Fig. 4B). The brain normicotine K_p ratio from 2 to 8 hours was 3.2 in the EA_(s.c.) group and 2.3 in the AD_(s.c.) group, indicative of higher partitioning of normicotine in the brain, as seen for nicotine (Fig. 4C). The brain nicotine-1'-N-oxide levels were relatively low




Fig. 4. Brain cotinine levels and AUC_(2-8h) were lower while brain normicotine levels and AUC_(2-8h) were higher in the EA_(s.c.) group following s.c. nicotine administration. (A) Brain cotinine, normicotine, and nicotine-1'-N-oxide levels measured over time. (B) Brain cotinine (COT), normicotine (NNIC), and nicotine-1'-N-oxide (NNO) AUC_(2-8h). (C) The brain-to-plasma partition coefficient (K_p) was < 1 for COT and NNO in both age groups. In contrast, the K_p for NNIC was > 1 in both age groups. * $P < 0.05$, ** $P < 0.01$ versus AD_(s.c.) group.

TABLE 4

Brain *trans*-3'-hydroxycotinine (3HC) and norcotinine (NCOT) levels in EA and AD rats following s.c. nicotine

Data are presented as mean (S.D.).

Time Point	Brain 3HC Level		Brain NCOT Level	
	EA	AD	EA	AD
	ng/g brain tissue			
2 h	< LOQ	< LOQ	4.3 (1.1)	3.1 (0.9)
4 h	1.0 (0.7)	1.4 (0.2)	4.9 (1.9)	5.1 (0.4)
8 h	< LOQ	1.2 (0.3)	2.7 (0.1)	3.5 (0.6)

and did not differ between the EA_(s.c.) and AD_(s.c.) animals (Fig. 4A); the K_p ratio from 2 to 8 hours was also <1, as seen for cotinine (Fig. 4C). Mean brain levels of *trans*-3'-hydroxycotinine and norcotinine were low, near the LOQ (1 ng/g), in both the EA_(s.c.) and AD_(s.c.) groups (Table 4).

Plasma and Brain Nicotine Levels Also Differed by Age Following i.v. Nicotine Administration. Plasma and brain nicotine levels were compared between the EA and AD age groups following an acute i.v. nicotine administration. The use of the i.v. route of administration, with 100% bioavailability, permits investigation of whether the differences in plasma and brain nicotine levels between the EA and AD age groups following acute s.c. nicotine resulted from differences in absorption, due perhaps to differences in body composition between these ages. Intravenous injection can also be used to examine the VD and CL.

Plasma nicotine levels were lower in the EA_(i.v.) compared with the AD_(i.v.) group at 30 minutes ($P < 0.05$), 1 hour ($P < 0.05$), 2 hours ($P < 0.001$), and 4 hours ($P < 0.05$) (Fig. 5A), resulting in a lower plasma nicotine AUC_(10min-4h) ($P < 0.001$) (Fig. 5B). Brain nicotine levels were not significantly different between the EA_(i.v.) and AD_(i.v.) groups at any time point (Fig. 6A); the mean brain nicotine AUC_(2-8h) trended toward being lower in the EA_(i.v.) group compared with the AD_(i.v.) group ($P = 0.09$) (Fig. 6B). The brain nicotine K_p ratio from 2 to 8 hours was 6.1 in the EA_(i.v.) group compared with 3.2 the AD_(i.v.)

group (Fig. 6C), indicating increased nicotine partitioning into the brain of EA rats, as observed following s.c. nicotine administration. Pharmacokinetic parameters for nicotine and metabolites following acute i.v. administration were similar to those seen following s.c. administration, with comparable age-related differences, and are reported in Supplementary Table 1. Due to the relatively flat elimination curve for nicotine from the brain (Fig. 6A), we examined the later time points of 18 and 24 hours. The only metabolite detected in the plasma beyond 8 hours was cotinine, detected in both age groups above the LOQ at 18 and 24 hours after nicotine administration. Nicotine and cotinine were detectable in the brain at 18 and 24 hours while other metabolites were not, suggesting a long residence time for these two compounds.

Volume of Distribution and Clearance Differ between Early Adolescent and Adult Rats. The EA_(i.v.) group had a mean VD that was ~1.6-fold larger than the AD_(i.v.) group (Fig. 7A). Furthermore, the estimated mean CL rate was >2-fold faster in the EA_(i.v.) group compared with that of the AD_(i.v.) group (Fig. 7B).

Discussion

There are nicotine-mediated behavioral differences between adolescent and adult rats (Belluzzi et al., 2004; Levin et al., 2007). We have shown that rat age (adolescent versus adult) substantially impacts nicotine pharmacokinetics, and consequently the plasma and brain levels of nicotine and its metabolites, which may contribute to these observed behavioral differences.

Pharmacokinetic parameters including absorption (Mangoni and Jackson, 2004), drug distribution (Turnheim, 1998; Molander et al., 2001), excretion (McLean and Le Coureur, 2004), and metabolism (Klinger, 2005) can be altered by age. Characterization of nicotine pharmacokinetics using s.c. nicotine can be influenced by absorption (Hukkanen et al., 2005). However, the pharmacokinetic differences between early adolescent and adult rats are not due to differences in nicotine absorption, as we obtained similar differences following i.v. nicotine administration, where bioavailability is 100% (Toutain and Bousquet-Mélou, 2004a). Throughout development, body mass




Fig. 5. Plasma nicotine levels and AUC_(10min-4h) were lower in the EA_(i.v.) group following i.v. nicotine administration. (A) Plasma nicotine levels measured over time. (B) Plasma nicotine AUC_(10min-4h). * $P < 0.05$, *** $P < 0.001$ versus AD_(i.v.) group.




Fig. 6. Brain nicotine levels were not significantly different in the EA_(i.v.) compared with the AD_(i.v.) group following i.v. nicotine administration. (A) Brain nicotine levels measured over time. (B) A trend toward a lower brain nicotine AUC_(2-8h) was noted in the EA_(i.v.) group ($P = 0.09$). (C) The K_p for brain nicotine was >1 in both age groups.

composition is altered in most mammalian species, with younger animals having a higher ratio of lean to fatty mass than older animals (Wolden-Hanson, 2010). As nicotine primarily distributes into lean mass, a change in this body mass composition could potentially impact nicotine's volume of distribution (Molander et al., 2001). As adolescent animals tend to have primarily lean body mass, they would be expected to have a larger volume of distribution, relative to their body weight, for nicotine, resulting in lower plasma nicotine levels (Tutka et al., 2005). Of note, the substantial change in volume of distribution (~ 1.6 times larger in EA) suggests that there are likely additional factors contributing to this difference.

Nicotine is excreted by glomerular filtration and tubular secretion with variable absorption, depending on urinary pH. Renal clearance accounts for $\sim 5\%$ of total nicotine clearance in humans (Benowitz et al., 2009). In vitro expression of rat organic cation transporters 1 and 2 indicated that they are basolaterally located transporters contributing to tubular secretion of cationic drugs, removing them from the plasma to be secreted in the urine (Urakami et al., 1998). In rats, renal transporter mRNA expression is increased in early adolescents (de Zwart et al., 2008). With developmental changes in transporter protein expression, it is predicted that the distribution and elimination of substrates would also change with age (de Zwart et al., 2008). The removal of a drug from the body is reflected by the plasma clearance, impacting plasma concentration (Toutain and Bousquet-Mélou, 2004b). The early adolescent animals have increased nicotine clearance, consistent with the lower plasma nicotine levels observed in this age group relative to the adult cohort.

In rats, nicotine is metabolized to cotinine by the hepatic enzyme CYP2B1, as the rat CYP2A family, chiefly responsible for nicotine metabolism in humans (Benowitz et al., 2006), is essentially inactive toward nicotine (Hammond et al., 1991; Nakayama et al., 1993). CYP2B expression changes through development; early adolescent rats had higher levels of hepatic CYP2B protein compared with adults (Yun et al., 2010). However, our data do not support a faster conversion of nicotine to cotinine in the early adolescent cohort, as both nicotine and cotinine levels are lower. Moreover, as nicotine is a high-extraction-ratio drug, its clearance is less affected by the expression and activity of hepatic enzymes (Benowitz et al., 2009).

This, and the faster nicotine clearance in younger animals, may be attributed in part to increased hepatic blood flow (Woodhouse and Wynne, 1992). Furthermore, most of the other metabolites were present at relatively lower levels in the adolescent compared with the adult cohort, suggesting that an increased rate of metabolism to these metabolites is not responsible, at least in full, for the lower nicotine levels in the adolescent cohort. Together, this suggests that the lower levels of nicotine in the adolescent cohort are consistent with an effect of volume of distribution and clearance rather than with a change in metabolic pathways.

Sex-dependent differences in drug metabolism and pharmacokinetics have been demonstrated in several species, including rats (Czerniak, 2001). In general, female rats have rates of hepatic drug metabolism 3- to 5-fold lower than those of males (Kyerematen et al., 1988a); the longer nicotine half-life and lower plasma cotinine levels observed in female versus male rats are consistent with lower recovery of urinary nicotine metabolites (Kyerematen et al., 1988a). Non-metabolic sex differences in nicotine pharmacokinetics include altered clearance and volume of distribution (Czerniak, 2001); female rats have a larger volume of distribution than males (Kyerematen et al., 1988a). However, despite the larger volume of distribution, and consistent with slower metabolism, female rats have higher plasma and brain nicotine levels compared with male rats, likely due to sex differences in nicotine distribution and metabolism (Rosecrans, 1972; Harrod et al., 2007). These sex differences in rat nicotine pharmacokinetics may also contribute to sex differences in nicotine-mediated behaviors, such as those seen for nicotine in the conditioned place preference paradigm (Torres et al., 2009), which may be related to the higher plasma and brain nicotine levels in females relative to males. Our findings of a potential role for differences in nicotine volume of distribution and clearance, and the known rodent sex differences in these parameters, suggest that differences between males and females of different ages may occur and should perhaps be considered when comparisons of nicotine-evoked behaviors are made.

Following acute nicotine administration, lower levels of nicotine were also observed in the brains of the adolescent animals. The lower brain nicotine levels may be important to the interpretation of differences in nicotine-mediated behaviors between adolescent and




Fig. 7. Age-related differences in volume of distribution and nicotine clearance may alter pharmacokinetics between the two age groups. (A) The initial VD was ~1.6-fold larger in the EA_(i.v.) versus the AD_(i.v.) group. (B) Nicotine CL was ~2-fold faster in the EA_(i.v.) group compared with the AD_(i.v.) group.

adult rats. Adolescent rats display increased rates of nicotine self-administration relative to adults (Levin et al., 2003, 2007), often attributed to an enhanced preference for nicotine. However, due to the lower brain nicotine levels observed in adolescent rats, it is possible that adolescent animals are self-administering more nicotine to attain rewarding brain levels comparable to those achieved by adults (Koob and Le Moal, 2001), effectively compensating for lower brain nicotine levels acquired per injection. We also observed a higher brain-to-plasma AUC_(2–8h) ratio in the early adolescents compared with the adults. Understanding the differing brain levels of nicotine measured directly, or by age-specific extrapolations from plasma levels, may be useful when interpreting animal behavioral data.

In contrast to the lower plasma and brain levels of nicotine, plasma normicotine levels were higher in early adolescent compared with adult rats. In both humans and rats, nicotine is metabolized to normicotine by *N*-demethylation (Cundy and Crooks, 1984; Yamanaka et al., 2005), accounting for a urinary recovery of 2–3% of a given nicotine dose in humans and ~8% in rats (Yamanaka et al., 2005). In this and former studies, normicotine represents a more substantial proportion of the nicotine dose in the brain (Plowchalk et al., 1992; Crooks et al., 1997). Normicotine has psychoactive properties and is self-administered by rats (Bardo et al., 1999); this metabolite also has a high affinity for nicotinic receptors and can evoke dopamine release in the striatum (Green et al., 2001). Normicotine desensitizes nicotinic receptor subtypes with a lower potency than nicotine, although cross-sensitization was observed between nicotine and normicotine, indicating common receptor subtypes (Dwoskin et al., 2001). We detected a significantly higher brain normicotine AUC_(2–8h) in the early adolescent cohort, indicating higher total brain normicotine exposure. In this age group, in which, for a given dose of nicotine, brain nicotine levels are lower and brain normicotine levels are higher than those in adults, the pharmacological actions of this metabolite may be contributing to observed age differences in nicotine-mediated behaviors.

Thus, given the same dose of nicotine, early adolescent rats have lower plasma and brain levels of nicotine, higher nicotine brain-to-plasma ratios, and differing levels of various nicotine metabolites in both plasma and brain compared with adult rats. Differences in volume of distribution and clearance are likely primarily responsible for these differences in nicotine levels. Additional contributing factors

include differences in renal and brain transport, hepatic blood flow, and, possibly, minor contributions from differences in metabolic pathways. Adolescence is a key period of development in which, in humans, the majority of initiation of nicotine use occurs (Breslau and Peterson, 1996). Animal models are useful for examining the underlying neurobiological vulnerability to dependence in adolescence in ways that cannot be easily investigated in humans (Caille et al., 2012). However, the rat may not be a suitable animal model of human nicotine pharmacokinetics given species-specific age differences in nicotine pharmacokinetics (Gourlay and Benowitz, 1996; Benowitz et al., 2009), differences in the enzyme primarily responsible for nicotine metabolism (Nakayama et al., 1993; Benowitz et al., 2006), and differences in the relative importance of various metabolic pathways of nicotine (Kyerematen et al., 1988b; Hukkanen et al., 2005). Overall, the results presented here highlight both the importance of age as a factor contributing to differences in nicotine pharmacokinetics in rats and the importance of taking these differences, as well as potential pharmacodynamic differences, into account when interpreting the results of animal model data and extrapolating them to interpretations of human smoking behaviors.

Acknowledgments

The authors thank Andy Z.X. Zhu and Reo Tanoshima for helpful discussions of the pharmacokinetics and statistical analyses.

Authorship Contributions

Participated in research design: Craig, Cui, Miksys, Tyndale.

Conducted experiments: Craig, Cui, Miksys, Zhao, Novalen.

Performed data analysis: Craig, Cui, Zhao.

Wrote or contributed to the writing of the manuscript: Craig, Tyndale.

References

- Al Koudsi N, Hoffmann EB, Assadzadeh A, and Tyndale RF (2010) Hepatic CYP2A6 levels and nicotine metabolism: impact of genetic, physiological, environmental, and epigenetic factors. *Eur J Clin Pharmacol* **66**:239–251.
- Bardo MT, Green TA, Crooks PA, and Dwoskin LP (1999) Normicotine is self-administered intravenously by rats. *Psychopharmacology (Berl)* **146**:290–296.
- Belluzzi JD, Lee AG, Oliff HS, and Leslie FM (2004) Age-dependent effects of nicotine on locomotor activity and conditioned place preference in rats. *Psychopharmacology (Berl)* **174**:389–395.
- Benowitz NL, Hukkanen J, and Jacob P, 3rd (2009) Nicotine chemistry, metabolism, kinetics and biomarkers, in *Nicotine Psychopharmacology* (Henningfield JE, London ED, and Pogun S eds) pp 29–60, Springer, Berlin.
- Benowitz NL, Swan GE, Jacob P, 3rd, Lessov-Schlaggar CN, and Tyndale RF (2006) CYP2A6 genotype and the metabolism and disposition kinetics of nicotine. *Clin Pharmacol Ther* **80**:457–467.
- Breslau N and Peterson EL (1996) Smoking cessation in young adults: age at initiation of cigarette smoking and other suspected influences. *Am J Public Health* **86**:214–220.
- Caille S, Clemens K, Stinus L, and Cador M (2012) Modeling nicotine addiction in rats. *Methods Mol Biol* **829**:243–256.
- Casey BJ and Jones RM (2010) Neurobiology of the adolescent brain and behavior: implications for substance use disorders. *J Am Acad Child Adolesc Psychiatry* **49**:1189–1201, quiz 1285.
- Crooks PA, Li M, and Dwoskin LP (1997) Metabolites of nicotine in rat brain after peripheral nicotine administration. Cotinine, normicotine, and norcotinine. *Drug Metab Dispos* **25**:47–54.
- Cundy KC and Crooks PA (1984) High-performance liquid chromatographic method for the determination of *N*-methylated metabolites of nicotine. *J Chromatogr A* **306**:291–301.
- Czerwiak R (2001) Gender-based differences in pharmacokinetics in laboratory animal models. *Int J Toxicol* **20**:161–163.
- Dempsey D, Jacob P, 3rd, and Benowitz NL (2000) Nicotine metabolism and elimination kinetics in newborns. *Clin Pharmacol Ther* **67**:458–465.
- de Zwart L, Scholten M, Monbaliu JG, Annaert PP, Van Houdt JM, Van den Wyngaert I, De Schaepe-drijver LM, Bailey GP, Coogan TP, and Coussement WC, et al. (2008) The ontogeny of drug metabolizing enzymes and transporters in the rat. *Reprod Toxicol* **26**:220–230.
- Doura MB, Gold AB, Keller AB, and Perry DC (2008) Adult and periadolescent rats differ in expression of nicotinic cholinergic receptor subtypes and in the response of these subtypes to chronic nicotine exposure. *Brain Res* **1215**:40–52.
- Dwoskin LP, Teng LH, and Crooks PA (2001) Normicotine, a nicotine metabolite and tobacco alkaloid: desensitization of nicotinic receptor-stimulated dopamine release from rat striatum. *Eur J Pharmacol* **428**:69–79.
- Gourlay SG and Benowitz NL (1996) The benefits of stopping smoking and the role of nicotine replacement therapy in older patients. *Drugs Aging* **9**:8–23.
- Green TA, Crooks PA, Bardo MT, and Dwoskin LP (2001) Contributory role for normicotine in nicotine neuropharmacology: normicotine-evoked [³H]dopamine overflow from rat nucleus accumbens slices. *Biochem Pharmacol* **62**:1597–1603.

- Hammond DK, Bjereke RJ, Langone JJ, and Strobel HW (1991) Metabolism of nicotine by rat liver cytochromes P-450. Assessment utilizing monoclonal antibodies to nicotine and cotinine. *Drug Metab Dispos* **19**:804–808.
- Harrod SB, Booze RM, and Mactutus CF (2007) Sex differences in nicotine levels following repeated intravenous injection in rats are attenuated by gonadectomy. *Pharmacol Biochem Behav* **86**:32–36.
- Hukkanen J, Jacob P, 3rd, and Benowitz NL (2005) Metabolism and disposition kinetics of nicotine. *Pharmacol Rev* **57**:79–115.
- Jacob P, 3rd, Yu L, Duan M, Ramos L, Yturralde O, and Benowitz NL (2011) Determination of the nicotine metabolites cotinine and trans-3'-hydroxycotinine in biologic fluids of smokers and non-smokers using liquid chromatography-tandem mass spectrometry: biomarkers for tobacco smoke exposure and for phenotyping cytochrome P450 2A6 activity. *J Chromatogr B Analyt Technol Biomed Life Sci* **879**:267–276.
- Khuder SA, Dayal HH, and Mutgi AB (1999) Age at smoking onset and its effect on smoking cessation. *Addict Behav* **24**:673–677.
- Klinger W (2005) Developmental pharmacology and toxicology: biotransformation of drugs and other xenobiotics during postnatal development. *Eur J Drug Metab Pharmacokinet* **30**:3–17.
- Koob GF and Le Moal M (2001) Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology* **24**:97–129.
- Kyerematen GA, Owens GF, Chattopadhyay B, deBethizy JD, and Vesell ES (1988a) Sexual dimorphism of nicotine metabolism and distribution in the rat. Studies in vivo and in vitro. *Drug Metab Dispos* **16**:823–828.
- Kyerematen GA, Taylor LH, deBethizy JD, and Vesell ES (1988b) Pharmacokinetics of nicotine and 12 metabolites in the rat. Application of a new radiometric high performance liquid chromatography assay. *Drug Metab Dispos* **16**:125–129.
- Levin ED, Lawrence SS, Petro A, Horton K, Rezvani AH, Seidler FJ, and Slotkin TA (2007) Adolescent vs. adult-onset nicotine self-administration in male rats: duration of effect and differential nicotinic receptor correlates. *Neurotoxicol Teratol* **29**:458–465.
- Levin ED, Rezvani AH, Montoya D, Rose JE, and Swartzwelder HS (2003) Adolescent-onset nicotine self-administration modeled in female rats. *Psychopharmacology (Berl)* **169**:141–149.
- Liang L, Chaloupka F, Nichter M, and Clayton R (2003) Prices, policies and youth smoking, May 2001. *Addiction* **98** (Suppl 1):105–122.
- Mangoni AA and Jackson SH (2004) Age-related changes in pharmacokinetics and pharmacodynamics: basic principles and practical applications. *Br J Clin Pharmacol* **57**:6–14.
- McCormick CM and Mathews IZ (2010) Adolescent development, hypothalamic-pituitary-adrenal function, and programming of adult learning and memory. *Prog Neuropsychopharmacol Biol Psychiatry* **34**:756–765.
- McLean AJ and Le Couteur DG (2004) Aging biology and geriatric clinical pharmacology. *Pharmacol Rev* **56**:163–184.
- Messina ES, Tyndale RF, and Sellers EM (1997) A major role for CYP2A6 in nicotine C-oxidation by human liver microsomes. *J Pharmacol Exp Ther* **282**:1608–1614.
- Micu AL, Miksys S, Sellers EM, Koop DR, and Tyndale RF (2003) Rat hepatic CYP2E1 is induced by very low nicotine doses: an investigation of induction, time course, dose response, and mechanism. *J Pharmacol Exp Ther* **306**:941–947.
- Molander L, Hansson A, and Lunell E (2001) Pharmacokinetics of nicotine in healthy elderly people. *Clin Pharmacol Ther* **69**:57–65.
- Nakajima M, Iwata K, Yamamoto T, Funae Y, Yoshida T, and Kuroiwa Y (1998) Nicotine metabolism in liver microsomes from rats with acute hepatitis or cirrhosis. *Drug Metab Dispos* **26**:36–41.
- Nakajima M and Yokoi T (2005) Interindividual variability in nicotine metabolism: C-oxidation and glucuronidation. *Drug Metab Pharmacokinet* **20**:227–235.
- Nakayama H, Okuda H, Nakashima T, Imaoka S, and Funae Y (1993) Nicotine metabolism by rat hepatic cytochrome P450s. *Biochem Pharmacol* **45**:2554–2556.
- Ochiai Y, Sakurai E, Nomura A, Itoh K, and Tanaka Y (2006) Metabolism of nicotine in rat lung microvascular endothelial cells. *J Pharm Pharmacol* **58**:403–407.
- Perna MK, Henderson YO, Bruner CL, and Brown RW (2011) An analysis of nicotine conditioned place conditioning in early postweanling and adolescent rats neonatally treated with quinpirole. *Behav Brain Res* **220**:254–261.
- Plowchalk DR, Andersen ME, and deBethizy JD (1992) A physiologically based pharmacokinetic model for nicotine disposition in the Sprague-Dawley rat. *Toxicol Appl Pharmacol* **116**:177–188.
- Rose JS, Lee CT, Dierker LC, Selya AS, and Mermelstein RJ (2012) Adolescent nicotine dependence symptom profiles and risk for future daily smoking. *Addict Behav* **37**:1093–1100.
- Rosecrans JA (1972) Brain area nicotine levels in male and female rats with different levels of spontaneous activity. *Neuropharmacology* **11**:863–870.
- Shapiro BH, Agrawal AK, and Pampori NA (1995) Gender differences in drug metabolism regulated by growth hormone. *Int J Biochem Cell Biol* **27**:9–20.
- Shram MJ, Siu EC, Li Z, Tyndale RF, and Lê AD (2008) Interactions between age and the aversive effects of nicotine withdrawal under mecamylamine-precipitated and spontaneous conditions in male Wistar rats. *Psychopharmacology (Berl)* **198**:181–190.
- Spear LP (2000) The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* **24**:417–463.
- Torres OV, Natividad LA, Tejada HA, Van Weelden SA, and O'Dell LE (2009) Female rats display dose-dependent differences to the rewarding and aversive effects of nicotine in an age-, hormone-, and sex-dependent manner. *Psychopharmacology (Berl)* **206**:303–312.
- Toutain PL and Bousquet-Mélou A (2004a) Bioavailability and its assessment. *J Vet Pharmacol Ther* **27**:455–466.
- Toutain PL and Bousquet-Mélou A (2004b) Plasma clearance. *J Vet Pharmacol Ther* **27**:415–425.
- Toutain PL and Bousquet-Mélou A (2004c) Volumes of distribution. *J Vet Pharmacol Ther* **27**:441–453.
- Turnheim K (1998) Drug dosage in the elderly. Is it rational? *Drugs Aging* **13**:357–379.
- Tutka P, Mosiewicz J, and Wielosz M (2005) Pharmacokinetics and metabolism of nicotine. *Pharmacol Rep* **57**:143–153.
- Urakami Y, Okuda M, Masuda S, Saito H, and Inui KI (1998) Functional characteristics and membrane localization of rat multispecific organic cation transporters, OCT1 and OCT2, mediating tubular secretion of cationic drugs. *J Pharmacol Exp Ther* **287**:800–805.
- US Department of Health and Human Services (2012) *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*, US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, Atlanta, GA.
- Vastola BJ, Douglas LA, Varlinskaya EI, and Spear LP (2002) Nicotine-induced conditioned place preference in adolescent and adult rats. *Physiol Behav* **77**:107–114.
- Vieira-Brock PL, Miller EI, Nielsen SM, Fleckenstein AE, and Wilkins DG (2011) Simultaneous quantification of nicotine and metabolites in rat brain by liquid chromatography-tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* **879**:3465–3474.
- Wolden-Hanson T (2010) Changes in body composition in response to challenges during aging in rats. *Interdiscip Top Gerontol* **37**:64–83.
- Woodhouse K and Wynne HA (1992) Age-related changes in hepatic function. Implications for drug therapy. *Drugs Aging* **2**:243–255.
- Yamanaka H, Nakajima M, Fukami T, Sakai H, Nakamura A, Katoh M, Takamiya M, Aoki Y, and Yokoi T (2005) CYP2A6 AND CYP2B6 are involved in normicotine formation from nicotine in humans: interindividual differences in these contributions. *Drug Metab Dispos* **33**:1811–1818.
- Yun KU, Oh SJ, Oh JM, Kang KW, Myung C-S, Song GY, Kim B-H, and Kim SK (2010) Age-related changes in hepatic expression and activity of cytochrome P450 in male rats. *Arch Toxicol* **84**:939–946.

Address correspondence to: Rachel F. Tyndale, Department of Pharmacology & Toxicology, University of Toronto, 1 King's College Circle, Room 4326, Medical Sciences Building, Toronto, Ontario M5S 1A8, Canada. E-mail: r.tyndale@utoronto.ca
