

Published in final edited form as:

Science. 2011 May 13; 332(6031): 855–858. doi:10.1126/science.1201618.

Natural microbe-mediated refractoriness to *Plasmodium* infection in *Anopheles gambiae*

Chris M. Cirimotich¹, Yuemei Dong¹, April M. Clayton¹, Simone L. Sandiford¹, Jayme A. Souza-Neto¹, Musapa Mulenga², and George Dimopoulos^{1,*}

¹W. Harry Feinstone Department of Molecular Microbiology and Immunology, Bloomberg School of Public Health, Johns Hopkins University, 615 N. Wolfe Street, Baltimore, MD 21205-2179, USA

²Malaria Institute at Macha, Choma, Zambia

Abstract

Malaria parasite transmission depends on the successful transition of *Plasmodium* through discrete developmental stages in the lumen of the mosquito midgut. Like the human intestinal tract, the mosquito midgut contains a diverse microbial flora, which may compromise the ability of *Plasmodium* to establish infection. We have identified an *Enterobacter* bacterium isolated from wild mosquito populations in Zambia that renders the mosquito 99% resistant to infection with the human malaria parasite *Plasmodium falciparum* by interfering with parasite development prior to invasion of the midgut epithelium. Phenotypic analyses showed that the anti-*Plasmodium* mechanism requires small populations of replicating bacteria and is mediated through a mosquito-independent interaction with the malaria parasite. We show that this anti-*Plasmodium* effect is largely caused by bacterial generation of reactive oxygen species.

Plasmodium parasites suffer considerable losses during their development in the mosquito midgut (1, 2), where they encounter a hostile environment of human blood-derived factors, mosquito innate immune responses, and resident microbiota. However, escape of only a few parasites is sufficient to ensure onward transmission. Midgut bacteria play a key role in modulating *Plasmodium* infection of the *Anopheles* mosquito vector (3–7), but the mechanism(s) of bacterially mediated parasite inhibition has not been described, although the mosquito's innate immune responses to the microbiota and parasite challenge have been implicated in this phenomenon (6–10).

In the present study, we isolated bacteria from southern Zambian populations of *A. arabiensis*, an important malaria vector, during two collecting trips in two consecutive years (11). A majority of the captured mosquito's midguts contained cultured bacteria, with an average concentration of 10⁴ bacteria per non-bloodfed gut, similar to that observed for

*To whom correspondence should be addressed: gdimopou@jhsph.edu.

Supporting Online Material

www.sciencemag.org

Materials and Methods

Figs. S1–S8

Tables S1–S3

References and notes

laboratory-reared *Anopheles* mosquitoes (6). Sixteen distinct bacterial strains were identified based on 16S ribosomal DNA sequence, with similar genera isolated from both collections (Table S1). Of these, some Gram-negative (G⁻) isolates reduced *P. falciparum* prevalence and intensity in the mosquito, while a Gram-positive (G⁺) isolate had no detectable effect on infection (Fig 1A), in accordance with previous reports involving other *Plasmodium-Anopheles* models (3–5). The G⁻ bacteria inhibited *P. falciparum* oocyst formation in the main African and Asian malaria vectors, *A. gambiae* and *A. stephensi*, respectively (Fig. 1A, S1). To investigate the temporal specificity of parasite inhibition by G⁻ bacteria, we monitored the development of the *Plasmodium* ookinetes in the presence of the field isolated bacteria. Significantly fewer ookinetes developed with G⁻ bacteria present, although the level of inhibition was bacterial species-dependent (Fig 1B). The *Enterobacter sp. (Esp_Z)* bacterium inhibited ookinete, oocyst, and sporozoite development of a highly virulent laboratory *Plasmodium* strain by 98%, 99% and 99%, respectively (Fig 1A–C), leading us to further investigate the mechanism of this inhibition.

The immune deficiency (IMD) innate immune pathway defends mosquitoes against *P. falciparum* in the gut tissue, and the microbiota has been shown to activate this pathway through the receptor protein peptidoglycan recognition protein-LC (PGRP-LC) (6–8); thus, we hypothesized that the refractory phenotype could be caused by *Esp_Z* activation of the IMD pathway. Two independent approaches showed that the mechanism of refractoriness is independent of the IMD pathway. First, although a general antibacterial response is mounted through the increased transcription of the antimicrobial peptide cecropin1 (CEC1), regulation of IMD pathway controlled genes, including several potent anti-*Plasmodium* effector genes [fibrinogen immunoelectin 9 (FBN9), leucine-rich repeat protein LRRD7, and thioester-containing protein 1 (TEP1) (12)] were similar in the midguts of mosquitoes challenged with *Esp_Z* or the non-inhibitory *Bacillus* bacterium (*Bpu*) (Fig 1D). Second, RNA interference-mediated depletion of the key pathway molecules, PGRP-LC, Imd, and REL2, did not rescue *P. falciparum* oocyst development in the presence of *Esp_Z* (Fig. 1E).

Hence, we investigated the potential for parasite inhibition exclusive of the mosquito. Although *P. falciparum* ookinete development was inefficient *in vitro*, co-culture of *Esp_Z* with gametocytes inhibited ookinete formation by 89% (Fig 1F). We observed a similarly strong *in vitro* inhibition of ookinete formation in the more-robust rodent malaria parasite *P. berghei* experimental model (Fig 1G).

Inhibition of *P. falciparum* by *Esp_Z* in the mosquito was dose-dependent, with a threshold of 10⁴ ingested bacteria providing near complete protection against parasite infection (Fig 2A). A remarkably low density of only 100 ingested bacteria was still able to significantly decrease oocyst intensity by 67% (Fig 2A). *In vitro* ookinete development of *P. berghei* was also inhibited in a dose-dependent manner (Fig. 2B). *Esp_Z* populations in the midgut expanded by 100–1,000 fold (Fig 2C) which is within the range of microbial proliferation that normally occurs in the midgut lumen after a blood meal (5, 6). The bacterial growth during the 24hr period immediately following blood ingestion correlates with the time period of parasite inhibition prior to ookinete formation (3–30hr after ingestion) (13). Negative correlations were also observed with the timing of bacterial replication and inhibition of *P. berghei* ookinete development *in vitro* (correlation coefficient= -0.95 for

10^6 *Esp_Z* and -0.94 for 10^7 *Esp_Z*) (Fig S2–3), and taken together, these results indicated that active replication of the bacteria was required for parasite inhibition. Mosquito exposure to heat-inactivated (HIA) *Esp_Z* upon feeding on a *P. falciparum* gametocyte culture did not result in the same level of refractoriness that was observed with exposure to live bacteria (Fig. 2D). Because of an overlap in antibacterial and anti-*Plasmodium* immune defenses in mosquitoes (6, 7, 9, 14), the observed decrease in oocyst numbers with high concentrations of HIA bacteria (Fig 2D) could be caused by the induction of an antibacterial response in the mosquito gut, while physical inhibition of parasite infection by the killed bacteria is also possible.

These observations, along with microscopy (Figure 2E), indicated that *Esp_Z* inhibition of *Plasmodium* did not involve direct association between the bacteria and parasite and was mediated by diffusible bacterial factors produced during replication, or bacterial sequestration of mosquito factors that are essential for *Plasmodium* development. In subsequent assays, we observed that *Plasmodium* inhibition was independent of bacterial fatty acid metabolism (Figure S4) (15), xanthurenic acid (XA) (Fig S5) and iron (Fig S6) utilization by the parasite.

Because the inhibitory effect did not appear to be dependent on the sequestration or acquisition of a molecule necessary for the parasite, we hypothesized that the bacteria were producing an anti-*Plasmodium* molecule. We tested this by examining the *in vitro* development of *P. berghei* first in co-culture with physically separated bacteria and second in filtered fresh supernatant of a bacterial culture. Separation of bacteria and parasites abolished bacteria-mediated inhibition of ookinete formation, except at very high bacterial concentrations (Fig 3A). When parasites were cultured in filtered fresh supernatant from an *Esp_Z* culture, there was a two-fold reduction in numbers of ookinetes formed (Fig 3B). Together, these data indicated that the inhibitory activity was mediated by a short-lived molecule in a concentration-dependent manner. Reactive oxygen species (ROS) have a short half-life, kill *Plasmodium* (16–19), and can be generated by bacteria (20), so we tested the hypothesis that bacteria were inhibiting parasite development by producing ROS.

Among the field mosquito –derived bacteria, those lacking ookinete inhibitory activity (Fig 1B) did not produce detectable levels of ROS, whereas cultures of the inhibitory *Esp_Z* did (Table S2). To determine whether ROS was involved in the parasite inhibition, we supplemented the *P. berghei* culture with an antioxidant to neutralize free radicals formed. The addition of vitamin C (vitC) to *in vitro* cultures of *P. berghei* gametocytes rescued development of ookinetes to untreated control levels when grown in filtered *Esp_Z* culture medium (Fig 3B), and parasite development in the presence of replicating *Esp_Z* was rescued by vitC in a dose-dependent fashion (Fig 3C). Furthermore, reduced glutathione, another potent antioxidant, also rescued *in vitro* ookinete formation in the presence of *Esp_Z* (Fig S7). We also showed that vertebrate leukocytes were not responsible for the observed *in vitro* ookinete inhibition (Fig S8). More importantly, supplementing an infectious bloodmeal with vitC did not impact parasite numbers in the absence of *Esp_Z* but rescued *P. falciparum* ookinete development two-fold in the lumen of *A. gambiae* midguts upon co-feeding with *Esp_Z* (Fig 3D). The significant, yet incomplete, rescue of ookinete development with higher bacterial concentrations could be attributed to a variety of factors

such as insufficient concentrations of antioxidant to neutralize the higher amount of bacteria produced ROS, excretion of significant amounts of antioxidant through mosquito diuresis, the intimate association between bacteria and parasites that may not enable detoxification of ROS prior to parasite inhibition, or the loss of antioxidant activity from prolonged exposure in the digestive environment of the midgut. Antioxidant concentrations higher than 10mM in the bloodmeal interfered with mosquito feeding propensity.

Genotypic analyses of laboratory and wild mosquito populations have suggested that a dominant refractory phenotype is associated with innate immunity and that *Plasmodium* infection is a result of immune failure (21–23). Our studies show a mechanism of *Plasmodium* inhibition that does not involve the mosquito-derived innate immune response, and they support the idea that the native microflora of *Anopheles* mosquitoes plays a crucial role in refractoriness to *Plasmodium* infection, and will therefore influence transmission success to humans.

Bacteria of the genus *Enterobacter* have been isolated from many anopheline mosquito species in diverse geographic regions (3, 5, 24). We show that mosquitoes do not become infected with *Plasmodium* parasites when exposed to an *Enterobacter* bacterium isolated from wild mosquito populations in southern Zambia, and we show that inhibition of parasite development can be mediated by bacterial generation of ROS. While *Esp_Z* was isolated from 25% of mosquitoes collected during one rainy season, it may be possible to manipulate the composition of the midgut microbial flora in wild mosquitoes to increase the prevalence of *Esp_Z* or other naturally inhibitory bacteria as part of an integrated malaria control strategy.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work has been supported by the National Institutes of Health/National Institute of Allergy and Infectious Disease R01AI061576 and a Johns Hopkins Malaria Research Institute Pilot Grant (to G.D.), the Calvin A. and Helen H. Lang fellowship (to C.M.C.), a JHMRI postdoctoral fellowship (to J.S.-N.), and a fellowship from the National Science Foundation (to A.M.C.). The authors thank the mosquito collection team at the Malaria Institute at Macha, Zambia, the Johns Hopkins Malaria Research Institute (JHMRI) Parasitology and Insectary Core facilities, Sanaria Inc., Dr. Eric Nelson (Cornell University) for providing mutant bacteria strains, and Dr. Deborah McClellan for editorial services. GenBank accession numbers generated for bacterial 16S rDNA sequences are listed in Table S1, SOM.

References and Notes

1. Alavi Y, et al. Int J Parasitol. 2003; 33:933. [PubMed: 12906877]
2. Vaughan JA, Noden BH, Beier JC. Am J Trop Med Hyg. 1994; 51:233. [PubMed: 8074258]
3. Gonzalez-Ceron L, Santillan F, Rodriguez MH, Mendez D, Hernandez-Avila JE. J Med Entomol. 2003; 40:371. [PubMed: 12943119]
4. Pumpuni CB, Beier MS, Nataro JP, Guers LD, Davis JR. Exp Parasitol. 1993; 77:195. [PubMed: 8375488]
5. Pumpuni CB, Demayo J, Kent M, Davis JR, Beier JC. Am J Trop Med Hyg. 1996; 54:214. [PubMed: 8619451]

6. Dong Y, Manfredini F, Dimopoulos G. PLoS Pathog. 2009; 5:e1000423. [PubMed: 19424427]
7. Meister S, et al. PLoS Pathog. 2009; 5:e1000542. [PubMed: 19662170]
8. Garver LS, Dong Y, Dimopoulos G. PLoS Pathog. 2009; 5:e1000335. [PubMed: 19282971]
9. Dong Y, et al. PLoS Pathog. 2006; 2:e52. [PubMed: 16789837]
10. Kumar S, Molina-Cruz A, Gupta L, Rodrigues J, Barillas-Mury C. Science. 2010; 327:1644. [PubMed: 20223948]
11. Materials and methods are available as supporting material on *Science* online.
12. Cirimotich CM, Dong Y, Garver LS, Sim S, Dimopoulos G. Dev Comp Immunol. 2010; 34:387. [PubMed: 20026176]
13. Vaughan JA, Noden BH, Beier JC. J Parasitol. 1992; 78:716. [PubMed: 1635032]
14. Meister S, et al. Proc Natl Acad Sci USA. 2005; 102:11420. [PubMed: 16076953]
15. van Dijk K, Nelson EB. Appl Environ Microbiol. 2000; 66:5340. [PubMed: 11097912]
16. Kumar S, et al. Proc Natl Acad Sci USA. 2003; 100:14139. [PubMed: 14623973]
17. Luckhart S, Vodovotz Y, Cui L, Rosenberg R. Proc Natl Acad Sci USA. 1998; 95:5700. [PubMed: 9576947]
18. Molina-Cruz A, et al. J Biol Chem. 2008; 283:3217. [PubMed: 18065421]
19. Peterson TML, Gow AJ, Luckhart S. Free Radic Biol Med. 2007; 42:132. [PubMed: 17157200]
20. Mai-Prochnow A, et al. J Bacteriol. 2008; 190:5493. [PubMed: 18502869]
21. Blandin S, et al. Science. 2009; 326:147. [PubMed: 19797663]
22. Niaré O, et al. Science. 2002; 298:213. [PubMed: 12364806]
23. Riehle MM, et al. Science. 2006; 312:577. [PubMed: 16645095]
24. Terenius O, et al. J Med Entomol. 2008; 45:172. [PubMed: 18283961]




Figure 1. Field bacteria-mediated inhibition of *Plasmodium* development

P. falciparum oocyst (A) and ookinete (B) loads in midguts of *A. gambiae* mosquitoes co-fed with parasites and field-isolated bacteria. Bpu, *Bacillus pumilus*; Asp, *Acinetobacter sp.*; Ppu, *Pseudomonas putida*; Esp_Z, *Enterobacter sp.* (C) *P. falciparum* sporozoite loads in salivary glands of *A. gambiae* mosquitoes co-fed with parasites and Esp_Z. For (A–C), circles represent the number of parasites from an individual mosquito and horizontal lines indicate the median number of parasites per tissue. (D) Midgut-specific transcript abundance of select genes at 8 h after bloodfeeding with equal quantities of either the Bpu or Esp_Z isolate. Each column and error bar represents the fold-change \pm standard deviation in transcript abundance when compared to PBS-fed controls. LC= PGRP-LC; CEC1=cecropin1; FBN9=fibrinogen immunolection 9; TEP1=thioester-containing protein 1; LRRD7=leucine-rich repeat-containing protein 7. (E) Oocyst loads in mosquitoes depleted of transcripts for IMD pathway molecules and co-challenged with Esp_Z and *P. falciparum*. The dsRNA and absence (–) or presence (+) of Esp_Z are indicated below each column. Circles represent the same as in (A). (F–G) *In vitro* development of *P. falciparum* (E) and *P. berghei* (F) ookinetes co-cultured with Esp_Z bacteria. Bars represent the mean \pm standard deviation in ookinetes. For all figures, statistical significance is represented by letters above each column, with different letters signifying distinct statistical groups ($p < 0.05$; Mann-Whitney test for (A–C, E); $p < 0.05$; unpaired t-test for (F–G)).




Figure 2. Phenotypic analyses of *Esp_Z* modulation of *Plasmodium* development

(A–B) Effect of *Esp_Z* dosage on *P. falciparum* oocyst formation in mosquitoes (A) and *P. berghei* ookinete formation *in vitro* (B). Circles in (A) represent the number of oocysts in an individual mosquito midgut, and the horizontal lines indicate the median number of parasites per midgut. Bars in (B) represent the mean \pm the standard deviation in percentage of the number of ookinetes formed in bacteria-treated groups as compared to PBS-treated controls. (C) Temporal replication of *Esp_Z* in mosquito midguts following bloodmeal administration of different inoculating doses of bacteria. CFU, colony-forming unit. Bars represent the mean \pm the standard deviation. (D) Effect of heat-inactivated *Esp_Z* bacteria on *P. falciparum* oocyst formation. Circles represent the same as in (A). For (A), (B) and (D), statistical significance is represented by letters above each column, with different letters signifying distinct statistical groups ($p < 0.05$; Mann-Whitney test for (A) and (D), unpaired t-test for (B)). (E) *In vivo* and *in vitro* *P. falciparum* development in the presence of *Esp_Z*. Scale bar = 10 μm .




Figure 3. Involvement of ROS generation by *Esp_Z* in inhibition of *Plasmodium* development
(A) Effect of physical separation of *Esp_Z* and parasites on *P. berghei* ookinete formation.
(B) Effect of filtered culture supernatant and addition of vitamin C on *P. berghei* ookinete formation. Sup=supernatant; Neg=supernatant from a bacteria-negative culture; VitC=vitamin C. For (A–B), bars represent the mean ± the standard deviation. **(C–D)** Effect of vitamin C addition on *P. berghei* ookinete formation *in vitro* (C) and *P. falciparum* ookinete formation in *A. gambiae* midguts (D). For (C), bars represent the mean ± the standard deviation in percentage of the number of ookinetes formed in bacteria-treated groups as compared to PBS-treated controls. For (D), circles represent the number of ookinetes from an individual mosquito and horizontal lines indicate the median number of parasites per midgut. For all figures, statistical significance is represented by letters above

each column, with different letters signifying distinct statistical groups ($p < 0.05$; unpaired t-test for (A–C); Mann-Whitney test for (D)).