

In-Vitro* Evaluation of Cytotoxic and Apoptogenic Properties of *Sophora Pachycarpa

Seyed Hadi Mousavi^a, Mahsa Motaez^a, Amir Zamiri-Akhlaghi^b,
Seyed Ahmad Emami^c and Zahra Tayarani-Najaran^{d*}

^aMedical Toxicology Research Centre, Mashhad University of Medical Sciences, Mashhad, Iran. ^bPharmacological Research Centre of Medicinal Plants, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. ^cDepartment of Pharmacognosy, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. ^dDepartment of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.

Abstract

Sophora pachycarpa Schrenk ex C.A.Mey. belongs to the family Fabaceae. Some species of the genus *Sophora* have shown to possess anti-proliferative and apoptosis-inducing activities in cancer cells. However, there is no available information addressing this effect in *S. pachycarpa*. Here, we investigated the cytotoxic effects of methanol extract and different fractions obtained from *S. pachycarpa* root on different cancer cell lines including A549, HeLa, HL-60, MCF-7, and PC3 cell lines and leukocytes as non-malignant cells. Apoptotic cells were determined using PI staining of DNA fragmentation by flow cytometry (sub-G1 peak). *S. pachycarpa* inhibited the growth of malignant cells in a dose-dependent manner. CH₂Cl₂ and EtOAc fractions showed the lowest IC₅₀ values ranging from 6 to 50 µg/mL in various cancer cell lines. HeLa cells as the most sensitive cells were chosen for further mechanistic studies. The sub-G1 peak in flow cytometry histogram of *S. pachycarpa* treated HeLa cells indicates apoptotic cell death in *S. pachycarpa*-induced toxicity. In conclusion, *S. pachycarpa* exerts cytotoxic effects in different cancer cell lines in which apoptosis plays an important role. Thus, *S. pachycarpa* could be considered as a potential chemotherapeutic agent in cancer treatment.

Keywords: Apoptosis; *Sophora pachycarpa*; Fabaceae; Cytotoxicity; Cancer.

Introduction

Plants have been served as the main sources of biologically active compounds in the prevention as well as treatment of many diseases including cancer. Molecular diversity of bio-active compounds in natural products provides diverse and wealthy source of phytochemical with recognized potential in drug discovery and

development (1, 2). Numerous cytostatic drugs with antitumor properties are plant derived components including alkaloids, phenolics, flavonoids, and terpenoids (3).

The genus *Sophora* (Fabaceae) is divided into two subgenera, *Sophora* and *Styphnolobium*. This genus has 52 species in the world and three of these species are found in Iran include: 1- *S. mollis* (Royle) Baker with three subspecies *mollis*, *griffithii* (Stocks) Ali and *sylves* M. Abbasi; 2- *S. alopecuroides* L. with two subspecies *alopecuroides* and *tomentosa* (Boiss) Yakovlev;

* Corresponding author:

E-mail: tayaraninz@mums.ac.ir

and 3- *S. pachycarpa* Schrenk ex C.A.Mey, and a hybrid *S. alopecroidese* × *S. pachycarpa* (4).

The dried roots of *Sophora* are one of the most popular and multi-purpose herbs used in China [*S. tonkinensis* (Pierre) Craib ex Hartwiss] and Japan [*S. flavescens* Aiton] (5).

The root of *S. flavescens* is a popular Chinese herbal medicine (Chinese name “Ku-Shen”) used in a variety of disorders including gastric disturbance and eczema, and also as anti-pyretic and anti-helminthic. This genus is an abundant source of peryl flavonoids (5-11), flavonoids (12) and alkaloids (13-17).

Previous studies on the chemical composition of *S. pachycarpa* showed the presence of quinolizidine alkaloids (18, 19), flavonoides and steroid glucosides (12). *Sophora* alkaloids have been found to be the chief active component of the plant including matrine, oxymatrine, sophocarpine, sophoramine, and sophoridine. Basic and clinical researches have confirmed that these alkaloids possess different pharmacological effects like anti-inflammatory (20,21), immunity-regulatory (22), antiviral (23), antibacterial (24), and anti-tumor properties (25, 26). From the published data mentioned above genera and families, it is possible to predict the type of compounds that might be present in a particular extract. This tentative prediction on possible identity of the classes of compounds may help choose suitable extraction and partitioning methods and appropriate solvents for extracting specific classes of compounds (27,28). In this study, the uses of solvents for screening and the isolation of active components were examined.

Despite several reports on the anti-cancer properties of some species of the genus, *S. pachycarpa* has not yet been studied for its cytotoxic potential.

Cancer is considered as the leading cause of death in developed countries and the second leading cause of death in developing countries. In order to evaluate new compounds and extracts for potential *in-vitro* activity in the primary screening model, employing diverse panel of human tumor cell lines representing major tumor types is recommended. Evaluations may carry out on compounds over a wide range of concentrations for cytotoxic or growth-inhibitory effects against each cell line comprising the panel. Lung cancer

(1.4 million deaths) is the most common cause of death in patients suffering from cancer worldwide (29). A549 is a cell line derived from human alveolar cell carcinoma (30) and has been used as model of both human primary alveolar epithelial cells (31) and human primary alveolar epithelial cells *in-vitro* (32). Over the course of a lifetime, 1 in 8 women will be diagnosed with breast cancer. MCF-7 is a human breast cancer cell line, extensively used in the study of breast cancer (33). Cervical cancer is the third most common type of cancer in women (34). HeLa cells are human epithelial cells studied widely in cervical cancer, the second most frequent malignant tumor in women (35). Prostate cancer is the most common cause of death from cancer in men over age 75 (36). PC3 is the classic human cell line of prostatic cancer (37). This cell line was used in studies evaluating the response of chemotherapeutic drugs on prostate cancer cells (38). The third most common childhood cancers is leukemia (34%) (39). The HL-60 (Human promyelocytic leukemia cells) cell line is a leukemic cell line which is used to study the apoptosis effect of the extracts on cells (40).

Here we analyzed for the first time the cytotoxic effects of *S. pachycarpa* root extract on different cancer cell lines including A549, HeLa, HL-60, MCF-7, PC3 cells, and human umbilical cord Leukocyte as non-malignant cells. The role of apoptosis in *S. pachycarpa*-induced cytotoxicity on cancer cell lines was explored as well.

Experimental

Reagents and chemicals

The fluorescent probe propidium iodide (PI), and Triton X-100 were purchased from Sigma (St Louis, MO, USA), MTS from Promega (Madison, WI, USA), RNase A and Proteinase K from Fermentas (Vilnius, Lithuania); RPMI-1640 and FCS from Gibco (Grand Island, USA).

Plant materials

Roots of *S. pachycarpa* were collected in June 2010 at the Ferdowsi University Campus, Mashhad, Razavi Khorasan province, northeast of Iran.

The plant was identified by Mr. M. R.




Figure 1. Partitioning scheme using immiscible solvents.

Joharchi from Ferdowsi University of Mashhad Herbarium (FUMH). A voucher specimen (No. 06-019-016) is deposited in the herbarium of the School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.

The dried root (380 g) was percolated with methanol (MeOH) at room temperature. The whole extract was filtered and the solvent was evaporated under reduced pressure at 40-45°C, to afford crude methanol extract. Methanol extract was then resolved in MeOH:H₂O 95:5 and partitioned successively between *n*-hexane, dichloromethane (CH₂Cl₂), ethylacetate (EtOAc), *n*-butanol (*n*-BuOH), and finally H₂O. CH₂Cl₂, EtOAc, and *n*-BuOH fractions were evaporated under vacuum and H₂O fraction was freeze dried to 0.4, 5.2, 3.5, and 5.9 g north of

respectively. Fractions were stored at 4°C until analysis. A partitioning scheme of *S. pachycarpa* methanol extract is presented in Figure 1 (27).

Sample preparation

To prepare the stock solutions (100 mg/mL), each extract was dissolved in DMSO. The concentrations of 3.5-250 µg/mL were then obtained by diluting these solutions with Roswell Park Memorial Institute-1640 (RPMI-1640) so that the final concentrations of DMSO did not exceed 0.25%. All dilutions were prepared fresh before addition to the cells.

Cell culture and Treatment

A549, HeLa, HL-60, MCF-7, and PC3 cells were obtained from Pasteur Institute (Tehran,

Iran) and maintained at 37°C in a humidified atmosphere (90%) containing 5% CO₂. Cell lines were cultured in RPMI 1640 supplemented with 10% (v/v) fetal bovine serum, 100 U/mL penicillin and 100 mg/mL streptomycin. Cells were seeded overnight and then incubated with various concentrations of different extracts for 48 h.

For MTS assay, cells were seeded at 10⁴ cell per well onto 96-well culture plates. For assay of apoptosis, cells were seeded at 10⁵ cell per well onto a 24-well plate.

For each concentration and time course study, there was a control sample which remained untreated and received an equal volume of the solvent. Paclitaxel (41) was used as positive control (350 nM).

Cell viability

The MTS assay (42) is based on the reduction of MTS by mitochondrial dehydrogenases in metabolically active cells, to form the colored, water-soluble formazan. Cells were seeded in each well of a 96-microwell plate and treated with various concentrations of methanol extract and different fractions obtained from *S. pachycarpa* root. After incubation for 48 h, CellTiter 96® Aqueous One Solution Reagent (Promega, Madison, WI, USA), which is composed of the novel tetrazolium compound MTS and an electron coupling reagent phenazine methosulfate (PES, a redox intermediary), was added to each well according to the manufacturer's instructions. After 1 h, the cell viability was determined by measuring the absorbance at 490 nm using an ELISA microplate reader (Awareness, Palm City, FL, USA).

Cytotoxicity was expressed as IC₅₀, which was calculated using Graph Pad prism 5 software and presented as mean ± SEM of three independent experiments with three replicates for each concentration.

Paclitaxel was used as a positive control.

Leukocyte culture

Human umbilical cord blood samples (50 ml) were collected from a fresh umbilical cord attached to the placenta by gravity flow in sterile 50 ml syringe containing citrate buffer as an anticoagulant. The sample was diluted with an

equal volume of Phosphate Buffered Saline (PBS), layered over Ficoll-Hypaque solution according to density gradient (1.077 g/mL), and centrifuged at 800 g for 20 min at room temperature. The mononuclear cell layer was removed, washed twice in PBS and resuspended in RPMI 1640 medium supplemented with 10% (v/v) fetal bovine serum, 100 U/mL penicillin and 100 mg/mL streptomycin. Leukocytes (5×10⁴ cells per well) were treated with different concentrations of each fraction of *S. pachycarpa* in 96-well plates, for 48 h. This study protocol was approved by the ethical committee of Mashhad University of Medical Sciences.

Apoptosis

Apoptotic cells were detected using PI staining of treated cells followed by flow cytometry to detect a sub-G1 peak (43,44).

It has been reported that DNA fragmentation creates small fragments of DNA that can be eluted following incubation in a hypotonic phosphate citrate buffer. When stained with a quantitative DNA-binding dye such as PI, cells that have lost DNA will take up less stain and will appear to the left of the G1 peak. Briefly, HeLa cells were cultured overnight in a 24-well plate and treated with various concentrations of CH₂Cl₂ extract for 48 h. Floating cells were harvested and incubated at 4°C overnight in the dark with 750 µL of a hypotonic buffer (50 µg/mL PI in 0.1% sodium citrate + 0.1% Triton X-100) before flow cytometric analysis using a Partec flow cytometer (GmbH, Münster, Germany) was conducted. Ten thousand events were acquired.

Statistics

One-way analysis of variance (ANOVA) and Bonferroni's post hoc were used for data analysis. All results were expressed as mean ± SEM and *p-values* < 0.05 were considered statistically significant.

Results

Inhibition of cell viability

The total methanol extract of *S. pachycarpa* decreased cell viability in a concentration-dependent manner. The toxicity was first observed at a concentration lower than 15 µg/mL




Figure 2. Dose-dependent growth inhibition of malignant cell lines by total methanol extract (15-500 µg/mL) after 48 h. Viability was quantitated by MTS assay. The toxicity started at a concentration as little as 15 µg/mL. Results are mean ± SEM (n =9). * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ compared to control.

(Figure 2). IC_{50} values for total methanol extract of *S. pachycarpa* in each cell line are presented in Table 1.

In order to compare the cytotoxicity of solvent fractions of *S. pachycarpa*, another MTS assay was carried out for different concentrations (0-250 µg/mL). Among all the fractions tested, the CH_2Cl_2 and EtOAc fractions were found to be more effective than the other fractions of the plant extract. CH_2Cl_2 fraction had the maximum inhibitory effect on the proliferation of HL-60 cells (Figure 3). In contrast, the H_2O fraction did not show any anti-proliferative effect. CH_2Cl_2 and EtOAc fractions effectively inhibited the growth of A549, HeLa, HL-60, MCF-7, as well as the PC3 cells in a concentration-dependent manner after 48 h. The IC_{50} values of solvent fractions of *S. pachycarpa* against malignant and non malignant cell lines are presented in Table 2.

In comparison, the cytotoxic effect of CH_2Cl_2 fraction on normal leukocyte proliferation isolated from peripheral blood was minimal (Figure 3).

We used 350 nM of Paclitaxel as a positive

Table 1. Doses of total methanol extract of *S. pachycarpa* inducing 50% cell growth inhibition (IC_{50}) against malignant cell lines. Cells incubated with different concentration of extracts for 48 h.

	Cell line		
	HeLa	HL-60	MCF-7
IC_{50}	84.07	15.07	52.33

control. Paclitaxel at this concentration decreased the viability of HeLa cells to 25.25% ± 5.58 (data not shown).

Together, these data point to the selective activity of CH_2Cl_2 and EtOAc fractions against tumor cells.

Apoptosis induction by CH_2Cl_2 fraction of *S. pachycarpa*

HeLa cells as the most sensitive cells were chosen for further mechanistic studies. To confirm that the CH_2Cl_2 fraction of *S. pachycarpa* induces apoptosis, we evaluated the percentage of apoptotic cells by PI staining and flow cytometry, to observe sub-G1 peak caused by DNA fragmentation. CH_2Cl_2 fraction treated cells showed sub-G1 peak in HeLa cells. This indicates that CH_2Cl_2 fraction induces cell death through apoptosis (Figure 4).

Discussion

Plants as the main source of the potent phytochemicals have been used to prevent and treat diseases including cancer for years. In this study, the cytotoxic and proapoptotic effects of methanol extract and its different fractions obtained from *S. pachycarpa* root extract were investigated on A549, HeLa, HL-60, MCF-7, and PC3 cells. Leukocytes were used as non-malignant cells while Paclitaxel (350 nM) was used as a positive control.




Figure 3. Dose-dependent growth inhibition of malignant cell lines by CH₂Cl₂ fraction (3.5-250 µg/mL) after 48 h. Viability was quantitated by MTS assay. Results are mean ± SEM (n=9). **p* < 0.05, ***p* < 0.01 and ****p* < 0.001 compared to control.

Table 2. Doses of solvent fractions of *S. pachycarpa* inducing 50% cell growth inhibition (IC₅₀) against malignant and non malignant cell lines. Cells incubated with different concentration of extracts for 48 h.

Cell line	Fraction			
	CH ₂ Cl ₂	EtOAc	<i>n</i> -butanol	H ₂ O
A549	31.15	40.53	189.3	>250
HeLa	14.54	26.62	86.20	>250
HL-60	10.52	6.651	35.29	>250
PC-3	43.22	233.2	72.17	>250
MCF-7	27.09	42.08	206.7	>250
Leukocyte (normal cells)	87.29	25.16	55.78	>250




Figure 4. Flow cytometry histograms of apoptosis assays by PI method in HeLa cells. Cells were treated with different concentration of CH_2Cl_2 fraction for 48 h. Sub-G1 peak as an indicative of apoptotic cells, was induced in CH_2Cl_2 fraction treated but not in control cells. CH_2Cl_2 fraction-treated cells exhibited a sub-G1 peak in HeLa cells in a concentration dependent manner that indicates the involvement of an apoptotic process in CH_2Cl_2 fraction-induced cell death.

To the authors' knowledge, this is the first report on *S. pachycarpa* induced cytotoxicity in cancer cell lines. Our data confirmed that *S. pachycarpa* extract has cytotoxic effect on various tumor cell lines tested.

Using MTS assay, the anti-proliferative effects of methanol extract and fractions of *S. pachycarpa* root extract on different cancer cell lines were compared. The effects of extracts and isolated fractions were not all anti-proliferative. Data showed that the CH_2Cl_2 fraction of *S. pachycarpa* has a significant cell growth inhibitory effect.

Similarly, a study to find bioactive compounds from plants, the CH_2Cl_2 fraction of the MeOH extract of the root of *S. flavescens*, showed significant cytotoxicity against human myeloid leukemia HL-60 cells (14,45).

Methanol extracts of *S. alopecuroides* and *S. pachycarpa* showed the LC_{50} values of 73.11 ± 1.20 and 56.73 ± 0.55 ($\mu\text{g}/\text{mL}$) in Brine shrimp lethality assay (46).

Herbal extract contains many components; different compounds may have opposing pharmacological activities. To ensure safe use of herbal medicine, these compounds with opposing effects should be removed from the extract.

In the present study, the purification by solvent extraction of *S. pachycarpa* was used and the potential antitumor activity of low-polar solvent fraction (CH_2Cl_2 , EtOAc) was compared to polar solvent fractions (*n*-BuOH and H_2O soluble).

The presence of compounds with synergistic or antagonistic interactions in crude extract may cause varying level of activity in total extract of the plant compared to its fractions. Differences in the activity of polar and non-polar fractions might be due to the presence of diverse constituents. The maximum activity of low-polar solvent fraction (CH_2Cl_2 , EtOAc) can serve to trace the highly active phytochemicals. Sharing various phytochemicals in different fractions might be responsible for the change in the activity either due to synergistic or antagonistic effects.

The chemo-physical properties of different compound in the extract can be exploited to initially divide them into various chemical groups using solvents of increasing polarity.

The CH_2Cl_2 fraction as the most cytotoxic fraction of *S. pachycarpa* extract was further analysed to establish its apoptotic activity in the HeLa and HL-60 cell lines.

Treatment of tumors is directed not only

on inhibition of cell proliferation, but also on induction of apoptosis of tumor cells.

It is well known that the earliest recognized morphological change in apoptosis is compaction and segregation of the nuclear chromatin, resulting in chromatin margination and cytoplasmic condensation. Condensation is accompanied by sinking followed by disintegration of the nucleus into discrete fragments, a characteristic feature of apoptosis involving endonucleases (47).

This characteristic oligonucleosomal DNA degradation (also known as low molecular weight (LMW) cleavage) and chromatin condensation are among the best known events of apoptotic execution and were initially assumed to be an essential part of the apoptotic pathway (48).

The CH₂Cl₂ fraction of *S. pachycarpa* possesses the ability to reduce the viability of HeLa cells via apoptosis induction confirmed by sub-G1 peak in the flow cytometry histogram of treated cells compared to control cells.

Our results showed that CH₂Cl₂ fraction of *S. pachycarpa* inhibits cell growth via apoptosis. On the basis of these findings, we predict that the CH₂Cl₂ and EtOAc fractions of *S. pachycarpa* would be good candidates to search for novel bioactive components and their application as anti-cancer potential.

Collectively, further studies are required to evaluate and understand the underlying molecular mechanisms, pathways involved and to identify active phyto-chemicals before *S. pachycarpa* being considered as a potential therapeutic agent against cancer.

Acknowledgements

The authors would like to thank Dr. H. Nasirli for her assistance in flow cytometry. This work was supported by grants (No. 89308) from Research Affairs of Mashhad University of Medical Sciences as a part of MD thesis.

References

- (1) Harvey A. Strategies for discovering drugs from previously unexplored natural products. *Drug Discov. Today* (2000) 5: 294-300.
- (2) Newman DJ, Cragg GM and Snader KM. The influence of natural products upon drug discovery. *Nat. Prod. Rep.* (2000) 172: 15-34.
- (3) Kintzios E. Terrestrial plant-derived anticancer agents and plant species used in anticancer research. *Crit. Rev. Plant Sci.* (2006) 25: 79-113.
- (4) Rechinger KH, Ali SI, Browicz K, Chrtkova-Zertova P, Heller D, Heyn CC, Thuling M and Vassilcznko IT. Flora Iranica No. 157. Papilionaceae. Akademische Druk-u. (1984) 15-20.
- (5) Ruangrunsi N, Iinuma M, Tanaka T, Ohyama M, Yokoyama J and Mizuno M. Three flavanones with a lavandulyl group in the roots of *Sophora exigua*. *Phytochem.* (1992) 35: 999-1000.
- (6) Iinuma M, Ohyama M, Kawasaka Y and Tanaka T. Flavonoid compounds in roots of *Sophora tetraptera*. *Phytochem.* (1995) 39: 667-672.
- (7) Iinuma M, Ohyama M, Tanaka T. Flavonoids in roots of *Sophora prostrata*. *Phytochem.* (1995) 38: 539-543.
- (8) Iinuma M, Ohyama M and Tanaka T. Six flavonostilbenes and a flavanone in roots of *Sophora alopecuroides*. *Phytochem.* (1995) 38: 519-525.
- (9) Iinuma M, Ohyama M, Tanaka T, Shirataki Y and Burandt CL. Isoflavonoids in roots of *Sophora secundiflora*. *Phytochem.* (1995) 39: 907-910.
- (10) Tanaka T, Iinuma M, Asai F, Ohyama M and Burandt CL. Flavonoids from the root and stem of *Sophora tomentosa*. *Phytochem.* (1997) 46: 1431-1437.
- (11) Tanaka T, Ohyama M, Iinuma M, Shirataki Y, Komatsu M and Burandt CL. Isoflavonoids from *Sophora secundiflora*, *S. arizonica* and *S. gypsophila*. *Phytochem.* (1998) 48: 1187-1193.
- (12) Emami SA, Amin-Ar-Ramimeh E, Ahi A, Bolourian-Kashy MR, Schneider B and Iranshahi M. Prenylated flavonoids and flavonostilbenes from *Sophora pachycarpa* Roots. *Pharm. Biol.* (2007) 45: 453-457.
- (13) Choudhary MI, Parvez K, Ahmed A and Akhtar F. Quinolizidine alkaloids from *Sophora alopecuroides*. *J. Nat. Prod.* (2000) 63: 190-192.
- (14) Izaddoost M. Alkaloid chemotaxonomy of the genus *Sophora*. *Phytochem.* (1975) 14: 203-204.
- (15) Kadooka, MM, Chang MY, Clardy J and Solheim BA. Mamanine and pohakuline. Two unprecedented quinolizidine alkaloids from *Sophora chrysophylla*. *Tetrahedron* (1979) 32: 919-924.
- (16) Keller WJ. Alkaloids from *Sophora secundiflora*. *Phytochem.* (1975) 14: 2305-2306.
- (17) Murakoshi I, Ito M, Haginiwa J, Ohmiya S, Otomasu H and Hirano RT. Lupine alkaloid from leaves of *Sophora secundiflora*. *Phytochem.* (1984) 23: 887-891.
- (18) Abdusalamov BA. Biosynthesis and metabolism of some matrine alkaloids in goebel. *Chem. Nat. Comp.* (1984) 20: 1-9.
- (19) Abdusalamov BA, Aslanov A and Sadycov AS. The biosynthesis of the alkaloids of *Goebelia pachycarpa*. *Chem. Nat. Comp.* (1997) 13: 458-460.
- (20) Jiang H, Hou CH, Zhang SB, Xie HY, Zhou WY, Jin QH, Cheng XD, Qian RL and Zhang XJ. Matrine upregulates the cell cycle protein E2F-1 and triggers apoptosis via the mitochondrial pathway in K562 cells. *Eur. J. Pharmacol.* (2007) 559, 98-108.
- (21) Liu SX and Chiou GC. Effects of Chinese herbal

- products on mammalian retinal functions. *J. Ocul. Pharmacol. Ther.* (1996) 12: 377-86.
- (22) Zhang JP, Zhang M, Jin C, Zhou B, Xie WF, Guo C, Zhang C and Qian DH. Matrine inhibits production and actions of fibrogenic cytokines released by mouse peritoneal macrophages. *Acta Pharmacol. Sin.* (2001) 22: 765-768.
- (23) Liu J, Zhu M, Shi R and Yang M. Radix *Sophorae flavescens* for chronic hepatitis B: a systematic review of randomized trials. *Am. J. Chin. Med.* (2003) 35: 337-354.
- (24) Fakhimi A, Iranshahi M, Emami SA, Amin-Ar-Ramimeh E, Zarrini G and Shahverdi AR. Sophoraflavanone G from *Sophora pachycarpa* enhanced the antibacterial activity of gentamycin against *Staphylococcus aureus*. *Z Naturforsch. C.* (2006) 65: 769-72.
- (25) MJ Zhang and J Huang. [Recent research progress of anti-tumor mechanism matrine]. *Zhongguo Zhong Yao Za Zhi.* (2004) 29: 115-118.
- (26) Ma L, Wen S, Zhan Y, He Y, Liu X and Jiang J. Anticancer effects of the Chinese medicine matrine on murine hepatocellular carcinoma cells. *Planta Med.* (2008) 74: 245-51
- (27) Otsuka H, Sarker SD, Latif Z and Gray AL. Purification by solvent extraction using partition coefficient. In: (eds) Natural Products Isolation. 2nd edn. Totowa New Jersey Humana Press. (2006) 269-273.
- (28) Tayarani-Najaran Z, Emami SA, Asili J, Mirzaei A and Mousavi SH. Analyzing cytotoxic and apoptogenic properties of *Scutellaria litwinowii* root extract on cancer cell lines. *Evid. Based Complement. Alternat. Med.* (2011) 2015: 1-9.
- (29) WHO (October 2010). «Cancer». World Health Organization. Retrieved 5 January 2011.
- (30) Lieber M, Smith B, Szakal A, Nelson-Rees W and Todaro GA. Continuous tumor-cell line from a human lung carcinoma with properties of type II alveolar epithelial cells. *Int. J. Cancer* (1976) 17: 62-70.
- (31) Shin S, Cha HJ, Lee EM, Lee SJ, Seo SK, Jin HO, Park IC, Jin YW and An S. Alteration of mRNA profiles by ionizing radiation in A549 human non-small cell lung cancer cells. *Int. J. Oncol.* (2009) 35: 81-86.
- (32) Mazzarella G, Ferraraccio F, Prati MV, Annunziata S, Bianco A, Mezzogiorno A, Liguori G, Angelillo IF and Cazzola M. Effects of diesel exhaust particles on human lung epithelial cells an *in-vitro* study. *Respir. Med.* (2007) 105: 1155-1162.
- (33) Simstein R, Burow M, Parker A, Weldon CA and Beckman B. Apoptosis, chemoresistance, and breast cancer: insights from the MCF-7 cell model system. *Exp. Biol. Med.* (2003) 228: 995-1003.
- (34) National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Cervical cancer. Version 2. 2013. Available from: URL: http://www.nccn.org/professionals/physician_gls/pdf/cervical.pdf. Accessed November (2012) 16.
- (35) Tavakol-Afshari J, Brook A and Mousavi SH. Study of cytotoxic and apoptogenic properties of saffron extract in human cancer cell lines. *Food Chem. Toxicol.* (2008) 463: 443-447.
- (36) Antonarakis ES and Eisenberger MA. Expanding treatment options for metastatic prostate cancer. *N. Engl. J. Med.* (2011) 364: 2055-2058
- (37) Alimirah F, Chen J, Basrawala Z, Xin H and Choubey D. «DU-145 and PC-3 human prostate cancer cell lines express androgen receptor: implications for the androgen receptor functions and regulation». *FEBS Lett.* (2006) 580: 2294-2300.
- (38) Kaighn ME, Narayan KS, Ohnuki Y, Lechner JF and Jones LW. Establishment and characterization of a human prostatic carcinoma cell line (PC-3). *Invest. Urol.* (1979) 17: 16-23.
- (39) Kaatsch P. Epidemiology of childhood cancer. *Cancer Treat. Rev.* (2010) 36: 277-85.
- (40) Sugimoto K, Yamada K, Egashira M, Yazaki Y, Hirai H, Kikuchi A and Oshimi K. «Temporal and spatial distribution of DNA topoisomerase II alters during proliferation, differentiation and apoptosis in HL-60 cells». *Blood* (1998) 95: 1407-1417.
- (41) Lee JI, Ha YW, Choi TW, Kim HJ, Kim SM, Jang HJ, Choi JH, Choi MH, Chung BC, Sethi G, Kim SH, Ahn KS, Choi SH, Shim BS and Ahn KS. Cellular uptake of ginsenosides in Korean white ginseng and red ginseng and their apoptotic activities in human breast cancer cells. *Planta Med.* (2011) 77: 133-140.
- (42) Malich G, Markovic B and Winder C. The sensitivity and specificity of the MTS tetrazolium assay for detecting the *in-vitro* cytotoxicity of 20 chemicals using human cell lines. *Toxicol.* (1997) 124: 179-192.
- (43) Nicoletti G, Migliorati MC, Pagliacci F, Grignani F and Riccardi C. rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. *J. Immunol. Methods.* (1991) 139: 271-279.
- (44) Parsaee H, Asili J, Mousavi SH, Soof H, Emami SA and Tayarani-Najaran Z. Apoptosis induction of *salvia chorassanica* root extract on human cervical cancer cell line. *Iran. J. Pharm. Res.* (2013) 12: 75-83.
- (45) Ko WG, Kang TH, Kim NY, Lee SJ, Kim YC, Ko GI, Ryu SY, Lee BH. Lavandulylflavonoids: a new class of *in vitro* apoptogenic agents from *Sophora flavescens*. *Toxicol. In-vitro.* (2000) 14: 429-433.
- (46) Khalighi-Sigaroodi F, Ahvazi M, Hadjiakhoondi A, Taghizadeh M, Yazdani D, Khalighi-Sigaroodi S and Bidel S. Cytotoxicity and antioxidant activity of 23 plant species of Leguminosae family. *Iran. J. Pharm. Res.* (2012) 11: 295-302.
- (47) Wyllie AH. Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature* (1980) 284: 555-556.
- (48) Samejima K and Earnshaw WC. Trashing the genome: the role of nucleases during apoptosis. *Nat. Rev. Mol. Cell Biol.* (2005) 6: 677-688.

Tell us if we are wrong?
Visit <http://www.ijpr.ir>
or
[http:// ijpr.sbm.ac.ir](http://ijpr.sbm.ac.ir)