

Published in final edited form as:

Cochrane Database Syst Rev. ; (6): CD004718. doi:10.1002/14651858.CD004718.pub3.

Early intervention for psychosis

Max Marshall¹ and John Rathbone²

¹University of Manchester, The Lantern Centre, Preston., UK

²HEDS, SchARR, The University of Sheffield, Sheffield, UK

Abstract

Background—Proponents of early intervention have argued that outcomes might be improved if more therapeutic efforts were focused on the early stages of schizophrenia or on people with prodromal symptoms. Early intervention in schizophrenia has two elements that are distinct from standard care: early detection, and phase-specific treatment (phase-specific treatment is a psychological, social or physical treatment developed, or modified, specifically for use with people at an early stage of the illness).

Early detection and phase-specific treatment may both be offered as supplements to standard care, or may be provided through a specialised early intervention team. Early intervention is now well established as a therapeutic approach in America, Europe and Australasia.

Objectives—To evaluate the effects of: (a) early detection; (b) phase-specific treatments; and (c) specialised early intervention teams in the treatment of people with prodromal symptoms or first-episode psychosis.

Search methods—We searched the Cochrane Schizophrenia Group Trials Register (March 2009), inspected reference lists of all identified trials and reviews and contacted experts in the field.

Selection criteria—We included all randomised controlled trials (RCTs) designed to prevent progression to psychosis in people showing prodromal symptoms, or to improve outcome for people with first-episode psychosis. Eligible interventions, alone and in combination, included: early detection, phase-specific treatments, and care from specialised early intervention teams. We accepted cluster-randomised trials but excluded non-randomised trials.

Copyright © 2011 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd

Contact address: Max Marshall, University of Manchester, The Lantern Centre, Vicarage Lane, Of Watling Street Road, Fulwood, Preston., Lancashire, UK. max.marshall@manchester.ac.uk. max.marshall@lancashirecare.nhs.uk.

CONTRIBUTIONS OF AUTHORS Max Marshall - designed the review, developed the search strategy, screened the search results, appraised the papers and extracted data, analysed and interpreted the data and wrote the final report.

John Rathbone - (2006/2011 update) screened the search results, appraised the papers and extracted data, analysed and interpreted the data and helped write the final report.

Editorial group: Cochrane Schizophrenia Group.

Publication status and date: New search for studies and content updated (conclusions changed), published in Issue 6, 2011.

Review content assessed as up-to-date: 3 June 2009.

DECLARATIONS OF INTEREST Max Marshall - received funding for the review from the UK Department of Health, which is committed to a policy of implementing Early Intervention teams across England and are in the early stages of developing a fidelity scale for early intervention teams. Max Marshall is Clinical Director of the Lancashire Early Intervention Service.

John Rathbone - none.

Data collection and analysis—We reliably selected studies, quality rated them and extracted data. For dichotomous data, we estimated relative risks (RR), with the 95% confidence intervals (CI). Where possible, we calculated the number needed to treat/harm statistic (NNT/H) and used intention-to-treat analysis (ITT).

Main results—Studies were diverse, mostly small, undertaken by pioneering researchers and with many methodological limitations (18 RCTs, total n=1808). Mostly, meta-analyses were inappropriate. For the six studies addressing prevention of psychosis for people with prodromal symptoms, olanzapine seemed of little benefit (n=60, 1 RCT, RR conversion to psychosis 0.58 CI 0.3 to 1.2), and cognitive behavioural therapy (CBT) equally so (n=60, 1 RCT, RR conversion to psychosis 0.50 CI 0.2 to 1.7). A risperidone plus CBT plus specialised team did have benefit over specialist team alone at six months (n=59, 1 RCT, RR conversion to psychosis 0.27 CI 0.1 to 0.9, NNT 4 CI 2 to 20), but this was not seen by 12 months (n=59, 1 RCT, RR 0.54 CI 0.2 to 1.3). Omega 3 fatty acids (EPA) had advantage over placebo (n=76, 1 RCT, RR transition to psychosis 0.13 CI 0.02 to 1.0, NNT 6 CI 5 to 96). We know of no replications of this finding.

The remaining trials aimed to improve outcome in first-episode psychosis. Phase-specific CBT for suicidality seemed to have little effect, but the single study was small (n=56, 1 RCT, RR suicide 0.81 CI 0.05 to 12.26). Family therapy plus a specialised team in the Netherlands did not clearly affect relapse (n=76, RR 1.05 CI 0.4 to 3.0), but without the specialised team in China it may (n=83, 1 RCT, RR admitted to hospital 0.28 CI 0.1 to 0.6, NNT 3 CI 2 to 6). The largest and highest quality study compared specialised team with standard care. Leaving the study early was reduced (n=547, 1 RCT, RR 0.59 CI 0.4 to 0.8, NNT 9 CI 6 to 18) and compliance with treatment improved (n=507, RR stopped treatment 0.20 CI 0.1 to 0.4, NNT 9 CI 8 to 12). The mean number of days spent in hospital at one year were not significantly different (n=507, WMD, -1.39 CI -2.8 to 0.1), neither were data for 'Not hospitalised' by five years (n=547, RR 1.05 CI 0.90 to 1.2). There were no significant differences in numbers 'not living independently' by one year (n=507, RR 0.55 CI 0.3 to 1.2). At five years significantly fewer participants in the treatment group were 'not living independently' (n=547, RR 0.42 CI 0.21 to 0.8, NNT 19 CI 14 to 62). When phase-specific treatment (CBT) was compared with befriending no significant differences emerged in the number of participants being hospitalised over the 12 months (n=62, 1 RCT, RR 1.08 CI 0.59 to 1.99).

Phase-specific treatment E-EPA oils suggested no benefit (n=80, 1 RCT, RR no response 0.90 CI 0.6 to 1.4) as did phase-specific treatment brief intervention (n=106, 1 RCT, RR admission 0.86 CI 0.4 to 1.7). Phase-specific ACE found no benefit but participants given vocational intervention were more likely to be employed (n=41, 1 RCT, RR 0.39 CI 0.21 to 0.7, NNT 2 CI 2 to 4). Phase-specific cannabis and psychosis therapy did not show benefit (n=47, RR cannabis use 1.30 CI 0.8 to 2.2) and crisis assessment did not reduce hospitalisation (n=98, RR 0.85 CI 0.6 to 1.3). Weight was unaffected by early behavioural intervention.

Authors' conclusions—There is emerging, but as yet inconclusive evidence, to suggest that people in the prodrome of psychosis can be helped by some interventions. There is some support for specialised early intervention services, but further trials would be desirable, and there is a question of whether gains are maintained. There is some support for phase-specific treatment focused on employment and family therapy, but again, this needs replicating with larger and longer trials.

Medical Subject Headings (MeSH)

*Psychotic Disorders [diagnosis; therapy]; *Schizophrenia [diagnosis; therapy]; Cognitive Therapy; Early Diagnosis; Randomized Controlled Trials as Topic; Suicidal Ideation; Time Factors

MeSH check words

Humans

BACKGROUND

Schizophrenia and other functional psychoses cause enormous suffering for individuals and their families, and are a financial burden to the NHS and other health services. The estimated total cost of schizophrenia in England was £6.7 billion in 2004/05; the direct cost of treatment and care was £2 billion, whilst the indirect cost to society was £4.7 billion, and the cost of informal care and private expenditure was £615 million (Mangalore 2007). Despite new medications and the development of community care, about one-third of people with schizophrenia have a poor long-term outcome (Mason 1997). An overview of studies investigating outcomes has shown that people with schizophrenia have a one-year relapse rate of 15% to 35%, rising to 80% within five years (Larsen 1998). Achievement of full remission becomes less likely after each relapse, and about 10% of sufferers eventually commit suicide (Wiersma 1998).

Description of the condition

Schizophrenia is a chronic, relapsing mental illness and has a worldwide lifetime prevalence of about 1% irrespective of culture, social class and race. Schizophrenia is characterised by positive symptoms such as hallucinations, delusions and jumbled thinking; and negative symptoms such as apathy, poverty of speech, and withdrawal from social activities.

Description of the intervention

Early intervention in psychosis has two elements that are distinct from standard care: early detection and phase-specific treatment. Early detection may be defined as either the identification of people thought likely to develop psychosis (i.e. those who display prodromal symptoms, but have never been psychotic (Schaffner 2001)) or the identification of people who are already psychotic, but have not yet received adequate treatment (Wyatt 2001).

Phase-specific treatments are defined as treatments (psychological, social or physical) that are especially targeted at people in the prodrome or early stages of schizophrenia (Miller 1999). Phase-specific treatments may be directed at preventing progression to psychosis (in people with prodromal symptoms), or at promoting recovery (in people who have recently experienced their first episode of psychosis).

Early detection and phase-specific treatments may be provided as supplements to standard psychiatric care, or they may be provided by means of a specialised early intervention team

(Garety 2000). Such teams provide care exclusively to people who have prodromal symptoms or are in early stages of schizophrenia (Edwards 2000). Prodromal patients are usually assessed by the attenuated psychotic symptom criteria, using either the criteria by Yung 2005 or the Scale of Psychotic Symptoms (SOPS, Miller 1999). A second method, is the detection of ‘basic symptoms’ developed in Germany (Schultze-Lutter 2007). When people are referred to as ‘ultra high risk’ they are using the Yung 2005 criteria. When they are referred to as early or late initial prodrome state, they are using basic symptoms.

How the intervention might work

Until recently, the orthodox approach to treating schizophrenia was to concentrate therapeutic resources on those people who developed severe and chronic disabilities (McGorry 1999). This approach has been challenged by proponents of early intervention, who have argued that greater investment of resources in the early stages of the disorder might substantially reduce the numbers of people developing chronic disabilities (Wyatt 1991). This argument has been strengthened by the observation that there may be an association between various outcome parameters and the duration of untreated psychosis (the time from the development of the first psychotic symptom to the receipt of adequate drug treatment) (Norman 2001). This has led to the proposition that untreated psychosis may be ‘toxic’ and that early intervention might prevent irreversible harm (Sheitman 1998).

Why it is important to do this review

The arguments in favour of early intervention have been so persuasive that early intervention teams are well-established in America, Europe and Australasia (Edwards 2002). In 2000, the UK government announced its intention to set up 50 early intervention teams in England to provide specialised care to all young people with a first episode of psychosis (DoH 2000). It remains unclear, however, how far these service developments are underpinned by evidence of effectiveness. There is particular concern over the ethics of early intervention with prodromal patients, when the benefits of early detection and treatment are unclear, and there is no certainty that they will go on to develop psychosis (Rosen 2000).

OBJECTIVES

To evaluate the effects of early intervention in the treatment of early psychosis.

The two specific objectives were to determine the following.

1. The effects of early detection and treatment of people with ‘prodromal’ symptoms, in terms of:
 - 1.1 prevention of progression to full-blown psychosis;
 - 1.2 clinical and social outcomes;
 - 1.3 process variables and costs.

2. To determine the effects of early detection and treatment of people in their first episode of psychosis, in terms of:

- 2.1 clinical and social outcomes;
- 2.2 prevention of relapse;
- 2.3 process variables and costs;
- 2.4 reduction in duration of untreated psychosis.

We defined ‘Treatment’ as including both phase-specific treatments and care from a specialised early intervention team. We are not concerned with evaluating the accuracy of methods of predicting who is likely to develop psychosis.

METHODS

Criteria for considering studies for this review

Types of studies—We included studies if they were randomised controlled trials (RCTs). We accepted cluster-randomised trials and listed non-randomised trials in the Characteristics of excluded studies table.

In broad terms we have included two types of trial in this review.

1. Trials to prevent the development of psychosis: These studies involved treatments and/or methods of management that are given to people who are believed to be showing prodromal (pre-psychotic) symptoms and are therefore considered at high risk of developing psychosis. The primary aim of such studies was to prevent progression to psychosis, and invariably the interventions they offered were combined with some method of early detection of people at risk.

2. Trials to improve outcome in first-episode psychosis: These studies involved treatments and/or methods of management designed for people in the early stages of psychosis. The primary aim of such studies was to improve the long term outcome. Early detection might be offered in addition to the treatments, with the aim of ensuring that the treatment was offered as early as possible after the onset of psychosis.

Types of participants—1. For trials to prevent the development of psychosis, we included people who were judged by the trialists to be in a prodromal phase of psychosis, on the basis of showing prodromal symptoms (however defined).

2. For trials to improve outcome in first-episode psychosis, we included people who were in their first episode of psychosis, or were in the process of recovering from their first episode. People with psychosis were defined as those presenting with any combination of delusions, hallucinations or thought disorder, or those who had been given a diagnosis of schizophrenia or schizophrenia-like disorder, bipolar disorder (manic episode i.e. with psychotic symptoms), or depression with psychotic features.

We excluded trials where the majority of participants were suffering from a learning disability or an organic psychosis. We did not exclude anyone for reasons such as age or type of psychosis (for example, affective psychosis). Where studies included both first and second episode participants, we excluded trials if more than 10% of the participants included in the study had experienced a second episode,

Types of interventions—In trials of early intervention there are many possible combinations of intervention and control condition. This depends on: the type of participant (prodromal or first episode); whether the trial involved early detection (which could involve the whole sample or just the treatment group); the type of intervention (phase-specific or specialised team); the nature of any phase-specific treatment (cognitive therapy, family therapy etc); and the type of control (no treatment, standard psychiatric care, care from a specialised team but not phase specific intervention, etc.). In this section the most likely combinations of intervention and control conditions are listed for trials to prevent the development of psychosis and trials to improve outcome in first-episode psychosis.

1. Trials to prevent the development of psychosis in prodromal patients: These trials require prodromal patients, and since such patients do not normally present to psychiatric services, the trials therefore require some form of early detection to be applied to the whole sample. The intervention may consist of: phase-specific treatment (medication, psychological treatment or other) or care from a specialised team (which might offer phase-specific treatments). The control condition may consist of no treatment, or standard psychiatric care, or care from a specialised team (in which case the intervention will consist of care from a specialised team plus a phase specific intervention). The various types of intervention and control condition are described in more detail below.

1.1 Phase-specific treatment: In the context of preventing psychosis, phase-specific treatments are discrete interventions including medication regimes, which have been specifically developed for use in patients experiencing prodromal symptoms. A phase-specific treatment could be offered by an individual therapist or provided in the context of receiving care from a specialised team (see 1.2 below). More than one phase-specific treatment might be offered at the same time (for example, medication regime and cognitive therapy).

1.2 Care from a specialised team: In the context of preventing psychosis, this is defined as a multi-disciplinary psychiatric team, specialising in the treatment of patients with prodromal symptoms. Such a team would normally provide comprehensive psychiatric care to its patients and would be an alternative, rather than an addition, to standard psychiatric care. In the context of a trial it is likely that any specialised team would also offer phase-specific interventions.

1.3 Control conditions: In the context of preventing psychosis, the common control conditions are: no treatment; non-specific supportive therapy or care from a specialised team (which did not offer phase-specific treatments to prevent onset of psychosis).

2. Trials to improve outcome in first-episode psychosis: The intervention may consist of: early detection; phase-specific treatments (medication, psychological intervention or other) or care from a specialised team (which might offer phase-specific treatments). The control condition may consist of standard psychiatric care or care from a specialised team (in which case the intervention will consist of care from a specialised team plus a phase-specific intervention). A 'no treatment' control group is not an ethically acceptable option in first-episode psychosis trials. The various types of intervention and control condition are described in more detail below.

2.1 Early detection: In trials to improve outcome in first-episode psychosis it is possible to use early detection as an intervention applied to the treatment group alone; this is in contrast to the situation in trials designed to prevent psychosis (see 1. above) where early detection must be applied to both treatment and control groups. The theoretical basis for using early detection as an intervention is that shortening the duration of untreated psychosis in itself improves outcome. In trials where early detection is the intervention being tested, the unit of randomisation must be a cluster (for example, general practices or catchment areas), since it is not possible to individually randomise patients who have not yet been diagnosed.

2.2 Phase-specific treatment: In the context of improving outcome in the first episode, phase-specific treatments are discrete treatments and include medication regimes which have been specifically developed for use in the early stages of psychosis. A phase-specific treatment can be offered by an individual therapist or provided in the context of receiving care from a specialised team (see 2.3 below). More than one phase-specific treatment might be offered at the same time (for example, medication regime and cognitive therapy).

2.3 Care from a specialised team: In the context of improving outcome in first episode, this is defined as a multi-disciplinary psychiatric team, specialising in the treatment of patients with first-episode psychosis. Such a team would normally provide comprehensive psychiatric care to its patients and is an alternative, rather than an addition, to standard psychiatric care. In the context of a trial, it is likely that any specialised team would also offer phase-specific interventions.

2.4 Control conditions: In the context of improving outcome in first episode, the common control conditions are standard care, or care from a specialised team (which does not offer the phase-specific treatment being provided in the treatment arm of the trial). Standard care would be the normal service for people with severe psychiatric illness in the region where the trial took place, and would normally consist of out-patient follow up, medication, and support from a community mental health team, but would not involve any phase-specific treatment or specialised team.

3. Excluded interventions: We considered treatment with low doses of neuroleptic medication (atypical or standard) a phase-specific treatment if given to prevent progression to psychosis, or in the context of a medication protocol designed specifically for treating patients in their first episode of psychosis. However, simple comparisons of atypical neuroleptic medication versus standard neuroleptics in first-episode patients were beyond the scope of this review.

Types of outcome measures

Primary outcomes: For trials to prevent the development of psychosis (i.e. prodromal participants) the primary outcomes were as follows.

1. General

1.1 Converting to psychosis during follow-up period: For trials to improve the outcome of first-episode psychosis the outcomes were as follows.

1. General

Secondary outcomes: For trials to prevent the development of psychosis (i.e. prodromal participants) the secondary outcomes were as follows.

1. General: 1.1 Overall functioning

1.2 Duration of hospital stay

1.3 Loss to follow up

1.4 Satisfaction with treatment - participant/carer

1.5 Remaining in contact

2. Mental state: 2.1 General symptoms

2.2 Specific symptoms

2.2.1 Positive symptoms (delusions, hallucinations, disordered thinking)

2.2.2 Negative symptoms (avolition, poor self-care, blunted affect)

2.2.3 Mood - depression

3. Behaviour: 3.1 General behaviour

3.2 Specific behaviours (for example, aggressive or violent behaviour)

3.2.1 Social functioning

3.2.2 Employment status during trial (employed/unemployed)

3.2.3 Occurrence of violent incidents (to self, others or property)

4. Adverse effects: 4.1 General

4.2 Specific

4.2.1 Death (suicide and non-suicide)

4.2.2 Movement disorders (extrapyramidal side effects, specifically tardive dyskinesia and neuroleptic malignant syndrome)

4.2.3 Sedation

4.2.4 Dry mouth

4.2.5 Weight gain

5. Economic: 5.1 Cost of care

6. Quality of life: 6.1 No substantial improvement in quality of life

For trials to improve the outcome of first-episode psychosis the secondary outcomes were:

1. General: 1.1 Overall functioning

1.2 Hospital readmission

1.3 Duration of hospital stay

1.4 Loss to follow-up

1.5 Satisfaction with treatment - participant/carer

1.6 Remaining in contact with services

2. Mental state: 2.1 General symptoms

2.2 Specific symptoms

2.2.1 Positive symptoms (delusions, hallucinations, disordered thinking)

2.2.2 Negative symptoms (avolition, poor self-care, blunted affect)

2.2.3 Mood - depression

3. Behaviour: 3.1 General behaviour

3.2 Specific behaviours (for example, aggressive or violent behaviour)

3.2.1 Social functioning

3.2.2 Employment status during trial (employed/unemployed)

3.2.3 Occurrence of violent incidents (to self, others or property)

4. Adverse effects: 4.1 General

4.2 Specific

4.2.1 Death (suicide and non-suicide)

4.2.2 Movement disorders (extrapyramidal side-effects, specifically tardive dyskinesia and neuroleptic malignant syndrome)

4.2.3 Sedation

4.2.4 Dry mouth

4.2.5 Weight gain

5. *Economic*: 5.1 Cost of care

Search methods for identification of studies

We applied no language restrictions within the limitations of the search.

Electronic searches

1. Cochrane Schizophrenia Group Trials Register (March 2009): The register was searched using the phrase:

[early* in title, abstract or keywords of REFERENCE] or [Early* in intervention or prodromal* or early** in Health Care Condition of STUDY]

This register is compiled by systematic searches of major databases, hand searches and conference proceedings (see Group Module).

2. Previous searches for earlier versions of this review: Please see Appendix 1.

Searching other resources

1. Reference lists: We inspected reference lists of all identified trials and reviews for additional trials.

2. Personal contact: We contacted experts in the field within the European First Episode Network (2003) to identify unpublished trials.

Data collection and analysis

Selection of studies—We (MM and AL) searched The Cochrane Schizophrenia Group's register. Working independently we examined the papers identified from the search strategy. We discarded obviously irrelevant publications and retained only those in which some form of early intervention had been compared against a control treatment, and obtained copies of papers relating to relevant trials. Once we had obtained these papers, we decided whether the trials were eligible. We resolved any disagreements by discussion. For the 2006 update we (MM and JR) independently inspected citations. Where disagreement occurred, we sought to resolve this by discussion, or where doubt remained, we acquired the full article for further inspection. Once we had obtained the full articles, we independently decided whether they met the review criteria. We resolved any disagreements that occurred by

discussion, and when this was not possible we added trials to the list of those awaiting assessment until we acquired further information. For the 2009 update we (MM and JR) inspected all study citations identified by the searches, and obtained full reports of the studies of agreed relevance.

Data extraction and management

1. Extraction: We (MM, AL) independently extracted and entered trial data into Review Manager (RevMan) twice, cross-checking for consistency (RevMan 2008). An initial analysis included all trials meeting inclusion criteria, whilst a second sensitivity analysis excluded all but the highest quality trials (Category A and B). For the 2006 and 2010 update, we (MM and JR) independently extracted and entered data into RevMan, cross-checking again for consistency. Where disputes arose, we attempted to resolve these by discussion. When this was not possible and further information was needed to resolve the dilemma, we did not enter the data, and added this outcome of the trial to the list of those awaiting assessment.

2. Management

2.1 Forms: We extracted the data onto standard, simple forms.

2.2 Direction of graphs: Where possible, we entered data into RevMan in such a way that the area to the left of the 'line of no effect' indicates a 'favourable' outcome for early intervention. Where this was not possible, (for example, scales that calculate higher scores=improvement) we inserted a minus sign into the data tables to reverse the graphical display in RevMan analyses so that the direction of effect was clear.

2.3 Scale-derived data: Unpublished scales are known to be subject to bias in trials of treatments for schizophrenia (Marshall 2000). Therefore we only included continuous data from rating scales were if the measuring instrument had been described in a peer-reviewed journal.

2.4 Skewed data: Continuous data on outcomes in trials relevant to mental health issues are often not normally distributed. To avoid the pitfall of applying parametric tests to non-parametric data we applied the following standards to continuous final value endpoint data before inclusion: (a) standard deviations and means were reported in the paper or were obtainable from the authors; (b) when a scale started from zero, the standard deviation, when multiplied by two, should be less than the mean (otherwise the mean is unlikely to be an appropriate measure of the centre of the distribution - Altman 1996); in cases with data that are greater than the mean we entered them into the 'Other data' table as skewed data. Where the skewed data are derived from a trial with ≥ 200 participants, the skewed data pose less of a problem when looking at means if the sample size is large and were entered into syntheses.

If a scale starts from a positive value (such as PANSS, which can have values from 30 to 210) the calculation described above in (b) should be modified to take the scale starting point into account. In these cases skewness is present if $2SD > (S - S_{min})$, where S is the mean

score and Smin is the minimum score. We reported non-normally distributed data (skewed) in the 'other data types' tables. For change data (mean change from baseline on a rating scale) it is impossible to tell whether data are non-normally distributed (skewed) or not, unless individual patient data are available. After consulting the ALLSTAT electronic statistics mailing list, we entered change data in RevMan analyses and reported the finding in the text to summarise available information. In doing this, we assumed either that data were not skewed or that the analysis could cope with the unknown degree of skew.

2.5 Final endpoint value versus change data: Where both final endpoint data and change data were available for the same outcome category, only final endpoint data were presented. We acknowledge that by doing this much of the published change data may be excluded, but argue that endpoint data is more clinically relevant and that if change data were to be presented along with endpoint data, it would be given undeserved equal prominence. Where studies reported only change data we contacted authors for endpoint figures.

2.6 Common measure: To facilitate comparison between trials, we converted variables (such as days in hospital) that could be reported in different metrics (mean days per year, per week or per month) to a common metric (for example, mean days per month).

2.7 Conversion of continuous to binary: Where possible, efforts were made to convert outcome measures to dichotomous data. This could be done by identifying cut-off points on rating scales and dividing participants accordingly into 'clinically improved' or 'not clinically improved'. It was generally assumed that if there had been a 50% reduction in a scale-derived score such as the Brief Psychiatric Rating Scale (BPRS, Overall 1962) or the Positive and Negative Syndrome Scale (PANSS, Kay 1986), this could be considered as a clinically significant response (Leucht 2005a; Leucht 2005b). If data based on these thresholds were not available, we used the primary cut-off presented by the original authors.

2.8 Summary of findings table: For the 2011 version of the review we had available to us the possibility of producing Summary of Findings tables. These should be considered before being biased by the results of analyses, but for us this is impossible. We have chosen to present two - but this choice is *post hoc*. We chose to present data from PACE-Australia and OPUS-Scandinavia as these are benchmark trials in this area and outcomes from these trials that we think to be clinically important.

- Progression to psychosis
- Compliance with treatment - treatment stopped in spite of need
- Leaving the study early
- Service use: 1. Average mean number of days per month in hospital
- Service use: 2. Not hospitalised
- Social outcomes: 1. Not living independently
- Social outcomes: 2. Not working or in education

Assessment of risk of bias in included studies—Again working independently, we assessed risk of bias using the tool described in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2008). This tool encourages consideration of how the sequence was generated, how allocation was concealed, the integrity of blinding at outcome, the completeness of outcome data, selective reporting and other biases. We would not have included studies where sequence generation was at high risk of bias or where allocation was clearly not concealed.

The categories are defined below.

- YES - low risk of bias
- NO - high risk of bias
- UNCLEAR - uncertain risk of bias

If disputes arose as to which category we should allocate a trial, again, we achieved resolution by discussion, after working with a third reviewer.

Earlier versions of this review used a different, less well-developed, means of categorising risk of bias (see Appendix 2).

Measures of treatment effect

1. Binary data: For binary outcomes we calculated an estimate of the risk ratio (RR) and its 95% (fixed-effect) confidence intervals (CI). RR is more intuitive (Boissel 1999) than odds ratios and odds ratios tend to be interpreted as RR by clinicians (Deeks 2000). This misinterpretation then leads to an overestimate of the impression of the effect. When the overall results were significant we calculated the number needed to treat/harm (NNT/NNH) using Visual Rx.

2. Continuous data: For continuous outcomes we estimated mean difference (MD) between groups. We preferred not to calculate effect size measures (standardised mean difference SMD). However, had scales of very considerable similarity been used, we would have presumed there was a small difference in measurement, and we would have calculated effect size and transformed the effect back to the units of one or more of the specific instruments.

Unit of analysis issues

1. Cluster trials: Studies increasingly employ cluster randomisation (such as randomisation by clinician or practice), but analysis and pooling of clustered data pose problems. Firstly, authors often fail to account for intra-class correlation in clustered studies, leading to a unit-of-analysis error (Divine 1992) whereby P values are spuriously low, confidence intervals unduly narrow and statistical significance overestimated. This causes Type I errors (Bland 1997; Gulliford 1999).

Where clustering had not been accounted for in primary studies, we presented the data in a table, with a (*) symbol to indicate the presence of a probable unit of analysis error. In subsequent versions of this review we will seek to contact first authors of studies to obtain

intra-class correlation co-efficients (ICCs) of their clustered data and to adjust for this using accepted methods (Gulliford 1999). Where clustering has been incorporated into the analysis of primary studies, we will also present these data as if from a non-cluster randomised study, but adjusted for the clustering effect.

We have sought statistical advice and have been advised that the binary data as presented in a report should be divided by a design effect. This is calculated using the mean number of participants per cluster (M) and the ICC (Design effect=1+ (M -1)*ICC) (Donner 2002). If the ICC is not reported we assumed it to be 0.1 (Ukoununne 1999). If cluster studies had been appropriately analysed taking into account ICCs and relevant data documented in the report, we synthesised these with other studies using the generic inverse variance technique.

2. Cross-over design: A major concern of cross-over trials is the carry-over effect. It occurs if an effect (for example, pharmacological, physiological or psychological) of the treatment in the first phase is carried over to the second phase. As a consequence, on entry to the second phase the participants can differ systematically from their initial state, despite a wash-out phase. For the same reason cross-over trials are not appropriate if the condition of interest is unstable (Elbourne 2002). As both effects are very likely in schizophrenia, we will only use data of the first phase of cross-over studies.

3. Studies with multiple treatment groups: We presented studies involving more than two treatment arms, if relevant, in comparisons.

Dealing with missing data

1. Overall loss of credibility: At some degree of loss to follow-up data must lose credibility (Xia 2007). We are forced to make a judgment where this is for the trials likely to be included in this review. Should more than 50% of data be unaccounted for by eight weeks, we did not reproduce these data or use them within analyses.

2. Intention to treat analysis

2.1 Binary data: We excluded data from studies where more than 50% of participants in any group were lost to follow-up (this did not include the outcome of 'leaving the study early'). In studies with less than 50% dropout rate, people leaving early were considered to have had the negative outcome, For example, those lost to follow-up for the outcome of relapse were treated in the analysis as having relapsed. Suicide was treated as relapse.

2.2 Continuous data

2.2.1 Attrition: In the case where attrition for a continuous outcome is between 0% and 50% and completer-only data were reported, we have reproduced these.

2.2.2 Standard deviations: We first tried to obtain the missing values from the authors. If not available, where there were missing measures of variance for continuous data but an exact standard error and confidence interval were available for group means, and either P value or T value were available for differences in mean, we noted these, and in future versions will calculate them according to the rules described in the *Handbook* (Higgins

2008): When only the standard error (SE) is reported, standard deviations (SDs) can be calculated by the formula $SD = SE * \text{square root } (n)$. Chapters 7.7.3 and 16.1.3 of the *Handbook* (Higgins 2008) present detailed formula for estimating SDs from P values, T or F values, confidence intervals, ranges or other statistics. If these formula do not apply, we, in the future will calculate SDs according to a validated imputation method which is based on the SDs of the other included studies (Furukawa 2006). Some of these imputation strategies can introduce error. The alternative would be to exclude a given study's outcome and thus to lose information. We will examine the validity of the imputations in a sensitivity analysis excluding imputed values.

2.2.3 Last observation carried forward: We anticipated that in some studies the method of last observation carried forward (LOCF) would be employed within the study report. As with all methods of imputation to deal with missing data, LOCF introduces uncertainty about the reliability of the results. Therefore, where LOCF data have been used in the trial, if less than 50% of the data had been assumed, we reproduced these data and indicated that they are the product of LOCF assumptions.

Assessment of heterogeneity

1. Clinical heterogeneity: We considered all included studies initially, without seeing comparison data, to judge clinical heterogeneity. We simply inspected all studies for clearly outlying situations or people which we had not predicted would arise. When such situations or participant groups arose, we would have fully discussed these.

2. Methodological heterogeneity: We considered all included studies initially, without seeing comparison data, to judge methodological heterogeneity. We simply inspected all studies for clearly outlying methods which we had not predicted would arise. Should such methodological outliers arise we would have fully discussed these.

3. Statistical heterogeneity

3.1 Visual inspection: We visually inspected graphs to investigate the possibility of statistical heterogeneity.

3.2 Employing the I^2 statistic: We investigated heterogeneity between studies by considering the I^2 method alongside the Chi^2 P value. The I^2 provides an estimate of the percentage of inconsistency thought to be due to chance (Higgins 2003). The importance of the observed value of I^2 depends on i. magnitude and direction of effects and ii. strength of evidence for heterogeneity (e.g. P value from Chi^2 test, or a confidence interval for I^2). We interpreted I^2 estimate greater than or equal to 50% accompanied by a statistically significant Chi^2 statistic as evidence of substantial levels of heterogeneity (Section 9.5.2 - Higgins 2008). When substantial levels of heterogeneity were found in the primary outcome, we explored reasons for heterogeneity (Subgroup analysis and investigation of heterogeneity).

Assessment of reporting biases—Reporting biases arise when the dissemination of research findings is influenced by the nature and direction of results. These are described in

section 10.1 of the *Handbook* (Higgins 2006). We are aware that funnel plots may be useful in investigating reporting biases but are of limited power to detect small-study effects (Egger 1997). We did not use funnel plots for outcomes where there were 10 or fewer studies, or where all studies were of similar sizes. In other cases, where funnel plots were possible, we sought statistical advice in their interpretation.

Data synthesis—Where possible we employed a fixed-effect model for analyses. We understand that there is no closed argument for preference for use of fixed-effect or random-effects models. The random-effects method incorporates an assumption that the different studies are estimating different, yet related, intervention effects. This does seem true to us; however, random-effects does put added weight onto the smaller of the studies - those trials that are most vulnerable to bias. For this reason we favour using the fixed-effect model.

Subgroup analysis and investigation of heterogeneity

1. Subgroup analyses: We did not anticipate subgroup analyses.

2. Investigation of heterogeneity: If inconsistency was high, we have reported this. First we investigated whether data had been entered correctly. Second, if data had been correct, we visually inspected the graph and successively removed studies outside of the company of the rest to see if homogeneity was restored. Should this occur with no more than 10% of the data being excluded, we have presented data. If not, we have not pooled data and have discussed relevant issues.

Should unanticipated clinical or methodological heterogeneity be obvious we simply stated hypotheses regarding these for future reviews or versions of this review. We did not anticipate undertaking analyses relating to these.

Sensitivity analysis—For the 2011 version of this review we did not anticipate undertaking any additional sensitivity analyses.

RESULTS

Description of studies

See: Characteristics of included studies; Characteristics of excluded studies; Characteristics of studies awaiting classification; Characteristics of ongoing studies.

For substantive descriptions of studies, please see Characteristics of included studies; Characteristics of excluded studies; Characteristics of ongoing studies.

Results of the search—The original search strategy identified 9279 abstracts, of which 184 referred to potentially eligible studies and 155 reviews of early intervention (Figure 1). From these we identified 100 relevant studies, of which 43 did not meet inclusion criteria. We were able to include three studies and the remainder are awaiting classification. For substantive descriptions of studies, please see the Characteristics of included studies and Characteristics of excluded studies tables.

For the 2006 update search we identified 159 new citations and were able to include four additional studies.

During the 2009 update search we identified 830 references (from 420 studies) and were able to include 11 additional studies (Alvarez-Spain; Amminger-Austria; Berger-Australia; Edwards-Australia; EIPS-Germany; Jackson-Australia; Killackey-Australia; Leavey-UK; LEO-CAT-UK; LIPS-Germany; Uzenoff-USA).

Included studies—We included 18 studies with 1808 participants.

1. Methods: For description of methods - please see Risk of bias in included studies.

2. Participants and setting

2.1 Participants with prodromal symptoms - and setting: Six trials (Amminger-Austria, EIPS-Germany, EDIE-UK, LIPS-Germany, PACE-Australia, PRIME-USA) were concerned with preventing the onset of psychosis.

EIPS-Germany undertook the research in community settings in Cologne, Bonn, Dusseldorf, and Munich. The participants had a mean age of 26 years and were judged at risk of psychosis because they met the criteria for the Early Initial Prodrome State according to the presence of Basic Symptoms. The LIPS-Germany study was undertaken in the same settings as EIPS-Germany, but included late prodrome participants who were at risk of psychosis according to the Basic Symptom criteria for Late Initial Prodrome State (where conversion to psychosis is considered more imminent than in the Early Initial Prodrome State).

Participants had a mean age of 25 years. EDIE-UK recruited participants from primary care teams (general practitioners, practice nurses and psychological therapists), student counselling services, accident and emergency departments, specialist services (community drug and alcohol teams, child and adolescent psychiatry and adult psychiatry services) and voluntary sector agencies. Participants had a mean age of 21 years and were judged to have an 'ultra high risk' of developing a first episode of psychosis (Yung's Criteria Yung 2005). PACE-Australia recruited participants referred to the Personal Assessment and Crisis Evaluation clinic, which is part of the EPPIC programme. Participants were from 14 to 30 years of age, and met Yung criteria for an 'ultra high risk' mental state (see included studies table for details). PRIME-USA recruited people from referrals and by participants responding to study advertisements. Participants were aged from 12 to 36 years with a diagnosis of being at risk of developing psychosis according to SOPS criteria (similar to Yung's criteria for the 'ultra high risk' mental state). Participants were recruited at four sites (three in the USA and one in Canada). Amminger-Austria recruited outpatients at the Vienna General Hospital who were at risk of developing first-episode psychosis according to Yung's criteria for the 'ultra high risk' mental state. Participants were eligible if they were aged from 13-24 years; the mean age of the participants was 16.4 years.

2.2 Participants with first-episode psychosis and setting: Twelve trials were concerned with improving outcome in first-episode psychosis (Alvarez-Spain; Berger-Australia; Edwards-Australia; Jackson-Australia; Killackey-Australia; Leavey-UK; LEO-CAT-UK; LifeSPAN-Australia; Linszen-Amsterdam; OPUS-Scandinavia; Uzenoff-USA; Zhang-China).

Alvarez-Spain included drug naïve participants with a DSM-IV diagnosis of psychosis, and a mean age of 26 years; the trial was set in the community. Edwards-Australia included participants with first-episode psychosis diagnosed as having a psychotic disorder using DSM-IV criteria. The study was undertaken at the EPPIC centre in Melbourne, Australia. Berger-Australia recruited participants in Melbourne, Australia (EPPIC Centre); participants had an average age of 20 years, and a mean antipsychotic treatment exposure prior to study of 17.5 days. Jackson-Australia included people with a mean onset of psychosis at 22 years; the settings used were at the participant's home, a neutral location or the EPPIC centre. Leavey-UK included first-episode psychosis patients who had been diagnosed within the last six months and were recruited from psychiatric services in North London. LEO-CAT-UK included participants with first-episode psychosis (Yung's criteria) with a mean age of 23 years. The study was undertaken in community settings within the borough of Lambeth, London, UK. LifeSPAN-Australia recruited participants from the Western region of Melbourne, Australia and is part of the EPPIC programme, which includes an early detection and crisis assessment team. Participants were aged 15 to 29 years, and were acutely suicidal. Linszen-Amsterdam recruited participants aged from 15 to 26 who were experiencing their first episode of schizophrenia and living in close contact with parents or relatives. All participants were recruited from an adolescent clinic and had to agree to an initial three months' inpatient programme before randomisation. Subsequent treatment took place on an outpatient basis. OPUS-Scandinavia recruited first-episode psychosis (ICD 10) patients from inpatient and outpatient departments in Denmark; participants were aged 18 to 45. Zhang-China recruited only men who had just been discharged from Suhoz Psychiatric Hospital in China, following their first admission for schizophrenia. The intervention and standard care were provided on an outpatient basis. Killackey-Australia enrolled participants from the EPPIC programme with first-episode psychosis; participants had a mean age of 21 years. Uzenoff-USA recruited participants with first episode schizophrenia in the USA.

3. Study size: OPUS-Scandinavia was the largest study, and had a sample size (n=547) which was arrived at using a pre-study power calculation. The other trials were small and, in ascending order of size, were: Uzenoff-USA (24), Killackey-Australia (41), Edwards-Australia (47), LifeSPAN-Australia (56), PACE-Australia (59), EDIE-UK (60), PRIME-USA (60), Alvarez-Spain (61), Jackson-Australia (62), Linszen-Amsterdam (76), Berger-Australia (80), Amminger-Austria (81), Zhang-China (83), Leavey-UK (106), LEO-CATUK (113), LIPS-Germany (124) and EIPS-Germany (128).

4. Intervention

4.1 Trials to prevent the onset of psychosis: Six trials (Amminger-Austria; EIPS-Germany; LIPS-Germany; EDIE-UK; PACE-Australia; PRIME-USA) were concerned with preventing the onset of psychosis.

Amminger-Austria compared omega-3 fatty acids (eicosapentaenoic acid and docosahexaenoic acid) with placebo over three months in adolescents at risk of first-episode psychosis according to the criteria of Yung 2005.

EIPS-Germany employed a complex intervention consisting of 12 months of cognitive behaviour therapy (CBT) delivered in group and individual therapy sessions, supplemented by cognitive remediation therapy. The participants were at risk of developing first episode of psychosis and met the early initial prodromal state criteria. The control group were given supportive therapy sessions which only provided minimal support involving psychoeducation and counselling.

LIPS-Germany employed a needs focused intervention combined with amisulpride (mean dose 118 mg/day) in participants judged at risk of psychosis because of the presence of prodromal symptoms. The control group also received the needs focused intervention but without amisulpride. The needs focused intervention included psychoeducation, crisis intervention, family counselling and assistance with education or work related difficulties according to the patient's need.

The Early Detection and Intervention Evaluation trial (EDIEUK) used cognitive therapy, limited to a maximum of 26 sessions over six months, following the principles developed by Beck 1976. The therapy was problem-orientated and time limited and was carried out by experienced cognitive therapists. Both control and treatment group received regular monitoring. Whilst participants (treatment and control) were not given medication, both treatment and control received elements of case management in order to resolve crises regarding social issues and mental health risk.

In PACE-Australia the intervention involved prescription of low dose risperidone (1-2 mg/day) combined with modified CBT, which aimed to enhance understanding and control of symptoms. Both the intervention and control groups also received case management from a PACE therapist. This involved supportive psychotherapy, assistance with accommodation and education/employment, and family support. Participants in the control and intervention groups received standard treatment if they developed psychosis, but control patients were not otherwise prescribed neuroleptics. Both groups could be prescribed antidepressants and benzodiazepines.

The Prevention through Risk Identification Management and Education study (PRIME-USA) randomised participants to olanzapine 5-15 mg/day (mean 8 mg/day) or placebo for one year, and then followed up for a further year without medication. Individual and family psychosocial interventions with supportive and psychoeducational components were available to all patients during the first year. The nature of the psychoeducational components varied across sites despite efforts to apply them in a uniform way. The psychosocial intervention available at the New Haven centre was modelled on the Problem Solving Training approach (D'Zurilla 1971; D'Zurilla 1986).

4.2 Trials to improve the outcome of first-episode psychosis: Twelve trials were concerned with improving outcome in first-episode psychosis (Alvarez-Spain; Berger-Australia; Edwards-Australia; Jackson-Australia; Killackey-Australia; Leavey-UK; LEO-CAT-UK; LifeSPAN-Australia; Linszen-Amsterdam; OPUS-Scandinavia; Uzenoff-USA; Zhang-China).

In Alvarez-Spain, drug naive first-episode participants were randomised to three different antipsychotics (risperidone, olanzapine, haloperidol) and then randomised to either Early Behavioural Intervention or the control group (routine clinical care).

In Berger-Australia, the first-episode psychosis participants were given ethyl-eicosapentaenoic acid oil (E-EPA) at a dose of 500 mg twice Daily, with a flexible dose of atypical antipsychotics. The control group received placebo capsules with a flexible dose of atypical antipsychotics.

In Edwards-Australia, the intervention group received a behavioural modification intervention, Cannabis and Psychosis Therapy (CAP). This consisted of weekly sessions of CBT provided by trained clinicians over three months. The aim of the CAP intervention was to reduce cannabis intake and to improve clinical and psychosocial functioning. CAP involved an assessment of engagement, followed by education about cannabis and psychosis and developing motivation to change. The focus of therapy was determined by the phase of commitment to change and could include further educational sessions, motivational interviewing, goal setting, and discussion about relapse prevention. An active control group was used which consisted of psychoeducation, which explained psychosis, medication and other treatments, and relapse prevention, but did not discuss cannabis.

In Jackson-Australia, the intervention group received CBT with 20 sessions provided for 45 minutes, plus antipsychotics. The control group were given a befriending service in addition to antipsychotics.

In Killackey-Australia, the intervention group received individual placement and support, which is an intervention designed to help people with mental illness to find and keep competitive employment. The support provided in the programme continued after employment was obtained, and was adapted to the needs of the individual. The control group received treatment as usual.

In Leavey-UK, the intervention group received a brief intervention and treatment as usual. The brief intervention was provided over seven sessions, lasting about one hour, and included: information gathering from the relative, plus sessions on: psychotic illness, symptoms and early warning signs, treatment, help seeking; coping strategies, problem solving and communication with the patient. The control group were given treatment as usual.

LEO-CAT-UK was a cluster-randomised trial in which primary care (GP) practices were randomly allocated to receive training in early detection of psychosis and direct access to LEO-CAT (a specialised treatment team for first-episode psychosis). The control group of General Practice clinics did not receive training in early detection and continued to refer new cases of psychosis to local mental health services who could then refer on to the LEO-CAT programme.

In LifeSPAN-Australia, the intervention group received standard clinical care plus LifeSPAN therapy which draws on the experience at EPPIC with Cognitive Orientated Therapy for Early Psychosis (COPE) and suicide manuals such as Choosing to Live and

Cognitive Therapy of Suicide Behaviour. Four phases are used for the intervention: (a) initial engagement, (b) suicide risk assessment/formulation, (c) cognitive modules and (d) final closure/handover.

In Linszen-Amsterdam the intervention was behavioural family therapy for one year. Eighteen family therapy sessions were held over a 12-month period. Each family was treated by two co-therapists, from a team of two psychologists and one social worker, all of whom had at least one year of experience in providing family interventions for schizophrenia. The intervention was based on the behavioural family management approach of Falloon 1984 and involved psychoeducation, communication training and development of problem solving skills. Both intervention and control groups also received care from a specialised first-episode team involving individual-oriented therapy consisting of maintenance medication and disease and stress management.

In OPUS-Scandinavia, participants received integrated treatment or standard care. Integrated treatment consisted of high fidelity assertive community treatment supplemented by behavioral family therapy and social skills training. Standard care consisted of care at a community mental health centre. All participants were offered antipsychotic drugs according to guidelines from the Danish Psychiatric Society, which recommends a low-dose atypical antipsychotic strategy for first episodes of psychotic illness. Each participant was usually in contact with a physician, community mental health nurse and in some cases a social worker. In a small proportion of cases, standard care also included psychosocial interventions such as training in social skills or Daily living activities, or supportive contacts with the family. Antipsychotics were given to both groups based on the psychiatrists' clinical assessment.

The Uzenoff-USA study provided participants with Adherence Coping Education (ACE) which consists of 14 sessions lasting between 30 and 45 minutes over six months. It is a manual-based psychotherapy grouped into four phases: (1) establishing therapeutic alliance; (2) promoting treatment adherence; (3) developing a plan for maintenance treatment and (4) rehabilitation. The control group received supportive therapy which involved (1) establishing the therapeutic relationship, and (2) providing emotional support plus discussion of non-illness issues or topics.

Zhang-China also used family therapy, but in the form of group and individual family sessions which were delivered on an outpatient basis over the 18-month follow-up period. Both intervention and control groups also received care from the outpatients department, (consisting of medication and review) but no regular appointments or community follow-ups were provided.

5. Trial duration: The six trials concerned with preventing the onset of psychosis reported data from between two months and two years: EIPS-Germany at 12 months; EDIE-UK at 12 months and 36 months; PACE-Australia at six and 12 months (the first six months being the period during which the intervention was received); PRIME-USA at two months, 12 months (first 12 months study intervention given) and 24 months (last 12 months without intervention); LIPS-Germany at three months (although the study is planned to last 24 months), and Amminger-Austria at three months.

The 12 trials concerned with improving outcome of first-episode psychosis also reported data at various time points (range three months to five years): Alvarez-Spain reported data at 13 weeks; Berger-Australia reported data at three months; Jackson-Australia followed participants for 12 months; Leavey-UK reported data at nine months. Four trials (Edwards-Australia; Killackey-Australia; LifeSPAN-Australia; Uzenoff-USA) reported data at six months. Linszen-Amsterdam reported at 12 months following an initial three-month inpatient admission and also reported a five-year follow-up, but data were provided for the whole sample only, not by group allocation. OPUS-Scandinavia reported data at 12 and 24 months, whereas Zhang-China reported at 18 months.

6. Outcomes

6.1 Non-scale data: We were able to report dichotomous data on suicide, death, leaving the study early, conversion to psychosis, adverse effects, hospital admission, days in hospital, compliance with medication, antipsychotic drug use, living independently and employment.

6.2 Scale derived data: Only details of the outcome scales that provided usable data are shown below. Reasons for exclusions of data are given under 'Outcomes' in Characteristics of included studies.

6.2.1 Global state scales: a. Global Assessment of Functioning - GAF (APA 1994)

This is an observer rated scale for measuring overall severity of functional impairment. GAF consists of nine behavioural descriptors. Patients are rated between 0 (most severe) and 90 (least severe) for each descriptor. PRIME-USA, PACE-Australia, OPUS-Scandinavia and LIPS-Germany reported data from this scale.

b. Clinical Global Impression - CGI (Guy 1970)

The CGI is a three-item scale commonly used in studies on schizophrenia that enables clinicians to quantify severity of illness and overall clinical improvement. The items are: severity of illness; global improvement and efficacy index. A seven-point scoring system is usually used with low scores indicating decreased severity and/or greater recovery. PRIME-USA reported data from this scale.

c. Knowledge About Psychosis Questionnaire - KAPQ (Birchwood 1992)

This questionnaire tests the patients understanding about psychosis and treatments. Data from this scale were reported by Edwards-Australia.

6.2.2 Mental state scales: a. Brief Psychopathological Rating Scale - BPRS (Overall 1962)

The BPRS is an 18-item scale measuring positive symptoms, general psychopathology and affective symptoms. The original scale has 16 items, but a revised 18-item scale is commonly used. Scores can range from 0-126. Each item is rated on a seven-point scale varying from 'not present' to 'extremely severe', with high scores indicating more severe symptoms. Data from this scale were reported by Edwards-Australia. In PACE-Australia the scale was used primarily to report severity of psychotic symptoms.

b. Positive and Negative Symptom Scale - PANSS (Kay 1987)

The Positive and Negative Symptom Scale was developed from the BPRS and the Psychopathology Rating Scale. It is used as a method for evaluating positive, negative and other symptom dimensions in schizophrenia. The scale has 30 items, and each item can be defined on a seven-point scoring system varying from one (absent) to seven (extreme). This scale can be divided into three sub-scales for measuring the severity of general psychopathology, positive symptoms (PANSS-P) and negative symptoms (PANSS-N). A low score indicates low levels of symptoms. EDIE-UK used this scale to determine transition to psychosis. PRIME-USA and LIPS-Germany reported data from the PANSS.

c. Scale of Psychotic Symptoms - SOPS (Miller 1999)

The SOPS scale was modelled on the PANSS scale and is designed to measure the presence/absence of prodromal states. It consists of five positive symptom items, six negative symptom items, four disorganisation symptoms items, and four general symptom items. Each has a severity rating from 0 (never, absent) to six (severe/extreme - and psychotic for the positive items). The severity of the prodromal state is based on the sum of the rating from the SOPS items and ranges between 0 and 114. PRIME-USA reported data from this scale.

d. Hamilton Rating Scale for Anxiety - HRSA (Hamilton 1959)

The Hamilton Anxiety Scale (HAMA) is a rating scale developed to quantify the severity of anxiety symptoms, often used in psychotropic drug evaluation. It consists of 14 items, each defined by a series of symptoms. Each item is rated on a five-point scale, ranging from 0 (not present) to 4 (severe). The 14 items consist of: anxious mood; tension; fears; insomnia; intellectual; depressed mood; somatic complaints (muscular); somatic complaints (sensory); cardiovascular symptoms; respiratory symptoms; gastrointestinal symptoms; genitourinary symptoms; autonomic symptoms and behaviour at Interview. Higher scores indicate greater anxiety. PACE-Australia reported data from this scale.

e. Hamilton Rating Scale for Depression - HRSD (Hamilton 1960)

This is an interviewer rated scale for measuring depression. It is used for quantifying the results of an interview and depends on the skill of the interviewer in eliciting the necessary information. It contains 17 variables measured on either a five point or a three-point rating scale. The variables include: depressed mood; suicide; employment and loss of interest; retardation; agitation; gastrointestinal symptoms; general somatic symptoms; hypochondriasis; loss of insight and loss of weight. Higher scores indicate more severe depression. PACE-Australia reported data from this scale.

f. Calgary Depression Rating Scale - CDRS (Addington 1990)

The Calgary Depression Scale for Schizophrenia is a nine-item scale (0=absent; 1=mild; 2=moderate; 3=severe.) that was specifically developed for assessment of depression in patients with schizophrenia. It has been evaluated in both relapsed and remitted patients, and

is provided as a semi-structured interview. High scores indicated worse outcome. Uzenoff-USA reported data from this scale.

g. Montgomery Asberg Depression Rating Scale - MADRS (Montgomery 1979)

This is a 65-item comprehensive psychopathology scale used to identify the 17 most commonly occurring symptoms in primary depressive illness. Ratings are based on 10 items, with higher scores indicating more symptoms. This scale was used by LIPS-Germany.

h. Beck Depression Inventory - BDI-SF (Beck 1961)

This is a 21-item self-rating scale for depression. Each item comprises four statements (rated 0-4) describing increasing severity of the abnormality concerned. The person completing the scale is required to read each group of statements and identify the one that best describes the way they have felt over the preceding week. A total of 12/13 is an indicative score for presence of significant depression. The short form of this scale was used by Edwards-Australia.

i. Presence of Psychosis Scale - POPS (Olsen 2006)

The Presence of Psychosis Scale (POPS), is part of the Structured Interview for Prodromal Syndromes scale (SIPS). It marks onset of psychosis by the presence of positive symptoms at the psychotic level of intensity and of sufficient frequency and duration. PRIME-USA reported data from this scale.

j. Scale for the Assessment of Negative Symptoms - SANS (Andreasen 1983)

This is also an interviewer rated scale for measuring the severity of negative symptoms of schizophrenia such as alogia, affective blunting, avolition-apathy, anhedonia-asociality and attention impairment. Items are rated on a six-point scale with higher scores indicating more symptoms. Edwards-Australia and PACE-Australia reported data from this scale.

k. Young Mania Scale - YMS (Young 1978)

Again an interviewer rated scale, but this time for measuring the severity of symptoms of mania. Higher scores indicate more severe symptoms. PRIME-USA and PACE-Australia reported data from this scale.

6.2.3 Social functioning: a. Social Functioning Scale II - SAS II (Weissman 1976)

The SAS II is an interviewer-administered scale adapted from the self-report Social Adjustment Scale for use with people with schizophrenia. It contains 52 questions that are administered in a semi-structured interview by a trained rater. The SAS II assesses current functioning with scores ranging from 1 to 5, with higher scores indicating worse functioning. EIPS-Germany reported data from this scale.

b. Social and Occupational Functioning Assessment Scale - SOFAS (APA 1994)

The SOFAS is a new instrument similar to the Global Assessment of Functioning is format, and attempts to assess the social and or occupational functioning independent of the overall severity of the illness. Higher scores indicate worse social functioning. Edwards-Australia and Jackson-Australia reported data from this scale.

6.2.4 Adverse effects: a. Simpson Angus Scale - SAS (Simpson 1970)

The SAS is a 10-item scale used to evaluate the presence and severity of drug-induced parkinsonian symptoms. The 10 items focus on rigidity rather than bradykinesia and do not assess subjective rigidity or slowness. The scale comprises a 10-item rating scale, each item rated on a five-point scale with zero meaning the complete absence of condition and four meaning the presence of condition in extreme. A low score indicates low levels of parkinsonism. PRIME-USA reported data from this scale,

b. Barnes Akathisia Rating Scale - BAS (Barnes 1989)

The Barnes Akathisia Rating Scale is a four-item scale to assess the presence and severity of drug-induced movement disorder akathisia. It is a widely used comprehensive rating scale for akathisia. Items include restless movements that characterise akathisia, the subjective awareness of restlessness and any distress associated with the condition. These items are rated from zero (normal) to three (severe). In addition, there is an item for rating the global severity that starts from zero (absent) to five (severe). A low score indicates low levels of akathisia. PRIME-USA reported data from this scale.

c. Abnormal Involuntary Movement Scale - AIMS (Guy 1976)

The Abnormal Involuntary Movement Scale has been used to assess abnormal involuntary movements associated with antipsychotic drugs, such as tardive dyskinesia and chronic akathisia, as well as 'spontaneous' motor disturbance related to the illness itself. Tardive dyskinesia is a long-term, drug-induced movement disorder. However, using this scale in short-term trials may also be helpful to assess some rapidly occurring abnormal movement disorders such as tremor. Scoring consists of rating movement severity in the anatomical areas (facial/oral, extremities, and trunk) on a five-point scale (0-4). A low score indicates low levels of dyskinesic movements. PRIME-USA reported data from this scale.

6.2.5 Quality of life: a. Quality of Life Scale - QLS (Heinrichs 1984)

This is a semi-structured interview administered and rated by trained clinicians. It contains 21 items rated on a seven-point scale based on the interviewer's judgement of patient functioning. Higher scores indicate better quality of life. PACE-Australia and Uzenoff-USA reported data from this scale.

6.2.6 Satisfaction with care: a. The Client Satisfaction Questionnaire - CSQ-8 (De-Wilde 2005)

The CSQ-8 is an eight-item self-report of global measure of patient satisfaction with services. The CSQ is substantially correlated with treatment dropout, number of therapy sessions attended, and with change in client-reported symptoms. The CSQ-8 consists of

eight items rated on a four-point Likert scale. The items are concerned with quality of services received, how well services met the client's needs and general satisfaction. The total score ranges from eight to 32. Higher scores indicate greater satisfaction of the responders OPUS-Scandinavia reported data from this scale.

6.2.7 Substance use: a. Cannabis and Substance Use Assessment Schedule - CASUAS (Wing 1990)

This scale measures the percentage of days using cannabis in the past four weeks and includes an index of severity of cannabis use. The scale is modified from the Schedule for Clinical Assessment on Neuropsychiatry and includes similar information to the Addiction Severity Index. Data from this scale were used by Edwards-Australia.

6.3 Redundant data: Some studies reported data only as P values or statements of significant or non-significant differences, and other continuous data could not be extracted because the number of participants was missing or standard deviations were not reported.

Excluded studies—There are currently 68 excluded studies. We have summarised reasons for exclusion in Table 1.

1. Awaiting classification: Thirty studies are awaiting assessment; 13 are brief reports where additional data are required; seven Chinese studies require further clarification; five studies are unclear regarding whether the status of participants is first episode or not; five studies are being sought. Ultimately, we will exclude studies where data are unobtainable.

2. Ongoing studies: We are awaiting data from 10 studies (see descriptions in Characteristics of ongoing studies table). This is an active area for research.

Risk of bias in included studies

Judgement of risks are illustrated in Figure 2.

Allocation—All 18 included studies were stated to be randomised. Only five described how randomisation had been performed, using computer-generated random numbers (Alvarez-Spain; Edwards-Australia; EIPS-Germany; Killackey-Australia; OPUS-Scandinavia). Attempts to conceal allocation were described in five studies which were judged to be adequately concealed. In the remaining studies allocation concealment was either not described or only briefly commented on and we were unable to determine in these instances if concealment is adequate.

Blinding—Blinding of participants and clinicians proved difficult in most studies. PRIME-USA blinded participants, investigators and dispensers to group assignment. Other studies used independent raters, some of whom were blind to allocation. PACE-Australia and OPUS-Scandinavia used raters who were independent of the study group, but were not blind to treatment allocation. In EDIE-UK, single blinding was attempted for the rater, but blinding was not maintained due to participants divulging information, or using language that suggested they were receiving cognitive therapy. Edwards-Australia and Leavey-UK

also used a single blind design. Zhang-China used independent raters blind to allocation, whereas the LifeSPAN-Australia study was described as only single blind. In Linszen-Amsterdam the status of the raters was unclear. EIPS-Germany did not state whether blinding had been attempted. Berger-Australia and Amminger-Austria used a double-blind design, whilst LIPS-Germany was an open-label study. In the LEO-CAT-UK study, the unit of randomisation were the GP practices and clinicians were not blinded to the intervention. Both Alvarez-Spain and Jackson-Australia used a single blinding. In Killackey-Australia blinding was not reported. Uzenoff-USA used blind raters. Overall, due to the nature of the intervention, blinding proved difficult in these studies and most studies are at risk of bias.

Incomplete outcome data—Follow-up rates where reported were quite high (see Table 2). Overall, study *attrition* did not suggest a risk of bias.

Outcomes were recorded, reported and analysed in many different ways. Even in this limited, relatively recent, research community, there is no indication of a consistency of approach and incomplete and selective reporting could easily be operating. PACE-Australia provided all outcomes on an intention-to-treat (ITT) basis. EDIE-UK used an ITT analysis but two people originally randomised to the treatment group were subsequently omitted from the analysis because they were found to be psychotic at the time of randomisation. (Because these exclusions are not compatible with an ITT analysis of relapse we counted such data as relapses and included these in the final analysis.) PRIME-USA also used an ITT analysis and reported scale data as change scores rather than endpoint scores. LifeSPAN-Australia only provided dichotomous data for leaving the study early and suicide. Linszen-Amsterdam only provided data on relapse at 12 months on an ITT basis. Berger-Australia only reported usable data on the number of participants not responding to treatment. Jackson-Australia did not report data on relapse or severity of illness, only data on hospitalisation, suicide and social functioning were usable. Killackey-Australia only reported usable data for employment and attrition. Leavey-UK used an ITT analysis. In the EIPS-Germany study 15 participants were not accounted for after randomisation. OPUS-Scandinavia used an ITT analysis. Zhang-China reported data on number of people readmitted and compliant on an ITT basis, but data on mental state and overall functioning were reported only for people who were not admitted to hospital. (This rendered data unusable.) Uzenoff-USA included only 24 participants and used a modified ITT analysis on 19 of this group. This small sample may have resulted in treatment effects being undetected. In Amminger-Austria, five participants were not accounted for after randomisation and is a potential source of bias to the outcome data. In LEO-CAT-UK much of the data were unusable due to the reporting of outcomes without the denominator; additionally, we divided binary data by a design effect to adjust for the excessive weight given to this cluster randomised trial. Edwards-Australia used the last observation carried forward method and all 47 participants were utilised in the reporting of outcome data.

Selective reporting—Outcome data from the SAPS scale were not reported by Alvarez-Spain. We did not identify overt under reporting of outcomes in the other included studies, although we did not have access to study protocols to check whether other data were recorded and not reported in the final papers.

Other potential sources of bias—Six of the 18 studies were undertaken at the pioneering EPPIC Centre in Melbourne, Australia (Berger-Australia; Edwards-Australia; Jackson-Australia; Killackey-Australia; LifeSPAN-Australia; PACE-Australia). This could lead to issues with applicability (see Overall completeness and applicability of evidence). However, as with the Australian studies, many of the trials were undertaken by leading figures in the world of early intervention who could have a vested interest in the findings - just as industry has in the outcomes for the drugs they manufacture. Early intervention studies are now less novel than a decade ago. The initial flourish of research has settled into a more steady stream and it will be interesting to see how findings average across time.

Effects of interventions

See: **Summary of findings for the main comparison PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM compared to SPECIALISED TEAM for psychosis**; **Summary of findings 2 SPECIALISED TEAM compared to STANDARD CARE for psychosis**

1. Obvious heterogeneity—Meta-analyses were often not possible because of the heterogeneity of the interventions and outcomes. In trials aiming to prevent onset of psychosis it was not possible to add together Amminger-Austria which used omega-3-fatty acids with other antipsychotic trials. PACE-Australia used risperidone and CBT and cannot be added to trials that used only antipsychotics. LIPS-Germany used amisulpride and a psychoeducation programme, whilst PRIME-USA used only olanzapine. Two studies (EDIE-UK, EIPS-Germany) were added together under the same comparison group, but the reported outcomes were dissimilar and we could not undertake meta-analysis. In trials aiming to improve outcome of first-episode psychosis Leavey-UK and Zhang-China were somewhat similar but it is not known what type of class of antipsychotic were given. In trials using cognitive therapy the focus of trials was different. Jackson-Australia targeted symptoms, Edwards-Australia targeted cannabis misuse, and LifeSPAN-Australia targeted suicidal ideas. We did perform an exploratory meta-analysis with three trials (Jackson-Australia; Leavey-UK; Uzenoff-USA), but this was limited by only few outcome measures being similar.

2. Trials to prevent the development of psychosis—Six studies addressed the question of prevention of psychosis by interventions for patients with prodromal symptoms.

2.1 COMPARISON 1. Phase specific intervention (olanzapine) + non specific supportive therapy versus placebo + non specific supportive therapy (all data from PRIME-USA)

2.1.1 Leaving the study early: In PRIME-USA we found the numbers of people leaving the study early by eight weeks (n=60, RR 1.29 CI 0.6 to 2.7) and also by 12 months to be equivocal (n=60, RR 1.59 CI 0.9 to 2.9).

2.1.2 Conversion to psychosis (POPS): By about 12 months the number of people converting to psychosis was 8/31 olanzapine group and 13/29 for the placebo group. This difference did not reach statistical significance (n=60, RR 0.58 CI 0.3 to 1.2).

2.1.3 Global state

2.1.3.1 Clinical Global Impression (CGI): We found the Clinical Global Impression change score 'severity of illness' equivocal by 12 months (n=59, WMD -0.23 CI -0.8 to 0.4).

2.1.3.2 Global Assessment of Functioning (GAF): We also found the Global Assessment of Functioning 'current' change score by 12 months to be equivocal (n=59, WMD 2.43 CI -4.8 to 9.6).

2.1.4 Mental state

2.1.4.1 Scale of Prodromal Symptoms (SOPS): PRIME-USA reported several outcomes as mean change scores from the SOPS at 12 months. We found the total score, positive score, negative score, disorganisation and general scores were not significantly different between the olanzapine and placebo group.

2.1.4.2 Positive and Negative Symptom Score (PANSS): We found the PANSS total (n=59, WMD 0.48 CI -10.7 to 11.7), PANSS positive (n=59, WMD -0.57 CI -3.8 to 2.6), PANSS negative (n=59, WMD 0.52 CI -2.6 to 3.6), and the PANSS general score (n=59, WMD 0.54 CI -5.4 to 6.5) were not significantly different between olanzapine and placebo groups.

2.1.4.3 Young Mania Rating Scale (YMRS): We found change scores by 12 months were equivocal (n=59, WMD -0.91 CI -3.8 to 2.0).

2.1.4.4 Montgomery and Asberg Depression Rating Scale (MADRS): We found depression change scores at 12 months were equivocal (n=59, WMD 0.68 CI -3.8 to 5.2).

2.1.5 Adverse effects

2.1.5.1 Simpson & Angus (SAS): Extrapyramidal symptoms were found to be equivocal between the olanzapine (mean 8 mg/day, range 5-15 mg/day) and the placebo group by eight weeks (n=59, WMD 0.10 CI -0.6 to 0.8).

2.1.5.2 Barnes Akathisia Scale (BAS): We also found change scores for akathisia to be equivocal by eight weeks (n=59, WMD 0.50 CI -0.6 to 1.6).

2.1.5.3 Abnormal Involuntary Movement Scale (AIMS): We found involuntary movement scores were not significantly different for those given low dose olanzapine by eight weeks compared with placebo (n=59, WMD 0.60 CI -0.3 to 1.5).

2.1.5.4 Weight change: We found the olanzapine group had a statistically significant increase in weight compared with the placebo group by 12 months (n=59, WMD 7.63 CI 4.0 to 11.2). We also found dichotomous data supported this finding, with the olanzapine group having significantly more weight gain (criteria not stated) than placebo by 12 months (n=60, RR 3.55, CI 1.5 to 8.3, NNH 3 CI 2 to 11).

2.1.5.5 Cardiovascular measures: Systolic and diastolic blood pressure values were measured sitting and standing (at eight weeks), and we found all data to be not statistically significantly different between olanzapine and placebo groups. Pulse rates were also measured (standing and sitting at eight weeks) and we again found data to be equivocal.

Twelve-month outcome data for change in pulse rates (sitting) did significantly favour the placebo group (n=58, WMD 8.31 CI 0.5 to 16.1), with the assumption that a lower pulse rate indicated an improvement. However, this significant finding was not replicated for pulse data recorded whilst standing, with data being non-significant.

2.1.5.6 Treatment emergent adverse events (CoStart terms): Somnolence, increased appetite, anxiety, nervousness, asthenia, joint disorder, abnormal thoughts were all equivocal by eight weeks. We found weight gain to be significantly higher in the olanzapine group (n=60, RR 10.29, CI 1.4 to 74.8, NNH 4 CI 2 to 70) by eight weeks compared with placebo.

2.1.5.7 Fatigue: We found the number of participants experiencing fatigue were significantly higher in the olanzapine group compared with the placebo control (n=60, RR 8.42 CI 1.1 to 62.4, NNH 4 CI 2 to 211).

2.2 COMPARISON 2. Phase specific intervention (cognitive behavioural therapy) + non-specific supportive therapy versus non specific supportive therapy (all data from EDIEUK and EIPS-Germany)

2.2.1 Leaving the study early: We found the numbers of people leaving the study early in EDIEUK were similar for both CBT (11/37) and control group (7/23) with no significant differences at 12 months (n=60, RR 0.98 CI 0.4 to 2.2). Two-year data also revealed no significant difference (n=60 RR 0.96 CI 0.6 to 1.5).

2.2.2 Transition to psychosis: The number of people who became psychotic during 12 months of observation were not significantly different for the CBT and monitoring groups (EDIE-UK, n=60, RR 0.50 CI 0.2 to 1.7).

2.2.3 Social functioning: SAS II: We found no significant differences for the outcomes of social activities, well-being, or employment. Global Social Adjustment scores also revealed no significant difference (EIPS-Germany, n= 69, WMD -0.10 CI -0.4 to 0.2).

2.3 COMPARISON 3. Phase specific intervention (risperidone + cognitive behavioural therapy) + specialised team versus specialised team (all data from PACE-Australia)

2.3.1 Leaving the study early: No participants were lost to follow-up at 12 months in either treatment (0/31) or control group (0/28).

2.3.2 Progression to psychosis (primary outcome): We found that participants with prodromal symptoms who received the intervention were significantly less likely to have developed psychosis at the six-month follow-up than controls (n=59, RR 0.27 CI 0.1 to 0.9, NNT 4 CI 2 to 20). However, this effect became non-significant by 12 months (n=59, RR 0.54 CI 0.2 to 1.3).

2.3.3 Global state: PACE-Australia used the GAF to rate overall functioning. At 12 months, data were skewed and no significant differences were found between the phase-specific treatment plus specialised team and the group receiving care from a specialised team (n=59, WMD 0.00 CI -5.2 to 5.2).

2.3.4 Mental state: There were no significant differences between intervention and control groups at six or 12 month follow up on any of the measures of mental state, but confidence intervals were generally wide. The BPRS results at both six and 12 months were equivocal and considerably skewed (n=59, WMD at 6 months -0.50 CI -2.3 to 1.3 ; WMD at 12 months 0.70 CI -1.0 to 2.4). This also applied to the SANS negative symptoms scores (n=59, WMD at six months -4.6 CI -12.7 to 3.5 ; WMD at 12 months -0.80 CI -7.9 to 6.3). Ratings of anxiety, depression and mania all had wide confidence intervals and data were skewed. No findings were statistically significant, either at six or 12 months.

2.3.5 Quality of life: We found no significant differences between intervention and control groups at six or 12 month follow up on the quality of life measure (n=59, WMD at 6 months -1.40 CI -13.6 to 10.8 ; WMD at 12 months 0.80 CI -10.2 to 11.8).

2.4 COMPARISON 4. Phase specific intervention (amisulpride + needs focused interventions - NFI) versus needs focused interventions (all data from LIPS-Germany)

2.4.1 Leaving the study early: Fewer participants dropped out of the amisulpride plus NFI group (n=124, RR 0.59 CI 0.4 to 0.9 , NNT 5 CI 4 to 34) compared with those given just Needs Focused Interventions.

2.4.2 Mental state: PANSS: The PANSS global scores and PANSS-positive scores favoured participants given amisulpride plus NFI (PANSS-G, n=102, WMD -3.40 CI -6.9 to 0.1), (PANSS +ve, n=102, WMD -2.10 CI -3.7 to -0.5) compared with the control group receiving NFI. We found no significant difference for PANSS negative symptoms (n=103, WMD -1.30 CI -3.3 to 0.7). Depression scores also revealed no significant differences (n=102, WMD -1.10 CI -4.5 to 2.3).

2.4.3 Global state: GAF: Global Assessment of Functioning scores favoured the amisulpride plus NFI (n=102, WMD -6.10 CI -11.8 to -0.5) compared with participants given Needs Focused Interventions alone.

2.5 COMPARISON 5. Omega 3 fatty acids (epa) versus PLACEBO (all data from Amminger-Austria)

2.5.1 Transition to psychosis: We found that participants who were given the essential fatty acid regime were significantly less likely to develop psychosis than the placebo group (n=76, RR 0.13 CI 0.02 to 1.0 , NNT 6 CI 5 to 96).

3. Trials to improve outcome in first-episode psychosis

3.1 COMPARISON 6. Phase-specific treatment (cognitive behavioural therapy for suicidality) + specialised team versus specialised team (all data are from a single study LifeSPAN-Australia)

3.1.2 Leaving the study early: We found the number of people leaving the study early by six months were not significantly different between the LifeSpan therapy group and those receiving standard care (n=56, RR 2.02 CI 0.7 to 5.7).

3.1.3 Suicide: Two people died from suicide during the six-month study, one from each intervention group.

3.2 COMPARISON 7. Phase-specific treatment (family therapy) + specialised team versus specialised team (all data from Linszen-Amsterdam)

3.2.1 Relapse (primary outcome): In Linszen-Amsterdam we found no significant difference between intervention and control groups at 12 months for the outcome of relapse (n=76, RR 1.05 CI 0.4 to 3.0).

3.3 COMPARISON 8. Phase-specific treatment (family therapy) + standard care versus standard care (all data from Zhang-China)

3.3.1 Leaving the study early: In Zhang-China we found only five people in a study of 83 participants were lost by 18 months. There were no significant differences in the number of people lost to follow-up for the two groups (n=83, RR 1.46 CI 0.3 to 8.3).

3.3.2 Admitted to hospital: We found that participants receiving the intervention were significantly less likely to be admitted to hospital at 18 months than people allocated to the standard care control group (n=83, RR 0.28 CI 0.1 to 0.6, NNT 3 CI 2 to 6).

3.3.3 Not compliant with medication: In both groups most people were compliant with medication. We found no significant difference in the number of people not compliant with medication at 18 months' follow-up, although the data suggested a trend favouring the intervention (p=0.06, n=83, RR 0.57 CI 0.3 to 1.0).

3.4 COMPARISON 9. Specialised team versus standard care (all data from OPUS-Scandinavia)

3.4.1 Leaving the study early: We found the numbers of people leaving early by one year were significantly lower in the integrated treatment group (n=547, RR 0.59 CI 0.4 to 0.8, NNT 9 CI 6 to 18) compared with the standard care group. By two years, numbers of people leaving the study early remained significantly lower in the integrated treatment group (n=547, RR 0.64 CI 0.5 to 0.8, NNT 7 CI 6 to 14). Five-year data, however were not significantly different (n=547, RR 1.01 CI 0.8 to 1.2).

3.4.2 Global state - Global Assessment of Functioning (GAF): We found GAF 'symptom' endpoint scores to significantly favour the integrated treatment group (n=419, WMD -3.71 CI -6.7 to -0.7) by one year. Two-year outcome data were, however, not significantly different (n=369, WMD -2.51 CI -5.7 to 0.7), and by five years follow-up data were equivocal. The GAF 'function' endpoint scores at 12 months were not significantly different (n=419, WMD -2.30 CI -5.25 to 0.6), but by two years results significantly favoured integrated treatment compared with standard care (n=369, WMD -4.03 CI -7.2 to -0.8), although by five years' follow-up functioning scores were equivocal.

3.4.3 User satisfaction - Client Satisfaction Questionnaire Score (CSQ-8): Overall satisfaction with levels of care were significantly better for the integrated treatment group (n=419, WMD -1.90 CI -3.1 to -0.7) compared with the standard care control at 12

months; this finding continued during the second year of follow-up (n=369, WMD -3.20 CI -4.1 to -2.3).

3.4.4 Compliance with treatment: We found 'treatment stopped in spite of need' - measured at one year significantly favoured the integrated treatment group (n=507, RR 0.20 CI 0.1 to 0.4, NNT 9 CI 8 to 12) compared with standard care. However, by two years we did not find any statistically significant differences between the treatment and control group (n=436, RR 0.66 CI 0.3 to 1.5).

3.4.5 Death/Suicide: Two people committed suicide (one from each treatment group) during the first year of the study. Also two people died in the control group, one an accidental death; the cause of the other death could not be ascertained.

3.4.6 Service use: We found the mean number of days spent in hospital at one year were not significantly different (n=507, WMD, -1.39 CI -2.8 to 0.1) and did indicate a trend favouring integrated treatment (p= 0.06), but by two and five years' follow-up the data revealed no significant differences. Also we found no significant difference in dichotomous data 'Not hospitalised' by five years (n=547, RR 1.05 CI 0.90 to 1.2).

3.4.7 Social outcomes: We found no significant differences in the numbers of people 'not living independently' by one year (n=507, RR 0.55 CI 0.3 to 1.2), and two-year data were also non-significant. However, by five years' follow-up, integrated treatment had significantly fewer participants 'not living independently' (n=547, RR 0.42 CI 0.21 to 0.8, NNT 19 CI 14 to 62) compared with the those receiving standard care. The numbers of participants who were either 'not working or in education' measured over one year showed no significant differences between the study groups, but by two years the integrated treatment group had significantly lower levels of not being in work or education (n=436, RR 0.72 CI 0.5 to 1.0, NNT 11 CI 7 to 99), compared with the control group. However by five years this advantage is not sustained and data were equivocal.

3.5 COMPARISON 10. Phase-specific treatment (CBT) + antipsychotics versus befriending + antipsychotics (all data from Jackson-Australia)

3.5.1 Leaving the study early: We found no significant difference in study attrition between treatment groups (n=62, 0.57 CI 0.2 to 1.8).

3.5.2 Hospitalised: No significant differences emerged in the number of participants being hospitalised over 12 months (n=62, RR 1.08 CI 0.59 to 1.99).

3.5.3 Suicide: Two people died due to suicide in the cognitive behavioural treatment group and none from the befriending group. This did not indicate a significant difference.

3.5.4 Social functioning: SOFAS: We found no significant difference in mean total endpoint scores from the SOFAS social functioning scale (n=62, RR 1.30 CI -6.3 to 8.9) by 12 months' assessment. Similarly, we found no significant differences in SOFRAS positive and negative symptoms scores.

3.6 COMPARISON 11. Phase-specific treatment E-EPA oils + atypicals versus placebo + atypical (all data from Berger-Australia)

3.6.1 Leaving the study early: For this short term study of 12 weeks we found study attrition to be equivocal (n=80, RR 0.83 CI 0.3 to 2.5).

3.6.2 Global state: not responded to treatment: We found that participants given the E-EPA oils in addition to antipsychotics had similar rates of non-response (18/40) to treatment compared with the control group (20/40) (n=80, RR 0.90 CI 0.6 to 1.4).

3.7 COMPARISON 12. Phase-specific treatment brief intervention + antipsychotics versus treatment as usual (all data from Leavey-UK)

3.7.1 Leaving the study: We found no significant differences in the number of people leaving the study early by nine months (n=106, RR 0.72 CI 0.3 to 1.5).

3.7.2 Hospital admission: Hospital admissions were reported before four months (n=106, RR 1.19 CI 0.9 to 1.6), and at up to four months (n=106, RR 0.75 CI 0.4 to 1.4) and between four and nine months (n=106, RR 0.86 CI 0.4 to 1.7); we found no significant difference between the intervention groups.

3.8 COMPARISON 13. Phase-specific treatment (ACE) + antipsychotics versus treatment as usual (all data from Uzenoff-USA)

3.8.1 Leaving the study early: We found no significant differences (n=24, RR 1.27 CI 0.3 to 6.3) at six months between ACE therapy and the control group.

3.8.2 Mental state: We found no significant differences for PANSS positive, negative, general or total scores (n=17, MD -1.57 CI -7.7 to 4.5). Depression rating from the CDRS also revealed no significant differences (n= 17, MD -1.46 CI -4.2 to 1.3).

3.8.3 Quality of life: We found quality of life scores as measured by the Heinrichs-Carpenter scale to be equivocal (n=16 MD -2.93 CI -25.6 to 19.7).

3.9 COMPARISON 14. Phase-specific treatment vocational intervention + TAU versus treatment as usual (all data from Killackey-Australia)

3.9.1 Not employed: We found participants who were given the vocational intervention were more likely to be employed (n=41, RR 0.39 CI 0.21 to 0.7, NNT 2 CI 2 to 4).

3.9.2 Leaving the study early: We found no significant differences in study attrition by six months (n=41, RR 0.21 CI 0.03 to 1.6).

3.10 COMPARISON 15. Phase-specific treatment (cannabis and psychosis therapy) + antipsychotics versus psychoeducation + antipsychotic (all data from Edwards-Australia)

3.10.1 Cannabis usage: We found no significant difference between the CAP intervention group and PE for use of cannabis at three months' assessment (n= 47, RR 1.04 CI 0.6 to 1.7). Data at nine months were also not significantly different (n=47, RR 1.30 CI 0.8 to 2.2).

The percentage of days using cannabis and the severity of cannabis usage were also reported by Edwards-Australia but data were skewed and are reported in 'other data' tables. Both outcomes were, however, non-significant.

3.10.2 Global state: KAPQ: We found no significant difference in the knowledge of psychosis and treatments between the participants at three months (n=47, WMD 0.80 CI -1.8 to 3.4) or nine months (n=47, WMD 0.90 -1.4 to 3.2).

3.10.3 Mental state

3.10.3.1 BPRS: We found average endpoint BPRS-extended scale scores at the three-month (n=47, WMD -3.60 CI -12.8 to 5.6) and nine-month assessment time point (n=47, WMD 0.80 CI -7.5 to 9.1) were not significantly different between the cannabis and psychosis therapy group, and those given psychoeducation. BPRS positive symptom scores were evaluated by Edwards-Australia, but data were skewed and are reported in 'other data' tables.

3.10.3.2 SANS: Data from the Scale for the Assessment of Negative Symptoms were skewed; the authors reported no significant differences in negative symptom scores between groups.

3.10.3.3 BDI: We found that the Beck Depression Inventory scores contained wide confidence intervals (skewed data) and have not reported these here.

3.10.4 Social functioning: SOFAS: We found no significant improvement in social functioning in the CAP group compared with PE at the three-month (n=47, WMD -0.80 CI -10 to 8.4) or the nine-month assessment (n=47, WMD -4.70 CI -14.5 to 5.1).

3.11 COMPARISON 16. Crisis assessment versus standard care (all data from LEO-CAT UK)

3.11.1 Hospitalisation: We found no significant difference in the number of participants being emitted to hospital between those who receiving crisis assessment and the control group who were given usual care (n=98, RR 0.85 CI 0.6 to 1.3).

3.11.2 Referred to mental health services: Crisis assessment did not result in significantly more participants being referred to mental health services, either by accident and emergency departments or emergency medical services (n=98, RR 0.85 CI 0.6 to 1.3).

3.12 COMPARISON 17. Early behavioural intervention versus routine care interval (all data from Alverez-Spain)

3.12.1 Weight: Measures of weight and body mass index were made. Data were skewed but we found no appreciable differences between the Early Behavioral Intervention and Routine Care Intervention groups in this small study (n=61).

3.13 COMPARISON 18. Phase-specific intervention versus control (Exploratory meta-analysis)

3.13.1 Leaving the study early: We pooled three studies in the meta-analysis (Jackson-Australia; Leavey-UK; Uzenoff-USA) and found no significant difference in the number of participants who left the study early (n=192, RR 0.72 CI 0.4 to 1.3).

3.13.2 Hospitalisation: Hospitalisation rates were equivocal between the phase-specific treatment and the control group (n=168, RR 0.97 CI 0.6 to 1.5) when we meta-analysed data from Jackson-Australia and Leavey-UK.

DISCUSSION

1. General

Studies were undertaken in the UK, Australia, Holland, Austria, Germany, Scandinavia, the USA and China. Six studies (Amminger-Austria; EDIE-UK; EIPS-Germany; LIPS-Germany; PACE-Australia; PRIME-USA) were concerned with preventing the development of psychosis in prodromal patients; 11 evaluated interventions for improving outcome in first-episode psychosis (Berger-Australia; Edwards-Australia; Jackson-Australia; Killackey-Australia; LEO-CAT-UK; Leavey-UK; LifeSPAN-Australia; Linszen-Amsterdam; OPUS-Scandinavia; Uzenoff-USA; Zhang-China).

Summary of main results

1. Trials to prevent the development of psychosis

1.1 COMPARISON 1. Phase specific intervention (olanzapine) + non-specific supportive therapy versus placebo + non-specific supportive therapy - PRIME-USA:

This study randomised a total of 60 people; we think this number is probably too small to detect a treatment effect. Attrition rates were not significantly different, although slightly more people did leave the olanzapine group by 12 months (65% by 12 months).

Olanzapine did not alter the numbers of people converting to psychosis over 12 months when compared with placebo. Also the Clinical Global Impression 'severity of illness' and the Global Assessment of Functioning 'current' change scores were both nonsignificant. Data were limited by study size, and larger groups may well have produced a different outcome. We did not find any significant outcome data from the SOPS scale during 12 months of evaluation. The PANSS, YMRS and MADRS scores were also equivocal, indicating that no real change in mental state occurred over a 12-month period for the olanzapine and placebo group.

For adverse effects, extrapyramidal symptoms were not more frequent in the olanzapine group compared with the placebo group (SAS, BAS and AIMS), even though olanzapine dosage levels were within the lower end of the normal dose range. Olanzapine did produce a statistically significant increase in weight compared with the placebo group. This limited data supports other recent reports of olanzapine's association with weight gain (Duggan 2005; Lieberman 2005). CoStart terms were also recorded and all were equivocal except for weight gain, with significantly more people gaining weight in the olanzapine group.

Cardiovascular measures were taken on blood pressure and pulse whilst sitting and standing over eight weeks with all data being equivocal. Twelve-month data (sitting) pulse rates were significantly lower in the placebo group, but this may have been a chance finding since all other data were non-significant. Fatigue was higher in the olanzapine group with a NNH of four.

1.2 COMPARISON 2. Phase specific intervention (cognitive behavioural therapy) + non-specific supportive therapy versus non-specific supportive therapy- EDIE-UK, EIPS-Germany):

Again there were few data to find (total n=188). We found no differences for numbers of participants leaving the study early by 12 months (n=60, RR 0.98 CI 0.4 to 2.2), and again at two years (n=60, RR 0.96 CI 0.6 to 1.5) and as a proxy measure of treatment acceptability, CBT did not enhance or worsen compliance in the (EDIE-UK) study.

The numbers of participants becoming psychotic over 12 months of observation were low and no significant differences between CBT (4/37) and the monitoring group (5/23) were found for this primary outcome (EDIE-UK).

Social functioning data were also non-significant (EIPS-Germany).

This approach, as with many others in this review, should be considered experimental.

1.3 COMPARISON 3. Phase-specific treatment (risperidone + cognitive behavioural therapy) + specialised team versus specialised team (PACE-Australia):

All participants (n=59) remained in the study for one year, which is unusual for randomised trials of this length (Summary of findings for the main comparison). The adherence to the study may have been due to participants being relatively well (i.e. prodromal) and also being cared for by a specialist team. This greatly limits applicability.

Initial findings from this comparison suggest that a phase-specific treatment combining risperidone and CBT can delay, but not prevent the onset of psychosis (Analysis 3.2). Whilst these findings are of interest, they are not definitive, as the single included trial (PACE-Australia) is substantially under-powered at the 12-month end-point. Moreover the use of a combination of phase-specific treatments makes it unclear how far each contributes to the outcome, though a sub-analysis by the trialists suggests that risperidone makes the primary contribution.

Global function (GAF) did not appear to have any effect on outcome in terms of global state, and again with such a small number of participants doubts will remain regarding efficacy. Delaying the onset of psychosis does not appear to have a substantial effect on medium-term outcome, in terms of mental state. This could be because the trial is under powered, or it may be that the benefits of delaying onset of psychosis are less than anticipated. It is also difficult to evaluate the benefits of delaying psychosis without more information on the impact of treatment from the perspective of service users and carers.

No improvements in quality of life occurred in the early intervention group, even though they were given CBT. However, they also received risperidone, which may have negated any gains in quality of life, due to its adverse effects profile.

1.4 COMPARISON 4. Phase specific intervention (amisulpride + needs focused interventions versus needs focused interventions (LIPS-Germany): Participants (total n=124) given both needs focused therapy and also amisulpride remained in the study more than the group given only needs focused intervention. This is not necessarily a beneficial outcome given the adverse effects of the amisulpride in a group of participants that are at risk of developing psychosis.

Global Assessment of Functioning scores favoured amisulpride plus NFI. Although the study reports data on global state of the prodromal participants, information on transition to psychosis would have been more useful in interpreting the value of such treatment combinations. Mental state measures in terms of PANSS global scores and PANSS-positive scores did favour amisulpride plus NFI. However, no significant differences were found for PANSS negative symptoms, or depression scores. Again, transition to psychosis data would have been more useful and are notable by their absence.

1.5 COMPARISON 5. Omega 3 fatty acids (EPA) versus placebo (Amminger-Austria): This study aimed to assess the effects of omega-3 fatty acids in preventing psychosis. The treatment group were significantly less likely to develop psychosis compared with the placebo group at 12 weeks. A NNT of 6 in a sample of 76 is an important result. This study should be replicated with a larger sample and participants followed for at least six months to determine whether a sustained treatment effect is present.

2. Trials to improve outcome in first-episode psychosis

2.1 COMPARISON 6. Phase-specific treatment (cognitive behavioural therapy for suicidality) + specialised team versus specialised team (LifeSPAN-Australia): The LifeSPAN study involved only 56 participants. Therapy did not affect the numbers of people leaving the study early over six months; larger sample sizes and perhaps longer study time may have produced less equivocal data. Two people committed suicide, one in each group. The study size is too small to determine whether LifeSPAN therapy can reduce suicide.

2.2 COMPARISON 7. Phase-specific treatment (family therapy) + specialised team versus specialised team (Linszen-Amsterdam): This was another important but small (n=76) study. Adding family therapy to care from a specialised team did not affect relapse rates, but Linszen-Amsterdam was substantially under powered, so that no definitive conclusions can be drawn. An unusual characteristic of the trial was that all participants had to consent to a three-month inpatient admission before randomisation. This may have limited the ability of the intervention to show an effect by excluding any differences in relapse rates occurring in the first three months after onset. It also limits applicability of findings to other early intervention services, which tend to be oriented towards reducing or avoiding admissions, rather than extending them (Edwards 2002).

2.3 COMPARISON 8. Phase-specific treatment (family therapy) + standard care versus standard care (Zhang-China): Retention of the 83 study participants over 18 months of care was good, with only 6% being lost to follow-up; the phase-specific intervention with family therapy did not prove either a benefit or hindrance to study attrition.

Family therapy in addition to standard care did reduce readmission rates and possibly helped compliance. Unfortunately other outcome data were not presented on an intention-to-treat basis and are impossible to use. The main limitation of this trial was the particular nature of the standard care given, which appeared to be a low-key form of outpatient treatment, with little continuity and no community follow-up. This makes it difficult to be certain how far the reduced admission rate in the intervention group was a non-specific effect of substantially increased contact with patients and their families, rather than a particular effect of family therapy. No data were available on how far the finding of fewer admissions was accompanied by improvements in outcome, or service user and carer satisfaction.

Family therapy did not appear, from limited participant numbers, to improve treatment compliance and larger sample sizes are needed to evaluate this outcome.

2.4 COMPARISON 9. Specialised team versus standard care (OPUS-Scandinavia):

This is the only large study in the review (n=547, Summary of findings 2). Integrated treatment significantly reduced the numbers of people leaving the study early by 12 months (NNT 9) and as a proxy measure of treatment acceptability was found to be more acceptable than placebo. By two years, attrition rates were still significantly favouring integrated treatment (NNT 7). It appears participants were prepared to remain in treatment longer when care was given from a specialist team, which for people with psychosis is an important outcome.

Global Assessment of Functioning 'symptom' scores significantly favoured integrated treatment by 12 months, but this was not sustained and two-year data were equivocal; GAF 'function' scores were equivocal at 12 months, but by two years did significantly favour integrated treatment. These outcomes seem to confound each other and more research is needed to adequately determine if this form of care can indeed improve global state.

Participants were significantly more satisfied with services in the integrated treatment group by 12 months, and this was sustained over two years. This result does fit with the positive findings for retention rates in the integrated treatment group. Overall, participants in the integrated care group were more compliant with treatment and outpatient visits than the standard care group. This effect was seen over one- and two-year time points, suggesting integrated treatment is more acceptable to people with first-episode psychosis than the standard care available to the control group. Again this result is consistent with attrition and user satisfaction outcomes.

Two people died from suicide, one person from each group.

We did not find any significant differences in the mean number of days per month participants spent in hospital and integrated treatment offered no advantages compared with the control group in terms of reducing the need for hospital care. Unfortunately, no data

were reported on relapse. Relapse is a primary outcome for this review and an important measure of treatment efficacy for people with first-episode psychosis, clinicians and health care managers.

We did not find integrated treatment to have any significant effect on participants' ability to live independently over one- and two-year assessments. Integrated treatment did significantly improve participants' employment and educational circumstances with 'not employed or in education' being lower by two years (NNT 11). However one-year data were non-significant, suggesting that two years of care are needed before benefits are obtained, although more data are needed to show this effect, especially as most outcome data were not significant.

OPUS-Scandinavia is an important study. It is a large study - but even it should be replicated.

2.5 COMPARISON 10. Phase-specific treatment cognitive behavioural treatment + antipsychotics versus befriending + antipsychotics (Jackson-Australia): Neither CBT or the befriending intervention managed to retain significantly more of the 62 participants over the 12 months of the study.

Although two deaths from suicide occurred, this did not suggest a significant difference between the two study groups.

We did not find advantage for CBT participants in their social functioning compared with the control befriending group.

2.6 COMPARISON 11. Specific intervention E-EPA oils + atypicals versus placebo + atypical (Berger-Australia): Neither intervention proved to be more advantageous in retaining the 80 participants within the study.

The number of people showing no response to treatment did not reveal any significant difference, in this short-term trial of 12 weeks' duration.

2.7 COMPARISON 12. Phase-specific treatment brief intervention + antipsychotics versus treatment as usual (Leavey-UK): At the end of the nine-month intervention, both treatments were similarly acceptable to the 106 participants, with no differences emerging in study attrition.

Over the course of nine months hospital admissions were not significantly lower in either intervention group.

2.8 COMPARISON 13. Phase-specific treatment (ACE) + antipsychotics versus treatment as usual (Uzenoff-USA): This was a study involving 24 participants. During the six months of the study, the interventions' study attrition was not significantly different.

We were able to report several mental state outcomes (PANSS positive, negative, general or total scores, depression rating scale scores), and all were non-significant. However, with

only 24 participants it is unlikely that any treatment effect would reach statistical significance.

Quality of life as measured using the Heinrichs-Carpenter Scale was equivocal.

2.9 COMPARISON 14. Phase-specific treatment vocational intervention + TAU versus treatment as usual (Killackey-Australia): The emphasis of this study of 41 people was the use of vocational-based interventions. Data revealed that the vocational training group was employed significantly more than the control group during the six months of evaluation.

No advantage emerged for either group in retaining participants within the study.

2.10 COMPARISON 15. Phase-specific treatment (cap) +antipsychotics versus psychoeducation (Edwards-Australia): The study (total n=47) aimed to minimise cannabis usage in people with first-episode psychosis. We found no significant difference between groups. The small sample size may have contributed to the lack of any treatment effect.

The participants understanding of psychosis, measured with the KAPQ questionnaire, did not reveal any differences at three and nine months' assessment. The use of an active control may have contributed to the non-significant finding. The BPRS-derived data on the positive symptoms of psychosis failed to reveal any significant benefits for CAP therapy compared with psychoeducation.

We found that neither CAP or psychoeducation improved social functioning over nine months in this small study; a larger scale trial would have increased the likelihood of finding statistically significant data.

2.11 COMPARISON 16. Crisis assessment versus standard care (LEO-CAT UK): This comparison involved 113 participants. From limited data, we found that hospital admissions were not increased or decreased for participants given crisis assessment. Similarly, participants given crisis assessment were no more likely to be referred to mental health services than the control group. Again, from such a small study it is possible that real effects of interventions are not being highlighted.

2.12 COMPARISON 17. Early behavioural intervention versus routine care interval (all data from Alvarez-Spain): As with most of the other trials, Alvarez-Spain was too small to really produce results with confidence. We are not sure if early behavioral intervention helps people whose weight is a problem (n=61).

2.13 COMPARISON 18. Phase-specific intervention versus control (Exploratory meta-analysis): Meta-analyses proved problematic because of the heterogeneity in the intervention and outcomes. No two studies were sufficiently alike to perform meta-analyses without reservations of their clinical meaningfulness. Only leaving the study early and hospitalisation data were pooled and in both cases no significant differences were found. An agreed set of core outcome measures in trials assessing early intervention for psychosis would increase the estimate of any treatment effects (COMET).

Overall completeness and applicability of evidence

From the comprehensive search, we included 18 studies, most of which have small sample sizes. OPUS-Scandinavia is the exception with over 500 people randomised. Additionally, only two studies were similar to be grouped together under the same comparison; the rest used different interventions or controls. These limitations are a source of uncertainty in our results.

1. Completeness—We had hoped to gather information from trials on a whole series of outcomes such as duration of hospital stay, employment status, quality of life and costs. We have identified a series of pioneering studies, none of which provide data on all outcomes, most of which are inappropriate to synthesise together and therefore data are incomplete for every outcome for every one of the many comparisons in this review.

2. Applicability—Six of the 18 studies were undertaken at the EPPIC Centre in Melbourne, Australia and work embedded in this region's excellent, pioneering and world-class services may not be widely applicable. However, and understandably, many of the other trials were undertaken by leading figures in the world of early intervention. Again, services within these studies may be better than is standard. This would serve to narrow any differences between intervention and control group.

Often in meta-analyses we have data from trials undertaken all around the world, in different cultures of care, synthesised. In these cases basically homogeneous findings would point to wide applicability of findings. In this review, however, meta-analyses have been very limited and, therefore, we have to remain very cautious regarding applicability.

Quality of the evidence

All studies were randomised, although in terms of allocation concealment, the quality of included studies was acceptable but not good, since precise details of the method of randomisation were lacking for most studies. One study (EDIE-UK) attempted to blind raters, but this proved difficult and was not adequately maintained as participants 'divulged' sufficient information to inform the rater which treatment participants were receiving. Edwards-Australia, Zhang-China and LifeSPAN-Australia also used raters who were blind to group allocation, but they did not report whether allocation concealment was maintained. One study (PRIME-USA) did use double-blind methodology, made possible by both groups receiving the same psychosocial intervention with the variable being medication (olanzapine or placebo), which was easier to blind for than non-pharmaceutical interventions. OPUS-Scandinavia, Linszen-Amsterdam and PACE-Australia used independent raters not blinded to treatment. The nature of these therapies is not suited to the raters being blind to treatment allocation and where this was attempted (as with EDIE-UK) it proved difficult to maintain throughout the study. In two trials (Linszen-Amsterdam; Zhang-China) key data were presented in a way that did not permit an intention-to-treat analysis on most outcomes. Rates of follow-up were particularly good in two trials (PACE-Australia; Zhang-China) and unclear (but probably acceptable) in one (Linszen-Amsterdam). Numbers of people lost to follow-up was not excessive in the other trials, but not good - EDIE-UK 70%, PRIME-USA 65%, OPUS-Scandinavia 77% and LifeSPAN-Australia 75%. One study did find significant

differences in follow-up rates between treatment and controls (OPUS-Scandinavia); the effects of this are unclear but may have had an impact on the findings of this review. A substantial omission from most trials was an attempt to capture the perspective of service users and their carers, by, for example, using satisfaction scales.

Potential biases in the review process

We have been able to identify only published reports; by doing this, we may be perpetuating reporting and publishing biases. It is possible that small studies have not been published, but we feel that large scale published studies would have been identified.

Also, we now have considerable knowledge of these trials and it is possible that we view the data in a way that others could consider biased or skewed in some way. We are keen to receive feedback via peer review before and after publication.

Agreements and disagreements with other studies or reviews

This review updates earlier versions (Marshall 2004).

AUTHORS' CONCLUSIONS

Implications for practice

The current update now includes 18 studies; however, due to the variety of interventions used we have added only two studies under the same comparison category, but because they did not measure the same outcomes we have not synthesised the data. We are, therefore, unable to make any firm recommendations for practice.

1. For people presenting with prodromal symptoms of psychosis—At the moment it is not clear whether treating people presenting with prodromal symptoms of schizophrenia provides benefits. There is inconclusive evidence on the personal and social consequences of providing treatment to people who will not necessarily become unwell. Further evidence is needed before recommendations can be given.

2. For people in their first episode of psychosis—There is some support for specialised early intervention services but again further evidence is needed. However, since such people do require treatment in some form, the ethical issues are less intense than for people presenting with prodromal symptoms. Moreover, there is also little evidence to support the 'standard care', which is the alternative to the employment of specialised first-episode teams (NICE 2002). The use of first-episode teams is therefore ethical even though there is not, as yet, strong evidence to support it.

Phase-specific treatments for people in their first episode of psychosis may help with employment and family therapy. Whilst this evidence is limited, it should be viewed in the broader context that family therapy is known to be effective for people with schizophrenia as a whole (Pharoah 2006). On this basis, it would seem reasonable to recommend family therapy to people experiencing their first episode of psychosis, but there is insufficient data to suggest that they should be given this intervention as a priority over people with established illness.

There is no evidence from clinical trials to support the benefits of early detection of patients in their first episode of psychosis.

3. For clinicians—Family intervention may be of value for people in their first episode of psychosis, as it may for people with longer established illnesses. It is important for clinicians to continue to keep up to date with this fast-expanding field.

4. For policy makers—It is premature to implement wide-spread treatment programmes for people with prodromal symptoms. Such treatment programmes should only be implemented within the context of a well-designed randomised study.

Implications for research

1. General—If CONSORT recommendations (Begg 1996; Moher 2001) had been followed by authors of the included studies and the editors of the journals in which those reports were published, the effects of early intervention for psychoses would be more evident.

2. Specific—This review has identified a discrepancy between the global rate of growth of early intervention services and the paucity of underpinning evidence. Whilst there is a compelling theoretical case for early intervention, much of the supporting evidence is circumstantial (based on the correlation between duration of untreated psychosis and outcome) rather than definitive (based on improved outcome in clinical trials). If this discrepancy persists, the obvious risk is that, eventually, early intervention will become routine practice, without its efficacy ever being definitively established. Whilst this review has found some evidence of a growing body of research in the field, there is no room for complacency over the amount of work that needs to be done. Possible combinations of early detection, type of participant (prodromal or first episode), type of control, and type of intervention (phase-specific or specialised team) generate at least 17 possible types of trial. However, the review has identified 18 included trials, most of which are clearly under-powered. The current substantial international interest in early intervention offers an opportunity to make major positive changes in psychiatric practice, but this opportunity may be missed without a concerted international programme of research to address key unanswered questions.

These key questions are:

- Can phase-specific treatments prevent people with prodromal symptoms from developing psychosis and, if so, do they or their carers benefit as a result?
- Can early detection reduce the duration of untreated psychosis, and if so, does this lead to improvements in outcome for service users and carers?
- Are there phase-specific treatments that improve outcome for people with first-episode psychosis, or for their carers?
- Do specialised early intervention teams offer improvements in outcome over and above those provided by phase-specific treatments alone?

These questions give rise to two important points, which if borne in mind at the design stage, might increase the value of future trials in the field. Firstly, a phase-specific treatment should not be a priority for investigation unless it is known to be substantially different from existing interventions that are already known to be helpful to people at all stages of schizophrenia. For example, there is little point in investigating the effects of behavioural family therapy with minor modifications for first-episode patients, when this intervention is known to be generally effective in schizophrenia (Pharoah 2006). Phase-specific treatments ought to be given priority for evaluation only when they are substantial departures from what would be considered standard care, or where there is evidence that they are likely to be more effective when offered in the early stages of the illness.

Secondly, great care must be taken in defining the characteristics and activities of specialised early intervention teams. The complexity of 'early intervention' makes it likely that no two specialised teams will be identical. Unless the essence of an early intervention team can be adequately characterised, it is inevitable that disappointing findings will lead to arguments over whether a particular specialised team was really practising early intervention. Years of research effort can be wasted in this way. Lessons should be learned from research which has already been undertaken with other specialised psychiatric teams (such as assertive outreach teams) and fidelity scales developed as an early priority.

Acknowledgments

Mark Fenton and Clive E Adams of the Cochrane Schizophrenia Group for conducting the initial database search. Many thanks to Sarah Uzenoff for providing additional data.

Many thanks to Austin Lockwood - who co-ordinated the review and collected data, screened the search results, organised the retrieval of papers, appraised the papers and extracted data, managed the data and entered data in RevMan, analysed and interpreted the data and wrote the final report, during the first publication of the review (Marshall 2004).

The Cochrane Schizophrenia Group produces and maintains a template Methods section. We have used this and adapted it to make it relevant for our review.

SOURCES OF SUPPORT

Internal sources

- Lancashire Care NHS Trust, UK.

External sources

- Department of Health, UK.
- Cochrane Incentive Scheme, UK.

CHARACTERISTICS OF STUDIES

Characteristics of included studies *[ordered by study ID]*

Alvarez-Spain

Methods	Allocation: randomised (by computer random numbers into 4 blocks). Blinding: single blind.
---------	---

Setting: community.
Duration: interventions offered for 3 months.

Participants	Diagnosis: drug-naive first-episode psychosis (DSM-IV). N=61. Sex: 46 M, 15 F. Age: mean 26 years. Inclusion criteria: 15 to 60 years, DSM criteria for schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, brief reactive psychosis, or psychosis not otherwise specified. Excluded: patients with neurological disease, head injury, mental retardation, and drug dependence. History: all patients experiencing their first episode of psychosis and had not received more than 6 weeks of adequate antipsychotic treatment
Interventions	1. Early Behavioural Intervention*. N=28. 2. Routine Care Intervention**. N=33. Interventions started when participants reached a minimum clinical stabilisation, defined by a global score of < 5 on the SAPS scale, or otherwise treatment began within 6 weeks of randomisation
Outcomes	Body weight. Body Mass Index. Unable to use - Leaving the study early- no data. Mental state: SAPS (no data reported).
Notes	* This package of care was specifically designed to teach strategies to enhance control over factors associated with antipsychotic-induced weight gain and consisted of 8 flexible intervention modules that incorporated behavioural interventions, nutrition and exercise. ** The control group were informed about potential weight gain and advised to increase their exercise and limit food intake. Participants were first randomised to 3 antipsychotics (olanzapine mean dose 13 mg/day, risperidone mean dose 4.2 mg/day or haloperidol mean dose 4.9 mg/day), then randomly assigned to either EBI or RCI

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Low risk	Randomised using computer generated random numbers
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Single
Incomplete outcome data addressed? All outcomes	High risk	Study attrition not reported
Free of selective reporting?	High risk	SAPS data not reported
Free of other bias?	Unclear risk	No details.

Amminger-Austria

Methods	Allocation: randomised, no further details. Blinding: double blind. Setting: Vienna, Austria. Inclusion criteria: not stated. Exclusion criteria: not stated. Follow up: 12 weeks.
Participants	Diagnosis: Adolescents at risk of 1st episode psychosis (Yung criteria). N=81*. Age: range 3-24 years; mean age 16.4. Sex: no details. History: no details.

Interventions	1. Omega-3 fatty acids: dose eicosapentaenoic acid 0.84 g/day; docosahexaenoic acid 0.7 g/day. N=38. 2. Placebo. N=38.
Outcomes	Transition to psychosis**. Unable to use - Leaving the study early Mental state: BPRS, PANSS (no usable data) Global state: GAF (no usable data). Adverse effects: UKU.
Notes	*5 participants not accounted for. **Operationally defined, based on Yung et al's criteria, using cut-off points on PANSS subscales, (4 or more on hallucinations, 4 or more on delusions, and 5 or more on conceptual disorganization), and the frequency of symptoms (at least several times a week) and their duration (more than one week)

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No described.
Blinding? All outcomes	Unclear risk	Double blind, untested.
Incomplete outcome data addressed? All outcomes	High risk	Study attrition not described.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	No details.

Berger-Australia

Methods	Allocation: randomised. Blinding: double blind. Setting: single centre Melbourne, Australia (EPPIC Centre). Inclusion criteria: age between 15-29 years, currently psychotic. Exclusion criteria: drug induced psychosis; first-episode mania; psychotic disorders due to organic illness. Follow-up: 12 weeks.
Participants	Diagnosis: psychosis (DSM-IV). N=80. Age: mean 20 years. Sex: male and female. History: first-episode psychosis; mean antipsychotic treatment prior to study 17.5 days
Interventions	1. Ethyl-Eicosapentaenoic Acid oil (E-EPA): dose 500 mg/bid, plus flexible dosage of atypical antipsychotics. N=40. 2. Placebo capsules: plus flexible dosage of atypical antipsychotics. N=40. Benzodiazepines, chlorpromazine or zuclopenthixol acetate allowed for behavioural control when indicated
Outcomes	Leaving the study early. Global state: Not responded to treatment. Unable to use - Mental state: BPRS, SANS, MADRAS (no data). Global state: CGI, GAF (no data). Social functioning: SOFAS (no data). Adverse events: SAS, UKU (no usable data). Social functioning: SAS II.
Notes	
Risk of bias	

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	Unclear risk	Double blind, untested.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Funded by Swiss National Science Foundation; Margaret and Walter Lichtenstein Foundation, National Health and Medical Research Council of Australia, the Colonial Foundation of Australia, and a National Alliance for Research on Schizophrenia and Depression

EDIE-UK

Methods	Allocation: randomised, stratified according to gender and genetic risk (independent clerical worker, sealed envelopes). Blinding: single blind (raters), attempts made to keep assessors blind*. Setting: community, Salford, Manchester. Inclusion criteria: based on an adaptation of the PACE criteria, and an age range between 16-36. Exclusion criteria: current of past receipt of antipsychotic medication. Follow-up: 1 year and 3 years. Evaluation: conducted by research assistants.
Participants	Diagnosis: ultra high risk of developing 1st episode of psychosis (Yung modified criteria). N=60. Age: mean 21 years. Sex: male and female. History: not reported.
Interventions	1. Cognitive therapy: dose maximum of 26 sessions, over 6 months. N=37 2. Monitoring control group. N=23.
Outcomes	Leaving the study early. Transition to psychosis (based on PANSS criteria). Unable to use - Transition to psychosis (3 year data, 53% lost to follow-up). Mental state: PANSS (no usable data). Global state: GAF, GHQ (no usable data). Sociotropy - Autonomy Scale (no usable data). Meta-Cognitions Questionnaire (no usable data). Oxford-Liverpool Inventory of Feelings and Experiences (no usable data)
Notes	*Blinding was not adequately maintained due to participants divulging information about their therapist, or used language that suggested they were receiving cognitive therapy. Three-year outcome data not used due to > 50% study attrition

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Low risk	By independent clerical worker, using sealed envelopes.
Blinding? All outcomes	High risk	Single blind.
Incomplete outcome data addressed?	Low risk	Study attrition reported.

All outcomes		
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Funded by North-West NHS Executive.

Edwards-Australia

Methods	Allocation: randomised, computer generated, placed in sealed envelopes. Blinding: single; attempts to maintain rater blindness included use of separate rooms and admin procedures for staff. Duration: 3 months intervention phase followed by 6 months of follow-up. Country: Melbourne, Australia. Setting: youth mental health service; early psychosis, prevention and intervention centre (EPPIC). Analyses: ITT analyses were used and last observation carried forward
Participants	Diagnosis: first-episode psychosis (DSM-IV). N=47. Age: 15-29 years. Sex: male and female. History: patients continuing to use cannabis after initial treatment for first episode psychosis. Inclusion criteria: DSM-IV diagnosis of a psychotic disorder (i.e. schizophrenia, schizophreniform, schizoaffective, delusional disorder, bipolar disorder, major depressive disorder with psychotic features, psychosis not otherwise stated, brief reactive psychosis). Exclusion criteria: only participants with at least 10 weeks continuous cannabis usage prior to study were eligible for study inclusion
Interventions	1. Cannabis and Psychosis Therapy: mean no. of CAP sessions 8. N=23 2. Psychoeducation: mean no. of sessions 8. N=24.
Outcomes	Behavioural: CASUAS. Mental state: BPRS, SANS, BDI-SF. Social functioning: SOFAS. Global state: KAPQ.
Notes	CAP therapy consisted of a cognitive-behavioural-orientated programme delivered in weekly sessions by trained clinicians over 3 months Psychoeducation was an active control.

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Low risk	Computer generated randomisation.
Allocation concealment?	Unclear risk	Insufficient details.
Blinding? All outcomes	High risk	Single blind.
Incomplete outcome data addressed? All outcomes	High risk	Study attrition not reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	No details.

EIPS-Germany

Methods	Allocation: randomised computer generated by block using sealed envelopes. Blinding: not stated. Setting: community, Cologne, Bonn, Dusseldorf, and Munich. Inclusion criteria: EIPS criteria.
---------	---

	Exclusion criteria: attenuated or brief limited intermittent psychotic symptoms; present or past diagnosis of schizophrenia; alcohol or drug dependence. Follow-up: 12 months therapy.
Participants	Diagnosis: people at risk of developing 1st episode of psychosis. N=128*. Age: mean age ~26 years. Sex: male and female. History: not reported.
Interventions	1. Cognitive Behavioural Therapy: sessions over 12 months, group therapy session n=15, individual therapy session n=25, plus cognitive remediation 12 sessions. N=54. 2. Supportive Therapy: sessions over 12 months, minimal support involving psychoeducation and counselling. N=59
Outcomes	Social functioning: SAS II. Unable to use - Leaving the study early - no usable data.
Notes	*15 participants not accounted for after randomisation. EIPS - early initial prodromal state

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Low risk	Randomised by computer generated numbers.
Allocation concealment?	Unclear risk	Using sealed envelopes.
Blinding? All outcomes	High risk	Not stated.
Incomplete outcome data addressed? All outcomes	High risk	Study attrition only part reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Funded by the German Federal Ministry of Education and Research

Jackson-Australia

Methods	Allocation: randomised (stratified according to affective and non-effective psychosis, by independent statistician). Blinding: single blind. Setting: participant's home, neutral location or EPPIC centre (Psychosis Prevention and Intervention Centre) Melbourne, Australia. Follow-up: one year. Inclusion criteria: Exclusion criteria: IQ < 70, psychosis due to a medical condition; exhibiting violent behaviour. Evaluation: research assistants blind to treatment.
Participants	Diagnosis: psychosis. N=62. Age: mean 22 years. Sex: 45 M, 17 F. History: mean onset age of psychosis 22 years.
Interventions	1. Cognitive Behavioural Therapy (ACE): dose 20 sessions* for 45 minutes, plus neuroleptics. N=31. 2. Befriending, plus neuroleptics. N=31.
Outcomes	Leaving the study early. Suicide. Hospital admission (not hospitalised). Social functioning: Social and Occupational Functioning Assessment Scale (SOFRAS).

Notes
*Participants could receive a maximum of 20 sessions over 14 weeks for approximately 45 minutes.
ACE - Active Cognitive Therapy for Early Psychosis.

Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Low risk	Randomised by independent statistician.
Blinding? All outcomes	High risk	Single, untested.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Funded by NH&MRC grant.

Killackey-Australia

Methods	Allocation: randomised, computer generated. Blinding: not stated. Setting: Early Psychosis Prevention and Intervention Centre (EPPIC) in Melbourne, Australia. Follow-up: 6 months. Inclusion criteria: Individuals were eligible for the study if they wanted to find work (including a different job if they currently held one) and had at least 6 months of care left at EPPIC (EPPIC is limited to providing 18 months of care). Exclusion criteria: lack of fluency in English.
Participants	Diagnosis: schizophrenia 1st episode. N=41. Age: mean 21 years. Sex: 33 M, 8 F. History: length of illness IPS 16 months, TAU 12 months.
Interventions	1. IPS plus TAU. N=20. 2. TAU. N=21.
Outcomes	Employment. Leaving the study early.
Notes	*IPS is focused on competitive employment; it is open to any person with mental illness who chooses to look for work and acceptance into the programme is not determined by measures of work-readiness or illness variables; the support provided in the programme is time-unlimited, continuing after employment is obtained, and is adapted to the needs of the individual

Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Low risk	Randomised, computer generated.
Allocation concealment?	Unclear risk	Not reported.
Blinding? All outcomes	Unclear risk	Not reported.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	No details

Leavey-UK

Methods	Allocation: randomised by block and placed into sealed envelopes, and drawn by individual with no connection to the study. Blinding: single blind. Setting: outpatients, London, UK. Follow-up: 9 months. Inclusion criteria: people with a 1st episode psychosis within the last 6 months. Exclusion criteria: organic disorders, or learning difficulties. Evaluation: assessment made by research assistants blind to allocation
Participants	Diagnosis: first-episode psychosis (ICD-9). N=106 patients and their carers*. Age: 16 + years. Sex: 68 M, 38 F. History: 1st episode psychosis, recruited from psychiatric services in North London
Interventions	1. Brief intervention and treatment as usual. N=57. 2. Treatment as usual with usual support from psychiatric services. N=49. Brief intervention was provided over seven sessions, lasting about one hour and included information gathering from the relative, an educational component on psychotic illness, symptoms and early warning signs, treatment, and help seeking; coping strategies, problem solving and communication with the patient
Outcomes	Leaving the study early. Hospital admission. Unable to use - Satisfaction: Verona Service Satisfaction Questionnaire (no usable data). Perceived severity of illness.
Notes	*Carers/relatives were blind to treatment allocation. ITT analysis used.

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Low risk	Using sealed envelopes, drawn by individuals with no connection to the study
Blinding? All outcomes	Low risk	Single, untested.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Funded by NHS Executive, London Research and Development Programme

LEO-CAT-UK

Methods	Allocation: cluster randomised with 46 GP practices. Blinding: open study. Setting: community health centre, London, UK. Follow-up: over 12 months.
Participants	Diagnosis: psychosis (CAARMS*) N=113. Age: mean age 23 years. Sex: 81 M, 32 F. Excluded: participants with a history of contact with mental health services for psychosis for more than 6 months or antipsychotic treatment for more than a month.

	History: first-episode psychosis.	
Interventions	1. GP education in early detection with access to LEO-CAT. N=23 practices, N=50 participants. 2. Usual care, with access to LEO-CAT. N=23 practices, N=63 participants	
Outcomes	Hospitalisation. Referred to Mental Health Services by Accident and Emergency or Emergency Health services	
Notes	*Comprehensive Assessment of At-Risk Mental States (CAARMS) (Yung 2005). LEO-CAT- Lambeth Early Onset Crisis Assessment Team. GP practices randomised into the intervention group received both the GP education training and direct access to the LEO CAT team for referrals.	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Open study.
Incomplete outcome data addressed? All outcomes	High risk	Study attrition not reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	No details.

LifeSPAN-Australia

Methods	Allocation: randomised (no further details). Blinding: single, no further details. Setting: Early Psychosis Prevention and Intervention Centre (EPPIC). Follow-up: 10 weeks and 6 months. Inclusion criteria: scoring from 4 to 7 on the expanded version of the BPRS suicidality subscore. Exclusion criteria: attended the EPPIC centre for more than one year. Evaluation: 'conducted blind to therapy'.
Participants	Diagnosis: first-episode psychosis and acutely suicidal. N=56. Age: range 15-29 years. Sex: not reported. History: not reported.
Interventions	1. LifeSPAN therapy: dose 8 to 10 sessions + standard clinical care. N=31 2. Standard care. N=25. LifeSPAN is a brief individual cognitively orientated therapy specifically designed for acutely suicidal youths with severe mental illness
Outcomes	Leaving the study early. Death from suicide. Unable to use - Mental state: BPRS, SANS: (no usable data). Global state: GAF: (no usable data). Quality of life: (no usable data). Beck Hopelessness Scale: (no usable data). Self Esteem Scale: (no usable data). Self Report Problem Solving Rating Scale: (no usable data). Suicide Ideation Questionnaire: (no usable data). Suicide Intent Scale: (no usable data). Reasons for Living Inventory: (no usable data).
Notes	
Risk of bias	

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Single blind.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Funded by the Government of Australia.

Linszen-Amsterdam

Methods	Allocation: 'randomly assigned'. Blinding: unclear if raters blind to treatment condition. Setting: In-patient unit for adolescents, Amsterdam, Netherlands. Follow-up: 12 months (following on from initial 3 month inpatient admission), then 5 years. Inclusion criteria: first episode of schizophrenia, age 15-26, living or in close contact with parents or other relatives, Dutch speakers, no primary drug problem. Evaluation: 'independent raters'.
Participants	Diagnosis: schizophrenia. N=76. Age: mean 20.6. Sex: M 53 F 23. History: mean DUP 5.4 months.
Interventions	1. Behavioural family therapy + individual-orientated therapy. N=37 2. Individual-orientated therapy alone. N=39.
Outcomes	Relapse. Unable to use - Mental state: BPRS: (data not reported). Compliance: (data not reported). Lost to follow up: (exact figures unclear, though no evidence of substantial loss)
Notes	First Episode Trial - care from a specialised team plus phase specific intervention versus care from specialised team Unclear when randomisation took place, possible the initial sample size was 97, in which case not intention to treat Five year data reported for whole sample, not by group allocation

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	No details.
Incomplete outcome data addressed? All outcomes	High risk	Study attrition not reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	Part funded by 'Praeventiefonds'.

LIPS-Germany

Methods	Allocation: randomised, no further details. Blinding: open label. Setting: Germany, outpatient clinic. Follow-up: two years. Evaluation: not blinded.
Participants	Diagnosis: prodromal at risk of psychosis. N=124. Age: mean age 25 years. Sex: M 70, F 54. Inclusion criteria: adapted ERraos criteria, age range restricted between 18-36 years. Exclusion criteria: any lifetime DSM-IV diagnosis of schizophrenia, bipolar disorder, any DSM-IV diagnosis of brief psychotic episode with a duration of more than 1 week; delirium, dementia, and other cognitive disorders, mental retardation, mental disorders due to a general medical condition or mental disturbances due to psychotropic substances, alcohol abuse or drugs within the past 3 months. History: no details.
Interventions	1. Needs Focused Intervention* with amisulpride mean dose 118 mg/day. N=65. 2. Needs Focused Intervention. N=59. Use of chloral hydrate or short-acting benzodiazepines were allowed to treat agitation or sleep disturbances, and biperiden permitted for EPS. Citalopram permitted for depression
Outcomes	Leaving the study early. Mental state: PANSS. Global state: GAF. Adverse events.
Notes	*Needs Focused Intervention included psychoeducation, crisis intervention, family counselling and assistance with education or work-related difficulties according to the patient's need

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Open label.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	High risk	Funded by German Federal Ministry for Education and research BMBF grant and Sanofi Synthelabo, Germany

OPUS-Scandinavia

Methods	Allocation: randomised (computer computer-generated ratio of 1:1 in blocks of 6, and stratified for each of 5 centres. In Aarhus, the researchers contacted a secretary by telephone when they had finished the entry assessment of each patient. The secretary then drew 1 lot from among 5 red and 5 white lots out of a black box. When the block of 10 was used, the lots were redrawn. Block sizes were unknown to the investigators). Blinding: raters not blind to treatment allocation, but independent of study group; raters at the 5-year follow-up were blinded to patients' previous treatment allocation. Setting: Copenhagen, Denmark, multicentre, 5 centres, participants visited in their homes or other places in the community, or at their primary team members office. Exclusion criteria: taking antipsychotics for more than 12 weeks. Duration: 2 years with 5 year follow-up.
---------	---

	Evaluations: by independent investigators, not blinded to treatment allocation
Participants	Diagnosis: first-episode schizophrenia spectrum disorder (ICD 10, codes in the F2 category). N=547. Age: range 18-45 years. Sex: M 256, F 291. History: participants included in and outpatients who had not received antipsychotics for more than 12 weeks continuously
Interventions	1. Integrated treatment. N=275. 2. Treatment as usual. N=272. Integrated treatment is an assertive community treatment, enhanced by better specific content via family involvement and social skill training. Treatment as usual consisted of care at a community mental health centre. The treatment was carried out for two years and patients follow up for a further 3 years All participants were offered antipsychotic drugs according to guidelines from the Danish Psychiatric Society, which recommends low dose, atypical antipsychotic strategy for 1st episodes of psychotic illness
Outcomes	Leaving the study early. Global state: GAF. Client Satisfaction: CSQ-8. Suicide attempts. Social outcome: Social Network Schedule, not living independently; no working or in education. Service utilisation: average number of days in hospital. Compliance with treatment. Unable to use - Mental state: SAPS, SANS: (no usable data).

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Low risk	Randomised, computer generated.
Allocation concealment?	Low risk	Secretary drew lots, which the investigators had no knowledge
Blinding? All outcomes	Low risk	Independent assessors aware of treatment allocation; 5-year follow-up assessors unaware of treatment allocation
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Low risk	Grants from the Danish Ministry of Health, Danish Ministry of Social Affairs, University of Copenhagen, Copenhagen Hospital Cooperation, Danish Medical Research Council, Wørners Foundation, and the Stanley Wada Research Foundation

PACE-Australia

Methods	Allocation: simple randomisation by study co-ordinator. Blinding: independent rater, not blind. Setting: PACE clinic (Personal Assessment and Crisis Evaluation), part of EPPIC program, Melbourne, Australia. Inclusion criteria: age 14-30, living in Melbourne, met one of 3 criteria for an Ultra High Risk mental state. Follow-up: 0, 6, 12 months.
Participants	Diagnosis: 'ultra high risk' of developing psychosis.* N=59. Age: mean 20.

	Sex: M 34, F 25. History: not reported.	
Interventions	1. Specific Preventive Intervention: dose risperidone 1-2 mg/day + cognitive behavioural therapy + needs-based case management + supportive psychotherapy. N=31 2. Needs based intervention alone. N=28.	
Outcomes	Progressing to psychosis. Mental state: BPRS, HRSA, HRSD, SANS, YMS. Quality of Life: QLS. Overall functioning: GAF.	
Notes	Prodromal Trial - Care from a specialised team plus a phase specific intervention versus care from a specialised team * Defined as either: family history of psychotic disorder & non specific symps & decrease in functioning on GAF of 30 points or more in last 12 ms, or attenuated psychotic symptoms sustained for at least 1 week, or brief episodes of psychotic symptoms not sustained beyond a week	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Not blinded.
Incomplete outcome data addressed? All outcomes	High risk	Study attrition not reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	Funding by the Commonwealth Government of Australia Research and Development Grants Advisory Committee, and Janssen-Cilag Pharmaceuticals; Australian Rotary Health Research Fund grant

PRIME-USA

Methods	Allocation: randomised. Blinding: double, pills dispensed in prepackaged packs, prelabelled by site number and sequential subject number within site. Setting: North America, 4 sites*, outpatient clinic. Inclusion criteria: age range 12-45 years. Exclusion criteria: past or current DSM-IV psychotic disorder; suffering from a psychiatric disorder that could account for the prodromal symptoms; judged to be suicidal or homicidal; prodromal symptoms due to drug or alcohol use. IQ less than 80; seizure disorders. Follow-up: one-year medication with one-year follow-up without medication. Evaluation: 'patient, investigator, prescriber and rater were maintained blind to group assignment throughout the study'
Participants	Diagnosis: prodromal at risk of psychosis (SIPS, SOPS). N=60. Age: range 12-36 years, mean 18 years. Sex: M 39, F 21. History: participants included those who had responded to advertisements, or were referred by clinicians. Exclusion criteria: past or current psychotic disorder, a treatable psychiatric disorder that could account for the prodromal symptoms, suicidal or homicidal ideation, or drug or alcohol use that could be responsible for their symptoms. Duration: 12 months.
Interventions	1. Olanzapine: dose 5-15 mg/day, mean 8 mg/day. N=31. 2. Placebo. N=29.

Olanzapine was adjusted within a range of 5-15 mg/day based on the clinician's judgement. Individual and family psychosocial interventions were available. Lorazepam (max 8 mg/day) diazepam (max 40 mg/day) and chloral hydrate (max 100 mg/day) were used for agitation and/or insomnia. Benztropine mesylate or biperiden up to 6 mg/day allowed to treat EPS. Nizatidine 300-600 mg/day for weight gain, beginning towards the end of the study

Outcomes	Leaving the study early. Progressing to psychosis: POPS scale. Mental state: PANSS, MADRS, YMRS, SOPS. Global state: CGI-S, GAF. Adverse effects: SAS, AIMS, BAS, weight gain, vital signs, CoStart terms Unable to use - Quality of life: no data. Resource utilisation: no usable data. Adverse effects: EPS (no usable data). Neurocognitive function: no data. Premorbid functioning: Cannon-Spoor Premorbid Adjustment Scale
Notes	*Yale University, New Haven, Connecticut; University of Toronto, Canada; University of North Carolina, USA; University of Carolina, Canada

Risk of bias

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Low risk	Pills dispensed in prepackaged packs, prelabelled by site number and sequential subject number within site
Blinding? All outcomes	Unclear risk	Double blind, untested.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	High risk	Funding Eli Lilly and Company.

Uzenoff-USA

Methods	Allocation: randomised. Blinding: assessors blind to treatment group. Setting: no details. Inclusion criteria: 16 years or older, DSM -IV schizophrenia, schizoaffective disorder or schizophreniform disorder and in treatment of a first episode of less than 12 months. Follow-up: 6 months. Evaluation: 3 and 6 months.
Participants	Diagnosis: schizophrenia first episode (DSM IV). N=24. Age: no details. Sex: men and women. History: no details.
Interventions	1. Adherence Coping Education* (ACE) with usual care. N=13. 2. Supportive therapy* with usual care. N=11.
Outcomes	Leaving the study early. Mental state: PANSS, CDPS, CDRS. Quality of life: Heinrichs-Carpenter.
Notes	*ACE 14 sessions lasting between 30-45 minutes over six months. It is a manual-based psychotherapy consisting of 4 phases: (1) establishing therapeutic alliance; (2) promoting treatment adherence; (3) developing a plan for maintenance treatment and (4) rehabilitation.

*Supportive therapy had two phases (1) establishment of the therapeutic relationship and (2) provide emotional support and discussion of non-illness issues or topics.
A modified ITT analysis was used based on 19 participants.

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Not blinded.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.
Free of other bias?	Unclear risk	No details.

Zhang-China

Methods	Allocation: 'randomly assigned'. Blinding: not reported. Setting: psychiatric hospital, Suzhou, China. Inclusion criteria: male, just discharged after first episode for schizophrenia, no other medical conditions. Follow-up: 18 months. Evaluation: by 'attending physicians' blind to allocation.
Participants	Diagnosis: schizophrenia (Chinese Medical Association Criteria). N=83. Age: mean 23.8. Sex: all male. History: mean DUP 34.6 months.
Interventions	1. Family psychoeducation in individual and group sessions plus standard out-patient care. N=42 2. Out-patient care. N=41.
Outcomes	Readmitted. Lost to follow-up. Compliant with medication. Unable to use - Chlorpromazine equivalent dosage of medication: (not a clinical or social outcome). Mental state: Chinese BPRS (excluded readmitted patients). Overall functioning: Chinese GAS (excluded readmitted patients)
Notes	First Episode Trial - phase-specific treatment plus standard care versus standard care

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Randomised, no further details.
Allocation concealment?	Unclear risk	No details.
Blinding? All outcomes	High risk	Not reported.
Incomplete outcome data addressed? All outcomes	Low risk	Study attrition reported.
Free of selective reporting?	Unclear risk	No details.

Free of other bias?	Unclear risk	No details.
CAARMS - Comprehensive Assessment of At-Risk Mental States		
COPS - Criteria of Prodromal Syndromes		
DSM - Diagnostic and Statistical Manual		
ICD-10 - International Classification of Diseases		
SIPS - Structured Interview for Prodromal Syndromes		
Rating Scales:		
Mental state		
BDI - Beck Depression Inventory		
BPRS - Brief Psychiatric Rating Scale		
CDRS - Calgary Depression Rating Scale		
HRSA - Hamilton Rating Scale for Anxiety		
HRSD - Hamilton Rating Scale for Depression		
MADRS - Montgomery and Asberg Depression Rating Scale		
PANSS - Positive and Negative Symptom Scale		
POPS - Presence of Psychosis Scale		
SANS - Schedule for the Assessment of Negative Symptoms		
SOPS - Scale of Prodromal Symptoms		
YMRS - Young Mania Rating Scale		
YMS - Young Mania Scale		
Global state		
CGI - Clinical Global Impression		
GAF - Global Assessment of Functioning		
GAS - Global Adjustment Scale		
Adverse effects		
AIMS - Abnormal and Involuntary Movement Scales		
BAS - Barnes Akathisia Scale		
SAS - Simpson & Angus		
User satisfaction		
CSQ-8 - Client Satisfaction Questionnaire-8		
Others		
APQ - Alcohol Problems Questionnaire		
DUP - Duration of untreated psychosis		
EM - Explanatory Model scale		
EIPS - early initial prodromal state		
GSI - General Symptom Index of the SCL-90-R		
IS/O - Integration/Sealing Over		
IPS - Individual Placement and Support		
ITT - Intention to Treat		
LOCF - Last Observation Carried Forward		
MCQ - Meta-Cognitions Questionnaire		
OLIFE - Oxford-Liverpool Inventory of Feelings and Experiences		
PACE - Personal Assessment and Crisis Evaluation		

QLS - Quality of Life Scale
 RCI - Reliable Change Index
 SAC - Sociotropy - Autonomy Scale
 TAU - Treatment as Usual
 UHR - Ultra high risk (of developing psychosis)

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Addington 1999	Allocation: not randomised, before and after design.
Agius 2007	Allocation: not randomised, no control group.
Alanen 1994	Allocation: not randomised, before and after design.
Albiston 1998	Allocation: not randomised, before and after design with historical control
Anonymous 1987	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: medication only (pimozide versus flupenthixol) at standard doses without specific early intervention protocol
Bao 2005	Allocation: randomised. Participants: people with psychosis. Interventions: no early intervention programme.
Birchwood 1989	Allocation: not randomised, service description. Participants: not first-episode patients (study of early detection of signs of relapse)
Clare 1994	Allocation: not randomised, service description. Participants: not first-episode patients (study of early signs of depression in long term patients)
COPE-Melbourne	Allocation: non-randomised quasi-experimental design, controls selected from a similar location to the experimental site
Craig 2004b	Allocation: randomised. Participants: people with 1st & 2nd episode psychosis.
Crow 1986	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: neuroleptic medication at standard doses versus no medication, no specific early intervention protocol
Culberg 1998	Allocation: not randomised - before and after design with historical controls
Davidson 2004	Allocation: randomised. Participants: people with early psychosis. Interventions: atypical versus conventional antipsychotic.
DeHaan 1997	Allocation: not randomised - before and after design.
Drury 2000	Allocation: randomised. Participants: not people with first-episode psychosis.
Emsley 1999	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: medication only (risperidone versus haloperidol), no specific early intervention protocol
Emsley 2004	Allocation: randomised. Participants: people with recent onset schizophrenia. Interventions: medication only (risperidone versus haloperidol)
Falloon 1992	Allocation: not randomised, no controls.
Fisher 2001	Allocation: not randomised - service description, outcome assessed by qualitative survey
Fitzgerald 1998	Allocation: not randomised - before and after design.
Fresan 2001	Allocation: not randomised - before and after study.
Gaebel 2004	Allocation: randomised. Participants: people with first-episode schizophrenia.

Study	Reason for exclusion
	Interventions: medication only (risperidone versus low-dose haloperidol)
Grawe 1998	Allocation: randomised. Participants: people with first-episode psychosis. Interventions: no early intervention programme.
Hartmann 1974	Allocation: not randomised - retrospective study.
Heydebrand 2004	Allocation: randomised. Participants: people with first-episode schizophrenia. Interventions: haloperidol versus risperidone.
Jenner 2001	Allocation: not randomised - no control group. Participants: not people with first-episode psychosis (adolescents, but on average in treatment for about 3 years)
Jenner 2004	Allocation: randomised. Participants: people with chronic psychosis.
Jolley 2003	Allocation: randomised. Participants: people with 1st & 2nd episode psychosis.
Jones 2005	Allocation: randomised. Participants: people with psychosis - not first episode.
Kadota 1992	Allocation: not randomised - uncontrolled follow-up study of response to neuroleptic treatment
Kauranen 2000	Allocation: not randomised - uncontrolled follow-up study.
Kavanagh 2004	Allocation: randomised. Participants: people with 1st and 2nd episode psychosis.
Keefe 2000	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: medication only (olanzapine versus haloperidol) at standard doses without specific early intervention protocol
Keshavan 1998	Allocation: not randomised - before and after study with historical control group
Kopala 2003	Allocation: randomised. Participants: people with recent onset schizophrenia. Interventions: risperidone versus haloperidol.
Kuipers 2004	Allocation: randomised. Participants: people with schizophrenia, 1st and 2nd or more episodes
Li 2004	Allocation: randomised. Participants: early schizophrenia, probably not 1st episode. Interventions: no early intervention program.
Lieberman 2005b	Allocation: randomised. Participants: people with first-episode psychosis and healthy volunteers. Interventions: medication only (haloperidol versus olanzapine)
Malla 2001	Allocation: not randomised - before and after study, no control group
McCay 2007	Allocation: quasi-randomised.
McGorry 1996	Allocation: not randomised - before and after study with historical controls
Mosher 1975	Allocation: not randomised - allocation on 'a consecutively admitted, space available basis' to 'Soteria' - a small home like facility in the community which acted as an alternative to admission for patients in their first-episode of schizophrenia
Mottaghipour 2000	Allocation: not randomised, no control group (participants compared with a group of families of long term patients)
Newton 2005	Allocation: none randomised study (before and after design).
Nuechterlein 2005	Allocation: randomised. Participants: people with first-episode psychosis. Interventions: no early intervention programme.
Parlato 1999	Allocation: non-randomised - a description of a service.
Perez 2003	Allocation: randomised. Participants: people with first-episode psychosis.

Study	Reason for exclusion
	Interventions: medication only (olanzapine versus risperidone versus haloperidol)
Power 2004	Allocation: randomised. Participants: people with 1st and 2nd episode psychosis.
Purdon 2000	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: medication only (olanzapine versus risperidone versus haloperidol) without specific early intervention protocol
Rund 1994	Allocation: not randomised, before and after study with historical control
Sanger 1999	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: medication only (olanzapine versus haloperidol) without specific early intervention protocol
Schooler 2003	Allocation: randomised. Participants: people with recent onset schizophrenia. Interventions: medication only (risperidone versus haloperidol)
SOCRATES-UK	Allocation: randomised. Participants: not first-episode patients, participants could be in either first or second admission, as long as second admission within 2 years of first admission (estimated that 61/309 participants were not first episode)
Szymanski 1994	Allocation: not randomised - before and after study without control group
Thomas 1979	Allocation: not randomised - no control group. Participants: not first-episode patients, although all were adolescents, some were experiencing an exacerbation of chronic schizophrenia
TIPS 2006	Allocation: not randomised.
Turetz 1997	Allocation: not randomised - no control group. Participants: probably not first-episode patients, although all participants were children, they were selected on the basis of treatment resistance and so probably not in the first episode
Ueland 2004	Allocation: randomised. Participants: people with psychosis, not first episode.
Walczewski 1998	Allocation: not randomised - a quasi-experimental design, patients receiving a psychosocial treatment program were compared with a group receiving an individual treatment programme
Wang 2000	Allocation: randomised. Participants: people with first-episode psychosis. Intervention: medication only (risperidone versus clozapine) without specific early intervention protocol
Welch 2000	Allocation: not randomised - service description.
Whitehorn 1998	Allocation: not randomised - before and after study without control
Whitwell 2000	Allocation: not randomised - service description.
Wieneke 2000	Allocation: not randomised - service description.
Wunderink 2003	Allocation: randomised. Participants: people with early onset psychosis. Intervention: drug trial - not early intervention.
Wykes 2007	Allocation: randomised. Participants: people with psychosis, but not first episode, (only early onset)
Yap 2001	Allocation: not randomised - before and after study without control group
Zhang-Wong 1999	Allocation: not randomised - prospective uncontrolled study to determine optimal dose of haloperidol

Characteristics of studies awaiting assessment [ordered by study ID]

Addington-2001

Methods	No details.
Participants	People with psychosis.
Interventions	Early psychosis programme.
Outcomes	No details.
Notes	

Alaghband-rad 2006a

Methods	Randomised.
Participants	People with psychosis.
Interventions	Early intervention.
Outcomes	No data.
Notes	

Berger 2006

Methods	No details.
Participants	People at risk of developing psychosis.
Interventions	Lithium.
Outcomes	No details.
Notes	

Cornblatt 2009

Methods	No details.
Participants	People at risk of developing psychosis.
Interventions	Risperidone versus sertraline.
Outcomes	No details.
Notes	

Dai

Methods	Randomised.
Participants	People with psychosis.
Interventions	Group or individual psychoeducation versus antipsychotic maintenance

Outcomes	No details.
Notes	

Deng 2006

Methods	Randomised.
Participants	People with schizophrenia.
Interventions	Early intervention and routine care versus routine care alone
Outcomes	No details.
Notes	

Doering 1998

Methods	No details.
Participants	People with psychosis.
Interventions	No details.
Outcomes	No details.
Notes	

Edwards 2003 EPPIC

Methods	Randomised.
Participants	People with first-episode psychosis.
Interventions	Clozapine versus CBT.
Outcomes	No details.
Notes	

Fillatre-1998

Methods	Unclear.
Participants	People with psychosis.
Interventions	Unclear.
Outcomes	Unclear.
Notes	

Furimsky 2005

Methods	No details.
Participants	No details.
Interventions	No details.
Outcomes	No details.
Notes	

Gleeson 2008

Methods	Controlled trial.
Participants	People with early psychosis.
Interventions	Cognitive and family therapy relapse prevention.
Outcomes	No details.
Notes	

Humphries 2005

Methods	No details.
Participants	People with early psychosis.
Interventions	No details.
Outcomes	No details.
Notes	

Johnson 2004

Methods	Cluster randomised trial.
Participants	People with early psychosis.
Interventions	Specialist team versus augmented community mental health teams
Outcomes	No details.
Notes	

Keshavan 2003

Methods	Controlled trial.
Participants	People with early psychosis.
Interventions	Psychoeducation and collaboration enhancement.

Outcomes	No details.
Notes	

Keshavan 2005

Methods	No details.
Participants	People with early psychosis.
Interventions	No details.
Outcomes	No details.
Notes	

Lecomte 2006

Methods	Randomised.
Participants	People with early episodes of psychosis.
Interventions	Group CBT versus skills training.
Outcomes	No details.
Notes	Recent onset psychosis i.e. early episode psychosis. Text is confusing as mentions first episode but is including people with 2 years established psychosis for example, having consulted for the first time a professional for psychotic symptoms in the past two years

Lee 2007

Methods	No details.
Participants	Adolescents with schizophrenia.
Interventions	No details.
Outcomes	No details.
Notes	

Li 2004b

Methods	Randomised.
Participants	People with first-episode schizophrenia.
Interventions	Early intervention programme.
Outcomes	Quality of life.
Notes	

Li 2007

Methods	No details.
Participants	People with schizophrenia.
Interventions	Psychotherapy.
Outcomes	No details.
Notes	

Richtand 2007

Methods	No details.
Participants	People with early schizophrenia.
Interventions	Omega-3 fatty acid.
Outcomes	No details.
Notes	

Schepp 1999

Methods	No details.
Participants	People with schizophrenia.
Interventions	Self-management therapy.
Outcomes	No details.
Notes	

Tao 2004

Methods	Randomised.
Participants	People with first-episode psychosis.
Interventions	Early intervention programme.
Outcomes	No details.
Notes	

Vinogradov 2008

Methods	No details.
Participants	People at risk of psychosis.
Interventions	Neuroadaptive cognitive training.

Outcomes	No details.
Notes	

Williams 2005

Methods	No details.
Participants	People with early psychosis.
Interventions	Psychosocial interventions.
Outcomes	No details.
Notes	

Woo 2009

Methods	No details.
Participants	People with schizophrenia.
Interventions	Tiagabine.
Outcomes	No details.
Notes	

Woods 2005

Methods	No details.
Participants	People with prodromal psychosis.
Interventions	Glycine.
Outcomes	No details.
Notes	

Woods 2008

Methods	No details.
Participants	People with prodromal psychosis.
Interventions	Ziprasidone.
Outcomes	No details.
Notes	

Xu 2003

Methods	No details.
Participants	People with first-episode schizophrenia.
Interventions	Psychotherapy.
Outcomes	No details.
Notes	

Yu 2005

Methods	Randomised.
Participants	People with first-episode schizophrenia.
Interventions	No details.
Outcomes	No details.
Notes	

Characteristics of ongoing studies [ordered by study ID]

Addington 2007

Trial name or title	A randomised controlled trial of individual therapy for first-episode psychosis
Methods	Randomised.
Participants	People with schizophrenia.
Interventions	CBT versus usual care.
Outcomes	Social Functioning Scale (SFS) Positive and Negative Syndrome Scale (PANSS) Psychotic Symptom Rating Scales (PSYRATS) Calgary Depression Scale for Schizophrenia (CDSS) The Time-Line Follow Back (TLFB) Alcohol and Drug Use Scale (AUS; DUS) Medication Event Monitoring System (MEMS) Rosenberg Self-Esteem Scale Maastricht Assessment of Coping Skills (MACS)
Starting date	June 2007.
Contact information	Jean_Addington@camh.net
Notes	

Arends 2006

Trial name or title	Prodromal symptoms and early intervention to prevent a relapse
Methods	No details.
Participants	People with schizophrenia.

Interventions	The Symptom Management Module (SMM) versus treatment as usual
Outcomes	Occurrence of a psychotic relapse: a worsening of at least two points on the CGI as assessed by psychiatrist and verified by researcher by a PANSS-interview within a week.
Starting date	No details.
Contact information	Johan.Arends@GGZDrenthe.nl
Notes	

EDIE-2 Morrison 2007

Trial name or title	Early detection and psychological intervention for individuals at high risk of psychosis (EDIE 2)
Methods	Randomised.
Participants	People with high risk of psychosis.
Interventions	CBT plus regular monitoring versus monitoring alone.
Outcomes	Transition to psychosis.
Starting date	No details.
Contact information	No details.
Notes	

Furimski 2005

Trial name or title	Access, detection and psychological treatments.
Methods	No details.
Participants	People at risk of developing psychosis.
Interventions	Psychological intervention in preventing or delaying the onset of a psychotic illness.
Outcomes	Psychosis.
Starting date	August 2004.
Contact information	No details.
Notes	

Gaebel 2005

Trial name or title	Maintenance treatment versus stepwise drug discontinuation in first-episode schizophrenia
Methods	Randomised.
Participants	People with schizophrenia.
Interventions	Maintenance treatment.
Outcomes	Relapse.
Starting date	November 2001.
Contact information	No details.

Notes

Heresco-Levy 2006

Trial name or title	Sarcosine preventive therapy for individuals at high risk for schizophrenia
Methods	Randomised.
Participants	People with the prodromal stage of schizophrenia.
Interventions	Sarcosine versus placebo.
Outcomes	Delay or prevention of illness.
Starting date	June 2006.
Contact information	No details.
Notes	

Lester 2006

Trial name or title	Birmingham early detection in untreated psychosis trial (REDIRECT)
Methods	Cluster randomised.
Participants	People with first-episode psychosis, aged 14-30 years.
Interventions	Early detection training versus detection as usual.
Outcomes	The primary outcome is the number of general practitioner referrals of young people with first-episode psychosis to early intervention services. Secondary outcomes are duration of untreated psychosis, time to recovery, use of the Mental Health Act, and general practitioner consultation rate
Starting date	2004.
Contact information	No details.
Notes	

McFarlane 2007

Trial name or title	Early detection and intervention prevention of psychosis.
Methods	Randomised, single blind (outcomes assessor).
Participants	People with a high risk of psychosis.
Interventions	Psychoeducational multifamily group treatment plus antipsychotics versus case management
Outcomes	Conversion to psychosis.
Starting date	No details.
Contact information	mcfarw@mmc.org
Notes	

Srihari 2006

Trial name or title	Specialized treatment early in psychosis (STEP): a pragmatic randomized controlled trial in the US public sector
Methods	Randomised.
Participants	People with early psychosis.
Interventions	Specialized treatment for early psychosis.
Outcomes	No details.
Starting date	No details.
Contact information	No details.
Notes	

Stain 2006

Trial name or title	The depth project: a multi site RCT for youths at risk for psychosis
Methods	Randomised.
Participants	People at risk of psychosis.
Interventions	Cognitive behavioural therapy and person centred therapy in ameliorating 'at risk mental states' for psychosis
Outcomes	No details.
Starting date	No details.
Contact information	No details.
Notes	

DATA AND ANALYSES

Comparison 1

PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early (for reasons other than psychosis)	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 by eight weeks	1	60	Risk Ratio (M-H, Fixed, 95% CI)	1.29 [0.60, 2.74]
1.2 by one year	1	60	Risk Ratio (M-H, Fixed, 95% CI)	1.59 [0.88, 2.88]
2 Converted to psychosis: POPS	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 over one year	1	60	Risk Ratio (M-H, Fixed, 95% CI)	0.58 [0.28, 1.18]
3 Global state: 1. Average total change score - by 1 month (CGI-severity of illness, high score=worse)	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.23 [-0.82, 0.36]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4 Global state: 2. Average total change score - by 12 months (GAF-current, high score=good)	1	59	Mean Difference (IV, Fixed, 95% CI)	2.43 [-4.77, 9.63]
5 Mental state: 1. Average total change score - by 12 months (SOPS, high score=worse)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
5.1 total score	1	59	Mean Difference (IV, Fixed, 95% CI)	-2.76 [-12.03, 6.51]
5.2 positive score	1	59	Mean Difference (IV, Fixed, 95% CI)	-2.73 [-6.18, 0.72]
5.3 negative	1	59	Mean Difference (IV, Fixed, 95% CI)	0.28 [-3.02, 3.58]
5.4 disorganisation	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.49 [-2.69, 1.71]
5.5 general	1	59	Mean Difference (IV, Fixed, 95% CI)	0.18 [-1.84, 2.20]
6 Mental state: 2. Average total change score - by 12 months (PANSS, high score=worse)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
6.1 total	1	59	Mean Difference (IV, Fixed, 95% CI)	0.48 [-10.69, 11.65]
6.2 positive	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.57 [-3.75, 2.61]
6.3 negative	1	59	Mean Difference (IV, Fixed, 95% CI)	0.52 [-2.60, 3.64]
6.4 general	1	59	Mean Difference (IV, Fixed, 95% CI)	0.54 [-5.44, 6.52]
7 Mental state: 3. Average total change score - by 12 months (YMRS, high score=worse)	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.91 [-3.77, 1.95]
8 Mental state: 4. Average total change score - by 12 months (MADRS, high score=worse)	1	59	Mean Difference (IV, Fixed, 95% CI)	0.68 [-3.81, 5.17]
9 Adverse effects: 1. Average total change score - by 8 weeks (SAS, high score=worse)	1	59	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.59, 0.79]
10 Adverse effects: 2. Average total change score - by 8 weeks (BAS, high score=worse)	1	59	Mean Difference (IV, Fixed, 95% CI)	0.5 [-0.58, 1.58]
11 Adverse effects: 3. Average total change score - by 8 weeks (AIMS, high score=worse)	1	59	Mean Difference (IV, Fixed, 95% CI)	0.60 [-0.34, 1.54]
12 Adverse effects: 4. Average total weight change score (kg) - by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	7.63 [4.04, 11.22]
13 Adverse effects: 5. Weight gain - by 12 months	1	60	Risk Ratio (M-H, Fixed, 95% CI)	3.55 [1.53, 8.28]
14 Adverse effects: 6. Average total change score - by 8 weeks (Cardiovascular)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
14.1 sitting systolic blood pressure	1	59	Mean Difference (IV, Fixed, 95% CI)	1.0 [-4.28, 6.28]
14.2 sitting diastolic blood pressure	1	59	Mean Difference (IV, Fixed, 95% CI)	0.70 [-4.43, 5.83]
14.3 standing systolic blood pressure	1	59	Mean Difference (IV, Fixed, 95% CI)	-2.80 [-9.18, 3.58]
14.4 standing diastolic blood pressure	1	59	Mean Difference (IV, Fixed, 95% CI)	0.20 [-4.96, 5.36]
14.5 sitting pulse rate	1	58	Mean Difference (IV, Fixed, 95% CI)	7.20 [-1.04, 15.44]
14.6 standing pulse rate	1	57	Mean Difference (IV, Fixed, 95% CI)	3.90 [-4.87, 12.67]
15 Adverse effects: 7. Average total change score - by 12 months (Pulse, BPM)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
15.1 sitting pulse rate	1	58	Mean Difference (IV, Fixed, 95% CI)	8.31 [0.53, 16.09]
15.2 standing pulse rate	1	57	Mean Difference (IV, Fixed, 95% CI)	2.86 [-6.69, 12.41]
16 Adverse effects: 8. Treatment emergent adverse events - by 8 weeks (CoStart Term)	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
16.1 somnolence	1	60	Risk Ratio (M-H, Fixed, 95% CI)	2.25 [0.90, 5.59]
16.2 weight gain	1	60	Risk Ratio (M-H, Fixed, 95% CI)	10.29 [1.42, 74.79]
16.3 increased appetite	1	60	Risk Ratio (M-H, Fixed, 95% CI)	1.87 [0.51, 6.80]
16.4 anxiety	1	60	Risk Ratio (M-H, Fixed, 95% CI)	4.68 [0.58, 37.68]
16.5 nervousness	1	60	Risk Ratio (M-H, Fixed, 95% CI)	1.87 [0.37, 9.46]
16.6 asthenia	1	60	Risk Ratio (M-H, Fixed, 95% CI)	3.74 [0.44, 31.55]
16.7 joint disorder	1	60	Risk Ratio (M-H, Fixed, 95% CI)	0.94 [0.20, 4.27]
16.8 abnormal thoughts	1	60	Risk Ratio (M-H, Fixed, 95% CI)	1.40 [0.25, 7.81]
17 Adverse effects: 9. Fatigue - by 12 months	1	60	Risk Ratio (M-H, Fixed, 95% CI)	8.42 [1.14, 62.40]

Comparison 2

PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 by 12 months	1	60	Risk Ratio (M-H, Fixed, 95% CI)	0.98 [0.44, 2.16]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.2 by 3 years	1	60	Risk Ratio (M-H, Fixed, 95% CI)	0.96 [0.60, 1.52]
2 Transition to psychosis - by 12 months	1	60	Risk Ratio (M-H, Fixed, 95% CI)	0.50 [0.15, 1.66]
3 Social Functioning: 1. SAS II endpoint data (long term, high score=worse, LOCF)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
3.1 global	1	67	Mean Difference (IV, Fixed, 95% CI)	0.40 [-0.05, 0.85]
3.2 social activities	1	67	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.28, 0.48]
3.3 well-being	1	67	Mean Difference (IV, Fixed, 95% CI)	0.10 [-0.22, 0.42]
3.4 work	1	67	Mean Difference (IV, Fixed, 95% CI)	-0.10 [-0.38, 0.18]

Comparison 3

PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early - by 12 months	1	59	Risk Ratio (M-H, Fixed, 95% CI)	0.0 [0.0, 0.0]
2 Progression to psychosis	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 by 6 months	1	59	Risk Ratio (M-H, Fixed, 95% CI)	0.27 [0.08, 0.89]
2.2 by 12 months	1	59	Risk Ratio (M-H, Fixed, 95% CI)	0.54 [0.23, 1.30]
3 Global state: Average endpoint score (GAF, high score=worse)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
3.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	4.20 [-2.57, 10.97]
3.2 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	0.0 [-5.21, 5.21]
4 Mental state: 1a. Average endpoint score (BPRS psychotic symptoms -general, high score=worse, skewed data)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
4.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	0.10 [-1.25, 1.45]
4.2 by 6 months	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.5 [-2.25, 1.25]
4.3 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	0.70 [-0.99, 2.39]
5 Mental state: 1b. Average endpoint score (SANS, psychotic symptoms - negative,	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
high score=worse, skewed data)				
5.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	-3.20 [-9.80, 3.40]
5.2 by 6 months	1	59	Mean Difference (IV, Fixed, 95% CI)	-4.60 [-12.72, 3.52]
5.3 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.80 [-7.87, 6.27]
6 Mental state: 2a. Average endpoint score anxiety (HRSA, high score=worse, skewed data)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
6.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.70 [-4.85, 3.45]
6.2 by 6 months	1	59	Mean Difference (IV, Fixed, 95% CI)	-1.10 [-4.81, 2.61]
6.3 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	0.60 [-4.18, 5.38]
7 Mental state: 2b. Average endpoint score depression (HRSD, high score=worse, skewed data)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
7.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	-1.0 [-5.51, 3.51]
7.2 by 6 months	1	59	Mean Difference (IV, Fixed, 95% CI)	-0.20 [-4.77, 4.37]
7.3 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	1.20 [-3.22, 5.62]
8 Mental state: 2c. Average endpoint score mania (YMS, high score=worse, skewed data)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
8.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	0.80 [-1.38, 2.98]
8.2 by 6 months	1	59	Mean Difference (IV, Fixed, 95% CI)	0.70 [-2.46, 3.86]
8.3 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	0.0 [-1.76, 1.76]
9 Quality of life: Average endpoint score (QLS, high score=worse)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
9.1 at baseline	1	59	Mean Difference (IV, Fixed, 95% CI)	-3.10 [-14.12, 7.92]
9.2 by 6 months	1	59	Mean Difference (IV, Fixed, 95% CI)	-1.40 [-13.63, 10.83]
9.3 by 12 months	1	59	Mean Difference (IV, Fixed, 95% CI)	0.80 [-10.15, 11.75]

Comparison 4**PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2 Mental state: 1. PANSS, endpoint score (by 12 weeks, higher scores=worse, LOCR)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 PANSS-G	1	102	Mean Difference (IV, Fixed, 95% CI)	-3.40 [-6.85, 0.05]
2.2 PANSS-P	1	102	Mean Difference (IV, Fixed, 95% CI)	-2.10 [-3.69, -0.51]
2.3 PANSS-N	1	102	Mean Difference (IV, Fixed, 95% CI)	-1.30 [-3.26, 0.66]
3 Mental state: 1. MADRS, endpoint score (by 12 weeks, higher scores=worse, LOCF)	1	102	Mean Difference (IV, Fixed, 95% CI)	-1.10 [-4.49, 2.29]
4 Global state: 1. GAF, endpoint score (by 12 weeks, higher scores=better)	1	102	Mean Difference (IV, Fixed, 95% CI)	-6.10 [-11.76, -0.44]

Comparison 5**OMEGA 3 FATTY ACIDS (EPA) versus PLACEBO**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Transition to psychosis	1	76	Risk Ratio (M-H, Fixed, 95% CI)	0.13 [0.02, 0.95]

Comparison 6**PHASE-SPECIFIC TREATMENT (CBT for SUICIDALITY) + SPECIALISED TEAM vs SPECIALISED TEAM**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early - by 6 months	1	56	Risk Ratio (M-H, Fixed, 95% CI)	2.02 [0.72, 5.66]
2 Suicide - by 6 months	1	56	Risk Ratio (M-H, Fixed, 95% CI)	0.81 [0.05, 12.26]

Comparison 7**PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + SPECIALISED TEAM vs
SPECIALISED TEAM**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Relapse by end of treatment - by 12 months	1	76	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.37, 2.98]

Comparison 8**PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs
STANDARD CARE**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early - by 18 months	1	83	Risk Ratio (M-H, Fixed, 95% CI)	1.46 [0.26, 8.31]
2 Readmitted to hospital - by 18 months	1	83	Risk Ratio (M-H, Fixed, 95% CI)	0.28 [0.13, 0.62]
3 Not compliant with medication	1	83	Risk Ratio (M-H, Fixed, 95% CI)	0.57 [0.31, 1.04]

Comparison 9**SPECIALISED TEAM vs STANDARD CARE**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 by one year	1	547	Risk Ratio (M-H, Fixed, 95% CI)	0.59 [0.43, 0.81]
1.2 by two years	1	547	Risk Ratio (M-H, Fixed, 95% CI)	0.64 [0.50, 0.82]
1.3 by five years	1	547	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.84, 1.21]
2 Global state: 1. Average endpoint score - by 12 and 24 months (GAF-symptom, high score=good)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 by one year	1	419	Mean Difference (IV, Fixed, 95% CI)	-3.71 [-6.69, -0.73]
2.2 by two years	1	369	Mean Difference (IV, Fixed, 95% CI)	-2.51 [-5.70, 0.68]
2.3 by 5 years	1	301	Mean Difference (IV, Fixed, 95% CI)	0.32 [-3.57, 4.21]
3 Global state: 2. Average endpoint score - by 12 and 24 months (GAF-function, high score=good)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.1 by one year	1	419	Mean Difference (IV, Fixed, 95% CI)	-2.30 [-5.15, 0.55]
3.2 by two years	1	369	Mean Difference (IV, Fixed, 95% CI)	-4.03 [-7.23, -0.83]
3.3 by five years	1	301	Mean Difference (IV, Fixed, 95% CI)	-1.20 [-5.23, 2.83]
4 User satisfaction: Average endpoint score - by 12 and 24 months (CSQ-8, high score=good)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
4.1 by one year	1	419	Mean Difference (IV, Fixed, 95% CI)	-1.90 [-3.07, -0.73]
4.2 by two years	1	369	Mean Difference (IV, Fixed, 95% CI)	-3.20 [-4.14, -2.26]
5 Compliance with treatment	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
5.1 treatment stopped in spite of need - by one year	1	507	Risk Ratio (M-H, Fixed, 95% CI)	0.20 [0.10, 0.42]
5.2 treatment stopped in spite of need - by two years	1	436	Risk Ratio (M-H, Fixed, 95% CI)	0.66 [0.29, 1.50]
6 Suicide: Death - by 12 months	1	506	Risk Ratio (M-H, Fixed, 95% CI)	0.93 [0.06, 14.81]
7 Death other than suicide - by 12 months	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
7.1 accident	1	506	Risk Ratio (M-H, Fixed, 95% CI)	0.31 [0.01, 7.59]
7.2 unexplained	1	507	Risk Ratio (M-H, Fixed, 95% CI)	0.31 [0.01, 7.56]
8 Service use: 1. Average mean number of days per month in hospital	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
8.1 by one year	1	507	Mean Difference (IV, Fixed, 95% CI)	-1.39 [-2.83, 0.05]
8.2 by two years	1	436	Mean Difference (IV, Fixed, 95% CI)	-0.67 [-1.88, 0.54]
8.3 by five years	1	547	Mean Difference (IV, Fixed, 95% CI)	-1.11 [-3.21, 0.99]
9 Service use: 2. Not hospitalised - by five years	1	547	Risk Ratio (M-H, Fixed, 95% CI)	1.05 [0.90, 1.22]
10 Social outcomes: 1. Not living independently	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
10.1 by one year	1	507	Risk Ratio (M-H, Fixed, 95% CI)	0.55 [0.25, 1.17]
10.2 by two years	1	436	Risk Ratio (M-H, Fixed, 95% CI)	0.74 [0.36, 1.53]
10.3 by five years	1	547	Risk Ratio (M-H, Fixed, 95% CI)	0.42 [0.21, 0.83]
11 Social outcomes: 2. Not working or in education	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
11.1 by one year	1	507	Risk Ratio (M-H, Fixed, 95% CI)	0.87 [0.65, 1.17]
11.2 by two years	1	436	Risk Ratio (M-H, Fixed, 95% CI)	0.72 [0.54, 0.97]
11.3 by five years	1	547	Risk Ratio (M-H, Fixed, 95% CI)	1.06 [0.92, 1.23]

Comparison 10

PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early by 12 months	1	62	Risk Ratio (M-H, Fixed, 95% CI)	0.57 [0.19, 1.76]
2 Hospitalised by 12 months	1	62	Risk Ratio (M-H, Fixed, 95% CI)	1.08 [0.59, 1.99]
3 Suicide by 12 months	1	62	Risk Ratio (M-H, Fixed, 95% CI)	5.0 [0.25, 100.08]
4 Social functioning: SOFRAS by 12 months (higher score=worse)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
4.1 total score	1	62	Mean Difference (IV, Fixed, 95% CI)	-1.30 [-8.86, 6.26]
4.2 positive symptoms	1	62	Mean Difference (IV, Fixed, 95% CI)	0.35 [-1.86, 2.56]
4.3 negative symptoms	1	62	Mean Difference (IV, Fixed, 95% CI)	4.89 [-1.58, 11.36]

Comparison 11

PHASE-SPECIFIC TREATMENT (E-EPA) + ATYPICALS vs PLACEBO + ATYPICALS

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early by 12 weeks	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.83 [0.28, 2.51]
2 Global state: Not responded to treatment by 12 weeks	1	80	Risk Ratio (M-H, Fixed, 95% CI)	0.9 [0.57, 1.43]

Comparison 12

PHASE-SPECIFIC TREATMENT (BRIEF INTERVENTION) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early by nine months	1	106	Risk Ratio (M-H, Fixed, 95% CI)	0.72 [0.34, 1.51]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2 Hospital admission: Hospitalised	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
2.1 before 4 months	1	106	Risk Ratio (M-H, Fixed, 95% CI)	1.19 [0.89, 1.58]
2.2 Up to 4 months	1	106	Risk Ratio (M-H, Fixed, 95% CI)	0.75 [0.41, 1.38]
2.3 between 4 months and 9 months	1	106	Risk Ratio (M-H, Fixed, 95% CI)	0.86 [0.43, 1.74]

Comparison 13

PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early (6 months)	1	24	Risk Ratio (M-H, Fixed, 95% CI)	1.27 [0.26, 6.28]
2 Mental state: 1. PANSS	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 total score	1	17	Mean Difference (IV, Fixed, 95% CI)	-4.80 [-18.42, 8.82]
2.2 positive score	1	17	Mean Difference (IV, Fixed, 95% CI)	-1.58 [-4.88, 1.72]
2.3 negative score	1	17	Mean Difference (IV, Fixed, 95% CI)	-1.64 [-8.05, 4.77]
2.4 general score	1	17	Mean Difference (IV, Fixed, 95% CI)	-1.57 [-7.65, 4.51]
3 Mental state: 2. Calgary Depression Rating Scale	1	17	Mean Difference (IV, Fixed, 95% CI)	-1.46 [-4.17, 1.25]
4 Quality of Life: 1. Heinrichs-Carpenter	1	16	Mean Difference (IV, Fixed, 95% CI)	-2.93 [-25.59, 19.73]

Comparison 14

PHASE-SPECIFIC TREATMENT (VOCATIONAL INTERVENTION) + TAU vs TREATMENT AS USUAL

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Not employed	1	41	Risk Ratio (M-H, Fixed, 95% CI)	0.39 [0.21, 0.71]
2 Leaving the study early	1	41	Risk Ratio (M-H, Fixed, 95% CI)	0.21 [0.03, 1.64]

Comparison 15

PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) +
ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Cannabis use: 1. Used cannabis in last 4 weeks	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 by 3 months - end of treatment	1	47	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [0.62, 1.74]
1.2 by 9 months - 6 months after end of treatment	1	47	Risk Ratio (M-H, Fixed, 95% CI)	1.30 [0.79, 2.15]
2 Cannabis use: 2. Percentage days used cannabis in last 4 weeks (skewed data)			Other data	No numeric data
2.1 by 3 months - end of treatment			Other data	No numeric data
2.2 by 9 months - 6 months after end of treatment			Other data	No numeric data
3 Cannabis use: 3. Severity of cannabis use (skewed data)			Other data	No numeric data
3.1 by 3 months - end of treatment			Other data	No numeric data
3.2 by 9 months - 6 months after end of treatment			Other data	No numeric data
4 Global state: Average score (KAPQ total endpoint, higher=good)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
4.1 by 3 months - end of treatment	1	47	Mean Difference (IV, Fixed, 95% CI)	0.80 [-1.78, 3.38]
4.2 by 9 months - 6 months after end of treatment	1	47	Mean Difference (IV, Fixed, 95% CI)	0.90 [-1.42, 3.22]
5 Mental state: 1. Average score (BPRS-E total endpoint, higher scores=poor)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
5.1 by 3 months - end of treatment	1	47	Mean Difference (IV, Fixed, 95% CI)	-3.60 [-12.81, 5.61]
5.2 by 9 months - 6 months after end of treatment	1	47	Mean Difference (IV, Fixed, 95% CI)	0.80 [-7.47, 9.07]
6 Mental state: 2. Average score (BPRS-PS total endpoint, higher scores=poor) (skewed data)			Other data	No numeric data
6.1 by 3 months - end of treatment			Other data	No numeric data
6.2 by 9 months - 6 months after end of treatment			Other data	No numeric data
7 Mental state: 3. Average negative symptom score			Other data	No numeric data

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
(SANS endpoint, higher scores=poor) (skewed data)				
7.1 by 3 months - end of treatment			Other data	No numeric data
7.2 by 9 months - 6 months after end of treatment			Other data	No numeric data
8 Mental state: 4. Average score (BDI-SF total endpoint , higher scores=poor) (skewed data)			Other data	No numeric data
8.1 by 3 months - end of treatment			Other data	No numeric data
8.2 by 9 months - 6 months after end of treatment			Other data	No numeric data
9 Social functioning: Average score (SOFAS total endpoint, higher scores=good)	1		Mean Difference (IV, Fixed, 95% CI)	Subtotals only
9.1 by 3 months - end of treatment	1	47	Mean Difference (IV, Fixed, 95% CI)	-0.80 [-9.95, 8.35]
9.2 by 9 months - 6 months after end of treatment	1	47	Mean Difference (IV, Fixed, 95% CI)	-4.70 [-14.52, 5.12]

Comparison 16

CRISIS ASSESSMENT versus STANDARD CARE (LEO-CAT)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Hospitalisation	1	98	Risk Ratio (M-H, Fixed, 95% CI)	0.85 [0.57, 1.27]
2 Referred to Mental Health Services by A & E or emergency medical services	1	98	Risk Ratio (M-H, Fixed, 95% CI)	0.85 [0.57, 1.27]

Comparison 17

EARLY BEHAVIOURAL INTERVENTION vs ROUTINE CARE INTERVAL (Alvarez-Spain)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Weight			Other data	No numeric data
1.1 Weight change at 13 weeks			Other data	No numeric data
1.2 Body Mass Index			Other data	No numeric data

Comparison 18

PHASE-SPECIFIC INTERVENTION versus CONTROL (Exploratory meta-analysis)


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Leaving the study early	3	192	Risk Ratio (M-H, Fixed, 95% CI)	0.72 [0.41, 1.29]
2 Hospitalisation	2	168	Risk Ratio (M-H, Fixed, 95% CI)	0.97 [0.61, 1.54]

Analysis 1.1. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 1 Leaving the study early (for reasons other than psychosis)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 1 Leaving the study early (for reasons other than psychosis)




Analysis 1.2. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 2 Converted to psychosis: POPS

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 2 Converted to psychosis: POPS




Analysis 1.3. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 3 Global state: 1. Average total change score - by 1 month (CGI-severity of illness, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 3 Global state: 1. Average total change score - by 1 month (CGI-severity of illness, high score=worse)




Analysis 1.4. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 4 Global state: 2. Average total change score - by 12 months (GAF-current, high score=good)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 4 Global state: 2. Average total change score - by 12 months (GAF-current, high score=good)




Analysis 1.5. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 5 Mental state: 1. Average total change score - by 12 months (SOPS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 5 Mental state: 1. Average total change score - by 12 months (SOPS, high score=worse)




Analysis 1.6. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 6 Mental state: 2. Average total change score - by 12 months (PANSS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 6 Mental state: 2. Average total change score - by 12 months (PANSS, high score=worse)




Analysis 1.7. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 7 Mental state: 3. Average total change score - by 12 months (YMRS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 7 Mental state: 3. Average total change score - by 12 months (YMRS, high score=worse)




Analysis 1.8. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC

SUPPORTIVE, Outcome 8 Mental state: 4. Average total change score - by 12 months (MADRS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 8 Mental state: 4. Average total change score - by 12 months (MADRS, high score=worse)




Analysis 1.9. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 9 Adverse effects: 1. Average total change score - by 8 weeks (SAS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 9 Adverse effects: 1. Average total change score - by 8 weeks (SAS, high score=worse)




Analysis 1.10. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO +

NON-SPECIFIC SUPPORTIVE, Outcome 10 Adverse effects: 2. Average total change score - by 8 weeks (BAS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 10 Adverse effects: 2. Average total change score - by 8 weeks (BAS, high score=worse)




Analysis 1.11. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 11 Adverse effects: 3. Average total change score - by 8 weeks (AIMS, high score=worse)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 11 Adverse effects: 3. Average total change score - by 8 weeks (AIMS, high score=worse)




Analysis 1.12. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO +

NON-SPECIFIC SUPPORTIVE, Outcome 12 Adverse effects: 4. Average total weight change score (kg) - by 12 months

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 12 Adverse effects: 4. Average total weight change score (kg) - by 12 months




Analysis 1.13. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 13 Adverse effects: 5. Weight gain - by 12 months

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 13 Adverse effects: 5. Weight gain - by 12 months




Analysis 1.14. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO +

NON-SPECIFIC SUPPORTIVE, Outcome 14 Adverse effects: 6. Average total change score - by 8 weeks (Cardiovascular)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 14 Adverse effects: 6. Average total change score - by 8 weeks (Cardiovascular)




Analysis 1.15. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 15 Adverse effects: 7. Average total change score - by 12 months (Pulse, BPM)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 15 Adverse effects: 7. Average total change score - by 12 months (Pulse, BPM)




Analysis 1.16. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 16 Adverse effects: 8. Treatment emergent adverse events - by 8 weeks (CoStart Term)

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 16 Adverse effects: 8. Treatment emergent adverse events - by 8 weeks (CoStart Term)




Analysis 1.17. Comparison 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE, Outcome 17 Adverse effects: 9. Fatigue - by 12 months

Review: Early intervention for psychosis

Comparison: 1 PHASE SPECIFIC TREATMENT (OLANZAPINE) + NON-SPECIFIC SUPPORTIVE THERAPY vs PLACEBO + NON-SPECIFIC SUPPORTIVE

Outcome: 17 Adverse effects: 9. Fatigue - by 12 months




Analysis 2.1. Comparison 2 PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY, Outcome 1 Leaving the study early

Review: Early intervention for psychosis

Comparison: 2 PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY

Outcome: 1 Leaving the study early




Analysis 2.2. Comparison 2 PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY, Outcome 2 Transition to psychosis - by 12 months

Review: Early intervention for psychosis

Comparison: 2 PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY

Outcome: 2 Transition to psychosis - by 12 months




Analysis 2.3. Comparison 2 PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY, Outcome 3 Social Functioning: 1. SAS II endpoint data (long term, high score=worse, LOCF)

Review: Early intervention for psychosis

Comparison: 2 PHASE SPECIFIC TREATMENT (CBT) + NON-SPECIFIC SUPPORTIVE THERAPY vs NON-SPECIFIC SUPPORTIVE THERAPY

Outcome: 3 Social Functioning: 1. SAS II endpoint data (long term, high score=worse, LOCF)




Analysis 3.1. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 1 Leaving the study early - by 12 months

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 1 Leaving the study early - by 12 months




Analysis 3.2. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 2 Progression to psychosis

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 2 Progression to psychosis




Analysis 3.3. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 3 Global state: Average endpoint score (GAF, high score=worse)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 3 Global state: Average endpoint score (GAF, high score=worse)




Analysis 3.4. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 4 Mental state: 1a. Average endpoint score (BPRS psychotic symptoms -general, high score=worse, skewed data)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 4 Mental state: 1a. Average endpoint score (BPRS psychotic symptoms -general, high score=worse, skewed data)




Analysis 3.5. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 5 Mental state: 1b. Average endpoint score (SANS, psychotic symptoms - negative, high score=worse, skewed data)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 5 Mental state: 1b. Average endpoint score (SANS, psychotic symptoms - negative, high score=worse, skewed data)




Analysis 3.6. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 6 Mental state: 2a. Average endpoint score anxiety (HRSA, high score=worse, skewed data)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 6 Mental state: 2a. Average endpoint score anxiety (HRSA, high score=worse, skewed data)




Analysis 3.7. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 7 Mental state: 2b. Average endpoint score depression (HRSD, high score=worse, skewed data)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 7 Mental state: 2b. Average endpoint score depression (HRSD, high score=worse, skewed data)




Analysis 3.8. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 8 Mental state: 2c. Average endpoint score mania (YMS, high score=worse, skewed data)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 8 Mental state: 2c. Average endpoint score mania (YMS, high score=worse, skewed data)




Analysis 3.9. Comparison 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 9 Quality of life: Average endpoint score (QLS, high score=worse)

Review: Early intervention for psychosis

Comparison: 3 PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 9 Quality of life: Average endpoint score (QLS, high score=worse)




Analysis 4.1. Comparison 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS, Outcome 1 Leaving the study early

Review: Early intervention for psychosis

Comparison: 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS

Outcome: 1 Leaving the study early




Analysis 4.2. Comparison 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS, Outcome 2 Mental state: 1. PANSS, endpoint score (by 12 weeks, higher scores=worse, LOCR)

Review: Early intervention for psychosis

Comparison: 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS

Outcome: 2 Mental state: 1. PANSS, endpoint score (by 12 weeks, higher scores=worse, LOCR)




Analysis 4.3. Comparison 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS

INTERVENTIONS, Outcome 3 Mental state: 1. MADRS, endpoint score (by 12 weeks, higher scores=worse, LOCF)

Review: Early intervention for psychosis

Comparison: 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS

Outcome: 3 Mental state: 1. MADRS, endpoint score (by 12 weeks, higher scores=worse, LOCF)




Analysis 4.4. Comparison 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS, Outcome 4 Global state: 1. GAF, endpoint score (by 12 weeks, higher scores=better)

Review: Early intervention for psychosis

Comparison: 4 PHASE-SPECIFIC TREATMENT (AMISULPRIDE) + NEEDS FOCUSED INTERVENTIONS vs NEEDS FOCUSED INTERVENTIONS

Outcome: 4 Global state: 1. GAF, endpoint score (by 12 weeks, higher scores=better)




Analysis 5.1. Comparison 5 OMEGA 3 FATTY ACIDS (EPA) versus PLACEBO, Outcome 1 Transition to psychosis

Review: Early intervention for psychosis

Comparison: 5 OMEGA 3 FATTY ACIDS (EPA) versus PLACEBO

Outcome: 1 Transition to psychosis




Analysis 6.1. Comparison 6 PHASE-SPECIFIC TREATMENT (CBT for SUICIDALITY) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 1 Leaving the study early - by 6 months

Review: Early intervention for psychosis

Comparison: 6 PHASE-SPECIFIC TREATMENT (CBT for SUICIDALITY) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 1 Leaving the study early - by 6 months




Analysis 6.2. Comparison 6 PHASE-SPECIFIC TREATMENT (CBT for SUICIDALITY) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 2 Suicide - by 6 months

Review: Early intervention for psychosis

Comparison: 6 PHASE-SPECIFIC TREATMENT (CBT for SUICIDALITY) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 2 Suicide - by 6 months




Analysis 7.1. Comparison 7 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + SPECIALISED TEAM vs SPECIALISED TEAM, Outcome 1 Relapse by end of treatment - by 12 months

Review: Early intervention for psychosis

Comparison: 7 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + SPECIALISED TEAM vs SPECIALISED TEAM

Outcome: 1 Relapse by end of treatment - by 12 months




Analysis 8.1. Comparison 8 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs STANDARD CARE, Outcome 1 Leaving the study early - by 18 months

Review: Early intervention for psychosis

Comparison: 8 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs STANDARD CARE

Outcome: 1 Leaving the study early - by 18 months




Analysis 8.2. Comparison 8 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs STANDARD CARE, Outcome 2 Readmitted to hospital - by 18 months

Review: Early intervention for psychosis

Comparison: 8 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs STANDARD CARE

Outcome: 2 Readmitted to hospital - by 18 months




Analysis 8.3. Comparison 8 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs STANDARD CARE, Outcome 3 Not compliant with medication

Review: Early intervention for psychosis

Comparison: 8 PHASE-SPECIFIC TREATMENT (FAMILY THERAPY) + STANDARD CARE vs STANDARD CARE

Outcome: 3 Not compliant with medication




Analysis 9.1. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 1 Leaving the study early

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 1 Leaving the study early




Analysis 9.2. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 2 Global state: 1. Average endpoint score - by 12 and 24 months (GAF-symptom, high score=good)

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 2 Global state: 1. Average endpoint score - by 12 and 24 months (GAF-symptom, high score=good)




Analysis 9.3. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 3 Global state: 2. Average endpoint score - by 12 and 24 months (GAF-function, high score=good)

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 3 Global state: 2. Average endpoint score - by 12 and 24 months (GAF-function, high score=good)




Analysis 9.4. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 4 User satisfaction: Average endpoint score - by 12 and 24 months (CSQ-8, high score=good)

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 4 User satisfaction: Average endpoint score - by 12 and 24 months (CSQ-8, high score=good)




Analysis 9.5. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 5 Compliance with treatment

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 5 Compliance with treatment




Analysis 9.6. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 6 Suicide: Death - by 12 months

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 6 Suicide: Death - by 12 months




Analysis 9.7. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 7 Death other than suicide - by 12 months

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 7 Death other than suicide - by 12 months




Analysis 9.8. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 8 Service use: 1. Average mean number of days per month in hospital

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 8 Service use: 1. Average mean number of days per month in hospital




Analysis 9.9. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 9 Service use: 2. Not hospitalised - by five years

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 9 Service use: 2. Not hospitalised - by five years




Analysis 9.10. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 10 Social outcomes: 1. Not living independently

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 10 Social outcomes: 1. Not living independently




Analysis 9.11. Comparison 9 SPECIALISED TEAM vs STANDARD CARE, Outcome 11 Social outcomes: 2. Not working or in education

Review: Early intervention for psychosis

Comparison: 9 SPECIALISED TEAM vs STANDARD CARE

Outcome: 11 Social outcomes: 2. Not working or in education




Analysis 10.1. Comparison 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS, Outcome 1 Leaving the study early by 12 months

Review: Early intervention for psychosis

Comparison: 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS

Outcome: 1 Leaving the study early by 12 months




Analysis 10.2. Comparison 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS, Outcome 2 Hospitalised by 12 months

Review: Early intervention for psychosis

Comparison: 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS

Outcome: 2 Hospitalised by 12 months




Analysis 10.3. Comparison 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS, Outcome 3 Suicide by 12 months

Review: Early intervention for psychosis

Comparison: 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS

Outcome: 3 Suicide by 12 months




Analysis 10.4. Comparison 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS, Outcome 4 Social functioning: SOFRAS by 12 months (higher score=worse)

Review: Early intervention for psychosis

Comparison: 10 PHASE-SPECIFIC TREATMENT (CBT) + ANTIPSYCHOTICS vs BEFRIENDING + ANTIPSYCHOTICS

Outcome: 4 Social functioning: SOFRAS by 12 months (higher score=worse)




Analysis 11.1. Comparison 11 PHASE-SPECIFIC TREATMENT (E-EPA) + ATYPICALS vs PLACEBO + ATYPICALS, Outcome 1 Leaving the study early by 12 weeks

Review: Early intervention for psychosis

Comparison: 11 PHASE-SPECIFIC TREATMENT (E-EPA) + ATYPICALS vs PLACEBO + ATYPICALS

Outcome: 1 Leaving the study early by 12 weeks




Analysis 11.2. Comparison 11 PHASE-SPECIFIC TREATMENT (E-EPA) + ATYPICALS vs PLACEBO + ATYPICALS, Outcome 2 Global state: Not responded to treatment by 12 weeks

Review: Early intervention for psychosis

Comparison: 11 PHASE-SPECIFIC TREATMENT (E-EPA) + ATYPICALS vs PLACEBO + ATYPICALS

Outcome: 2 Global state: Not responded to treatment by 12 weeks




Analysis 12.1. Comparison 12 PHASE-SPECIFIC TREATMENT (BRIEF INTERVENTION) + ANTIPSYCHOTICS vs TREATMENT AS USUAL, Outcome 1 Leaving the study early by nine months

Review: Early intervention for psychosis

Comparison: 12 PHASE-SPECIFIC TREATMENT (BRIEF INTERVENTION) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome: 1 Leaving the study early by nine months




Analysis 12.2. Comparison 12 PHASE-SPECIFIC TREATMENT (BRIEF INTERVENTION) + ANTIPSYCHOTICS vs TREATMENT AS USUAL, Outcome 2 Hospital admission: Hospitalised

Review: Early intervention for psychosis

Comparison: 12 PHASE-SPECIFIC TREATMENT (BRIEF INTERVENTION) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome: 2 Hospital admission: Hospitalised




Analysis 13.1. Comparison 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL, Outcome 1 Leaving the study early (6 months)

Review: Early intervention for psychosis

Comparison: 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome: 1 Leaving the study early (6 months)




Analysis 13.2. Comparison 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL, Outcome 2 Mental state: 1. PANSS

Review: Early intervention for psychosis

Comparison: 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome: 2 Mental state: 1. PANSS




Analysis 13.3. Comparison 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL, Outcome 3 Mental state: 2. Calgary Depression Rating Scale

Review: Early intervention for psychosis

Comparison: 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome: 3 Mental state: 2. Calgary Depression Rating Scale




Analysis 13.4. Comparison 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL, Outcome 4 Quality of Life:

1. Heinrichs-Carpenter

Review: Early intervention for psychosis

Comparison: 13 PHASE-SPECIFIC TREATMENT (ACE) + ANTIPSYCHOTICS vs TREATMENT AS USUAL

Outcome: 4 Quality of Life: 1. Heinrichs-Carpenter




Analysis 14.1. Comparison 14 PHASE-SPECIFIC TREATMENT (VOCATIONAL INTERVENTION) + TAU vs TREATMENT AS USUAL, Outcome 1 Not employed

Review: Early intervention for psychosis

Comparison: 14 PHASE-SPECIFIC TREATMENT (VOCATIONAL INTERVENTION) + TAU vs TREATMENT AS USUAL

Outcome: 1 Not employed




Analysis 14.2. Comparison 14 PHASE-SPECIFIC TREATMENT (VOCATIONAL INTERVENTION) + TAU vs TREATMENT AS USUAL, Outcome 2 Leaving the study early

Review: Early intervention for psychosis

Comparison: 14 PHASE-SPECIFIC TREATMENT (VOCATIONAL INTERVENTION) + TAU vs TREATMENT AS USUAL

Outcome: 2 Leaving the study early




Analysis 15.1. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 1 Cannabis use: 1. Used cannabis in last 4 weeks

Review: Early intervention for psychosis

Comparison: 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS

Outcome: 1 Cannabis use: 1. Used cannabis in last 4 weeks



Analysis 15.2. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 2 Cannabis use: 2. Percentage days used cannabis in last 4 weeks (skewed data)

Cannabis use: 2. Percentage days used cannabis in last 4 weeks (skewed data)

Study	Intervention	Mean	SD	N
by 3 months - end of treatment				
Edwards-Australia	CAP	30.4	41.8	23
Edwards-Australia	PE	18.8	30.6	24
by 9 months - 6 months after end of treatment				
Edwards-Australia	CAP	32.4	44.9	23
Edwards-Australia	PE	19.3	30.4	24

Analysis 15.3. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 3 Cannabis use: 3. Severity of cannabis use (skewed data)

Cannabis use: 3. Severity of cannabis use (skewed data)

Study	Intervention	Mean	SD	N
by 3 months - end of treatment				
Edwards-Australia	CAP	1.4	1.4	23


Study	Intervention	Mean	SD	N
Edwards-Australia	PE	1.3	1.4	24
by 9 months - 6 months after end of treatment				
Edwards-Australia	CAP	1.4	1.4	23
Edwards-Australia	PE	1.3	1.5	24

Analysis 15.4. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 4 Global state: Average score (KAPQ total endpoint, higher=good)

Review: Early intervention for psychosis

Comparison: 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS

Outcome: 4 Global state: Average score (KAPQ total endpoint, higher=good)




Analysis 15.5. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 5 Mental state: 1. Average score (BPRS-E total endpoint, higher scores=poor)

Review: Early intervention for psychosis

Comparison: 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS

Outcome: 5 Mental state: 1. Average score (BPRS-E total endpoint, higher scores=poor)



Analysis 15.6. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 6 Mental state: 2. Average score (BPRS-PS total endpoint, higher scores=poor) (skewed data)

Mental state: 2. Average score (BPRS-PS total endpoint, higher scores=poor) (skewed data)

Study	Intervention	Mean	SD	N
by 3 months - end of treatment				
Edwards-Australia	CAP	8.9	4.8	23
Edwards-Australia	PE	9.5	5.4	24
by 9 months - 6 months after end of treatment				
Edwards-Australia	CAP	9.4	4.6	23
Edwards-Australia	PE	8.8	4.8	24

Analysis 15.7. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 7 Mental state: 3. Average negative symptom score (SANS endpoint, higher scores=poor) (skewed data)

Mental state: 3. Average negative symptom score (SANS endpoint, higher scores=poor) (skewed data)

Study	Intervention	Mean	SD	N
by 3 months - end of treatment				

Study	Intervention	Mean	SD	N
Edwards-Australia	CAP	21.8	14.9	23
Edwards-Australia	PE	23.5	14.0	24
by 9 months - 6 months after end of treatment				
Edwards-Australia	CAP	23.7	17.2	23
Edwards-Australia	PE	19.4	13.5	24

Analysis 15.8. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 8 Mental state: 4. Average score (BDI-SF total endpoint , higher scores=poor) (skewed data)

Mental state: 4. Average score (BDI-SF total endpoint , higher scores=poor) (skewed data)


Study	Intervention	Mean	SD	N
by 3 months - end of treatment				
Edwards-Australia	CAP	6.2	5.9	23
Edwards-Australia	PE	7.8	8.1	24
by 9 months - 6 months after end of treatment				
Edwards-Australia	CAP	7.5	6.3	23
Edwards-Australia	PE	6.3	7.2	24

Analysis 15.9. Comparison 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS, Outcome 9 Social functioning: Average score (SOFAS total endpoint, higher scores=good)

Review: Early intervention for psychosis

Comparison: 15 PHASE-SPECIFIC TREATMENT (CANNABIS AND PSYCHOSIS THERAPY) + ANTIPSYCHOTICS vs PSYCHOEDUCATION + ANTIPSYCHOTICS

Outcome: 9 Social functioning: Average score (SOFAS total endpoint, higher scores=good)




Analysis 16.1. Comparison 16 CRISIS ASSESSMENT versus STANDARD CARE (LEO-CAT), Outcome 1 Hospitalisation

Review: Early intervention for psychosis

Comparison: 16 CRISIS ASSESSMENT versus STANDARD CARE (LEO-CAT)

Outcome: 1 Hospitalisation




Analysis 16.2. Comparison 16 CRISIS ASSESSMENT versus STANDARD CARE (LEO-CAT), Outcome 2 Referred to Mental Health Services by A&E or emergency medical services

Review: Early intervention for psychosis

Comparison: 16 CRISIS ASSESSMENT versus STANDARD CARE (LEO-CAT)

Outcome: 2 Referred to Mental Health Services by A&E or emergency medical services



Analysis 17.1. Comparison 17 EARLY BEHAVIOURAL INTERVENTION vs ROUTINE CARE INTERVAL (Alvarez-Spain), Outcome 1 Weight

Weight


Study	Intervention	Mean endpoint score	SD	N
Weight change at 13 weeks				
Alvarez-Spain	Early Behavioral Intervention	4.10	3.99	28
Alvarez-Spain	Routine Care Intervention	6.98	4.50	33
Body Mass Index				
Alvarez-Spain	Early Behavioral Intervention	1.40	1.34	28
Alvarez-Spain	Routine Care Intervention	2.39	1.53	33

Analysis 18.1. Comparison 18 PHASE-SPECIFIC INTERVENTION versus CONTROL (Exploratory meta-analysis), Outcome 1 Leaving the study early

Review: Early intervention for psychosis

Comparison: 18 PHASE-SPECIFIC INTERVENTION versus CONTROL (Exploratory meta-analysis)

Outcome: 1 Leaving the study early




Analysis 18.2. Comparison 18 PHASE-SPECIFIC INTERVENTION versus CONTROL (Exploratory meta-analysis), Outcome 2 Hospitalisation

Review: Early intervention for psychosis

Comparison: 18 PHASE-SPECIFIC INTERVENTION versus CONTROL (Exploratory meta-analysis)

Outcome: 2 Hospitalisation



Appendix 1. Previous searches for earlier versions of this review

1. Electronic search for update (March 2006)

We searched The Cochrane Schizophrenia Group Trials Register (July 2003 to March 2006) using the phrase: [early* in title, abstract or keywords of REFERENCE] or [Early* in intervention or 'prodromal or early*' in HealthCare Condition of STUDY]

This register is compiled by systematic searches of major databases, hand searches and conference proceedings (see Group Module).

2. Details of previous searches

We generated a list of relevant papers from our personal databases. On the basis of the indexing of these papers, we developed the following searches:

2.1 Electronic searches

2.1.1 We searched The Cochrane Schizophrenia Group's Register (July 2003) using the following phrase:

[Early* in intervention or 'prodromal or early*' in Health Care Condition of STUDY] or [early* in title, abstract or keywords of REFERENCE]

2.1.2 We searched CINAHL (1982 to November 2002, Ovid online) using the following phrase:

1. exp SCHIZOPHRENIA/ or exp SCHIZOPHRENIA, CATATONIC/ or exp SCHIZOPHRENIA, CHILDHOOD/ or exp SCHIZOPHRENIA, DISORGANIZED/ or exp SCHIZOPHRENIA, PARANOID/
 2. exp Paranoid Disorders/
 3. (schizo\$ or psychotic\$ or psychosis or psychoses or hebephreni\$ or oligophreni\$).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
 4. ((CHRONIC\$ or SEVER\$) adj5 MENTAL\$ adj5 (ILL\$ or DISORDER\$)).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
 5. 1 or 2 or 3 or 4
 6. ((risk\$ adj3 schiz\$) or (screen\$ adj3 schiz\$)).mp.
 7. ((duration or length) adj3 untreat\$).mp.
 8. ((first or initial or primary) adj3 (admission\$ or hospital\$ or episod\$ or breakdown\$)).mp.
 9. (early adj3 (intervent\$ or treat\$ or recogni\$ or detect\$)).mp.
 10. (delay\$ adj3 treat\$).mp.
 11. (' (DUP) ' or premorbid\$ or prodrom\$).mp.
 12. 6 or 7 or 8 or 9 or 10 or 11
 13. 12 and 5
 14. (animal not human).mp.
 15. 13 not 14
- 2.1.3 We searched The Cochrane Controlled Trials Register (November 2001) using the following phrase:

1. exp SCHIZOPHRENIA/ or exp SCHIZOPHRENIA, CATATONIC/ or exp SCHIZOPHRENIA, CHILDHOOD/ or exp SCHIZOPHRENIA, DISORGANIZED/ or exp SCHIZOPHRENIA, PARANOID/
2. exp Paranoid Disorders/
3. (schizo\$ or psychotic\$ or psychosis or psychoses or hebephreni\$ or oligophreni\$).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
4. ((CHRONIC\$ or SEVER\$) adj5 MENTAL\$ adj5 (ILL\$ or DISORDER\$)).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
5. 1 or 2 or 3 or 4
6. ((risk\$ adj3 schiz\$) or (screen\$ adj3 schiz\$)).mp.
7. ((duration or length) adj3 untreat\$).mp.
8. ((first or initial or primary) adj3 (admission\$ or hospital\$ or episod\$ or breakdown\$)).mp.
9. (early adj3 (intervent\$ or treat\$ or recogni\$ or detect\$)).mp.
10. (delay\$ adj3 treat\$).mp.
11. (‘ (DUP) ’ or premorbid\$ or prodrom\$).mp.
12. 6 or 7 or 8 or 9 or 10 or 11
13. 12 and 5
14. (animal not human).mp.
15. 13 not 14

21.4 We searched Embase (1966 to November 2002, Ovid online) using the following phrase:

1. exp SCHIZOPHRENIA/ or exp SCHIZOPHRENIA, CATATONIC/ or exp SCHIZOPHRENIA, CHILDHOOD/ or exp SCHIZOPHRENIA, DISORGANIZED/ or exp SCHIZOPHRENIA, PARANOID/
2. exp Paranoid Disorders/
3. (schizo\$ or psychotic\$ or psychosis or psychoses or hebephreni\$ or oligophreni\$).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]

4. ((CHRONIC\$ or SEVER\$) adj5 MENTAL\$ adj5 (ILL\$ or DISORDER\$)).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
5. 1 or 2 or 3 or 4
6. ((risk\$ adj3 schiz\$) or (screen\$ adj3 schiz\$)).mp.
7. ((duration or length) adj3 untreat\$).mp.
8. ((first or initial or primary) adj3 (admission\$ or hospital\$ or episod\$ or breakdown\$)).mp.
9. (early adj3 (intervent\$ or treat\$ or recogni\$ or detect\$)).mp.
10. (delay\$ adj3 treat\$).mp.
11. (' (DUP) ' or premorbid\$ or prodrom\$).mp.
12. 6 or 7 or 8 or 9 or 10 or 11
13. 12 and 5
14. (animal not human).mp.
15. 13 not 14

2.1.5 We searched Medline (1966 to November 2002, Ovid online) using the phrase:

1. exp SCHIZOPHRENIA/ or exp SCHIZOPHRENIA, CATATONIC/ or exp SCHIZOPHRENIA, CHILDHOOD/ or exp SCHIZOPHRENIA, DISORGANIZED/ or exp SCHIZOPHRENIA, PARANOID/
2. exp Paranoid Disorders/
3. (schizo\$ or psychotic\$ or psychosis or psychoses or hebephreni\$ or oligophreni\$).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
4. ((CHRONIC\$ or SEVER\$) adj5 MENTAL\$ adj5 (ILL\$ or DISORDER\$)).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
5. 1 or 2 or 3 or 4
6. ((risk\$ adj3 schiz\$) or (screen\$ adj3 schiz\$)).mp.
7. ((duration or length) adj3 untreat\$).mp.
8. ((first or initial or primary) adj3 (admission\$ or hospital\$ or episod\$ or breakdown\$)).mp.
9. (early adj3 (intervent\$ or treat\$ or recogni\$ or detect\$)).mp.

10. (delay\$ adj3 treat\$).mp.
11. (' (DUP) ' or premorbid\$ or prodrom\$).mp.
12. 6 or 7 or 8 or 9 or 10 or 11
13. 12 and 5
14. (animal not human).mp.
15. 13 not 14

2.1.6 We searched PsychINFO (1872 to November 2002, Ovid online) using the following phrase:

1. exp SCHIZOPHRENIA/ or exp SCHIZOPHRENIA, CATATONIC/ or exp SCHIZOPHRENIA, CHILDHOOD/ or exp SCHIZOPHRENIA, DISORGANIZED/ or exp SCHIZOPHRENIA, PARANOID/
2. exp Paranoid Disorders/
3. (schizo\$ or psychotic\$ or psychosis or psychoses or hebephreni\$ or oligophreni\$).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
4. ((CHRONIC\$ or SEVER\$) adj5 MENTAL\$ adj5 (ILL\$ or DISORDER\$)).mp. [mp=title, abstract, subject headings, drug trade name, original title, device manufacturer, drug manufacturer name]
5. 1 or 2 or 3 or 4
6. ((risk\$ adj3 schiz\$) or (screen\$ adj3 schiz\$)).mp.
7. ((duration or length) adj3 untreat\$).mp.
8. ((first or initial or primary) adj3 (admission\$ or hospital\$ or episod\$ or breakdown\$)).mp.
9. (early adj3 (intervent\$ or treat\$ or recogni\$ or detect\$)).mp.
10. (delay\$ adj3 treat\$).mp.
11. (' (DUP) ' or premorbid\$ or prodrom\$).mp.
12. 6 or 7 or 8 or 9 or 10 or 11
13. 12 and 5
14. (animal not human).mp.
15. 13 not 14

Appendix 2. Risk of Bias

Assessment of methodological quality

We assessed the methodological quality of included trials in this review using the criteria described in the Cochrane Handbook (Higgins 2005) and the Jadad scale (Jadad 1996). The former is based on the evidence of a strong relationship between allocation concealment and direction of effect (Schulz 1995). We allocated non-randomised studies (of early detection only, see above) to Category C. We performed a sensitivity analysis excluding trials in randomisation Category C, and trials with a follow up rate of less than 80%. The categories are defined below:

- A. Low risk of bias (adequate allocation concealment)
- B. Moderate risk of bias (some doubt about the results)
- C. High risk of bias (inadequate allocation concealment). For the purpose of the analysis in this review, we excluded trials if they met the Cochrane Handbook criteria A or B.

The Jadad Scale measures a wider range of factors that impact on the quality of a trial. The scale includes three items:

1. Was the study described as randomised?
2. Was the study described as double-blind?
3. Was there a description of withdrawals and drop outs?

Each item receives one point if the answer is positive. In addition, a point can be deducted if either the randomisation or the blinding/masking procedures described are inadequate. For this review we used a cut-off of two points on the Jadad scale to check the assessment made by the Handbook criteria. However we did not use the Jadad Scale was to exclude trials.

HISTORY

Protocol first published: Issue 1, 2004

Review first published: Issue 2, 2004

Date	Event	Description
23 August 2006	New citation required and conclusions have changed	Substantive amendment

WHAT'S NEW

Last assessed as up-to-date: 3 June 2009.

Date	Event	Description
11 March 2011	New citation required and conclusions have changed	conclusions changed after addition of 11 new studies
16 February 2010	New search has been performed	Updated, 11 new studies added, conclusions changed.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

For the 2009 update we used a more conservative estimate of $I^2 > 50\%$ to indicate heterogeneity. We have extensively reformatted this review but not substantively changed any methods.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON

PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM compared to SPECIALISED TEAM for psychosis

Patient or population: patients with psychosis

Settings: Melbourne, Australia

Intervention: PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM

Comparison: SPECIALISED TEAM

Outcomes	Illustrative comparative risks* (95% CI)	Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk				
	Corresponding risk				
	SPECIALISED TEAM				
	PHASE SPECIFIC TREATMENT (RISPERIDONE + CBT) + SPECIALISED TEAM				
Leaving the study early - by 12 months Follow-up: 12 months	See comment	Not estimable	59 (1 study ²)	⊕⊕⊕⊕ moderate ¹	No participant left early: this would be unusual for most situations
Progression to psychosis - by 6 months Follow-up: 6 months	Low risk population ³ 200 per 1000 54 per 1000 (16 to 178)	RR 0.27 (0.08 to 0.89)	59 (1 study ²)	⊕⊕⊕⊕ moderate ¹	This important study was undertaken in a centre of excellence, received funds from industry, and has generated much debate and research in this area
	Medium risk population ³ 350 per 1000 95 per 1000 (28 to 311)				
	High risk population ³ 500 per 1000 135 per 1000 (40 to 445)				
Progression to psychosis - by 12 months Follow-up: 12 months	Low risk population ³ 200 per 1000 108 per 1000 (46 to 260)	RR 0.54 (0.23 to 1.3)	59 (1 study ²)	⊕⊕⊕⊕ moderate ¹	
	Medium risk population ³ 350 per 1000 189 per 1000 (81 to 455)				
	High risk population ³ 500 per 1000 270 per 1000 (115 to 650)				

The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Limitations in design: rated - 'serious'. Randomisation generation not clear, sequence concealment not clear, outcomes not blindly rated, attrition not reported, unclear if selective reporting.

² Benchmark study.

³ Medium control risk is that of the control group of the trial.

ADDITIONAL SUMMARY OF FINDINGS

SPECIALISED TEAM compared to STANDARD CARE for psychosis

Patient or population: patients with psychosis

Settings: Scandinavia

Intervention: SPECIALISED TEAM

Comparison: STANDARD CARE

Outcomes	Illustrative comparative risks* (95% CI)	Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk			
	STANDARD CARE	SPECIALISED TEAM			
Compliance with treatment - treatment stopped in spite of need - by one year Follow-up: 12 months	Low risk population ¹		507 (1 study)	⊕⊕⊕○ moderate ²	
	100 per 1000	20 per 1000 (10 to 42)			
	Medium risk population ¹				
	150 per 1000	30 per 1000 (15 to 63)			
	High risk population ¹				
200 per 1000	40 per 1000 (20 to 84)				
Compliance with treatment - treatment stopped in spite of need - by two years Follow-up: 24 months	Low risk population ¹		436 (1 study)	⊕⊕⊕○ moderate ²	
	20 per 1000	13 per 1000 (6 to 30)			
	Medium risk population ¹				
	60 per 1000	40 per 1000 (17 to 90)			
	High risk population ¹				
100 per 1000	66 per 1000 (29 to 150)				
Service use: 1. Average mean number of days per month in hospital - by 5 years Follow-up: 5 years			547 (1 study)	⊕⊕⊕○ moderate ²	
		The mean Service use: 1. Average mean number of days per month in hospital - by 5 years in the intervention groups was 1.11 lower (3.21 lower to 0.99 higher)			

Outcomes	Illustrative comparative risks* (95% CI)	Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk				
	Corresponding risk				
	STANDARD CARE				
	SPECIALISED TEAM				
Service use: 2. Not hospitalised - by five years Follow-up: 5 years	Low risk population [†]	RR 1.05 (0.9 to 1.22)	547 (1 study)	⊕⊕⊕○ moderate ²	
	300 per 1000	315 per 1000 (270 to 366)			
	Medium risk population [†]				
	500 per 1000	525 per 1000 (450 to 610)			
	High risk population [†]				
	700 per 1000	735 per 1000 (630 to 834)			
Social outcomes: 1. Not living independently - by 5 years Follow-up: 5 years	Low risk population [†]	RR 0.42 (0.21 to 0.83)	547 (1 study)	⊕⊕⊕○ moderate ²	
	50 per 1000	21 per 1000 (10 to 41)			
	Medium risk population [†]				
	100 per 1000	42 per 1000 (21 to 83)			
	High risk population [†]				
	150 per 1000	63 per 1000 (31 to 124)			
Social outcomes: 2. Not working or in education - by 5 years Follow-up: 5 years	Low risk population [†]	RR 1.06 (0.92 to 1.23)	547 (1 study)	⊕⊕⊕○ moderate ²	
	200 per 1000	212 per 1000 (184 to 246)			
	Medium risk population [†]				
	500 per 1000	530 per 1000			

Outcomes	Illustrative comparative risks* (95% CI)	Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk				
	Corresponding risk				
	STANDARD CARE				
	SPECIALISED TEAM				
	(460 to 615)				
	High risk population /				
	800 per 1000				
					848 per 1000

Outcomes	Illustrative comparative risks* (95% CI)	Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk				
	Corresponding risk				
	STANDARD CARE				
	SPECIALISED TEAM				
	(736 to 984)				

The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR**: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Medium control risk is that of the control group of the trial.

² Limitations in design: rated 'Serious'. There were limitations but rating 'serious' may be harsh (but limited choice). Attempts were made to address these all though the study. Blinding not clear. Selective reporting possible.

References to studies included in this review

- Alvarez-Spain {published data only} . Alvarez-Jimenez M, Gonzalez-Blanch C, Vazquez-Barquero JL, Perez-Iglesias R, Martinez-Garcia O, Perez-Pardal T, Ramirez-Bonilla ML, Crespo-Facorro B. Attenuation of antipsychotic-induced weight gain with early behavioral intervention in drug-naive first-episode psychosis patients: a randomized controlled trial. *Journal of Clinical Psychiatry*. 2006; 67(8):1253–60. [EMBASE: 2006468531]. [PubMed: 16965204]
- Alvarez M, Gonzalez-Blanch C, Perez-Iglesias R, Perez-Pardal T, Martinez-Garcia O, Crespo-Facorro B, Vazquez-Barquero JL. Early intervention in antipsychotic - induced weight gain in first episode psychosis. *Schizophrenia Bulletin*. 2005; 31:518.
- Amminger-Austria {published data only} . Amminger G, Schaefer MR, Papageorgiou K, Becker J, Mossaheb N, Harrigan SM, McGorry PD, Berger GE. Omega-3 fatty acids reduce the risk of early transition to psychosis in ultra-high risk individuals: a double blind randomized, placebo-controlled treatment study. *Schizophrenia Bulletin*. 2007; 33(2):418–9. Amminger GP. Indicated prevention with omega-3 fatty acids in adolescents with ‘at-risk-mental-state’ for psychosis: a randomised, double blind, placebo-controlled treatment trial. 2006 <http://www.clinicaltrials.gov>.
- Amminger GP, Schafer MR. Indicated prevention with omega-3 fatty acids in adolescents at ultra-high risk for psychosis - rationale, methods, and 3-months outcome. *Schizophrenia Research*. 2006; 86(Suppl 1):S97–8. Amminger GP, Schafer MR. Is it feasible to conduct a RCT in ultra-high risk individuals at a child and adolescent psychiatric service? *Schizophrenia Research*. 2006; 86(Suppl 1):S98. Schafer MR, Klier CM, Papageorgiou K, Friedrich MH, Amminger GP. Early detection of psychotic disorders. *Neuropsychiatrie*. 2007; 21(1):37–44. [PubMed: 17555006]
- Berger-Australia {published data only} . Berger G, Wood S, Proffitt T, McConchie M, Khan A, O'Donnell C, Yuen H, Smith D, Horrobin D, McGorry P. Ethyl-eicosapentaenoic acid (e-epa) supplementation in early psychosis. A double-blind randomized add on standard therapy study in 80 drug- native or early treated first episode psychosis patients. *Schizophrenia Research*. 2004; 70(1):41. [: ISI:000224551100105]. Berger GB, Proffitt TM, McConchie MA, Wood SJ, Yuen HP, Smith D, Horrobin D, McGorry PD. Ethyleicosapentaenoic acid (e-epa) supplementation in early psychosis. *Schizophrenia Research*. 2004; 67(1):7–8. [: ISI 000188788100014]. Berger GE, Proffitt T-M, McConchie M, Yuen H, Wood SJ, Amminger GP, Brewer W, McGorry PD. Ethyl-eicosapentaenoic acid in first-episode psychosis: a randomized, placebo-controlled trial. *Journal of Clinical Psychiatry*. 2007; 68(12):1867–75. [MEDLINE: 18162017]. [PubMed: 18162017]
- Berger GE, Proffitt TM, McConchie MA, Wood SJ, Yuen HP, McGorry PD. Ethyl - eicosapentaenoic acid (e - epa) supplementation in early psychosis. A double - blind, randomized, placebo - controlled trial (RCT) comparing 2g e - epa versus placebo add - on therapy in 80 drug - naive or early treated first - episode psychos. *Schizophrenia Bulletin*. 2005; 31:475. Berger GE, Wood SJ, Wellard RM, Proffitt TM, McConchie M, Amminger GP, Jackson GD, Velakoulis D, Pantelis C, McGorry PD. Ethyl-eicosapentaenoic acid in first-episode psychosis. A 1H-MRS study. *Neuropsychopharmacology*. 2008; 33(10):2467–73. [MEDLINE: 18199999]. [PubMed: 18199999] McConchie MA, Berger GE, Proffitt TM, Yuen HP, Wood S, Smith D, Horrobin D, McGorry PD. Effect of diagnostic heterogeneity on response to ethyl-eicosapentaenoic acid (epa) in first-episode psychosis. *Schizophrenia Research*. 2004; 67(1):147–8. [: ISI 000188788100352].
- EDIE-UK {published data only} . French P, Shryane N, Bentall RP, Lewis SW, Morrison AP. Effects of cognitive therapy on the longitudinal development of psychotic experiences in people at high risk of developing psychosis. *British Journal of Psychiatry - Supplementum*. 2007; 51:s82–7. Lewis S. EDIE - Early detection and intervention for psychosis. *National Research Register*. 2001; Vol. 3. Morrison A. Early detection and intervention for psychosis in primary care. *National Research Register*. 2003. Morrison A. Follow-up of prodromal symptoms. *National Research Register*. 2004; Vol. 3. Morrison AP. Cognitive therapy for the prevention of psychosis in people at ultra-high risk: results of a randomised controlled trial. *Schizophrenia Research*. 2006; 86(Suppl 1):s59. Morrison AP, Bentall RP, French P, Walford L, Kilcommons A, Knight A, Kreutz M, Lewis SW. Randomised controlled trial of early detection and cognitive therapy for preventing transition to psychosis in high-risk individuals. Study design and interim analysis of transition rate and psychological risk factors. *British Journal of Psychiatry*. 2002; 43:s78–84. Morrison AP, French P, Parker S, Roberts M, Stevens H, Bentall RP, Lewis SW. Three-year

follow-up of a randomized controlled trial of cognitive therapy for the prevention of psychosis in people at ultrahigh risk. *Schizophrenia Bulletin*. 2007; 33(3):682–7. [EMBASE: 2007428663; MEDLINE: 16973786; PSYCINFO: 2007–08119–011]. [PubMed: 16973786] *Morrison AP, French P, Walford L, Lewis SW, Kilcommons A, Green J, Parker S, Bentall RP. Cognitive therapy for the prevention of psychosis in people at ultra-high risk: randomised controlled trial. *British Journal of Psychiatry*. 2004; 185:291–7. [PubMed: 15458988] Morrison T. Early detection and intervention for psychosis in primary care. *National Research Register*. 2001; Vol. 1 Renton J, Morrison AP. Effectiveness of cognitive therapy for psychosis and implications for early intervention. *Schizophrenia Research*. 2004; 70(1):142. [ISI: 000224551100421].

Edwards-Australia {published data only} . Edwards J, Elkins K, Hinton M, Harrigan SM, Donovan K, Athanasopoulos O, McGorry PD. Randomized controlled trial of a cannabis-focused intervention for young people with first-episode psychosis. *Acta Psychiatrica Scandinavica*. 2006; 114:109–17. [PubMed: 16836598]

EIPS-Germany {published data only} . Bechdorf A. Development and pilot evaluation of modified cognitive behavioural therapy for adolescents with EARLY onset psychosis. 2007 <http://www.clinicaltrials.gov>. Bechdorf A. Psychological intervention for persons at risk of psychosis in the early initial prodromal state. 2005 <http://www.clinicaltrials.gov>. Bechdorf A, Buhler B, Berning J, Wagner M, Stamm E, Streit M. Cognitive behavioural therapy in the early initial prodromal state of psychosis: first results. *Schizophrenia Research*. 2004; 67(1):202. [ISI 000188788100483]. Bechdorf A, Klosterkötter J. Cognitive-behavioural treatment (CBT) in the early initial prodromal state of psychosis: concept and practical approach. *Schizophrenia Research*. 2004; 70(1):52. [ISI:000224551100137]. Bechdorf A, Ruhrmann S, Janssen B, Bottlender R, Wagner M, Maurer K, Hafner H, Maier W, Klosterkötter J. Early recognition and intervention for people at risk of schizophrenia [FRUHERKENNUNG UND – INTERVENTION BEI PERSONEN MIT ERHOHEMTEM PSYCHOSERISIKO]. *Psychoneuro*. 2004; 30(11):606–14. [EMBASE 2004516569]. Bechdorf A, Veith V, Berning J, Stamm E, Decker P, Janssen B, Bottlender R, Wagner M, Klosterkötter J. Cognitive behavioral therapy (CBT) in the early initial prodromal state of psychosis: first results of a randomized trial. *Schizophrenia Research*. 2004; 70(1):62–3. [ISI: 000224551100168]. Bechdorf A, Wagner M, Veith V, Ruhrmann R, Janssen B, Bottlender R, et al. A randomised controlled multicenter trial of cognitive behaviour therapy in the early initial prodromal state of psychosis. *Schizophrenia Research*. 2006; 86(Suppl 1):S8. Bechdorf A, Wagner M, Veith V, Ruhrmann S, Pukrop R, Brockhaus-Dumke A, Berning J, Stamm E, Janssen B, Decker P, Bottlender R, Moller H-J, Gaebel W, Maier W, Klosterkötter J. Randomized controlled multicentre trial of cognitive behaviour therapy in the early initial prodromal state: effects on social adjustment post treatment. *Early Intervention in Psychiatry*. 2007; 1(1):71–8. [PSYCINFO: 2008–07770–011]. [PubMed: 21352110] Klosterkötter J. Early pharmacological and psychological intervention for late prodromal states of psychosis. <http://www.clinicaltrials.gov>. Lammertink, M. FETZ the first German early recognition and intervention centre for schizophrenic psychoses - how it works and what it aims for. *Proceedings of the 2nd International Conference on Early Psychosis*; New York, USA. 2000 Mar 31-Apr 2; 2000. [ISI 000188788100381]

Jackson-Australia {published data only} . Jackson HJ, Killackey E, Bendall S, Allott K, Dudgeon P, Harrigan S. A randomised controlled trial of CBT for early psychosis with 1 year follow-up. *Australian and New Zealand Journal of Psychiatry*. 2005; 39(Suppl 2):A42. [CINAHL: 2009775290; PSYCINFO: 2007–19131–008]. *Jackson HJ, McGorry PD, Killackey E, Bendall S, Allott K, Dudgeon P, Gleeson J, Johnson T, Harrigan S. Acute-phase and 1-year follow-up results of a randomized controlled trial of CBT versus befriending for first-episode psychosis: the ACE project. *Psychological Medicine*. 2008; 38(5):725–35. [EMBASE: 2008165197; MEDLINE: 18005494]. [PubMed: 18005494] Killackey E, Jackson H, McGorry P, Bendall S, Allott K, Johnson T, Gleeson J, Harrigan S. The ACE project: a randomised controlled trial of CBT versus befriending for acute first episode psychosis: acute phase results. *Schizophrenia Bulletin*. 2005; 31:526–27. [ISI: 114097].

Killackey-Australia {published data only} . Killackey E, Jackson H, McGorry P. Vocational intervention in a specialist early psychosis service: first results of a pilot randomised controlled trial. *Schizophrenia Research*. 2006; 86(Suppl 1):S28–S29. Killackey E, Jackson HJ, McGorry PD. Vocational intervention in first-episode psychosis: individual placement and support versus treatment as usual. *British Journal of Psychiatry*. 2008; 193(2):114–20. [PubMed: 18669993]

- Leavey-UK {published data only} . Leavey G, Gulamhussein S, Papadopoulos C, Johnson-Sabine E, Blizard BKM. A randomized controlled trial of a brief intervention for families of patients with a first episode of psychosis. *Psychological Medicine*. 2004; 34(3):423–31. [MEDLINE: 15259827]. [PubMed: 15259827]
- LEO-CAT-UK {published data only} . Power P, Craig T, McGuire P, Iacoponi E, Garety P, Russell M. A randomised controlled trial of an early detection team in first episode psychosis: the leo cat trial. *Schizophrenia Research*. 2004; 67(1):36. [ISI 000188788100082]. Power P, Iacoponi E, Reynolds N, Fisher H, Russell M, Garety P, McGuire PK, Craig T. The Lambeth early onset crisis assessment team crisis assessment study: general practitioner education and access to an early detection team in first-episode psychosis. *British Journal of Psychiatry*. 2007; (Supp 51):s133–9. Power P, Iacoponi E, Russell M, Fisher H, McGuire P, Garety P, Valmaggio L, Craig T. A randomised controlled trial of an early detection team in first- episode psychosis: provisional findings of the leo cat study. *Schizophrenia Research*. 2004; 70(1):131. [ISI: 000224551100386].
- LifeSPAN-Australia {published data only} . Power, P.; Bell, R.; Mills, R.; Herrmann-Doig, T.; Davern, M.; Henry, L.; McGorry, P.; Khademy-Deljo, A. A randomized controlled trial of suicide prevention therapy for young people with first episode psychosis. *Proceedings of the Second International Conference on Early Psychosis; New York, USA*. 2000 March 31-2; 2000. Power P, Bell R, Mills R, Herrmann-Doig T, Davern M, Yuen H, McGorry P. A randomised controlled trial of a suicide preventative cognitive oriented psychotherapy for suicidal young people with first episode psychosis. *Schizophrenia Research*. 1999; 36:332. *Power PJ, Bell RJ, Mills R, Herrman-Doig T, Davern M, Henry L, Yuen HP, Khademy-Deljo A, McGorry PD. Suicide prevention in first episode psychosis: the development of a randomised controlled trial of cognitive therapy for acutely suicidal patients with early psychosis. *Australian and New Zealand Journal of Psychiatry*. 2003; 37(4):414–20. [MEDLINE: 12873325]. [PubMed: 12873325]
- Linszen-Amsterdam {published data only} . Lenior ME, Dingemans PM, Schene AH, Linszen DH. Predictors and five-year outcome in early-onset schizophrenia. A path analysis. *Schizophrenia Research*. 2003; 60:292. Lenior ME, Dingemans PMAJ, Schene AH, Linszen DH. Predictors of the early 5-year course of schizophrenia: a path analysis. *Schizophrenia Bulletin*. 2005; 31(3): 781–91. [PubMed: 16123531] Linszen D. Early and critical period intervention in first episode schizophrenia: relapse, chronicity, early stabilisation, predictors over 4 years and new research. *Schizophrenia Research*. 2004; 70(1):66. [ISI:000224551100177]. *Linszen D, Dingemans P, Lenior ME. Early intervention and a five-year follow-up in young adults with a short duration of untreated psychosis: ethical implications. *Schizophrenia Research*. 2001; 51:55–61. [PubMed: 11479066] Linszen D, Dingemans P, Van Der Does JW, Nugter A, Scholte P, Lenior R, Goldstein MJ. Treatment, expressed emotion and relapse in recent onset schizophrenic disorders. *Psychological Medicine*. 1996; 26(2):333–42. [PubMed: 8685289] Linszen D, Dingemans PM, Lenior ME, Scholte WF, Goldstein M. Early family and individual interventions and relapse in recent-onset schizophrenia and related disorders. *Italian Journal of Psychiatry & Behavioural Sciences*. 1998; 8:77–84. Linszen D, Dingemans PM, Scholte WF, Lenior ME, Goldstein M. Early recognition, intensive intervention and other protective and risk factors for psychotic relapse in patients with first psychotic episodes in schizophrenia. *International Clinical Psychopharmacology*. 1998; 13(Suppl):S7–S12. Linszen D, Lenior M, De Haan L, Dingemans P, Gersons B. Early intervention, untreated psychosis and the course of early schizophrenia. *British Journal of Psychiatry - Supplement*. 1998; 172(Suppl):84–9. [PubMed: 9764132] Linszen D, Wouters L, Dingemans P, De Haan L, Nieman D. Early and 3-year sustained intervention in first episode schizophrenia: relapse, stabilization and its predictors. *Schizophrenia Research*. 2004; 67(1):18. [ISI 000188788100038]. Linszen DH, De Haan L, Dingemans P, Van Bruggen M, Hofstra N, Van Engelsdorp H, Smitten M. Treatment reluctance in first episode schizophrenia: Lack of insight, non-compliance and cannabis abuse predict bad outcome after eighteen months intervention. *Schizophrenia Research*. 2003; 60:325. [EMBASE: 2005343078; MEDLINE: 12777272].
- LIPS-Germany {published data only} . Bechdolf A, Ruhrmann S, Wagner M, Kuhn KU, Janssen B, Bottlender R, Wieneke A, Schulze-Lutter F, Maier W, Klosterkötter J. Interventions in the initial prodromal states of psychosis in Germany: concept and recruitment. *British Journal of Psychiatry Supplementum*. 2005; 187(Suppl 48):s45–8. [MEDLINE: 16055807]. Klosterkötter J. Secondary prevention of schizophrenia: a randomised controlled trial. *Current Controlled Trials*.

- Ruhrmann S, Bechdorf A, Kuhn KU, Wagner M, Schultze-Lutter F, Janssen B, Maurer K, Hafner H, Gaebel W, Moller HJ, Maier W, Klosterkotter J, LIPS study group. Acute effects of treatment for prodromal symptoms for people putatively in a late initial prodromal state of psychosis. *British Journal of Psychiatry Supplementum*. 2007; 51:s88–95. [MEDLINE: 18055944].
- Ruhrmann S, Hoppmann B, Theysohn S, Picker H, Kuhn KU, Schultze-Lutter F, Wagner M, Bechdorf A, Moller HJ, Gaebel W, Maier W, Klosterkotter J. Acute symptomatic treatment effects in persons clinically at risk for psychosis. *Schizophrenia Research*. 2006; 86(Suppl 1):S8.
- OPUS-Scandinavia {published data only} . Bertelsen M, Jeppesen P, Petersen L, Thorup A, Ohlenschlaeger J, le Quach P, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. Five-year follow-up of a randomized multicenter trial of intensive early intervention vs standard treatment for patients with a first episode of psychotic illness: the OPUS trial. *Archives of General Psychiatry*. 2008; 65(7):762–71. [MEDLINE: 18606949; PSYCHINFO: 2008–18369–003]. [PubMed: 18606949] Bertelsen M, Thorup A, Petersen L, Jeppesen P, Oehlenschlaeger J, Joergensen P, Le Quach P, Krarup G, Mortensen P, Nordentoft M. The OPUS trial: results from the five-year follow-up. *Schizophrenia Research*. 2006; 86(Suppl 1):S43. Jeppesen P, Petersen L, Thorup A, Abel MB, Oehlenschlaeger J, Christensen TO, Krarup G, Hemmingsen R, Jorgensen P, Nordentoft M. Integrated treatment of first-episode psychosis: effect of treatment on family burden: OPUS trial. *British Journal of Psychiatry*. 2005; 48:s85–90. Jeppesen P, Petersen L, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. The association between pre-morbid adjustment, duration of untreated psychosis and outcome in first-episode psychosis. *Psychological Medicine*. 2008; 8:1157–66. [PubMed: 18447961] Jorgensen P, Nordentoft M, Abel M, Gouliaev G, Jeppesen P, Kassow P. Early detection and assertive community treatment of young psychotics: the opus study rationale and design of the trial. *Social Psychiatry & Psychiatric Epidemiology*. 2000; 35:283–7. [PubMed: 11016522] Jorgensen P, Nordentoft M, Abel MB, Gouliaev G, Jeppesen P, Kassow P. Early detection and assertive community treatment of young psychotics: the opus study rationale and design of the trial. *Social Psychiatry Psychiatric Epidemiology*. 2000; 35(7):283–7. [PubMed: 11016522] Nordentoft M, Jeppesen P, Abel M, Kassow P, Petersen L, Thorup A, Krarup G, Hemmingsen R, Jorgensen P. OPUS study: suicidal behaviour, suicidal ideation and hopelessness among patients with first-episode psychosis. One-year follow-up of a randomised controlled trial. *British Journal of Psychiatry*. 2002; 43:s98–106. Nordentoft M, Jeppesen P, Abel M, Kassow P, Petersen L, Thorup A, Krarup G, Hemmingsen R, Jorgensen P. OPUS study: suicidal behaviour, suicidal ideation and hopelessness among patients with first-episode psychosis. One-year follow-up of a randomised controlled trial. *British Journal of Psychiatry Supplementum*. 2002; 181(Suppl 43):S98–106. [MEDLINE: 12271808; MEDLINE: 22233038]. Nordentoft, M.; Jeppesen, P.; Abel, M.; Petersen, L.; Thorup, A.; Christensen, T.; Øhlenschläger, J.; Jorgensen, P. Opus-project: a randomised controlled trial of integrated psychiatric treatment in first-episode psychosis - clinical outcome improved. *Proceeding of the 3rd International Conference on Early Psychosis; Copenhagen, Denmark*. 2002 Sep 25–28; 2002. p. 56 Nordentoft, M.; Jeppesen, P.; Jørgensen, P.; Abel, MB.; Kassow, P.; Reisby, N.; Hemmingsen, R. Opus - project : a randomised controlled trial of first episode psychotic patients better compliance. *Proceedings of the 2nd International Conference on Early Psychosis; New York, New York, USA*. 2000 Mar 31 - Apr 2; 2000. Nordentoft M, Jeppesen P, Kassow P, Abel M, Petersen L, Thorup A, Cristensen T, Øhlenschläger J, Jørgensen P. Opus-project: a randomised controlled trial of integrated psychiatric treatment in first-episode psychosis-clinical outcome improved. *Schizophrenia Research*. 2002; 53(3 Suppl 1):51. [MEDLINE: 11016522; MEDLINE: 20468737]. Nordentoft, M.; Jeppesen, P.; Petersen, L.; Thorup, A.; Abel, M. Opus project: a randomised controlled trial of integrated psychiatric treatment in first episode psychosis. *Proceedings of the 9th International Congress on Schizophrenia Research; Colorado Springs, USA*. 2003 Mar 29 - Apr 2; 2003. [CINAHL: 2009245707; MEDLINE: 16141449][CINAHL: 2009245707; MEDLINE: 16141449] Nordentoft M, Jeppesen P, Petersen L, Thorup A, Jorgensen P. Duration of untreated psychosis predicts psychotic symptoms but not negative symptoms. *Schizophrenia Bulletin*. 2005; 31:234. Nordentoft M, Petersen L, Jeppesen P, Thorup AA, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P. OPUS: a randomised multicenter trial of integrated versus standard treatment for patients with a first-episode psychosis--secondary publication. *Ugeskr Laeger*. 2006; 168(4):381–4. [PubMed: 16436240] Nordentoft, M.; Reisby, N.; Jeppesen, P.; Abel, M-B.; Kassow, P.; Jørgensen, P. Opus-project: differences in treatment outcome of a

randomised controlled trial of integrated psychiatric treatment of first-episode psychotic patients. Proceedings of the 11th World Congress of Psychiatry; Hamburg, Germany. 1999 Aug 6-11; 1999. p. 165

Nordentoft M, Thorup A, Petersen L, Ohlenschlaeger J, Melau M, Christensen TO, Krarup G, Jorgensen P, Jeppesen P. Transition rates from schizotypal disorder to psychotic disorder for first-contact patients included in the opus trial. A randomized clinical trial of integrated treatment and standard treatment. *Schizophrenia Research*. 2006; 83(1):29–40.

*Petersen L, Jeppesen P, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P, Nordentoft M, Petersen L, Jeppesen P, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. A randomised multicentre trial of integrated versus standard treatment for patients with a first episode of psychotic illness. *BMJ*. 2005; 331(7517):602. [PubMed: 16141449]

Petersen L, Jeppesen P, Thorup A, Ohlenschlaeger J, Christensen T, Krarup G, Jorgensen P, Nordentoft M. Substance abuse in first-episode schizophrenia-spectrum disorders. *Schizophrenia Research*. 2006; 86(Suppl 1):S44.

Petersen L, Jeppesen P, Thorup A, Ohlenschlaeger J, Krarup G, Ostergard T, Jorgensen P, Nordentoft M. Substance abuse and first-episode schizophrenia-spectrum disorders The Danish opus trial. *Early Intervention in Psychiatry*. 2007; 1(1):88–96. [PubMed: 21352112]

Petersen L, Nordentoft M, Jeppesen P, Ohlenschlaeger J, Thorup A, Christensen TO, Krarup G, Dahlstrom J, Haastrup B, Jorgensen P. Improving 1-year outcome in first-episode psychosis: Opus trial. *British Journal of Psychiatry*. 2005; 48(Suppl):s98–103. [MEDLINE: 16055817].

Petersen L, Nordentoft M, Thorup A, Ohlenschlaeger J, Jeppesen P, Christensen T, Krarup G, Joergensen P. The opus trial: a randomised multi-centre trial of integrated versus standard treatment for 547 first-episode psychotic patients. *Schizophrenia Bulletin*. 2005; 31:531.

Thorup A, Petersen L, Jeppesen P, Ohlenschlaeger J, Christensen T, Krarup G, Jorgensen P, Nordentoft M. Integrated treatment ameliorates negative symptoms in first episode psychosis - results from the Danish opus trial. *Schizophrenia Research*. 2005; 79(1):95–105. [PubMed: 16122909]

Thorup A, Petersen L, Jeppesen P, Ohlenschlaeger J, Christensen T, Krarup G, Jorgensen P, Nordentoft M. Social network among young adults with first-episode schizophrenia spectrum disorders: results from the Danish opus trial. *Social Psychiatry and Psychiatric Epidemiology*. 2006; 41(10):761–70. [MEDLINE: 16900304]. [PubMed: 16900304]

PACE-Australia {published data only} . McGorry PD, Yung AR, Phillips LJ, Yuen HP, Francey S, Cosgrave EM, Germano D, Bravin J, McDonald T, Blair A, Adlard S, Jackson H. Randomized controlled trial of interventions designed to reduce the risk of progression to first-episode psychosis in a clinical sample with subthreshold symptoms. *Archives of General Psychiatry*. 2002; 59(10):921–8. [PubMed: 12365879]

Phillips LJ, Leicester SB, O'Dwyer LE, Francey SM, Koutsogiannis J, Abdel-Baki A, Kelly D, Jones S, Vay C, Yung AR, McGorry PD. The PACE Clinic: identification and management of young people at 'ultra' high risk of psychosis. *Journal of Psychiatric Practice*. 2002; 8(5):255–69.

Phillips LJ, McGorry PD, Yuen HP, Ward J, Donovan K, Kelly D, Francey SM, Yung AR. Medium term follow-up of a randomized controlled trial of interventions for young people at ultra high risk of psychosis. *Schizophrenia Research*. 2007; 96(1-3):25–33. [PubMed: 17611080]

Phillips LJ, Yung AR, Yuen HP, Pantelis C, McGorry PD. Prediction and prevention of transition to psychosis in young people at incipient risk for schizophrenia. *American Journal of Medical Genetics*. 2002; 114(8):929–37. [PubMed: 12457389]

PRIME-USA {published data only} . Hawkins KA, Addington J, Keefe R, Christensen B, Woods S, Zipursky R, Perkins D, McGlashan T. Neuropsychological functioning in the first episode prodrome and early psychosis. *Schizophrenia Research*. 2004; 70(1):101. [ISI: 000224551100287]. [PubMed: 15246469]

Hawkins, KA.; Addington, J.; Keefe, RS.; Christensen, B.; Woods, SW.; Zipursky, RB.; Perkins, DO.; McGlashan, TH. Effect of olanzapine versus placebo on the neuropsychological status of prodromal subjects. Proceedings of the 12th Biennial Winter Workshop on Schizophrenia; Davos, Switzerland. 2004 Feb 7-13; 2004.

Hawkins KA, Keefe RSE, Christensen BK, Addington J, Woods SW, Callahan J, Zipursky RB, Perkins DO, Tohen M, Breier A, McGlashan TH. Neuropsychological course in the prodrome and first episode of psychosis: findings from the prime North America double blind treatment study. *Schizophrenia Research*. 2008; 105(1-3):1–9. [EMBASE: 2008459795; PSYCHINFO: 2008–14504–002]. [PubMed: 18774696]

Hoffman RE, Woods S, Preda A, Tohen M, Breier A, Glist J, Addington J, Perkins DO, Hawkins K, McGlashan TH. Excessive top-down perceptual processing and reduced real-world investment exhibited by prodromal patients predict

- subsequent conversion to schizophrenia. *Schizophrenia Research*. 2006; 86(s1):S46. Hoffman RE, Woods SW, Hawkins KA, Pittman B, Tohen M, Preda A, Breier A, Glist J, Addington J, Perkins DO, McGlashan TH. Extracting spurious messages from noise and risk of schizophrenia-spectrum disorders in prodromal population. *British Journal of Psychiatry*. 2007; 191(2):355–6. [PubMed: 17906248] McGlashan, TH. Intervention in the prodrome to first psychosis. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 March 31-2; 2000. McGlashan, TH.; Miller, TJ.; Zipursky, RB.; Woods, SW.; Perkins, DO.; Hawkins, KA.; Addington, JM. Intervention in the schizophrenic prodrome: the prevention through risk identification, management, and education initiative. Proceeding of the 156th Annual Meeting of the American Psychiatric Association; San Francisco, California, USA. 2003 May 17-22; 2003. McGlashan TH, Vaglum P, Friis S, Johannessen JO, Simonsen E, Larsen TK, Melle I, Haahr U, Opjordsmoen S, Zipursky R, Perkins D, Addington J, Miller T, Woods S, Hoffman R, Preda A, Epstein I, Addington D, Lindborg S, Trzaskoma Q, Tohen M, Breier A. Early detection and intervention in first episode psychosis: empirical update of the tips and prime projects. *Schizophrenia Bulletin*. 2005; 31:496.*McGlashan TH, Zipursky RB, Perkins D, Addington J, Miller T, Woods SW, Hawkins KA, Hoffman RE, Preda A, Epstein I, Addington D, Lindborg S, Trzaskoma Q, Tohen M, Breier A. Randomized, double-blind trial of olanzapine versus placebo in patients prodromally symptomatic for psychosis. *American Journal of Psychiatry*. 2006; 163(5):790–9. [PubMed: 16648318] McGlashan TH, Zipursky RB, Perkins D, Addington J, Miller T, Woods SW, Hawkins KA, Hoffman RE, Preda A, Epstein L, Addington D, Lindborg S, Trzaskoma Q, Tohen M, Breier A. Randomized, double-blind trial of olanzapine versus placebo in patients prodromally symptomatic for psychosis. *American Journal of Psychiatry*. 2006; 163(5):790–9. [CINAHL: 2009175479; MEDLINE: 16648318]. [PubMed: 16648318] McGlashan TH, Zipursky RB, Perkins D, Addington J, Miller TJ, Woods SW, Hawkins KA, Hoffman R, Lindborg S, Tohen M, Breier A. The prime North America randomized double-blind clinical trial of olanzapine versus placebo in patients at risk of being prodromally symptomatic for psychosis. I. Study rationale and design. *Schizophrenia Research*. 2003; 61(1):7–18. [MEDLINE: 12648731]. [PubMed: 12648731] McGlashan TH, Zipursky RB, Perkins DO, Addington J, Woods SW, Miller TJ, Lindborg S, Marquez E, Hawkins K, Hoffman RE. Olanzapine versus placebo treatment of the schizophrenia prodrome: One year results. *Schizophrenia Research*. 2003; 60:295. [EMBASE: 2008459795; PSYCHINFO: 2008–14504–002]. McGlashan, TH.; Zipursky, RB.; Perkins, DO.; Addington, JM.; Woods, SW.; Lindborg, S.; Breier, AF. Olanzapine versus pbo for the schizophrenic prodrome: one-year results. Proceedings of the 156th Annual Meeting of the American Psychiatric Association; San Francisco, California, USA. 2003 May 17-22; 2003.
- Uzenoff-USA {published data only} . Uzenoff SR, Perkins DO, Hamer RM, Wiesen CA, Penn DL. A preliminary trial of adherence-coping-education (ACE) therapy for early psychosis. *Journal of Nervous and Mental Disease*. 2008; 196(7):572–5. [CINAHL: 2009976831; MEDLINE: 18626299; PSYCINFO: 2008–10312–009]. [PubMed: 18626299]
- Zhang-China {published data only} . Zhang M, Wang M, Li J, Phillips MR. Randomised-control trial of family intervention for 78 first-episode male schizophrenic patients: an 18-month study in Suzhou, Jiangsu. *British Journal of Psychiatry*. 1994; 165(Suppl 24):96–102.

References to studies excluded from this review

- Addington 1999 {published data only} . Addington J, Jones B, Ko T, Addington D. Family intervention in an early psychosis program. *Psychiatric Rehabilitation Skills*. In press. Addington, J.; Jones, B.; Ko, T.; Addington, D. Intervention strategies for families of first episode patients. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000. Addington J, McCleery A, Addington D. Three-year outcome of family work in an early psychosis program. *Schizophrenia Research*. 2005; 79(1):107–16. [PubMed: 15907375]
- Agius 2007 {published data only} . Agius M, Shah S, Ramkisson R, Murphy S, Zaman R. Three-year outcomes of an early intervention for psychosis service as compared with treatment as usual for first psychotic episodes in a standard community mental health team - final results. *Psychiatria Danubina*. 2007; 19(3):130–8. [MEDLINE: 17914313]. [PubMed: 17914313]

- Alanen 1994 {published data only} . Alanen, YO.; Lehtinen, V.; Lehtinen, K.; Aaltonen, J.; Rakkolainen, V. The Finnish Integrated Model for Early Treatment of Schizophrenia and Related Psychoses. Martindale, B.; Bateman, A., editors. Gaskell/Royal College of Psychiatrists; London, England: 2000. p. 235-65. Alanen, YO.; Ugelstad, E.; Armelius, BA.; Lehtinen, K.; Rosenbaum, B.; Sjostrom, R. Early Treatment for Schizophrenic Patients: Scandinavian Psychotherapeutic Approach. Scandinavian University Press; Oslo: 1994. Lehtinen K, Tuori T, Jaaskelainen NJ. Using psychosis teams in early intervention. A follow-up study of the Finnish schizophrenia project. *Nordic Journal of Psychiatry*. 1998; 52(Suppl):74. Lehtinen V, Aaltonen J, Koffert T, Rakkolainen V, Syvalahti E. Two-year outcome in first-episode psychosis treated according to an integrated model. Is immediate neuroleptisation always needed? *European Psychiatry*. 2000; 15:312–20. [PubMed: 10954876]
- Albiston 1998 {published data only} . Albiston D, Francey SM, Harrigan SM. Group programmes for recovery from early psychosis. *British Journal of Psychiatry*. 1998; 172(33):117–21.
- Anonymous 1987 {published data only} . Anonymous. The Scottish first episode schizophrenia study. ii. Treatment: pimozide versus flupenthixol. *British Journal of Psychiatry*. 1987; 150:334–8. [PubMed: 2889495]
- Bao 2005 {published data only} . Bao W-Q, Sun X-J, Wang M-L. Research on home intervention to community schizophreniform by applying PDCA. *Journal of Practical Nursing*. 2005; 21(3A): 9–11.
- Birchwood 1989 {published data only} . Birchwood M, Smith J, Macmillan F, Hogg B, Prasad R, Harvey C. Predicting relapse in schizophrenia: the development and implementation of an early signs monitoring system using patients and families as observers, a preliminary investigation. *Psychological Medicine*. 1989; 19:649–56. [PubMed: 2798634]
- Clare 1994 {published data only} . Clare L, Singh K. Preventing relapse in psychotic illness: a psychological approach to early intervention. *Journal of Mental Health*. 1994; 3:541–50.
- COPE-Melbourne {published data only} . Jackson, H.; McGorry, PD.; Edwar, SJ. Cognitively oriented psychotherapy for early psychosis: theory, praxis, outcomes, and challenges. In: Corrigan, PW.; Penn, DL., editors. *Social Cognition and Schizophrenia*. American Psychological Association; Washington, DC, US: 2001. Jackson H, McGorry P, Edwards J, Hulbert C, Henry L, Francey S, Maude D, Cocks J, Power P, Harrigan SDP. Cognitively-oriented psychotherapy for early psychosis (cope). Preliminary results. *British Journal of Psychiatry*. 1998; 172(Suppl): 93–100. Jackson H, McGorry P, Edwards J, Hulbert C, Henry L, Harrigan S, Dudgeon P, Francey S, Cocks J, Killackey E. A randomised controlled trial of cognitively-orientated psychotherapy for early psychosis (cope) with 5-year follow-up relapse data. *Schizophrenia Research*. 2004; 70(1):61. [: ISI:000224551100163]. Jackson H, McGorry P, Henry L, Edwards J, Hulbert C, Harrigan S, Dudgeon P, Francey S, Maude D, Cocks J, Power P. Cognitively oriented psychotherapy for early psychosis (cope): a 1-year follow-up. *British Journal of Clinical Psychology*. 2001; 40:57–70. [PubMed: 11317949] Jackson, HJ.; Edwards, J.; McGorry, PD.; Hulbert, C. The recognition and management of early psychosis: a preventive approach. In: McGorry, PD.; Jackson, HJ., editors. *Recovery From Psychosis: Psychological Interventions*. Cambridge University Press; Cambridge: 1999. p. 213-35. Jackson, HJ.; McGorry, PD.; Edwards, J.; Hulbe, TC. Cognitively oriented psychotherapy for early psychosis (cope). In: Cotton; Peter; Jackson; Henry, editors. *Early Intervention & Prevention in Mental Health*. Australian Psychological Society Ltd; Carlton South VIC, Australia: 1996.
- Craig 2004b {published data only} . Craig T, Garety P, Power P, Rahaman N, Colbert S, Fornells-Ambrojo M. Lambeth early onset service: a randomised controlled trial. *Schizophrenia Research*. 2004; 70(1):145–6. [: ISI:000224551100433]. Craig TKJ. Brixton early psychosis project. National Research Register. 2000. Craig TKJ, Garety P, Power P, Rahaman N, Colbert S, Fornells-Ambrojo M, Dunn G. The Lambeth early onset (LEO) Team: randomised controlled trial of the effectiveness of specialised care for early psychosis. *BMJ*. 2004; 329(7474):1067–70. [MEDLINE: 15485934; : CINAHL 2005085280]. [PubMed: 15485934]
- Crow 1986 {published data only} . Crow TJ, MacMillan J, Johnson AL, Johnstone EC. The Northwick Park study of first episodes of schizophrenia: ii. A randomised controlled trial of prophylactic neuroleptic treatment. *British Journal of Psychiatry*. 1986; 148:120–7. [PubMed: 2870753]

- Cullberg 1998 {published data only} . Cullberg J. Integrating intensive psychosocial therapy and low dose medical treatment in a total material of first episode psychotic patients compared to 'treatment as usual': a 3-year follow-up. *Medicinski Arhiv*. 1999; 53:167–70. [PubMed: 10546453] Cullberg J, Levander S, Stefans C. Symposium 7: early detection and intervention in psychosis. 'the parachute project' - a Swedish multicenter project for first episode psychosis. *Nordic Journal of Psychiatry*. 1998; 52(Suppl):73–4. Cullberg, J.; Thoren, G.; Abb, S.; Mesterton, A.; Svedb, B. Integrating intensive psychosocial and low-dose neuroleptic treatment: a three-year follow-up. In: Martindale, B.; Bateman, A., et al., editors. *Psychosis: Psychological Approaches and Their Effectiveness*. Gaskell/Royal College of Psychiatrists; London: 2000. p. 200-9.
- Davidson 2004 {published data only} . Davidson, M. Reducing the risk of early transition to psychosis: using long-acting atypical antipsychotics in young patients. Proceedings of the Thematic Conference of the World Psychiatric Association on 'Treatments in Psychiatry: An Update'; Florence, Italy. 2004 Nov 10-13; 2004.
- DeHaan 1997 {published data only} . De Haan L, Linsze D, Gorsira R. Early intervention, social functioning, psychotic relapse and suicide of patients with recent onset schizophrenia and other psychotic disorders [Vroegtijdige intensieve interventie, sociaal functioneren, psychoserecidief en suicide bij patienten met recent ontstane schizofrenie en verwante stoornissen]. *Tijdschrift voor Psychiatrie*. 1997; 39:24–36. De Haan L, Linsze D, Gorsira R. Early intervention, social functioning and psychotic relapse of patients with recent-onset schizophrenic disorders. *International Clinical Psychopharmacology*. 1998; 13(suppl1):S63–66.
- Drury 2000 {published data only} . Drury V, Birchwood M, Cochrane R. Cognitive therapy and recovery from acute psychosis: a controlled trial. 3. Five-year follow-up. *British Journal of Psychiatry*. 2000; 177:8–14. [PubMed: 10945081] Drury V, Birchwood M, Cochrane R, Macmillan F. Cognitive therapy and recovery from acute psychosis: a controlled trial. I. Impact on psychotic symptoms. *British Journal of Psychiatry*. 1996; 169:593–601. Drury V, Birchwood M, Cochrane R, Macmillan F. Cognitive therapy and recovery from acute psychosis: a controlled trial. II. Impact on recovery time. *British Journal of Psychiatry*. 1996; 169:602–7. [PubMed: 8932889]
- Emsley 1999 {published data only} . Emsley R, Lambert T, McGrath J, Power P, Schweitzer I, Bosma G, Mertens C, De Nayer A, De Wilde J. Risperidone in the treatment of first-episode psychotic patients: a double-blind multicenter study. *Schizophrenia Bulletin*. 1999; 25:721–9. [PubMed: 10667742]
- Emsley 2004 {published data only} . Emsley R, Davidson M, Rabinowitz J. Risk for akathisia in patients with recent onset schizophrenia treated with risperidone and haloperidol and its association with suicidality. *Schizophrenia Research*. 2004; 67(1):183. [ISI 000188788100438].
- Falloon 1992 {published data only} . Falloon I. Early intervention for first episodes of schizophrenia: a preliminary exploration. *Psychiatry*. 1992; 55:4–15. [PubMed: 1557469] Falloon I. General practice recruitment for people at risk of schizophrenia: the Buckingham experience. *Australian & New Zealand Journal of Psychiatry*. 2000; 34(Suppl):S131–6. discussion S140-4. [PubMed: 11129297]
- Fisher 2001 {published data only} . Fisher A, Savin-Bade M. The benefits to young people experiencing psychosis, and their families, of an early intervention programme: evaluating a service from the consumers' and the providers' perspectives. *British Journal of Occupational Therapy*. 2001; 64:58–65.
- Fitzgerald 1998 {published data only} . Fitzgerald P, Kulkarni J. Home-oriented management programme for people with early psychosis. *British Journal of Psychiatry Supplement*. 1998; 172:39–44. [PubMed: 9764125] Kulkarni, J. Home-based treatment of first-episode psychosis. In: McGorry, PD.; Jackson, HJ., editors. *The Recognition and Management of Early Psychosis: A Preventive Approach*. Cambridge University Press; New York, NY, US: 1999. p. 206-25.
- Fresan 2001 {published data only} . Fresan A, Apiquian R, Ulloa R, Loyzaga C, Garcia-Anaya M, Gutierrez D. Family environment and psychoeducation in first-episode schizophrenia: preliminary results [Ambiente familiar y psicoeducacion en el primer episodio de esquizofrenia: resultados preliminares]. *Salud Mental*. 2001; 24:36–40.
- Gaebel 2004 {published data only} . Gaebel W, Moller HJ, Buchkremer G, Ohmann C, Riesbeck M, Wolwer WWM, Bottlender R, Klingberg S. Pharmacological long-term treatment strategies in first episode schizophrenia--study design and preliminary results of an ongoing RCT within

the German research network on Schizophrenia. *European Archives of Psychiatry and Clinical Neuroscience*. 2004; 254(2):129–40. [EMBASE: 2004421362; MEDLINE: 15146342]. [PubMed: 15146342] Gaebel W, Riesbeck M, Von Wilmsdorff M, Zielasek J. Pharmacological long-term treatment strategies in first episode schizophrenia: preliminary results of an ongoing randomised clinical trial within the german research network on schizophrenia. *Schizophrenia Bulletin*. 2005; 31:483.

- Grawe 1998 {published data only} . Grawe R, Widen JH. Result of two years optimal out-patient treatment of first episode schizophrenia: a controlled study. *Nordic Journal of Psychiatry Supplement*. 1998; 52:76.
- Hartmann 1974 {published data only} . Hartmann W, Muller P. The therapy of first time hospitalized schizophrenics [Zur stationaren behandlung erstmals hospitalisierter schizophrener]. *Fortschritte der Neurologie Neurologie und Medizinische Psychologie*. 1974; 42:601–14.
- Heydebrand 2004 {published data only} . Heydebrand G, Weiser M, Rabinowitz J, Hoff AL, DeLisi LE, Csernansky JG. Correlates of cognitive deficits in first episode schizophrenia. *Schizophrenia Research*. 2004; 68(1):1–9. [EMBASE: 2004130672]. [PubMed: 15037334]
- Jenner 2001 {published data only} . Jenner J, Van de Willige G. Hit, hallucination focused integrative treatment as early intervention in psychotic adolescents with auditory hallucinations: a pilot study. *Acta Psychiatrica Scandinavica*. 2001; 103(2):148–52. [PubMed: 11167318]
- Jenner 2004 {published data only} . Jenner J, Van De Willige G. Results of integrated treatment (hit) in early psychoses: a pilot study. *Schizophrenia Research*. 2004; 70(1):16. [ISI: 000224551100038].
- Jolley 2003 {published data only} . Jolley S, Garety P, Craig T, Dunn G, White J, Aitken M. Cognitive therapy in early psychosis: a pilot randomized controlled trial. *Behavioural and Cognitive Psychotherapy*. 2003; 31(4):473–8.
- Jones 2005 {published data only} . Fowler D, Hodgekins J, Painter M, Reilly T, Crane C, Macmillan I, Mugford M, Croudace T, Jones PB. Cognitive behaviour therapy for improving social recovery in psychosis: a report from the ISREP MRC trial platform study (Improving social recovery in early psychosis). *Psychological Medicine*. 2009; 39:1627–36. [PubMed: 19335932] Jones PB. Improving social recovery in early affective and non-affective psychosis: a randomised controlled trial of social recovery orientated cognitive behaviour (SRCBT). *National Research Register*.
- Kadota 1992 {published data only} . Kadota K. A study of clinical response to different kinds of neuroleptics in first time medicated schizophrenics. *Japanese Journal of Psychiatry & Neurology*. 1992; 46:127–53. [PubMed: 1353124]
- Kauranen 2000 {published data only} . Kauranen A, Seikkula J, Alakar B. The change of social network of first episode psychotic patients in a network oriented treatment [Akuutin psykoosipotilaan sosiaalinen verkosto ja sen muutos verkostokeskeisessae hoidossa]. *Psykologia*. 2000; 35:403–15.
- Kavanagh 2004 {published data only} . Kavanagh DJ, Young R, White A, Saunders JB, Wallis J, Shockley NJL, Clair A. A brief motivational intervention for substance misuse in recent-onset psychosis. *Drug and Alcohol Review*. 2004; 23(2):151–5. [MEDLINE: 15370020]. [PubMed: 15370020]
- Keefe 2000 {published data only} . Keefe, RSE. Treatment of neurocognitive deficits with olanzapine or haloperidol in first episode psychosis. *Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000*.
- Keshavan 1998 {published data only} . Keshavan M, Schooler NR, Sweeney JA, Haas GL, Pettegrew JW. Research and treatment strategies in first-episode psychoses. *The Pittsburgh experience*. *British Journal of Psychiatry Supplement*. 1998; 172:60–5.
- Kopala 2003 {published data only} . Kopala L, Rabinowitz J, Davidson M. Extra-pyramidal signs and symptoms (eps) in recent onset schizophrenia: a comparison of risperidone and haloperidol. *Journal of the European College of Neuropsychopharmacology*. 2003; 13(4):S338.
- Kuipers 2004 {published data only} . Kuipers E, Holloway F, Rabe-Hesketh S, Tennakoon L. An RCT of early intervention in psychosis: Croydon outreach and assertive support team (COAST). *Social Psychiatry and Psychiatric Epidemiology*. 2004; 39(5):358–63. [PubMed: 15133591]

- Li 2004 {published data only} . Li X-H, Wan J. Effects of the nursing care of mutual participation model on the rehabilitation of inpatients with early schizophrenia. *Zhongguo Linchuang Kangfu*. 2004; 8(24):4958–9.
- Lieberman 2005b {published data only} . Lieberman JA, Tollefson GD, Charles C, Zipursky R, Sharma T, Kahn RSKRS, Green AI, Gur RE, McEvoy J, Perkins D, Hamer RM, Gu H, Tohen M, HGDH Study Group. Antipsychotic drug effects on brain morphology in first-episode psychosis. *Archives of General Psychiatry*. 2005; 62(4):361–70. [EMBASE: 2005157308; MEDLINE: 15809403]. [PubMed: 15809403]
- Malla 2001 {published data only} . Malla, A.; Norman, R.; McLean, T.; Manchanda, R.; Townsend, L.; Cortese, L.; Harricharan, R.; Takhar, J. Development of a community focussed early intervention program for psychosis: combining service, research and public education. *Proceedings of the Second International Conference on Early Psychosis*; New York, USA. 2000 31st March - 2nd April; 2000. Malla A, Norman RM, McLean TS, McIntosh E. Impact of phase-specific treatment of first episode of psychosis on wisconsin quality of life index (client version). *Acta Psychiatrica Scandinavica*. 2001; 103(5):355–61. [PubMed: 11380305]
- McCay 2007 {published data only} . McCay E, Beanlands H, Leszcz M, Goering P, Seeman MV, Ryan K, Johnston N, Vishnevsky T. A group intervention to promote healthy self-concepts and guide recovery in first episode schizophrenia: a pilot study. *Psychiatric Rehabilitation Journal*. 2006; 30(2):105–11. [PubMed: 17076053] McCay E, Beanlands H, Zipursky R, Roy P, Leszcz M, Landeen J, Ryan K, Conrad, Romano D, Francis D, Hunt J, Costantini L, Chan E. A randomised controlled trial of a group intervention to reduce engulfment and self-stigmatisation in first episode schizophrenia. *Australian E-Journal for the Advancement of Mental Health*. 2007; 6(3):1–9. [CINAHL: 2009775290; PSYCINFO: 2007–19131–008].
- McGorry 1996 {published data only} . Francey, SM. The role of day programmes in recovery in early psychosis. In: McGorry, PD.; Jackson, HJ., editors. *The Recognition and Management of Early Psychosis: A Preventive Approach*. Cambridge University Press; New York, NY, US: 1999. p. 407–37. McGorry P, Edwards J, Mihalopoulos C, Harrigan SM, Jackson HJ. Eppic: an evolving system of early detection and optimal management. *Schizophrenia Bulletin*. 1996; 22:305–26. [PubMed: 8782288] Power P, Elkins K, Adlard S, Curry C, McGorry P, Harrigan S. Analysis of the initial treatment phase in first-episode psychosis. *British Journal of Psychiatry Supplement*. 1998; 172:71–6. [PubMed: 9764130] Yung AR, McGorry PD, McFarlane CA, Jackson HJ, Patton GC, Rakkar A. Monitoring and care of young people at incipient risk of psychosis. *Schizophrenia Bulletin*. 1996; 22:283–303. [PubMed: 8782287] Yung, AR.; Phillips, LJ.; Drew, LT. Promoting access to care in early psychosis. In: McGorry, PD.; Jackson, HJ., editors. *The Recognition and Management of Early Psychosis: A Preventive Approach*. Cambridge University Press; New York, NY, US: 1999. p. 80–114.
- Mosher 1975 {published data only} . Bola JR. Evaluation of treatment in early episode acute psychosis: a secondary analysis of the soteria study. *Dissertation Abstracts International (Humanities and Social Sciences)*. 1999; Vol. 60 Hirschfeld RM, Matthews SM, Mosher LR, Menn AZ. Being with madness: personality characteristics of three treatment staffs. *Hospital Community Psychiatry*. 1977; 28(4):267–73. [PubMed: 844816] Matthews SM, Roper MT, Mosher LR, Menn AZ. A nonneuroleptic treatment for schizophrenia: analysis of the two-year postdischarge risk of relapse. *Schizophrenia Bulletin*. 1979; 5(2):322–33. [PubMed: 37598] Mosher LR. Soteria and other alternatives to acute psychiatric hospitalization: a personal and professional review. *Journal of Nervous and Mental Diseases*. 1999; 187(3):142–9. Mosher LR, Menn A. Soteria: an alternative to hospitalization for schizophrenia. *Current Psychiatric Therapy*. 1975; 15:287–96. Mosher LR, Menn A. Soteria: an alternative to hospitalization for schizophrenics. *Current Psychiatric Therapy*. 1982; 21:189–203. Mosher LR, Menn A, Matthew SM. Soteria: Evaluation of a home-based treatment for schizophrenia. *American Journal of Orthopsychiatry*. 1975; 45(3):455–67. [PubMed: 238399] Mosher LR, Menn AZ. Community residential treatment for schizophrenia: two-year follow-up. *Hospital Community Psychiatry*. 1978; 29(11):715–23. [PubMed: 700610] Mosher LR, Menn AZ. Soteria House: one-year outcome data. *Psychopharmacology Bulletin*. 1977; 13(2):46–8. [PubMed: 859995] Mosher LR, Vallone R, Menn A. The treatment of acute psychosis without neuroleptics: six-week psychopathology outcome data from The Soteria Project. *International Journal of Social Psychiatry*. 1995; 41(3):157–73. [PubMed: 8847197]

- Mottaghipour 2000 {published data only} . Mottaghipour, Y.; Woodland, L.; Sara, G. Efficacy and effectiveness in early psychosis family education group program. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000.
- Newton 2005 {published data only} . Newton E, Landau S, Smith P, Monks P, Shergill S, Wykes T. Early psychological intervention for auditory hallucinations: an exploratory study of young people's voices groups. *Journal of Nervous and Mental Disease*. 2005; 193(1):58–61. [MEDLINE: 2005088980]. [PubMed: 15674136]
- Nuechterlein 2005 {published data only} . Nuechterlein KH, Subotnik KL, Ventura J, Gitlin MJ, Green MF, Wallace CJ, Becker DR, Liberman RP, Drake RE, Mintz J. Advances in improving and predicting work outcome in recent - onset schizophrenia. *Schizophrenia Bulletin*. 2005; 31:31.
- Parlato 1999 {published data only} . Parlato L, Lloyd C, Basse TJ. Young occupations unlimited: an early intervention programme for young people with psychosis. *British Journal of Occupational Therapy*. 1999; 62:113–6.
- Perez 2003 {published data only} . Perez R, Gonzalez-Blanch C, Sierra-Biddle D, Martinez I, Vazquez-Barquero JL, Crespo-Facorro B. Efficacy and safety of olanzapine, risperidone and haloperidol in acute treatment of patients with first episode psychosis. *Schizophrenia Research*. 2003; 60:298–9. [EMBASE: 2005343078; MEDLINE: 12777272].
- Power 2004 {published data only} . Garety PA, Craig TKJ, Dunn G, Fornells-Ambrojo M, Colbert S, Rahaman N, Reed J, Power P. Specialised care for early psychosis: symptoms, social functioning and patient satisfaction: randomised controlled trial. *British Journal of Psychiatry*. 2006; 188:37–45. [PubMed: 16388068]
- Purdon 2000 {published data only} . Purdon S, Jones BD, Stip E, Labelle A, Addington D, David SR, Breier A, Tollefson GD. Neuropsychological change in early phase schizophrenia during 12 months of treatment with olanzapine, risperidone, or haloperidol. The Canadian collaborative group for research in schizophrenia. *Archives of General Psychiatry*. 2000; Vol. 57:249–58. [PubMed: 10711911]
- Rund 1994 {published data only} . Rund B, Moe L, Sollien T, Fjell A, Borchgrevink T, Hallert M, Naess PO. The psychosis project: outcome and cost-effectiveness of a psychoeducational treatment programme for schizophrenic adolescents. *Acta Psychiatrica Scandinavica*. 1994; Vol. 89:211–8. [PubMed: 7909978]
- Sanger 1999 {published data only} . Lieberman, J.; Sanger, T.; Tohen, M.; The First Episode Collaborative Study Group. Olanzapine and haloperidol treatment of first episode schizophrenia and schizoaffective disorder: 12 week outcome of a two year randomized double blind trial. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; Sanger, T. Olanzapine versus haloperidol treatment in first episode psychosis [poster]. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; Sanger, T. Treatment of neurocognitive deficits with olanzapine or haloperidol in first episode psychosis. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; Sanger T, Lieberman JA, Tohen M, Grundy S, Beasley C Jr, Tollefson GD. Olanzapine versus haloperidol treatment in first-episode psychosis. *American Journal of Psychiatry*. 1999; 156:79–87. [PubMed: 9892301]
- Schooler 2003 {published data only} . Schooler N, Davidson M, Kopala L. Reduced relapse rates in recent onset schizophrenia patients treated with risperidone vs. haloperidol. *Journal of the European College of Neuropsychopharmacology*. 2003; 13(4):S337.
- SOCRATES-UK {published data only} . Drake RJ, Haley CJ, Akhtar S, Lewis SW. Causes and consequences of duration of untreated psychosis in schizophrenia. *British Journal of Psychiatry*. 2000; 177:511–5. [PubMed: 11102325] Lewis, S.; Tarrier, N.; Haddock, G.; Bentall, R.; Kinderman, P.; Kingdon, D.; Siddle, R.; Everitt, J.; Benn, A.; Leadley, K.; Glazebrook, K.; Drake, R.; Haley, C.; Akhtar, S.; Faragher, B. A randomised, controlled trial of cognitive-behaviour therapy in acute, early schizophrenia: the SOCRATES trial summary. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000. Lewis SW, Tarrier N, Haddock G, Bentall R, Kinderman P, Kingdon D, Siddle R, Drake R, Everitt J, Leadley K, Benn A, Grazebrook K, Haley C, Akhtar S, Davies L, Palmer

- S, Faragher B, Dunn G. A randomised controlled trial of cognitive behaviour therapy in early schizophrenia: acute phase outcomes. *British Journal of Psychiatry*. 2002; 181(Suppl 43):s91–97. Tarrier N, Lewis S, Haddock G, Bentall R, Drake R, Kinderman P, Kingdon D, Siddler R, Everitt J, Leadley K, Benn A, Grazebrook K, Haley C, Akhtar S, Davies L, Palmer S, Dunn G. Cognitive-behavioural therapy in first-episode and early schizophrenia: 18-month follow-up of a randomised controlled trial. *British Journal of Psychiatry*. 2004; 184:231–9. [EMBASE: 2004121890]. [PubMed: 14990521]
- Szymanski 1994 {published data only} . Szymanski S, Masiar S, Mayerhoff D, Loebel A. Clozapine response in treatment-refractory first-episode schizophrenia. *Biological Psychiatry*. 1994; 35:278–80. [PubMed: 8186332]
- Thomas 1979 {published data only} . Thomas JL. Loxapine oral liquid concentrate in the treatment of young adult patients with acute schizophrenic symptoms. *Current Therapeutic Research*. 1979; 25:371–7.
- TIPS 2006 {published data only} . Melle I, Johannesen JO, Friis S, Haahr U, Joa I, Larsen TK, Opjordsmoen S, Rund BR, Simonsen E, Vaglum P, McGlashan T. Early detection of the first episode of schizophrenia and suicidal behavior. *American Journal of Psychiatry*. 2006; 163(5): 800–4. [MEDLINE: 16648319]. [PubMed: 16648319] Melle I, Larsen TK, Haahr U, Friis S, Johannesen JO, Opjordsmoen S, Simonsen E, Rund BR, Vaglum P, McGlashan T. Reducing the duration of untreated first-episode psychosis: effects on clinical presentation. *Archives of General Psychiatry*. 2004; 61(2):143–50. [PubMed: 14757590] Opjordsmoen, S.; Brunsvik, S.; Melle, I.; Dahl, A.; Friis, S.; Haahr, U.; Hustoft, K.; Johannesen, JO.; Larsen, TK.; McGlashan, TH.; Simonsen, E.; Vaglum, P. A comparison between novel and traditional antipsychotics as first-line medication in early psychosis. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000. Simonsen, E. A randomized comparative trial of an early detection program and treatment as usual in the duration of untreated psychosis. Stanley Foundation Research Programs; Denmark: 2000.
- Turetz 1997 {published data only} . Turetz M, Mozes T, Toren P, Chernauzan N, Yoran-Hegesh R, Mester R, Wittenberg N, Tyano SWA. An open trial of clozapine in neuroleptic-resistant childhood-onset schizophrenia. *British Journal of Psychiatry*. 1997; 170:507–10. [PubMed: 9330014]
- Ueland 2004 {published data only} . Ueland T, Rund BR. A controlled randomized treatment study: the effects of a cognitive remediation program on adolescents with early onset psychosis. *Acta Psychiatrica Scandinavica*. 2004; 109(1):70–4. [MEDLINE: 14674961]. [PubMed: 14674961]
- Walczewski 1998 {published data only} . Walczewski K, Wojciechowski A. Social networks of schizophrenic patients three years after the first psychiatric hospitalization. Differences between a group treated in psychosocial program and patients treated in an individual treatment program. *Psychiatria Polska*. 1998; 32(1):59–67. [PubMed: 9594584]
- Wang 2000 {published data only} . Wang, L.; Wang, Z. Risperidone versus clozapine treatment in first-episode psychosis. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000.
- Welch 2000 {published data only} . Welch, M. Early intervention in first episode psychosis in a rural mental health service. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; Welch M, Garland G. The safe way to early intervention: an account of the SAFE (Southern area first episode) project. *Australasian Psychiatry*. 2000; 8:243–8.
- Whitehorn 1998 {published data only} . Whitehorn, D. Establishing an early psychosis program in Nova Scotia. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; 2000. Whitehorn D, Lazier L, Kopala L. Psychosocial rehabilitation early after the onset of psychosis. *Psychiatric Services*. 1998; 49(9):1135–7. 1147. [PubMed: 9735953]
- Whitwell 2000 {published data only} . Whitwell, D. Early intervention on a shoestring. Proceedings of the Second International Conference on Early Psychosis; New York, USA. 2000 31st March - 2nd April; Whitwell D. Service innovations: early intervention in psychosis as a core task for general psychiatry. *Psychiatric Bulletin*. 2001; 25:146–8.

- Wieneke 2000 {published data only} . Wieneke A, Bechdorf A, Schultze-Lutter F. Early intervention in increased risk for schizophrenic illnesses. Multimodal psychotherapy - antipsychotic drugs only in treatment failure [Fruhintervention bei erhöhtem risiko für schizophrene erkrankungen]. *Fortschritte der Medizin*. 2000; 142(10):30–2. [PubMed: 10748600]
- Wunderink 2003 {published data only} . Wunderink L, Nienhuis FJ, Wiersma D, Van Den Bosch RJ, Bruggeman R, Faber G, Van Der Linde J, Noorthoorn E, Slooff CJ, Vlaminc P. Medication strategies in first onset schizophrenia: A randomized trial of effectiveness of short versus sustained antipsychotic treatment and quality of life in first episode psychosis preliminary findings: Incidence, compliance and early course. *Schizophrenia Research*. 2003; 60:307. [EMBASE: 2005343078; MEDLINE: 12777272].
- Wykes 2007 {published data only} . Wykes T, Newton E, Landau S, Rice C, Thompson N, Frangou S. Cognitive remediation therapy (CRT) for young early onset patients with schizophrenia: an exploratory randomized controlled trial. *Schizophrenia Research*. 2007; 94(1-3):221–30. [EMBASE: 2007327184; MEDLINE: 17524620]. [PubMed: 17524620] Wykes T, Newton E, Landau S, Rice C, Thompson N, Frangou S. Remediation in adolescent onset results of a randomised control trial. *Schizophrenia Bulletin*. 2007; 33(2):610.
- Yap 2001 {published data only} . Yap H, Mahendran R, Lim D, Liow PH, Lee A, Phang S, Tiong A. Risperidone in the treatment of first episode psychosis. *Singapore Medical Journal*. 2001; 42:170–3. [PubMed: 11465317]
- Zhang-Wong 1999 {published data only} . Zhang-Wong J, Zipursk R, Beiser M, Bean G. Optimal haloperidol dosage in first-episode psychosis. *Canadian Journal of Psychiatry*. 1999; 44:164–7.

References to studies awaiting assessment

- Addington-2001 {published data only} . Addington J, Addington D. Impact of an early psychosis program on substance use. *Journal of Psychiatric Rehabilitation*. 2001; 25(1):60–7.
- Alagband-rad 2006a {published data only} . Alagband-rad J, Shahriyar Z, Mahmoodi J, Salesian N. First episode psychoses among Iranian adolescents. *Schizophrenia Research*. 2006; 86(Suppl 1):S65. Alagband-Rad J, Sharifi V, Amini H, Shahriyar Z, Mottaghipour Y, Mahmoodi J, Seddigh A, Salesian N, Ali-Malayeri A, Tabatabaee M. Management of first episode psychoses in Iran: unique features and challenges. *Schizophrenia Research*. 2006; 86(Suppl 1):S42.
- Berger 2006 {published data only} . Berger, G. Lithium in patients at ultra high risk of developing a first psychotic episode. *Stanley Foundation Research Programs*; 2006.
- Cornblatt 2009 {published and unpublished data} . Cornblatt, B. Risperidone vs sertraline for prodromal schizophrenia. *Stanley Foundation Research Programs*; 2009. NCT00169988. Sertraline alone vs in combination with risperidone in the treatment of attenuated positive and negative symptoms. 2005 <http://www.clinicaltrials.gov>.
- Dai {published data only} . Dai M, Liu F, Fan J. A study on influence of early comprehensive interventions on prognosis of incipient schizophrenia patients. *Chinese Nursing Research*. 2007; 21:2393–4.
- Deng 2006 {published data only} . Deng Q-Y, Huang Z-L, Xie Z-Y. Research the effect of early intervention of first episode schizophrenia. *Medical Journal of Chinese Civil Administration*. 2006; 18(9):794–6.
- Doering 1998 {published data only} . Doering S, Muller E, Kopcke W, Pietzcker A, Gaebel W, Linden M, Muller P, Muller-Spahn F, Tegeler J, Schussler G. Prädiktoren für rezidiv und rehospitalisierung bei schizophrenie und schizoaffektiver psychose. *Psychiatria Danubina*. 1998; 10(4):419–30.
- Edwards 2003 EPPIC {published data only} . Edwards, J. Early psychosis service developments: becoming real. *Proceedings of the 3rd International Conference on Early Psychosis*; Copenhagen, Denmark. 2002 Sep 25-28; 2003. Edwards J, Wong L, Burnett P, Harrigan SM, McGorry PD, Wade D, Murphy B, Drew L, Albiston D. Enduring positive symptoms in first episode psychosis: A randomised controlled trial of clozapine and CBT. *Schizophrenia Research*. 2003; 60:321. [EMBASE: 2005343078; MEDLINE: 12777272]. [PubMed: 12591594]

- Fillatre-1998 {published data only} . Fillatre M, Fahs H, Rose-Reinhardt H, Degiovanni A, Marc ID. First acute psychotic disorder of the young [Premier episode psychotique de l'adolescent et de l'adulte jeune]. *Annales de Psychiatrie*. 1998; 13:256–61.
- Furimsky 2005 {published data only} . NCT00260273. Access, detection and psychological treatments. 2005 <http://www.clinicaltrials.gov>.
- Gleeson 2008 {published data only} . Gleeson J, Wade D, Castle D, Gee D, Crisp K, Pearce T, et al. The EPISODE II trial of cognitive and family therapy for relapse prevention in early psychosis: rationale and sample characteristics. *Journal of Mental Health*. 2008; 17(1):19–32. [CINAHL: 2009815127; EMBASE: 2008078812].
- Humphries 2005 {published data only} . Humphries B, Honer W. V3: Vancouver-Victoria valacyclovir trial for early psychosis. <http://www.clinicaltrials.gov>.
- Johnson 2004 {published data only} . Johnson S. Cluster randomised trial comparing outcomes of early psychosis care by a specialist team and augmented community mental health teams (CMHTs). 2005 <http://www.controlled-trials.com/>. Johnson, S. Establishing a framework for long-term research in early psychosis: use for research of routine outcome measures. National Research Register;
- Keshavan 2003 {published data only} . Keshavan, M. A 6-month controlled study of psychoeducation and collaboration enhancement (PEACE) in early psychosis to improve treatment compliance. Stanley Foundation Research Programs; 2003.
- Keshavan 2005 {published data only} . Keshavan MS, Hogarty GE. Rehabilitation, brain function and early schizophrenia. <http://www.clinicaltrials.gov>.
- Lecomte 2006 {published data only} . Lecomte T, Leclerc C, Wykes T, Wallace CJ, Spidel A, Corbiere M. Group CBT vs skills training for first episodes of psychosis - results of a RCT. *Schizophrenia Research*. 2006; 86(Suppl 1):S45–6.
- Lee 2007 {published data only} . Lee H. Protective and risk factors in adolescents with schizophrenia. Dissertation Abstracts International: Section B: The Sciences and Engineering.
- Li 2004b {published data only} . Li T-Y, Li Y-Y, Wu T-C, Yan W-Y, Chen Y-H, An Y, Lu C-Y, Zhu H-L, Jiang C-X. Effect of early intervention on quality of life in patients with first episode schizophrenia. *Zhongguo Linchuang Kangfu*. 2004; 8(18):3464–5. [CENTRAL: CN-00516341]. Li YY, Wu TC, Li M. Efficacy of early intervention on first-episode schizophrenic patients. *Journal of Clinical Psychological Medicine*. 2004; 14(1):24–6.
- Li 2007 {published data only} . Li G, Zong L, Tian S. Effect of early psychotherapy on post-schizophrenia depression. *Chinese Journal of Health Psychology*. 2007; 15(1):24–5.
- Richtand 2007 {published data only} . Richtand N. Omega-3 fatty acid deficiency replacement in early schizophrenia. <http://www.clinicaltrials.gov>.
- Schepp 1999 {published data only} . Schepp KG. Self-management therapy for youth with schizophrenia. <http://www.clinicaltrials.gov>.
- Tao 2004 {published data only} . Tao YL, Yong YL, Tian CW, Wei YY, Yu HC, Yuan A, Chang YL, Hong LZ, Cai XJ. Effect of early intervention on quality of life in patients with first episode schizophrenia. *Chinese Journal of Clinical Rehabilitation*. 2004; 8(18):3464–65.
- Vinogradov 2008 {published data only} . Vinogradov S, May L, Loewy R, Moua K, Carter C. Effectiveness of neuroadaptive cognitive training in adolescents at risk for psychosis. <http://www.clinicaltrials.gov>.
- Williams 2005 {published data only} . Williams, S. Effectiveness trial of systematic psychosocial interventions in early psychosis. National Research Register;
- Woo 2009 {published data only} . Woo W-U. Addition of tiagabine to second-generation antipsychotics in the treatment of recent-onset schizophrenia by modification of developmental reorganization of the prefrontal cortex. <http://www.clinicaltrials.gov>. Woo, TU. Tiagabine for schizophrenia. Stanley Research Programmes;
- Woods 2005 {published data only} . Woods SW. Glycine treatment of prodromal symptoms. <http://www.clinicaltrials.gov>.
- Woods 2008 {published data only} . Woods SW, Saksa JR, De Silva S, Cannon TD, Daley M, Shafer K, Cadenhead K, Haroun N, Walsh B, McGlashan T, Gelernter J, Brasfield J, Walker E, Compton M, Groves J, Seidman L, Frazier JA, Atencio J, Cornblatt BA, Ather A, Smith C,

Correll C, Cook E, Graham K, Perkins D, Anderson G, Addington D, Addington J, Raedler T. Ziprasidone in the psychosis prodrome. <http://www.clinicaltrials.gov>.

Xu 2003 {published data only} . Xu L, Zhang F, Meng Q. Early recovery psychotherapy and first episode schizophrenics. *Journal of Nursing Science*. 2003; 18(10):723–6.

Yu 2005 {published data only} . Yu EL, Xiu YL, Zhao XZ. One-year follow-up study of systemic early intervention to first episode schizophrenia. *Chinese Journal of Clinical Rehabilitation*. 2004; 8(36):8178–81.

References to ongoing studies

Addington 2007 {published data only} . Addington J, Christensen B, Remington G, Lau M, Monette G. A randomized controlled trial of individual therapy for first episode psychosis. <http://www.clinicaltrials.gov>.

Arends 2006 {published data only} . Arends J. Prodromal symptoms and early intervention to prevent a relapse. 2006 <http://www.controlled-trials.com/>.

EDIE-2 Morrison 2007 {published data only} . Morrison AP. EDIE-2: early detection and psychological intervention for individuals at high risk of psychosis (2). *Current Controlled Trials*. 2007Morrison T. Early detection and psychological intervention for individuals at high risk of psychosis. EDIE-2. Data on file 2006.

Furimski 2005 {published data only} . Furimski I, Mcmillan S, Zipursky RB, Addington J, Dewa C, Boydell K, Noh S. Access, detection and psychological treatments. <http://www.clinicaltrials.gov>.

Gaebel 2005 {published data only} . Gaebel W. Prodrome-based early intervention with antipsychotics vs benzodiazepine in patients with first-episode schizophrenia after one year maintenance treatment under further maintenance treatment vs stepwise discontinued drugs. <http://www.clinicaltrials.gov>. Gaebel W. Maintenance treatment vs stepwise drug discontinuation after one year of maintenance treatment in first-episode schizophrenia. <http://www.clinicaltrials.gov>.

Heresco-Levy 2006 {published data only} . Heresco-Levy U, Lerer B, Kohn Y, Nashim E. Sarcosine (n-methylglycine) trial for individuals at risk for developing schizophrenia and related disorders. 2006 <http://www.clinicaltrials.gov>.

Lester 2006 {published data only} . Lester H. Birmingham early detection in untreated psychosis trial (REDIRECT). *Current Controlled Trials*. 2004Lester H. Birmingham early detection in untreated psychosis trial (REDIRECT). *National Research Register*. 2004; Vol. 1Lester HE, Birchwood M, Tait L, Freemantle N. Redirect: evaluating the effectiveness of an educational intervention about first episode psychosis in primary care. *Schizophrenia Research*. 2006; 86(Suppl 1):S39–40.Tait L, Lester H, Birchwood M, Freemantle N, Wilson S. Design of the Birmingham early detection in untreated psychosis trial (REDIRECT): cluster randomised controlled trial of general practitioner education in detection of first episode psychosis [ISRCTN87898421]. *BMC Health Services Research*. 2005; 5(1):19. [MEDLINE: 15755321]. [PubMed: 15755321]

McFarlane 2007 {published data only} . NCT00531518. Early detection and intervention for the prevention of psychosis, a multisite study. 2007 <http://www.clinicaltrials.gov>.

Srihari 2006 {published data only} . Srihari VH, Woods SW, Walsh B, Saksa JR, Pollard J, Hyman L, Walsh K, Cartier S. Specialized treatment early in psychosis (STEP): a pragmatic randomized controlled trial in the US public sector. *Schizophrenia Research*. 2006; 86(Suppl 1):S165.

Stain 2006 {published data only} . Stain HJ, Startup M, Carr V, Baker A, Schall U. The depth project: a multisite RCT for youths at risk for psychosis. *Schizophrenia Research*. 2006; 86(Suppl 1):S51–2.

Additional references

Addington 1990 . Addington D, Addington J, Schissel B. A depression rating scale for schizophrenics. *Schizophrenia Research*. 1990; 3:247–25. [PubMed: 2278986]

- Altman 1996 . Altman DG, Bland JM. Detecting skewness from summary information. *BMJ*. 1996; 313:1200. [PubMed: 8916759]
- Andreasen 1983 . Andreasen, NC. Schedule for the Assessment of Negative Symptoms. University of Iowa; Iowa City, Iowa: 1983.
- APA 1994 . American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (DSM-IV). 4th Edition. APA; Washington, DC: 1994.
- Barnes 1989 . Barnes TR. A rating scale for drug-induced akathisia. *British Journal of Psychiatry*. 1989; 154:672–6. [PubMed: 2574607]
- Beck 1961 . Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Archives of General Psychiatry*. 1961; 4:53–63.
- Beck 1976 . Beck, AT. *Cognitive Therapy and the Emotional Disorders*. International Universities Press; New York: 1976.
- Begg 1996 . Begg C, Cho M, Eastwood S, Horton R, Moher D, Olkin I, Pitkin R, Rennie D, Schulz KF, Simel D, Stroup DF. Improving the quality of randomized controlled trials. The CONSORT statement. *JAMA*. 1996; 276:637–9. [PubMed: 8773637]
- Birchwood 1992 . Birchwood M, Smith J, Cochrane R. Specific and nonspecific effects of educational intervention for families living with schizophrenia. A comparison of three methods specific and non-specific effects of educational intervention for families living with schizophrenia. A comparison of three methods. *British Journal of Psychiatry*. 1992; 160:806–14. [PubMed: 1617364]
- Bland 1997 . Bland JM, Kerry SM. Statistics notes. Trials randomised in clusters. *BMJ*. 1997; 315:600.
- Boissel 1999 . Boissel JP, Cucherat M, Li W, Chatellier G, Gueyffier F, Buyse M, Boutitie F, Nony P, Haugh M, Mignot G. The problem of therapeutic efficacy indices. 3. Comparison of the indices and their use. *Therapie*. 1999; 54(4):405–11. [PubMed: 10667106]
- D’Zurilla 1971 . D’Zurilla TJ, Goldfried MR. Problem solving and behavior modification. *Journal of Abnormal Psychology*. 1971; 78:107–26. [PubMed: 4938262]
- D’Zurilla 1986 . D’Zurilla, TJ. *Problem-solving therapy: A social competence approach to clinical intervention*. Springer; New York: 1986.
- De-Wilde 2005 . De Wilde EF, Hendriks VM. The client satisfaction questionnaire: Psychometric properties in a Dutch addict population Dutch addict population psychometric properties in a Dutch addict population. *European Addiction Research*. 2005; 11:157–62. [PubMed: 16110221]
- Deeks 2000 . Deeks, J. Issues in the selection for meta-analyses of binary data. Proceedings of the 8th International Cochrane Colloquium; Cape Town. 2000 Oct 25–28th; 2000.
- Divine 1992 . Divine GW, Brown JT, Frazer LM. The unit of analysis error in studies about physicians’ patient care behavior. *Journal of General Internal Medicine*. 1992; 7:623–9. [PubMed: 1453246]
- DoH 2000 . Department of Health. *The NHS plan: a summary*. Department of Health; London: 2000.
- Donner 2002 . Donner A, Klar N. Issues in the meta-analysis of cluster randomized trials. *Statistics in Medicine*. 2002; 21:2971–80. [PubMed: 12325113]
- Duggan 2005 . Duggan L, Fenton M, Rathbone J, Dardennes R, El-Dosoky A, Indran S. Olanzapine for schizophrenia. *Cochrane Database of Systematic Reviews*. 2005; (Issue 2) [DOI: %3Chtml%3E%3Cbody%3E%3C!--StartFragment--%3E%3Cspan class=%22Apple-style-span%22 style=%22border-collapse: separate; color: rgb(0, 0, 0); font-family: %27Times New Roman%27; font-size: medium; font-style: normal; font-variant: normal; font-weight: normal; letter-spacing: normal; line-height: normal; orphans: 2; text-align: auto; text-indent: 0px; text-transform: none; white-space: normal; widows: 2; word-spacing: 0px; -webkit-border-horizontal-spacing: 0px; -webkit-border-vertical-spacing: 0px; -webkit-text-decorations-in-effect: none; -webkit-text-size-adjust: auto; -webkit-text-stroke-width: 0px; %22%3E%3Cspan class= %22Apple-style-span%22 style=%22font-family: sans-serif; font-size: 13px; -webkit-border-horizontal-spacing: 2px; -webkit-border-vertical-spacing: 2px; %22%3E10.1002/14651858.CD001359.pub2%3C/span%3E%3C/span%3E%3C!--EndFragment--%3E%3C/body%3E%3C/html%3E].

- Edwards 2000 . Edwards, J.; McGorry, PD.; Pennell, K. Models of early intervention in psychosis: an analysis of service approaches. In: Birchwood, M.; Fowler, D.; Jackson, C., editors. *Early Intervention in Psychosis*. Wiley; Chichester: 2000. p. 281-314.
- Edwards 2002 . Edwards, J.; McGorry, P. Multi-component early intervention - models of good practice. In: Edwards, J.; McGorry, PD., editors. *Implementing Early Intervention in Psychosis*. Martin Dunitz; London: 2002. p. 63-84.
- Egger 1997 . Egger M, Davey G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997; 315:629–34. [PubMed: 9310563]
- Elbourne 2002 . Elbourne DR, Altman DG, Higgins JPT, Curtin F, Worthington HV, Vailancourt JM. Meta-analyses involving cross-over trials: methodological issues. *International Journal of Epidemiology*. 2002; 31:140–49. [PubMed: 11914310]
- Falloon 1984 . Falloon, IRH.; Boyd, JL.; McGill, CW. *Family Care of Schizophrenia*. Guildford Press; New York: 1984.
- Furukawa 2006 . Furukawa TA, Barbui C, Cipriani A, Brambilla P, Watanabe N. Imputing missing standard deviations in meta-analyses can provide accurate results. *Journal of Clinical Epidemiology*. 2006; 59(7):7–10. [PubMed: 16360555]
- Garety 2000 . Garety P, Jolley S. Early intervention in psychosis. *Psychiatric Bulletin*. 2000; 24:321–3.
- Gulliford 1999 . Gulliford MC, Ukoumunne OC, Chinn S. Components of variance and intraclass correlations for the design of community-based surveys and intervention studies: data from the Health Survey for England 1994. *American Journal of Epidemiology*. 1999; 149:876–83. [PubMed: 10221325]
- Guy 1970 . Guy, W.; Bonato, RR., editors. *Manual for the ECDEU Assessment Battery 2*. National Institute of Mental Health; 1970. Revised
- Guy 1976 . Guy, W. *ECDEU Assessment Manual for Psychopharmacology (DOTES: Dosage Record and Treatment Emergent Symptom Scale)*. National Institute of Mental Health; Rockville: 1976.
- Hamilton 1959 . Hamilton M. The assessment of anxiety states by rating. *British Journal of Psychiatry*. 1959; 32:50–5.
- Hamilton 1960 . Hamilton M. A rating scale for depression. *Journal of Neurology, Neurosurgery and Psychiatry*. 1960; 23:56–62.
- Heinrichs 1984 . Heinrichs DW, Hanlon TE, Carpenter WT. The quality of life scale: an instrument for rating the schizophrenic deficit syndrome. *Schizophrenia Bulletin*. 1984; 10:389–98.
- Higgins 2003 . Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003; 327:557–60. [PubMed: 12958120]
- Higgins 2005 . Higgins, JPT.; Green, S., editors. *Cochrane Handbook for Systematic Reviews of Interventions 4.2.5* [updated May 2005]. The Cochrane Collaboration; 2005.
- Higgins 2006 . Higgins, JPT.; Green, S., editors. *Cochrane Handbook for Systematic Reviews of Interventions 4.2.6* [updated September 2006]. The Cochrane Collaboration; 2006.
- Higgins 2008 . Higgins, JPT.; Green, S., editors. *Cochrane Handbook for Systematic Reviews of Interventions*. John Wiley & Sons; Chichester: 2008.
- Jadad 1996 . Jadad AR, Moore A, Carroll D, Jenkinson C, Reynolds JM, Gavaghan DJ, McQuay HJ. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Controlled Clinical Trials*. 1996; 17:1–12. [PubMed: 8721797]
- Kay 1986 . Kay, SR.; Opler, LA.; Fiszbein, A. *Positive and negative syndrome scale (PANSS) manual*. Multi-Health Systems; North Tonawanda, NY: 1986.
- Kay 1987 . Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*. 1987; 13:261–76. [PubMed: 3616518]
- Larsen 1998 . Larsen TK, Johannssem JO, Opjordsmoen S. First-episode schizophrenia with long duration of untreated psychosis. Pathways to care. *British Journal of Psychiatry*. 1998; 172(Suppl 33):45–52.
- Leucht 2005a . Leucht S, Kane JM, Kissling W, Hamann J, Etschel E, Engel R. Clinical implications of Brief Psychiatric Rating Scale scores. *British Journal of Psychiatry*. 2005; 187:366–71. [PubMed: 16199797]

- Leucht 2005b . Leucht S, Kane JM, Kissling W, Hamann J, Etschel E, Engel RR. What does the PANSS mean? *Schizophrenia Research*. 2005; 79(2-3):231–8. [PubMed: 15982856]
- Lieberman 2005 . Lieberman JA, Stroup TS, McEvoy JP, Swartz MS, Rosenheck RA, Perkins DO, Keefe RS, Davis SM, Davis CE, Lebowitz BD, Severe J, Hsiao JK. Effectiveness of antipsychotic drugs in patients with chronic schizophrenia. *New England Journal of Medicine*. Sep 22; 2005 353(12):1209–23. [PubMed: 16172203]
- Mangalore 2007 . Mangalore R, Knapp M. Cost of Schizophrenia in England. *Journal of Mental Health Policy and Economics*. 2007; 109:23–41. [PubMed: 17417045]
- Marshall 2000 . Marshall M, Lockwood A, Adams C, Bradley C, Joy C, Fenton M. Unpublished rating scales - a major source of bias in randomised controlled trials of treatments for schizophrenia? *British Journal of Psychiatry*. 2000; 176:249–52. [PubMed: 10755072]
- Mason 1997 . Mason P, Harrison G, Glazebrook C, Medley I, Dalkin T, Croudace TJ. Characteristics of outcome in schizophrenia at 13 years. *British Journal of Psychiatry*. 1997; 167:596–603. [PubMed: 8564314]
- McGorry 1999 . McGorry, PD. 'A stitch in time' the scope for preventive strategies in early psychosis. In: McGorry, PD.; Jackson, HJ., editors. *The Recognition and Management of Early Psychosis*. Cambridge University Press; Cambridge: 1999. p. 3-23.
- Miller 1999 . Miller R, Mason S. Phase-specific psychosocial interventions for first-episode schizophrenia. *Bulletin of the Menninger Clinic*. 1999; 63:499–519. [PubMed: 10589141]
- Moher 2001 . Moher D, Altman D, Schulz KF. The CONSORT statement: Revised recommendations for improving the quality of reports parallel-group randomized trials. *Journal of the American Medical Association*. 2001; 285:1987–91. [PubMed: 11308435]
- Montgomery 1979 . Montgomery SA, Asberg M. Montgomery asberg depression rating scale. *British Journal of Psychiatry*. 1979; 134:382–89. [PubMed: 444788]
- NICE 2002 . National Institute for Clinical Excellence. *Schizophrenia: Core Interventions in the Treatment and Management of Schizophrenia in Primary and Secondary Care*. National Institute for Clinical Excellence; London: 2002.
- Norman 2001 . Norman R, Malla AK. Duration of untreated psychosis: a critical examination of the concept and its importance. *Psychological Medicine*. 2001; 31(3):381–400. [PubMed: 11305847]
- Olsen 2006 . Olsen KA, Rosenbaum B. Prospective investigations of the prodromal state of schizophrenia: assessment instruments state of schizophrenia: assessment instruments. *Acta Psychiatrica Scandinavica*. 2006; 113:273–82. [PubMed: 16638071]
- Overall 1962 . Overall JE, Gorham DR. The brief psychiatric rating scale. *Psychological Report*. 1962; 10:799–812.
- Pharoah 2006 . Pharoah FM, Rathbone J, Mari JJ. Family intervention for schizophrenia. *Cochrane Database of Systematic Reviews*. 2006; (Issue 4) [DOI: 10.1002/14651858.CD000088.pub2].
- RevMan 2008 . The Nordic Cochrane Centre, The Cochrane Collaboration. *Review Manager (RevMan)*. 5.0. The Nordic Cochrane Centre, The Cochrane Collaboration; Copenhagen: 2008.
- Rosen 2000 . Rosen A. Ethics of early prevention in schizophrenia. *Australian & New Zealand Journal of Psychiatry*. 2000; 34(Suppl):S208–12. [PubMed: 11129311]
- Schaffner 2001 . Schaffner K, McGorry PD. Preventing severe mental illnesses - new prospects and ethical challenges. *Schizophrenia Research*. 2001; 51(1):3–15. [PubMed: 11479061]
- Schultze-Lutter 2007 . Schultze-Lutter F, Ruhrmann S, Pickler H, Von Reventlow HG, Brockhaus-Dumke A, Klosterkötter J. Basic symptoms in early psychotic and depressive disorders. *British Journal of Psychiatry Supplement*. 2007; 51:s31–7. [PubMed: 18055935]
- Schulz 1995 . Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias: dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA*. 1995; 273:408–12. [PubMed: 7823387]
- Sheitman 1998 . Sheitman B, Lieberman JA. The natural history and pathophysiology of treatment resistant schizophrenia. *Journal of Psychiatric Research*. 1998; 32:143–50. [PubMed: 9793867]
- Simpson 1970 . Simpson GM, Angus JWS. A rating scale for extrapyramidal side effects. *Acta Psychiatrica Scandinavica Supplementum*. 1970; 212:11–9. [PubMed: 4917967]

- Ukoumunne 1999 . Ukoumunne OC, Gulliford MC, Chinn S, Sterne JAC, Burney PGJ. Methods for evaluating area-wide and organisation-based intervention in health and health care: a systematic review. *Health Technology Assessment*. 1999; 3(5):1–75.
- Weissman 1976 . Weissman MM, Bothwell S. Assessment of social adjustment by patient self report. *Archives of General Psychiatry*. 1976; 33:1111–15. [PubMed: 962494]
- Wiersma 1998 . Wiersma D, Nienhuis FJ, Slooff CJ, Giel R. Natural course of schizophrenic disorders: a 15-year follow up of a Dutch incidence cohort. *Schizophrenia Bulletin*. 1998; 24(1): 75–85. [PubMed: 9502547]
- Wing 1990 . Wing, Jk; Barbo, T.; Brugha, T.; Burke, J.; Cooper, J.; Giel, R.; Jablenski, A.; Regier, D.; Sartorius, N. Schedules for clinical assessment in neuropsychiatry. *Archives of General Psychiatry*. 1990; 47:589–93. [PubMed: 2190539]
- Wyatt 1991 . Wyatt R. Early intervention with neuroleptics may decrease the long-term morbidity of schizophrenia. *Schizophrenia Research*. 1991; 5(3):201–2. [PubMed: 1684719]
- Wyatt 2001 . Wyatt R, Henter I. Rationale for the study of early intervention. *Schizophrenia Research*. 2001; 51(1):69–76. [PubMed: 11479068]
- Xia 2007 . Xia, J.; Adams, CE.; Bhagat, N.; Bhagat, V.; Bhoopathi, P.; El-Sayeh, H.; Pinfeld, V.; Takriti, Y. The Leeds outcomes stakeholders survey (LOSS) study. *Proceedings of the 15th Cochrane Colloquium; Sao Paulo, Brazil*. 2007 October 23-27; 2007.
- Young 1978 . Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. *British Journal of Psychiatry*. 1978; 133:429–35. [PubMed: 728692]
- Yung 2005 . Yung AR, Yuen HP, McGorry PD, Phillips LJ, Kelly D, Dell'Olio M, Francey SM, Cosgrave EM, Killackey E, Stanford C, Godfrey K, Buckby J. Mapping the onset of psychosis: the comprehensive assessment of at-risk mental states. *Australian and New Zealand Journal of Psychiatry*. 2005; 39(11-12):964–71. [PubMed: 16343296]

References to other published versions of this review

- Marshall 2004 . Marshall M, Lockwood A. Early Intervention for psychosis. *Cochrane Database of Systematic Reviews*. 2004; (Issue 2) [DOI: 10.1002/14651858.CD004718].

* *Indicates the major publication for the study*

PLAIN LANGUAGE SUMMARY

Early Intervention for psychosis

Schizophrenia typically begins in young adulthood and may lead to disability that lasts a lifetime. The onset of psychosis is usually preceded by a period of non-psychotic symptoms, known as prodromal symptoms. The symptoms of full-blown schizophrenia include hallucinations, delusions, disordered thinking and emotional withdrawal. There is some evidence that a delay in receiving adequate treatment reduces the chances or the extent of recovery.

In broad terms, early intervention has two objectives: the first is to prevent the onset of schizophrenia in people with prodromal symptoms; the second is to provide effective treatment to people in the early stages of schizophrenia, with the goal of reducing the ultimate severity of the illness. Early intervention services are now widespread in America, Europe and Australia.

We sought to review all trials that involved early intervention for people with prodromal symptoms, or a first episode of psychosis. We identified 18 studies, most were underpowered and at present we have insufficient data to draw any definitive conclusions, although further trials are expected.




Figure 1.
PRISMA diagram

	Adequate sequence generation?	Allocation concealment?	Blinding?	Incomplete outcome data addressed?	Free of selective reporting?	Free of other bias?
Alvarez-Spain	+	?	-	-	-	?
Amminger-Austria	?	?	?	-	?	?
Berger-Australia	?	?	?	+	?	+
EDIE-UK	?	+	-	+	?	+
Edwards-Australia	+	?	-	-	?	?
EIPS-Germany	+	?	-	-	?	+
Jackson-Australia	?	+	-	+	?	+
Killackey-Australia	+	?	?	+	?	?
Leavey-UK	?	+	+	+	?	+
LEO-CAT-UK	?	?	-	-	?	?
LifeSPAN-Australia	?	?	-	+	?	+
Linszen-Amsterdam	?	?	-	-	?	?
LIPS-Germany	?	?	-	+	?	-
OPUS-Scandinavia	+	+	+	+	?	+
PACE-Australia	?	?	-	-	?	?
PRIME-USA	?	+	?	+	?	-
Uzenoff-USA	?	?	-	+	?	?
Zhang-China	?	?	-	+	?	?

Figure 2. Methodological quality summary: review authors' judgements about each methodological quality item for each included study.

Table 1

Reasons for excluding studies

Totals	Reasons	Totals	Reasons	Totals	References
68	Randomised	32	not first episode	13	Craig 2004b, Drury 2000, Grawe 1998, Jenner 2004, Jones 2005, Jolley 2003, Keshavan 1998, Kuipers 2004, Li 2004, Nuechterlein 2005, Power 2004, SOCRATESUK, Ueland 2004, Wykes 2007
			drug studies	16	Anonymous 1987, Crow 1986, Davidson 2004, Emsley 2004, Emsley 1999, Gaebel 2004, Heydebrand 2004, Keefe 2000, Kopala 2003, Lieberman 2005b, Sanger 1999, Perez 2003, Purdon 2000, Schooler 2003, Wang 2000, Wunderink 2003
Not randomised	36	descriptions of services	7	Bao 2005, Birchwood 1989, Clare 1994, Fisher 2001, Parlato 1999, Welch 2000, Whitwell 2000, Wieneke 2000	
		before and after studies	12	Addington 1999, Alanen 1994, Albiston 1998, Culberg 1998, DeHaan 1997, Fitzgerald 1998, Fresan 2001, Newton 2005, Rund 1994, Szymanski 1994, Whitehorn 1998, Yap 2001	
		quasi-experimental studies	3	COPE-Melbourne, TIPS 2006, Walczewski 1998	
		inadequate randomisation	2	McCay 2007, Mosher 1975	
		uncontrolled studies	13	Agius 2007, Falloon 1992, Hartmann 1974, Jenner 2001, Kadota 1992, Kauranen 2000, Keshavan 1998, Malla 2001, McGorry 1996, Mottaghypour 2000, Thomas 1979, Turetz 1997, Zhang-Wong 1999	

Table 2

Percentage followed up

Study	Duration of follow up (months)			
	6	12	18	24
Amminger-Austria	No data	No data	No data	No data
Berger-Australia		86%		
EDIE-UK		60%		
Edwards-Australia	No data	No data	No data	No data
EIPS-Germany	No data	No data	No data	No data
Jackson-Australia		82%		
Killackey-Australia	75%			
Leavey-UK	79% (9 months)			
LEO-CAT-UK	No data	No data	No data	No data
LifeSPAN-Australia	75%			
Linszen-Amsterdam*		unclear		
LIPS-Germany				61%
OPUS-Scandinavia		77%		67%
PACE-Australia		100%		
PRIME-USA		84%		72%
Uzenoff-USA	79%			
Zhang-China			94%	

Loss to follow-up did not appear to be substantial