

Evaluation of the efficacy of 2% curcumin gel in the treatment of experimental periodontitis

Rajesh Ramesh Hosadurga, S. N. Rao¹, Jobin Jose², Narayana Charyulu Rompicharla², Moidin Shakil³, R. Shashidhara⁴

Departments of Periodontics, ¹Pharmacology and ³Oral and Maxillofacial Pathology, Yenepoya Dental College, Yenepoya University, ²Department of Pharmaceutics, NGSM Institute of Pharmaceutical Sciences, Paneer, Deralakatte, Mangalore, ⁴Department of Oral and Maxillofacial Pathology, Coorg Institute of Dental Sciences, Maggula, Virajpet, Karnataka, India

Submitted: 24-04-2014

Revised: 04-06-2014

Published: 06-08-2014

ABSTRACT

Context: Local drug delivery (LDD) systems have been proposed for the treatment of periodontitis. Curcumin could be a suitable agent as LDD for the treatment of periodontitis.

Aim: To formulate, evaluate the anti-inflammatory activity and to assess the duration of the action and the efficacy of 2% curcumin gel in the treatment of experimental periodontitis in Wistar albino rat model. **Settings and Design:** Twenty-one Wistar albino rats were randomly assigned to three groups. Periodontitis was induced using ligature model. Group 1: Control; group 2: Plain gel, and group 3: 2% curcumin gel. **Materials and Methods:** About 2% curcumin gel was prepared. The anti-inflammatory activity and duration of action was assessed. Silk ligature 5-0 was used to induce periodontitis. Gingival index (GI) and probing pocket depth (PPD) were measured. Treatment was done. The rats were sacrificed. Morphometric analysis was performed using stereomicroscope and ImageJ software. **Statistical Analysis Used:** Analysis of variance followed by Bonferroni's test, Wilcoxon's test for inter-group comparison, Mann-Whitney test for *P* value computation was used. The observations are mean \pm standard deviation and standard error of the mean. *P* < 0.01 when compared to control was considered as statistically significant. **Results:** About 2% curcumin gel showed 42.98% inhibition of edema and peak activity was noted at 24 h. There was statistically significant change in the GI and PPD. Morphometric analysis did not show any significant difference between groups. No toxic effects were seen on oral administration of 2000 mg/kg of curcumin. **Conclusions:** About 2% curcumin gel was effective in the treatment of experimental periodontitis.

Key words: Anti-inflammatory activity, curcumin gel, experimental periodontitis, local drug delivery, morphometric analysis

INTRODUCTION

Periodontitis is one of the most common diseases affecting the teeth. It is seen in individuals who are more susceptible to the disease.^[1]

The remnants of plaque and calculus can cause failure in periodontal treatment.^[2] Chemotherapeutic agents can be used to enhance the results achieved by mechanical instrumentation or to improve outcomes at sites not responsive to conventional

therapy.^[3] Several research studies have established their efficacy in the treatment of periodontal diseases.^[4]

MATERIALS AND METHODS

Institutional Animal Ethical Committee, approval was obtained. Dosage of the gel was selected based on the results of acute oral toxicity study of *Curcuma longa* in Wistar albino rats carried out according to Organization for Economic Cooperation and Development Guidelines No. 420. The rats were housed in polypropylene cages under standard conditions with access to food and water ad-libitum.

The aim of this study was to formulate, evaluate the anti-inflammatory activity, to assess the duration of action

Access this article online

Website:

www.phcogres.com

DOI: 10.4103/0974-8490.138287

Quick Response Code:



Address for correspondence:

Dr. Rajesh Ramesh Hosadurga, Department of Periodontics, Yenepoya Dental College, Yenepoya University, Mangalore - 575 018, Karnataka, India.
E-mail: drsharaj@gmail.com

and the efficacy of 2% curcumin gel in the treatment of experimentally induced periodontitis in Wistar albino rat model.

Preparation of 2% curcumin gel by simple dispersion method

Carbopol-940 was soaked in purified water containing 0.2% w/v sodium benzoate overnight. Using tissue homogenizer hydroxypropyl methylcellulose (HPMC) solution was mixed in propylene glycol. 2 mg of curcumin (Rajesh Chemicals, Mumbai) was transferred into HPMC solution and homogenized. This drug solution was transferred to carbopol solution and homogenized. Triethanolamine was added quantity sufficient (q.s.) to neutralize the pH. Then, distilled water was added to make q.s. to 100 ml. Control gel was prepared in the same manner. The gel was stored at ambient temperature. The formulation of 2% curcumin gel was prepared by Department of Pharmaceutics, NGSM Institute of Pharmaceutical Sciences [Figure 1, Table 1]. Shavetha *et al.* showed in an *in-vitro* study that this particular combination showed increased bioavailability and a higher percentage of drug diffusion and good rheological and texture properties.^[5]

Evaluation of physicochemical parameters of 2% curcumin gel

The formulations were subjected to tests such as homogeneity, spreadability, grittiness, extrudability, pH measurement, drug content, and percentage drug release.

Spreadability

Spreadability was measured by modifying an apparatus suggested by Mutimer *et al.*^[6]

Homogeneity

Homogeneity was graded. Grades were allotted as +++ good, ++ fair, and + poor.



Figure 1: Curcumin gel

Extrudability

The formulation was filled in a clean, lacquered aluminum collapsible one ounce tube with a nasal tip of 5 mm opening. The extrudability was then determined by measuring the amount of gel extruded through the tip when a constant load of 1 kg was placed. The extruded gel was collected and weighed. The percentage of gel extruded was calculated and grades were allotted.^[7]

Determination of pH

About 5 ± 0.1 g of gel is taken in a 100 ml in a beaker and the gel was dispersed in 45 ml of water. pH meter was used to determine the pH.

Determination of drug content uniformity

A total of 5 g of the prepared formulation was subjected for analytical assay using UV spectrophotometer at λ_{\max} 430 nm.

Drug release studies

Curcumin release from the gel was studied using permeation apparatus.

Evaluation of anti-inflammatory activity of 2% curcumin gel

A total of 18 healthy Wistar albino rats of either sex were randomly allocated to test (2% curcumin gel), standard (Voveron[®] gel), and control group (plain gel) with six animals ($n = 6$) in each group. The anti-inflammatory activity was assessed by carrageenan induced paw edema method.

Duration of anti-inflammatory activity of 2% curcumin gel

A total of 36 healthy Wistar albino rats of either sex were divided into six groups ($n = 6$) based on the time duration at which 2% curcumin gel was applied prior to injection of 0.1% carrageenan. In groups 1-6, 50 mg of 2% curcumin gel was applied at 0, 2, 4, 6, 12, 24, and 48 h prior to administration of carrageenan.

About 50 mg of the 2% gel was divided into two equal parts of 25 mg. The first part of 25 mg gel was applied on the plantar surfaces of their left hind paw surface

Table 1: Formula used to prepare 2% curcumin gel

Ingredients	Quantity
Carbapol	2 g
Polymer (HPMC)	2 g
Curcumin	2 g
Propylene glycol	5 ml
Sodium benzoate	0.2 ml
Triethanolamine	q.s.
Distilled water	q.s. to make 100 ml

HPMC=Hydroxypropyl methylcellulose; q.s.=Quantity sufficient

by gentle rubbing with the index finger approximately 50 times until no gel was seen or felt on the skin. After 5 min, 25 mg gel was applied in a similar manner at 2, 4, 6, 12, 24, and 48 h prior to injection of 0.1 ml of 1% carrageenan solution.^[8,9] Three hours after the injection the 3rd h reading was noted. The percentage inflammation and the percent inhibition of edema at each dosing interval were calculated. In the control group ($n = 6$) the plain gel and standard (Voveron[®] gel) group ($n = 6$) gel was applied 3 h prior to the carrageenan injection.

Experimental periodontitis

A total of 21 Wistar albino rats (5-10 weeks old) weighing between 150 and 250 g were used in the study. Five rats weighing 150-250 g were taken in a poly propylene cage each day of the experiment for 9 days. The rats were anesthetized with ketamine anesthesia. Preperiodontal examination was done and the upper second molars were ligated using a sterile braided silk suture (5-0) [Figure 2]. Soft tissue indicators were measured.

Treatment of experimental periodontitis

Four weeks after ligature placement the rats were divided into three groups. The rats in the control group did not receive any treatment, group 2 received plain gel, group 3 received 2% curcumin gel. The gels were applied with a tuberculin syringe with a blunt tip. The application was done every alternate day for 6 days. The soft tissue indicators were measured prior to euthanizing the rats.^[10]

Soft tissue indicators of periodontitis

Gingival index

Gingival index (GI) was recorded on maxillary second molar on four surfaces: Mesial, buccal, distal, and palatal surface. GI was recorded 1 week after ligature placement and 1 week after treatment.^[11]



Figure 2: Ligature placed

Probing pocket depth

Probing pocket depth (PPD) was taken from the gingival margin to the bottom of the pocket using a modified graduated silver cone with the round-ended tip of approximately 0.4 mm in diameter. Six values were recorded and averaged (mesiobuccal, midbuccal, distobuccal, mesiopalatal, midpalatal, and distopalatal). PPD was examined 1 week after ligature placement and 1 week after treatment.^[10]

Sacrificial indicator

Morphometric analysis of alveolar bone loss

After the rats were euthanized with ketamine overdose the specimens were dissected carefully to maintain their integrity then, immersed in sodium hypochlorite for 4 h and manual scavenging of the remaining tissue was done. The specimens were stained with methylene blue dye (1 g/100 mL, Sigma-Aldrich, Saint Louis, MO, USA) for 1 min to demarcate the cemento-enamel junction (CEJ) and examined under a stereomicroscope. In order to ensure reproducibility of the alignment of the image, the buccal cusp tip of the first and second molars were placed such that they superimposed on the corresponding lingual/palatal cusp tip. Photographs were obtained with a 6.1-megapixel digital camera (Nikon D100, Ayutthaya, Thailand). Measurements were made on the maxillary second molar in a blinded fashion, 3 times using Java based image processing software, NIH, USA, and the mean values were used in statistical analysis.^[12] The distance method applied by Crawford, Taubman, and Smith on digitalized images was used to perform linear measurements from the CEJ to the alveolar bone crest, on half of each root following the axis. Six measurements were obtained for the maxillary second molar [Figures 3 and 4]. The data was subjected to statistical analysis.

RESULTS

All statistical analysis was performed using InStat-GraphPad software [GraphPad software Inc. CA, USA]. $P < 0.001$ was considered as highly significant and $P < 0.05$ was considered as significant.




Figure 3: Morphometric analysis: Buccal view

Evaluation of physicochemical parameters of the 2% curcumin gel

The gels showed optimum spreadability of 15 s, homogeneity of +++ grade, good extrudability. The pH was within the acceptable range of 6.9-7.2 even at the end of 30 days. Curcumin release was 61% [Table 2].

Evaluation of anti-inflammatory action of 2% curcumin gel

Percentage inflammation was calculated using the formula:

$$\text{Percentage inflammation} = \{V - V_i / V_i\} \times 100$$

where $V_i = 0$ h reading and $V = 3$ h reading. The average paw thickness in the drug treated group was compared within the control group.

Percent inhibition of the edema was calculated using the formula:

$$\frac{\text{Percentage swelling of drug treated group } (V_t)}{\text{Percentage swelling of control group } (V_c)} \times 100$$

The mean weight of the control group was 238.33 ± 9.83 g, the test group was 237.50 ± 22.30 g and the standard was 227.33 ± 62.19 g. Analysis of variance was used for multiple comparisons. *F* value was 0.151. *P* value was 0.861. There was no statistically significant difference between the groups.

The mean percentage of inflammation as measured using change in paw thickness in the control, standard,




Figure 4: Morphometric analysis: Palatal view

and test group was 68.38 ± 25.05 mm, 15.74 ± 4.63 mm, and 38.99 ± 22.20 mm, respectively. Percent inhibition of edema in the standard group was 76.97% and in the test group was 42.98%. The test group showed moderate anti-inflammatory activity [Tables 3-5].

Duration of the anti-inflammatory activity of 2% curcumin gel

The gel base was used as control, 1% Voveron® gel was used as standard and the test formulation was 2% curcumin gel. The test groups 1–6, the mean percentage of inflammation was 36.65 ± 11.98 mm, 23.05 ± 4.79 , 22.46 ± 6.63 , 21.78 ± 7.91 , 19.88 ± 14.69 , and 48.79 ± 15.72 mm. Percent inhibition of edema was 46.39%, 66.28%, 67.14%, 68.15%, 70.91%, and 28.65%, respectively [Table 6].

Soft tissue indicators of periodontitis

Gingival index

The mean GI at the end of 7 days was in the control, plain gel, and curcumin gel group was 2.19 ± 0.33 , 2.21 ± 0.33 , and 2.67 ± 0.49 . The mean GI at the end of 35 days in the control, plain gel, and curcumin gel group was 3.08 ± 0.34 , 3.08 ± 0.36 , and 1.23 ± 0.18 . The mean percentage change was -40.35% , -39.62% , and 53.75% . There was a statistically significant change in the GI at the end of 35 days, *P* value was 0.001 [Table 7].

Probing pocket depth

Mean PPD in control group, plain gel group, curcumin group, and Tulsi group, 1 week after placement of ligature, was 2.75 ± 0.121 , 2.73 ± 0.120 and 2.681 ± 0.098 mm, respectively. Mean PPD in control group, plain gel group 1 week after placement of ligature, in the control group was, plain gel group and curcumin group was 2.79 ± 0.137 , 2.73 ± 0.120 and 1.79 ± 0.094 , respectively. On comparison between

Table 2: Physicochemical characters of 2% curcumin gel

Physicochemical parameters	
Homogeneity	++
Grittiness	-
Extrudability	++
Spreadability (s)	15.0
pH	7.0
Drug content	98.1
Percentage release	61

Table 3: Weight of the animals selected for the study

Weight (g)	n	Mean	SD	SE	95% CI for mean		ANOVA F	P value
					Lower bound	Upper bound		
Standard	6	227.3	62.19	25.39	162.06	292.6	1.51	0.86 NS
Test	6	237.5	22.30	9.10	214.09	260.9		
Control	6	238.3	9.832	4.01	228.02	248.6		
Total	18	234.3	36.593	8.625	216.19	252.5		

SD=Standard deviation; SE=Standard error; CI=Confidence interval; ANOVA=Analysis of variance; NS=Not significant

control and plain gel and curcumin gel using Tukey's multiple comparison 1 week after ligature placement, mean difference was 0.018, 0.072, and 0.027, respectively. There was no statistical significant difference between groups [Tables 8 and 9].

Sacrificial indicators

Morphometric analysis

The mean bone loss at mesiobuccal, midbuccal, distobuccal, mesiopalatal, midpalatal, and distopalatal site treated with plain gel was 0.30 ± 0.05 mm, 0.28 ± 0.05 , 0.22 ± 0.05 , 0.37 ± 0.06 , 0.29 ± 0.03 , and 0.29 ± 0.05 mm respectively.

The mean bone loss at mesiobuccal, midbuccal, distobuccal, mesiopalatal, midpalatal, and distopalatal site treated with 2% curcumin gel was 0.25 ± 0.04 mm, 0.20 ± 0.05 , 0.14 ± 0.03 , 0.35 ± 0.17 , 0.25 ± 0.12 , and 0.25 ± 0.12 mm respectively.

On comparison there was no statistically significant bone loss between the various groups at the mesiobuccal,

midbuccal, distobuccal, mesiopalatal, midpalatal, and distopalatal site.

Bonferroni's multiple comparisons test was used to compare the effect of control, plain gel and 2% curcumin gel at mesiobuccal, midbuccal, mesiopalatal, midpalatal, and distopalatal sites. There was no statistically significant difference between the mean bone loss at mesiobuccal, midbuccal, mesiopalatal, midpalatal, and distopalatal sites treated with control, plain gel, and 2% curcumin gel [Tables 10-16].

Acute oral toxicity

All the animals survived for the period of 14 days. They appeared healthy throughout the study. All the animals appeared to gain weight during the observation period of 14 days. There were no signs of gross toxicity, adverse pharmacological events or changes in behavior. Gross necropsy findings did not show any abnormalities.

DISCUSSION

Periodontitis is a chronic inflammatory disease caused by bacterial infection of the supporting tissues around the teeth.^[13,14] The concept of locally delivering chemotherapeutic agents to the periodontal pocket as a method to treat periodontal disease has been studied for over few decades. Several herbal drugs have been an area of interest in the treatment of periodontal diseases.^[4]

Periodontal disease can be induced in the rats by tying a ligature of 2-0-5-0 braided silk around the cervix of the maxillary or mandibular molars or by injecting lipopolysaccharides into the papilla or combination of both.^[15] Souza *et al.* used 4 weeks for periodontitis induction in the maxilla. It was similar to the study period used in our study.^[16]

Table 4: Percentage inflammation in control, standard, and test (curcumin) group

	Control	Standard	Test (curcumin)
Mean percentage of inflammation±SEM	68.387±25.05	15.74±4.634	38.992±22.20

SEM=Standard error of mean

Table 5: Percent inhibition of edema

Formulation	Dose (mg/paw)	Number of rats	Percent swelling (mm)	Percent inhibition of edema
Control	50	6	38.99±42.98	-
Standard	50	6	15.74±4.63	76.975
Test	50	6	38.99±23.22	42.983

Table 6: Duration of anti-inflammatory action of 2% curcumin gel

Curcumin	2 h	4 h	6 h	12 h	24 h	48 h
Mean percentage of inflammation±SEM	36.65±11.98	23.05±4.79	22.46±6.63	21.78±7.918	19.88±14.69	48.791±15.72
Percent inhibition	46.39	66.28	67.14	68.15	70.91	28.65

SEM=Standard error of mean

Table 7: Comparison of GI before and after treatment

Group	n	Minimum	Maximum	Mean	SD	Median	Wilcoxon signed rank test value	P value
Control group								
GI at 7 days	12	2.00	3.00	2.19	0.33	2.00	3.24	0.001** HS
GI at 35 days	12	2.50	3.50	3.08	0.34	3.00		
Plain gel								
GI at 7 days	12	2.00	3.00	2.21	0.33	2.00	3.11	0.002** HS
GI at 35 days	12	2.50	3.50	3.08	0.36	3.00		
2% curcumin gel								
GI at 7 days	12	2.00	3.00	2.67	0.49	3.00	1.43	0.002** HS
GI at 35 days	12	1.00	1.50	1.23	0.18	1.20		

GI=Gingival index; SD=Standard deviation; **HS=Highly significant

Table 8: Probing pocket depth

Mean probing depth	Mean±SD	
	7 days postligature placement	At the end of the treatment
Control	2.75±0.121	2.79±0.137
Plain	2.73±0.120	2.73±0.120
Curcumin gel	2.681±0.098	1.79±0.094

SD=Standard deviation

Table 9: Comparison between control and treatment group at the end of treatment

Turkey's multiple comparisons test	Mean difference	95% CI of difference	Summary
Control versus plain gel	0.05455	-0.1094 to 0.2185	NS
Control versus curcumin gel	0.6818	0.5178 to 0.8458	HS**

CI=Confidence interval; NS=Not significant; **HS=Highly significant

Table 10: Mean bone level at sites treated with control, plain gel, and 2% curcumin gel

Site	Mean±SD		
	Control	Plain gel	Curcumin gel
Mesiobuccal	0.30±0.056	0.30±0.055	0.25±0.046
Midbuccal	0.28±0.056	0.24±0.064	0.20±0.055
Distobuccal	0.22±0.057	0.17±0.036	0.14±0.031
Mesiopalatal	0.37±0.063	0.42±0.203	0.35±0.171
Midpalatal	0.29±0.030	0.33±0.152	0.25±0.129
Distopalatal	0.29±0.053	0.31±0.153	0.25±0.126

SD=Standard deviation

Table 11: Comparison of the effect of control, plain gel, and 2% curcumin gel on bone loss (mesiobuccal site)

Bonferroni's multiple comparisons test	Mean difference	95% CI of difference	Statistical significance
Mesiobuccal			
Control versus plain gel	0.0492	-0.05154 to 0.1499	NS
Control versus curcumin gel	0.0534	-0.04734 to 0.1541	NS
Plain gel versus curcumin gel	0.0042	-0.09654 to 0.1049	NS

CI=Confidence interval; NS=Not significant

Studies have supported the positive role of anti-inflammatory agents in the treatment of periodontal disease.^[17] The anti-inflammatory properties of curcumin are mediated by the modulating the activity of signaling pathways and transcription factors, especially nuclear factor kappa-β, activating protein-1 and mitogen-activated protein kinases. Curcumin suppresses the expression of interleukin (IL)-6, IL-1β, tumor necrosis factor-α, prostaglandins, matrix metalloproteinase 2 (MMP-2) and MMP-9.^[18-23] Curcumin was also shown to improve wound healing by increasing

Table 12: Comparison of the effect of control, plain gel, and 2% curcumin gel on bone loss (midbuccal site)

Bonferroni's multiple comparisons test	Mean difference	95% CI of difference	Statistical significance
Midbuccal			
Control versus plain gel	0.0734	-0.02734 to 0.1741	NS
Control versus curcumin gel	0.0793	-0.02144 to 0.1800	NS
Plain gel versus curcumin gel	0.0059	-0.09484 to 0.1066	NS

CI=Confidence interval; NS=Not significant

Table 13: Comparison of the effect of control, plain gel and 2% curcumin gel on bone loss (distobuccal site)

Bonferroni's multiple comparisons test	Mean difference	95% CI of difference	Statistical significance
Distobuccal			
Control versus plain gel	0.0804	-0.02034 to 0.1811	NS
Control versus curcumin gel	0.0814	-0.01934 to 0.1821	NS
Plain gel versus curcumin gel	0.001	-0.09974 to 0.1017	NS

CI=Confidence interval; NS=Not significant

Table 14: Comparison of the effect of control, plain gel, and 2% curcumin gel on bone loss (mesiopalatal site)

Bonferroni's multiple comparisons test	Mean difference	95% CI of difference	Statistical significance
Mesiopalatal			
Control versus plain gel	0.0169	-0.08384 to 0.1176	NS
Control versus curcumin gel	0.0197	-0.08104 to 0.1204	NS
Plain gel versus curcumin gel	0.0028	-0.09794 to 0.1035	NS

CI=Confidence interval; NS=Not significant

Table 15: Comparison of the effect of control, plain gel, and 2% curcumin gel on bone loss (mid palatal site)

Bonferroni's multiple comparisons test	Mean difference	95% CI of difference	Statistical significance
Midpalatal			
Control versus plain gel	0.0191	-0.08164 to 0.1198	NS
Control versus curcumin gel	0.0417	-0.05904 to 0.1424	NS
Plain gel versus curcumin gel	0.0226	-0.07814 to 0.1233	NS

CI=Confidence interval; NS=Not significant

Table 16: Comparison of the effect of control, plain gel, and 2% curcumin gel on bone loss (distopalatal site)

Bonferroni's multiple comparisons test	Mean difference	95% CI of difference	Statistical significance
Distopalatal			
Control versus plain gel	0.0367	-0.06404 to 0.1374	NS
Control versus curcumin gel	0.0461	-0.05464 to 0.1468	NS
Plain gel versus curcumin gel	0.0094	-0.09134 to 0.1101	NS

CI=Confidence interval, NS=Not significant

collagen deposition, angiogenesis and the density of fibroblasts, reducing the radiation-induced delay in wound repair.¹²⁴¹ These pharmacological properties make it an effective drug to treat periodontal disease and hence, we fabricated a 2% curcumin gel by simple dispersion method.

The gel showed a standard physicochemical profile with a drug release of 61% inhibition of edema was 42.98%. Maximum inhibition was seen at 24 h. The anti-inflammatory activity lasted for 48 h. Hence, the gel was applied on alternate days to treat experimental periodontitis.

The GI scores in the control group and plain gel group increased indicating that the inflammation progressed. In the 2% curcumin gel group, a highly significant mean percentage decrease was noted. This showed that the curcumin gel had good anti-inflammatory effect.

Morphometric analysis did not show any significant difference between the groups. Whether or not the local drug delivery (LDD) system results in bone regeneration is controversial. In our study, we did not notice any significant difference in the residual periodontal bone level among various groups.

Limitations of the study

- Pressure sensitive probe and stent needs to be used
- No scaling and root planning was done
- Microbial profiling, biochemical, and immuno-histochemical parameters have to be evaluated
- It is worthwhile to do chronic toxicity study. But, the animals were monitored closely during the entire study period for any signs of toxicity. No signs of chronic toxicity were noted.

CONCLUSION

About 2% curcumin gel has shown good anti-inflammatory effect for 24-48 h despite the limitations in our study. The

2% curcumin gel showed significant reduction in gingival inflammation and pocket depth. The periodontal bone loss was not statistically significant. This is consistent with the use of LDD systems as they help in disease limitation and not significant bone regeneration. Hence, 2% curcumin can be used as useful adjunct to enhance the results of standard periodontal therapy.

ACKNOWLEDGMENTS

I would like to acknowledge the technical and intellectual support provided by Dr. B. H. Sripathi Rao, Principal and Dean, Dr. A. Shashikanth Hegde, Prof. and Head, Dr. K. S. Rajesh, Sr. Professor, Department of Periodontics, Dr. Maji Jose, Prof. and Head, Department of Oral and Maxillofacial Pathology, Yenepoya Dental College, Mangalore, Mr. E. P. Rejeesh, Department of Pharmacology, Yenepoya Medical College, Dr. Sucharita Suresh, statistician, Father Mullers College, Mangalore.

REFERENCES

1. Ryan ME. Nonsurgical approaches for the treatment of periodontal diseases. *Dent Clin North Am* 2005;49:611-36. vii.
2. Rabbani GM, Ash MM Jr, Caffesse RG. The effectiveness of subgingival scaling and root planing in calculus removal. *J Periodontol* 1981;52:119-23.
3. Greenstein G, Polson A. The role of local drug delivery in the management of periodontal diseases: A comprehensive review. *J Periodontol* 1998;69:507-20.
4. Hirasawa M, Takada K, Makimura M, Otake S. Improvement of periodontal status by green tea catechin using a local delivery system: A clinical pilot study. *J Periodontal Res* 2002;37:433-8.
5. Sharma S, Shwetha P, Upendra KJ. Development and evaluation of topical Curcumin from different combination of polymers formulation and evaluation of herbal gel. *Int J Pharm Pharm Sci* 2012;4:4452-6.
6. Mutimer MN, Riffkin C, Hill JA, Glickman ME, Cyr GN. Modern ointment base technology. II. Comparative evaluation of bases. *J Am Pharm Assoc Am Pharm Assoc (Baltim)* 1956;45:212-8.
7. Vijaybhaskar D, Kumar NS, Rao KP, Pratima S. Histopathological effect of mucoadhesive herbal gel on tobacco carcinogenicity. *Int J Pharm Biol Sci Arch* 2012;3:394-9.
8. Brinker F. *Herb Contraindications and Drug Interactions*. Sandy, OR: Eclectic Institute Publishers; 1998. p. 33.
9. Pereira SL, de Oliveira JW, Angelo KK, da Costa AM, Costa F. Clinical effect of a mouth rinse containing *Ocimum gratissimum* on plaque and gingivitis control. *J Contemp Dent Pract* 2011;12:350-5.
10. Liu R, Li N, Liu N, Zhou X, Dong ZM, Wen XJ, et al. Effects of systemic ornidazole, systemic and local compound ornidazole and pefloxacin mesylate on experimental periodontitis in rats. *Med Sci Monit* 2012;18:BR95-102.
11. Löe H. The gingival index, the plaque index and the retention index systems. *J Periodontol* 1967;8 6 Suppl: 610-6.
12. Amer M, Elverdin JC, Fernández-Solari J, Medina VA, Chiarenza AP, Vacas MI. Reduced methacholine-induced submandibular salivary secretion in rats with experimental periodontitis. *Arch Oral Biol* 2011;56:421-7.

13. Evans AS. Epidemiological concepts. In: Evans AS, Brachmen PS, editors. *Bacterial Infections of Humans Epidemiology and Control*. New York: Plenum; 1991. p. 1-58.
14. Page RC, Offenbacher S, Schroeder HE, Seymour GJ, Kornman KS. Advances in the pathogenesis of periodontitis: Summary of developments, clinical implications and future directions. *Periodontol 2000* 1997;14:216-48.
15. Struillou X, Boutigny H, Soueidan A, Layrolle P. Experimental animal models in periodontology: A review. *Open Dent J* 2010;4:37-47.
16. Souza DM, Prado Fde A, Prado Mde A, Rocha RF, Carvalho YR. Evaluation of two morphometric methods of bone loss percentages caused by periodontitis in rats in different locations. *J Appl Oral Sci* 2010;18:493-7.
17. Drisko CH. Non-surgical pocket therapy: Pharmacotherapeutics. *Ann Periodontol* 1996;1:491-566.
18. Kim GY, Kim KH, Lee SH, Yoon MS, Lee HJ, Moon DO, *et al.* Curcumin inhibits immunostimulatory function of dendritic cells: MAPKs and translocation of NF-kappa B as potential targets. *J Immunol* 2005;174:8116-24.
19. Gulcubuk A, Altunatmaz K, Sonmez K, Haktanir-Yatkin D, Uzun H, Gurel A, *et al.* Effects of curcumin on tumour necrosis factor-alpha and interleukin-6 in the late phase of experimental acute pancreatitis. *J Vet Med A Physiol Pathol Clin Med* 2006;53:49-54.
20. Mun SH, Kim HS, Kim JW, Ko NY, Kim do K, Lee BY, *et al.* Oral administration of curcumin suppresses production of matrix metalloproteinase (MMP)-1 and MMP-3 to ameliorate collagen-induced arthritis: Inhibition of the PKCdelta/JNK/c-Jun pathway. *J Pharmacol Sci* 2009;111:13-21.
21. Swarnakar S, Ganguly K, Kundu P, Banerjee A, Maity P, Sharma AV. Curcumin regulates expression and activity of matrix metalloproteinases 9 and 2 during prevention and healing of indomethacin-induced gastric ulcer. *J Biol Chem* 2005;280:9409-15.
22. Chandrasekaran CV, Sundarajan K, Edwin JR, Gururaja GM, Mundkinajeddu D, Agarwal A. Immune-stimulatory and anti-inflammatory activities of *Curcuma longa* extract and its polysaccharide fraction. *Pharmacognosy Res* 2013;5:71-9.
23. Ghodasara J, Pawar A, Deshmukh C, Kuchekar B. Inhibitory effect of rutin and curcumin on experimentally-induced calcium oxalate urolithiasis in rats. *Pharmacognosy Res* 2010;2:388-92.
24. Hanai H, Sugimoto K. Curcumin has bright prospects for the treatment of inflammatory bowel disease. *Curr Pharm Des* 2009;15:2087-94.

Cite this article as: Hosadurga RR, Rao SN, Jose J, Rompicharla NC, Shakil M, Shashidhara R. Evaluation of the efficacy of 2% curcumin gel in the treatment of experimental periodontitis. *Phcog Res* 2014;6:326-33.

Source of Support: Nil, **Conflict of Interest:** None declared.