J. Anat. (2014) 225, pp376

doi: 10.1111/joa.12217

Corrigenda

Morphometrics and inertial properties in the body segments of chimpanzees (Pan troglodytes)

Kirsten Schoonaert, Kristiaan D'Août and Peter Aerts

DOI: 10.1111/j.1469-7580.2007.00720.x

Growth of segment parameters and a morphological classification for children between 15 and 36 months

M. Van Dam, A. Hallemans and P. Aerts


DOI: 10.1111/j.1469-7580.2008.01016.x

Our group has published two articles in the Journal of Anatomy that contain a similar error (Schoonaert et al., 2007; Van Dam et al., 2009). Both these papers report on segment inertial data. One of the reported variables, the location of the Center of Mass, is erroneously stated as being a percentage relative to the proximal joint. This is wrong: the reported values are relative to the distal joint. Values relative to the proximal joint can be obtained by subtracting the reported values from 100%.

Journal of Anatomy

We sincerely apologise for this error, and can provide the full corrected data set in electronic form upon request.

For Schoonaert et al. (2007), CoM values in Table 2 should be subtracted from 100, and we provide a corrected Figure 3 for this article.



The conclusions from both articles are not affected.

References

Schoonaert K, D'Août K, Aerts P (2007) Morphometrics and inertial properties in the body segments of chimpanzees (Pan troglodytes). J Anat 210, 518–531.

Van Dam M, Hallemans A, Aerts P (2009) Growth of segment parameters and a morphological classification for children between 15 and 36 months. J Anat 214, 79–90.