



Published in final edited form as:

J Prim Care Community Health. 2014 October ; 5(4): 234–241. doi:10.1177/2150131914529307.

Psychosocial Barriers to Follow-up Adherence After an Abnormal Cervical Cytology Test Result Among Low-Income, Inner-City Women

Siu-ken Azor Hui¹, Suzanne M. Miller¹, Kuang-Yi Wen¹, Zhu Fang², Tianyu Li¹, Joanne Buzaglo³, and Enrique Hernandez⁴

¹Fox Chase Cancer Center/Temple University Health System, Philadelphia, PA, USA

²Everest Clinical Research Services, Little Falls, NJ, USA

³Research and Training Institute of Cancer Support Community, Philadelphia, PA, USA

⁴Temple University School of Medicine, Philadelphia, PA, USA

Abstract

Objectives—Low-income, inner-city women bear a disproportionate burden of cervical cancer in both incidence and mortality rates in the United States, largely because of low adherence to follow-up recommendations after an abnormal cervical cytology result in the primary care setting. The goals of the present study were to delineate the theory-based psychosocial barriers underlying these persistent low follow-up rates and their sociodemographic correlates.

Methods—Guided by a well-validated psychosocial theory of health behaviors, this cross-sectional, correlational study assessed the barriers to follow-up adherence among underserved women (N = 210) who received an abnormal cervical cytology result. Participants were recruited through an inner-city hospital colposcopy clinic, and were assessed by telephone prior to the colposcopy appointment.

Results—Participants were largely of African American race (82.2%), lower than high school completion education (58.7%), single, never married (67.3%), and without full-time employment (64.1%). Knowledge barriers were most often endorsed (68%, M = 3.22), followed by distress barriers (64%, M = 3.09), and coping barriers (36%, M = 2.36). Forty-six percent reported more than one barrier category. Less education and being unemployed were correlated with higher knowledge barriers ($P < .0001$ and $P < .01$, respectively) and more coping barriers ($P < .05$ and $P < .05$, respectively). Women who were younger than 30 years displayed greater distress barriers ($P < .05$).

Conclusion—In the primary care setting, assessing and addressing knowledge and distress barriers after feedback of an abnormal cervical cytology result may improve adherence to follow-

© The Author(s) 2014

Corresponding Author: Suzanne M. Miller, Fox Chase Cancer Center/Temple Health, 333 Cottman Avenue, Robert C. Young Pavilion, Philadelphia, PA 19111, USA., suzanne.miller-halegoua@fccc.edu.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

up recommendations. The use of structured counseling protocols and referral to navigational and other resources may facilitate this process and thereby reduce disparities in cervical cancer.

Keywords

cervical risk; psychosocial barriers; low-income women; adherence

Cervical cancer disparities persist in the United States despite the availability of the well-established screening test, the cervical cytology, and resultant prevention and early treatment of precancerous lesions. Women of racial and ethnic minorities suffer the highest incidence rates in the United States, and therefore bear a disproportionate burden of the disease.¹ Indeed, mortality from cervical cancer among African American women is almost twice that among White women.¹ Persistent infection with oncogenic strains of human papillomavirus (HPV) plays a major role in the development of cervical cancer, and the prevalence of HPV infection is highest among women with minority race/ethnicity, low education, and low income.^{2–5} Although the recent availability of the HPV vaccine has the potential to lower the rates of HPV infection,⁶ the uptake of HPV vaccination remains low.⁷

There are about 55 million cervical cytology tests performed each year in the United States, mainly in primary care settings. Approximately 3.5 million (~6.4%) are abnormal and require medical follow-up,^{8–12} which generally entails colposcopy (ie, microscopic evaluation of the cervix) and biopsy of suspicious areas.¹³ Although cervical cancer is preventable, adherence to colposcopy and follow-up recommendations is less than optimal, with the lowest adherence rates occurring among low-income, inner-city African American women,^{14–19} generally in the 30% to 40% range,^{19–24} as well as among women who are younger and less educated.^{25,26} These populations not only experience access barriers (eg, insurance, language, place of residence) but also psychosocial barriers (eg, low knowledge and high anxiety) that undermine adherence.^{18,19,27–32}

Despite the fact that adherence remains a persistent problem, only a few studies have focused on psychosocial barriers to follow-up testing and management after an abnormal cytology result, particularly in the most vulnerable populations.^{19,30,33,34} This study builds on previous work^{19,30} to provide a more comprehensive and systematic assessment of barriers in an inner-city, underserved population, guided by the Cognitive-Social Health Information Processing (C-SHIP) model,^{19,30,35–37} which highlights 5 cognitive-affective constructs that can undermine adherence (ie, cancer-risk knowledge and perceptions, cancer-risk beliefs and expectancies, cancer-risk distress, cancer-risk goals and values, and cancer-risk coping skills).^{19,30,35–37} In the current study, we addressed 2 research issues among underserved women scheduled for an initial colposcopy: (a) to delineate the profile of cognitive-affective barriers of an inner-city, predominantly African American population and (b) to describe the relations between these barriers and sociodemographic factors to identify strategies to improve follow-up adherence for use in the primary care setting.

Methods

This is a cross-sectional, correlational study that assessed barriers to follow-up adherence among low-income, minority women, who were notified of an abnormal cytological test

result and received a scheduled colposcopy appointment. This design was chosen in order to assess psychosocial barriers and their sociodemographic correlates prior to the actual follow-up appointment to capture women during the anticipatory phase of feedback impact. The data for this study were collected as part of a precolposcopy baseline assessment for a larger parent study, which is a randomized controlled trial that evaluated the efficacy of a tailored, telephone-delivered barriers counseling intervention program in the colposcopy clinic (National Institutes of Health Grant R01 CA104979; Suzanne M. Miller, Principal Investigator). For ease of communication, the parent study design and outcomes are reported in another article.³⁸ The study was approved by the institutional review boards of Fox Chase Cancer Center (FCCC) and Temple University Hospital (TUH).

Participants

Patients who were scheduled for an initial colposcopy were recruited from the Temple University School of Medicine Women's Care Center Colposcopy Clinic in North Philadelphia, Pennsylvania, which serves a predominately low-income African-American population, from May 2006 to June 2010. The final sample size was 210.

Procedure

Patients with an initial abnormal cervical cytology test result were mailed a notification letter informing them of their result and the need to follow-up with a diagnostic colposcopy, a scheduled appointment date, and colposcopy clinic contact phone numbers. The TUH research nurse reviewed the Colposcopy Clinic schedules and identified eligible patients. Approximately 2 to 4 weeks before the initial colposcopy appointment, eligible patients (N = 343, 100%) were contacted by the TUH research nurse by telephone. On contact, the patient's upcoming colposcopy appointment was confirmed and the research nurse informed the patient about the study opportunity and invited her to be transferred to a FCCC study staff to learn more. Patients who provided verbal HIPAA (Health Insurance Portability and Accountability Act) authorization and informed consent were then telephonically transferred to an FCCC study staff member.

For patients who were transferred, study staff provided further information about the study and confirmed verbal consent. Verbally consenting participants (N = 324, 94.5%) were administered the demographic and barriers assessments, and were sent a written informed consent document in the mail for them to sign and return in a prestamped envelope. Although the barriers assessment was administered to all verbally consented participants, only those who returned the written consent form were included in the final sample (N = 210, 61.2%).

Measures

Sociodemographic Assessment—Variables assessed included age, race/ethnicity, level of education, marital status, and employment status. The sociodemographic characteristics of the study sample (N = 210) are presented in Table 1.

Psychosocial Barriers Assessment—The barriers assessment instrument used in the current study was developed from our prior work,^{19,30} formative evaluation, and our guiding

theory. Participants were asked about the 5 categories of psychosocial barriers on a Likert-type scale of 1 = *not at all*, 2 = *slightly*, 3 = *somewhat*, 4 = *very much*, and 5 = *extremely*. The Likert-type scale rating format was used to capture the extent to which a given barrier was operative, in order to increase the rigor of the results and their application to intervention. All items were scored in the direction that higher ratings indicated greater barriers. Reverse-scored items are indicated with an “R” below. A rating of 3 or higher was considered endorsing the item as a barrier to adherence. The 5 psychosocial barrier categories posited by the C-SHIP model were operationalized into the following:

Knowledge/risk perceptions: Three items assessed how well the participant understood (1) the meaning of an abnormal Pap smear [R]; (2) what a colposcopic examination includes [R]; and (3) what the human papilloma virus (HPV) is and its link to cervical cancer [R].

Expectancies and beliefs: Five items assessed the participant’s extent of beliefs in (1) having an abnormal Pap smear means having cancer; (2) having a colposcopy will be helpful [R]; (3) there is nothing that can be done that would change whether or not she gets cancer; (4) being able to keep her colposcopy appointment [R]; and (5) being able to follow what the doctor tells her to do after the colposcopy [R].

Distress: Five items assessed the participant’s worry about (1) the possibility of having cancer; (2) having pain or discomfort from the colposcopy or biopsy; (3) treatments that might be needed after colposcopy; (4) sexual problems after colposcopy; and (5) transmitting the virus that causes cancer to sexual partners.

Values and goals: Five items assessed the importance to the participant of (1) keeping the colposcopy appointment and following through with the doctor’s medical management recommendations [R]; (2) keeping a good body image and feeling good about herself [R]; (3) [if age appropriate] being able to become pregnant [R]; (4) being able to have a healthy sex life [R]; and (5) not being embarrassed or feeling ashamed during the colposcopy appointment.

Coping skills: Five items assessed whether the participant had ways to manage the following challenges: (1) distress associated with the appointment [R]; (2) paying the co-pay for the colposcopy appointment and getting referral letters that might be needed for insurance [R]; (3) remembering the appointment [R]; (4) managing responsibilities such as getting childcare, eldercare, and coverage at work [R]; and (5) ability to get transportation.

Results

The final study sample characteristics are presented in Table 1. More than half of the participants had completed education of high school, trade school, or GED (General Education Development) or below ($n = 122$, 58.65%). The majority were single, never married ($n = 136$, 67.33%), and a significant subset were unemployed ($n = 76$, 36.36%). Some participants did not provide complete demographic information; therefore, the numbers of the demographic groups are smaller than the total sample size. Women who had high school completion or below were more likely to be unemployed ($P < .0001$).

Regarding barriers to adherence, the frequencies and mean ratings of the barrier categories are displayed in Table 2. Overall, 81.43% of study participants reported having at least one barrier to adherence, and almost half (45.71%) endorsed 2 or more barriers.

In examining the sociodemographic correlates to each C-SHIP barrier category, we focused on those correlating with the knowledge, distress, and coping skills barriers because they were the most frequently endorsed and had the highest mean ratings. Bivariate relations between sociodemographic characteristics and barrier category ratings are presented in Table 3. Women with less education (high school completion or below) reported significantly higher knowledge ($P < .0001$) and coping barriers ($P < .05$). Women who were unemployed also reported higher ratings of knowledge ($P < .01$) and coping ($P < .05$). Women who were younger than 30 years reported significantly higher distress barriers ($P < .05$). Please see Table 4 for a summary of major findings of this study.

Discussion

Although cervical cancer is highly preventable, inner-city, low-income minority women continue to be at higher risk for the disease, due to disparities in adherence to follow-up regimens after an abnormal cervical cytology result.^{14–19,39,40} Building on existing literature and theory, this study delineated the psychosocial barriers to adherence, and their sociodemographic correlates, in an underserved population. The study sample had a large proportion of African Americans (82.2%), who are at the highest risk of cervical cancer morbidity and mortality^{1–5,41} and have been found to have the lowest adherence rates to follow-up care after an abnormal cervical cytology.^{14–19} Educational level and employment status were representative of a low socioeconomic status sample, who suffers the greatest disparities in health care access and uptake.

We found that knowledge and distress barriers were the most frequently endorsed, suggesting that these factors require attention among this vulnerable population. Of interest, coping barriers also characterized a significant proportion of the participants. Even within this underserved group, less education, being unemployed, and age younger than 30 years were associated with greater adherence barriers. Less educated and unemployed women may not have sufficient cognitive or material resources to understand their cervical cancer risk and follow through with adaptive actions. Younger women may also be emotionally less able to manage the worries and distress associated with cervical cancer risk and related diagnostic/ treatment procedures.

The results show that cognitive–affective barriers vary by individual characteristics, and hence it is important to develop personalized interventions to assess and address barriers that are most relevant to the individual. Theory-based barriers assessment is a critical first step for identifying women who are at high risk for nonadherence, followed by delivering tailored counseling messages. This process can be facilitated by employing refined barriers assessment instruments, such as the one used in the current study, which offer advantages in terms of precision and sophistication of tailoring algorithms for specific individuals. As knowledge barriers are the most commonly endorsed, it would be worthwhile to explore whether a protocol for assessing and addressing barriers could be integrated within primary

care services. Primary care staff who communicate with patients about the feedback of test results, and/or who prepare patients for the receipt of test results, could reinforce educational messages in a supportive fashion and provide referrals to community navigators, colposcopy clinic support services, and other resources.

In addition, based on best practice principles in health communication,^{42,43} the messages delivered to patients need to be carefully crafted for low health literacy levels to maximize understanding of the meaning and personal relevance of an abnormal cervical cytology result. Patients would also seem to benefit from messages that correct unrealistic fears and worries about disease and diagnostic/treatment procedures, especially since follow-up regimens require sustained adherence over time. Finally, given the importance of coping skills for enabling the patient to manage distress, distress management skills and reinforcement for adherence behaviors should be components of a comprehensive approach to barriers reduction.

While telephone-based intervention has been shown to be efficacious in the colposcopy clinic,^{19,30} it can be time and service provider intensive. Focusing on the most common barriers at the time of feedback in a protocolized and automated fashion should refine and streamline the process and better prepare patients. Furthermore, there is growing evidence supporting the role of patient navigators in addressing barriers and promoting adherence to abnormal follow-up in underserved patient populations, such as the Screening Adherence Follow-Up (SAFe) program for Latino women³⁹ and other patient navigation programs.^{34,44–49} Embedding state-of-the-science psychosocial barriers counseling into clinical- and community-based patient navigation programs may work synergistically to address cognitive–affective and access barriers.^{50,51} This integrated approach has high potential to improve adherence to follow-up among inner-city, minority women, and hence reduce cervical cancer disparities. In future research, it will be important to replicate these findings with a larger and more culturally diverse underserved sample. Furthermore, it will also be important to link reported patterns of barriers with adherence outcomes, particularly over time.

Acknowledgments

We also thank Mary Anne Ryan and Gem Roy for their technical assistance on this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by National Institutes of Health grants, R01 CA104979, RC1 CA145063, R01 CA076644, R01 HG017666, Department of Defense grant DAMD17-01-1-0238, American Cancer Society grant TURGS-02227, and the Fox Chase Cancer Center Behavioral Research Core Facility P30-CA06927 to Dr Miller.

References

1. American Cancer Society. [Accessed March 10, 2014] Cancer Facts & Figures 2013. <http://www.cancer.org/acs/groups/content/@epidemiologysurveillance/documents/document/acspc-036845.pdf>
2. Dunne EF, Unger ER, Sternberg M, et al. Prevalence of HPV infection among females in the United States. *JAMA*. 2007; 297:813–819. [PubMed: 17327523]

3. Hariri S, Unger ER, Sternberg M, et al. Prevalence of genital human papillomavirus among females in the United States, the National Health and Nutrition Examination Survey, 2003–2006. *J Infect Dis.* 2011; 204:566–573. [PubMed: 21791659]
4. Kahn JA, Lan D, Kahn RS. Sociodemographic factors associated with high-risk human papillomavirus infection. *Obstet Gynecol.* 2007; 110:87–95. [PubMed: 17601901]
5. Dunne EF, Sternberg M, Markowitz LE, et al. Human papillomavirus (HPV) 6, 11, 16, and 18 prevalence among females in the United States—National Health and Nutrition Examination Survey, 2003–2006: opportunity to measure HPV vaccine impact? *J Infect Dis.* 2011; 204:562–565. [PubMed: 21791658]
6. Markowitz LE, Hariri S, Lin C, et al. Reduction in human papillomavirus (HPV) prevalence among young women following HPV vaccine introduction in the United States, National Health and Nutrition Examination Surveys, 2003–2010. *J Infect Dis.* 2013; 208:385–393. [PubMed: 23785124]
7. Laz TH, Rahman M, Berenson AB. An update on human papillomavirus vaccine uptake among 11–17-year-old girls in the United States: National Health Interview Survey, 2010. *Vaccine.* 2012; 30:3534–3540. [PubMed: 22480927]
8. Forhan SE, Gottlieb SL, Sternberg MR, et al. Prevalence of sexually transmitted infections among female adolescents aged 14 to 19 in the United States. *Pediatrics.* 2009; 124:1505–1512. [PubMed: 19933728]
9. Miller, SM.; Roussi, P. Cervical cancer screening. In: Holland, JC.; Breitbart, WS.; Jacobsen, PB.; Lederberg, MS.; Loscalzo, MJ.; McCorkle, R., editors. *Psycho-oncology.* 2nd ed.. New York, NY: Oxford University Press; 2010. p. 84–89.
10. National Cancer Institute. [Accessed April 27, 2012] Fact Sheet 5.16, Pap Test. 2010. http://www.cancer.gov/cancertopics/factsheet/detection/Fs5_16.pdf
11. Tiltman AJ. The pathology of cervical tumours. *Best Pract Res Clin Obstet Gynaecol.* 2005; 19:485–500. [PubMed: 16150389]
12. Wright TC Jr, Massad LS, Dunton CJ, Spitzer M, Wilkinson EJ, Solomon D. 2006 Consensus guidelines for the management of women with cervical intraepithelial neoplasia or adenocarcinoma in situ. *J Low Genit Tract Dis.* 2007; 11:223–239. [PubMed: 17917567]
13. Massad LS, Einstein MH, Huh WK, et al. 2012 Updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors. *Obstet Gynecol.* 2013; 121:829–846. [PubMed: 23635684]
14. Benard VB, Lawson HW, Ehemann CR, Anderson C, Helsel W. Adherence to guidelines for follow-up of low-grade cytologic abnormalities among medically underserved women. *Obstet Gynecol.* 2005; 105:1323–1328. [PubMed: 15932824]
15. Doyle JP, Parker RM, Jacobson TA, McNagny SE. Breast and cervical cancer screening in an inner-city medical walk-in clinic: taking advantage of an often missed opportunity. *Am J Prev Med.* 1996; 12:345–350. [PubMed: 8909645]
16. Engelstad LP, Stewart SL, Nguyen BH, et al. Abnormal Pap smear follow-up in a high-risk population. *Cancer Epidemiol Biomarkers Prev.* 2001; 10:1015–1020. [PubMed: 11588126]
17. Jones BA, Novis DA. Follow-up of abnormal gynecologic cytology: a College of American Pathologists Q-Probes study of 16 132 cases from 306 laboratories. *Arch Pathol Lab Med.* 2000; 124:665–671. [PubMed: 10782145]
18. Marcus AC, Crane LA, Kaplan CP, et al. Improving adherence to screening follow-up among women with abnormal Pap smears: results from a large clinic-based trial of three intervention strategies. *Med Care.* 1992; 30:216–230. [PubMed: 1538610]
19. Miller SM, Siejak KK, Schroeder CM, Lerman C, Hernandez E, Helm CW. Enhancing adherence following abnormal Pap smears among low-income minority women: a preventive telephone counseling strategy. *J Natl Cancer Inst.* 1997; 89:703–708. [PubMed: 9168185]
20. Engelstad LP, Stewart S, Otero-Sabogal R, Leung MS, Davis PI, Pasick RJ. The effectiveness of a community outreach intervention to improve follow-up among underserved women at highest risk for cervical cancer. *Prev Med.* 2005; 41:741–748. [PubMed: 16125761]
21. Fox P, Arnsberger P, Zhang X. An examination of differential follow-up rates in cervical cancer screening. *J Community Health.* 1997; 22:199–209. [PubMed: 9178119]

22. Hartz LE, Fenaughty AM. Management choice and adherence to follow-up after colposcopy in women with cervical intraepithelial neoplasia 1. *Obstet Gynecol.* 2001; 98:674–679. [PubMed: 11576587]
23. Kaplan CP, Bastani R, Belin TR, Marcus A, Nasser K, Hu MY. Improving follow-up after an abnormal pap smear: results from a quasi-experimental intervention study. *J Womens Health Gend Based Med.* 2000; 9:779–790. [PubMed: 11025870]
24. McKee MD, Schechter C, Burton W, Mulvihill M. Predictors of follow-up of atypical and ASCUS papanicolaou tests in a high-risk population. *J Fam Pract.* 2001; 50:609. [PubMed: 11485710]
25. Abercrombie PD. Improving adherence to abnormal Pap smear follow-up. *J Obstet Gynecol Neonatal Nurs.* 2001; 30:80–88.
26. Jean-Pierre P, Hendren S, Fiscella K, et al. Understanding the processes of patient navigation to reduce disparities in cancer care: perspectives of trained navigators from the field. *J Cancer Educ.* 2011; 26:111–120. [PubMed: 20407860]
27. Crane LA. Social support and adherence behavior among women with abnormal Pap smears. *J Cancer Educ.* 1996; 11:164–173. [PubMed: 8877577]
28. Eger RR, Peipert JF. Risk factors for noncompliance in a colposcopy clinic. *J Reprod Med.* 1996; 41:671–674. [PubMed: 8887192]
29. Eggleston KS, Coker AL, Das IP, Cordray ST, Luchok KJ. Understanding barriers for adherence to follow-up care for abnormal pap tests. *J Womens Health (Larchmt).* 2007; 16:311–330. [PubMed: 17439377]
30. Lerman C, Hanjani P, Caputo C, et al. Telephone counseling improves adherence to colposcopy among lower-income minority women. *J Clin Oncol.* 1992; 10:330–333. [PubMed: 1732434]
31. Marcus AC, Kaplan CP, Crane LA, et al. Reducing loss-to-follow-up among women with abnormal Pap smears. Results from a randomized trial testing an intensive follow-up protocol and economic incentives. *Med Care.* 1998; 36:397–410. [PubMed: 9520963]
32. Nelson K, Geiger AM, Mangione CM. Effect of health beliefs on delays in care for abnormal cervical cytology in a multi-ethnic population. *J Gen Intern Med.* 2002; 17:709–716. [PubMed: 12220368]
33. McKee MD, Lurio J, Marantz P, Burton W, Mulvihill M. Barriers to follow-up of abnormal Papanicolaou smears in an urban community health center. *Arch Fam Med.* 1999; 8:129–134. [PubMed: 10101983]
34. Tejeda S, Darnell JS, Cho YI, Stolley MR, Markossian TW, Calhoun EA. Patient barriers to follow-up care for breast and cervical cancer abnormalities. *J Womens Health (Larchmt).* 2013; 22:507–517. [PubMed: 23672296]
35. Miller SM, Shoda Y, Hurley K. Applying cognitive-social theory to health-protective behavior: breast self-examination in cancer screening. *Psychol Bull.* 1996; 119:70–94. [PubMed: 8559860]
36. Miller SM, Mischel W, O'Leary A, Mills M. From human papillomavirus (HPV) to cervical cancer: psychological processes in infection, detection, and control. *Ann Behav Med.* 1996; 18:219–228. [PubMed: 18425667]
37. Miller, SM.; Rodoletz, M.; Buzaglo, J.; Gray, T.; Sherman, K. Monitoring attentional style among low-income minority women at risk for cervical cancer: cognitive social determinants of adjustment and adherence; Poster presented at: 12th Annual Meeting of the Society of Behavioral Medicine; 2001.
38. Miller SM, Hui SA, Wen KY, et al. Tailored telephone counseling to improve adherence to follow-up regimens after an abnormal Pap smear among minority, underserved women. *Patient Educ Couns.* 2013; 93:488–495. [PubMed: 24007767]
39. Ell K, Vourlekis B, Muderspach L, et al. Abnormal cervical screen follow-up among low-income Latinas: Project SAFE. *J Womens Health Gend Based Med.* 2002; 11:639–651. [PubMed: 12396896]
40. Carey P, Gjerdingen DK. Follow-up of abnormal Papanicolaou smears among women of different races. *J Fam Pract.* 1993; 37:583–587. [PubMed: 8245810]
41. National Cancer Institute. [Accessed June 14, 2013] Surveillance Epidemiology and End Results (SEER) Stat Fact Sheets: Cervix Uteri. 2013. <http://seer.cancer.gov/statfacts/html/cervix.html>

42. Institute of Medicine. [Accessed June 12, 2013] Speaking of health: assessing health communication strategies for diverse populations. 2002. http://www.nap.edu/catalog.php?record_id=10018
43. National Cancer Institute. [Accessed March 10, 2014] Making Health Communication Programs Work. 2002. <http://www.cancer.gov/cancertopics/cancerlibrary/pinkbook>
44. Battaglia TA, Bak SM, Heeren T, et al. Boston Patient Navigation Research Program: the impact of navigation on time to diagnostic resolution after abnormal cancer screening. *Cancer Epidemiol Biomarkers Prev.* 2012; 21:1645–1654. [PubMed: 23045539]
45. Freund KM, Battaglia TA, Calhoun E, et al. National Cancer Institute Patient Navigation Research Program: methods, protocol, and measures. *Cancer.* 2008; 113:3391–3399. [PubMed: 18951521]
46. Markossian TW, Darnell JS, Calhoun EA. Follow-up and timeliness after an abnormal cancer screening among underserved, urban women in a patient navigation program. *Cancer Epidemiol Biomarkers Prev.* 2012; 21:1691–1700. [PubMed: 23045544]
47. Paskett ED, Harrop JP, Wells KJ. Patient navigation: an update on the state of the science. *CA Cancer J Clin.* 2011; 61:237–249. [PubMed: 21659419]
48. Paskett ED, Katz ML, Post DM, et al. The Ohio Patient Navigation Research Program: does the American Cancer Society patient navigation model improve time to resolution in patients with abnormal screening tests? *Cancer Epidemiol Biomarkers Prev.* 2012; 21:1620–1628. [PubMed: 23045536]
49. Wells KJ, Battaglia TA, Dudley DJ, et al. Patient navigation: state of the art or is it science? *Cancer.* 2008; 113:1999–2010. [PubMed: 18780320]
50. Yabroff KR, Kerner JF, Mandelblatt JS. Effectiveness of interventions to improve follow-up after abnormal cervical cancer screening. *Prev Med.* 2000; 31:429–439. [PubMed: 11006069]
51. Radecki Breitkopf C, Dawson L, Grady JJ, Breitkopf DM, Nelson-Becker C, Snyder RR. Intervention to improve follow-up for abnormal Papanicolaou tests: a randomized clinical trial [published online June 3, 2013]. *Health Psychol.*

Biographies

Siu-kuen Azor Hui, PhD, is a Research Assistant Professor at Fox Chase Cancer Center, Cancer Prevention and Control Program. She is a health psychologist studying community-based interventions to promote cancer preventive behaviors, particularly in underserved populations, and their linkage to primary care services.

Suzanne M. Miller, PhD, is Professor and Director of the Psychosocial and Behavioral Medicine Department in the Cancer Prevention and Control Program at Fox Chase Cancer Center/Temple Health. She is also an Adjunct Professor at the University of Pennsylvania and Temple University. Her funded research is focused at the interface of psychology and oncology, particularly with a view to applying personalized biobehavioral and health communication principles to bridge the gap between the availability of groundbreaking cancer prevention and control technologies and the cancer populations they are meant to serve. Most recently, Dr. Miller's work has focused on the use of traditional and new technologies (web-based, texting, tweeting) to improve decision making, adherence and quality of life across the cancer continuum including cancer risk, diagnosis, management, and survivorship.

Kuang-Yi Wen, PhD, is an Assistant Professor at Fox Chase Cancer Center, Cancer Prevention and Control Program. She is a health systems engineer with interests and experience in developing, evaluating, and implementing consumer health informatics interventions.

Fang Zhu, PhD, is a senior biostatistician. She received her PhD in Biostatistics from University of Pittsburgh in 2008. She spent 5 years working as Biostatistician at Fox Chase Cancer Center and have collaborated in the publication of numerous papers.

Tianyu Li, MS, is a senior research statistician at the biostatistics and bioinformatics facility at Fox Chase Cancer Center. Her research interest is predicting the survival outcome and treatment preference of various cancer patients using advanced statistical methodologies

Joanne Buzaglo, PhD, is Vice President of Research & Training at the Cancer Support Community, a non-profit organization that provides evidence-based support and education to cancer patients and their families across an international affiliate network and online community. She is a licensed clinical psychologist and behavioral researcher with a focus on the development, evaluation and dissemination of tailored interventions designed to address the psychosocial needs of cancer patients and their families.

Enrique Hernandez, MD, is the Abraham Roth Professor and Chairman of the department of Obstetrics and Gynecology at Temple University School of Medicine and a member of Temple's Fox Chase Cancer Center. He is also director of gynecologic oncology and professor of pathology. His research interest includes the prevention, early detection, and treatment of cervical cancer.

Table 1

Baseline Demographic Characteristics of the Total Sample (N = 210).

Age in years, <i>M</i> (<i>SD</i>)	30.00 (10.67)
Race/ethnicity; n (%)	
White	9 (4.46)
Black	166 (82.18)
Hispanic	25 (12.38)
Other	2 (0.99)
Education (highest level completed); n (%)	
Below high school	11 (5.29)
Some high school	27 (12.98)
High school, trade school, or GED	84 (40.38)
Vocational school	22 (10.58)
Some college	55 (26.44)
Undergraduate degree	8 (3.85)
Graduate courses	1 (0.48)
Marital status; n (%)	
Widowed	7 (3.47)
Single, never married	136 (67.33)
Single, living with significant others	37 (18.32)
Married	13 (6.44)
Divorced	9 (4.46)
Employment status; n (%)	
Employed full time	50 (23.92)
Employed part time	37 (17.70)
Disabled	21 (10.05)
Retiree	1 (0.48)
Unemployed	76 (36.36)
Student	24 (11.48)

Table 2

Descriptive Statistics of Barriers Ratings by Cognitive-Social Health Information Processing (C-SHIP) Constructs (N = 210).

	Mean (SD)	Proportion of Women Endorsing Any Barrier Item in the Category (%)
Knowledge/risk perceptions	3.22 (1.17)	67.72
Beliefs and expectancies	1.93 (0.52)	25.10
Distress	3.09 (1.04)	64.01
Values and goals	2.17 (0.65)	28.49
Coping skills	2.36 (0.70)	36.00

Table 3
 Bivariate Relations Between Sociodemographic Characteristics and Cognitive-Social Health Information Processing (C-SHIP)-Based Barriers Ratings (N = 210).

Barrier category	Age			Race/Ethnicity				Education			Employment Status				
	<30 Years, Mean (SD)	30 Years, Mean (SD)	P	White, Mean (SD)	African American, Mean (SD)	Hispanic, Mean (SD)	Other, Mean (SD)	P	High School, Mean (SD)	>High School, Mean (SD)	P	Student, Mean (SD)	Unemployed, Mean (SD)	Employed, Mean (SD)	P
Knowledge/risk perception	3.15 (1.15)	3.29 (1.17)	.39	2.41 (1.16)	3.27 (1.14)	2.83 (1.21)	4.00 (0.47)	.04	3.46 (1.19)	2.84 (1.01)	<.0001	3.01 (1.16)	3.49 (1.15)	2.96 (1.10)	.005
Beliefs and expectancies	1.91 (0.50)	1.91 (0.50)	.97	1.64 (0.54)	1.91 (0.50)	2.08 (0.49)	1.90 (0.14)	.17	1.94 (0.49)	1.89 (0.51)	.46	1.91 (0.40)	1.91 (0.50)	1.93 (0.52)	.98
Distress	3.27 (1.06)	2.92 (0.88)	.02	2.76 (0.81)	3.09 (1.02)	3.50 (0.88)	4.60 (0.57)	.03	3.19 (1.01)	3.07 (1.02)	.38	3.19 (0.96)	3.20 (0.97)	3.07 (1.08)	.65
Values and goals	2.04 (0.54)	2.30 (0.57)	<.001	2.29 (0.78)	2.12 (0.54)	2.13 (0.66)	2.00 (0.57)	.84	2.22 (0.56)	1.99 (0.55)	<.005	1.78 (0.47)	2.22 (0.56)	2.11 (0.55)	.002
Coping skills	2.32 (0.64)	2.31 (0.65)	.99	1.98 (0.60)	2.32 (0.65)	2.35 (0.65)	2.80 (0.57)	.31	2.41 (0.66)	2.19 (0.61)	.01	2.3 (0.60)	2.45 (0.68)	2.18 (0.58)	.02

Table 4

Summary of Major Findings (N = 210).

Number of psychosocial barrier categories endorsed by patients	<ul style="list-style-type: none"> • At least one barrier category endorsed (81.43%) • Two or more barrier categories endorsed (45.71%)
Psychosocial barrier categories most frequently endorsed by patients	<ol style="list-style-type: none"> 1 Knowledge/risk perceptions (67.72%, mean rating = 3.22) 2 Distress (64.01%, mean rating = 3.09) 3 Coping skills (36.00%, mean rating = 2.36)
Significant sociodemographic correlations with the most frequently endorsed psychosocial barriers	<ol style="list-style-type: none"> 1 Higher knowledge/risk perceptions barriers are associated with lower education level (ie, high school completion or below) and unemployment 2 Higher distress barriers are associated with younger age (ie, younger than 30 years) 3 Higher coping skills barriers are associated with lower education level (ie, high school completion or below) and unemployment