

Early Evolution of the Eukaryotic Ca²⁺ Signaling Machinery: Conservation of the CatSper Channel Complex

Xinjiang Cai,^{*1} Xiangbing Wang,¹ and David E. Clapham^{*,2,3}

¹Division of Endocrinology, Metabolism and Nutrition, Department of Medicine, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ

²Howard Hughes Medical Institute, Department of Cardiology, Boston Children's Hospital, Boston, MA

³Department of Neurobiology, Harvard Medical School, Boston, MA

*Corresponding author: E-mail: caixinj@gmail.com; dclapham@enders.tch.harvard.edu.

Associate editor: Joel Dudley

Abstract

Calcium signaling is one of the most extensively employed signal transduction mechanisms in life. As life evolved into increasingly complex organisms, Ca²⁺ acquired more extensive and varied functions. Here, we compare genes encoding proteins that govern Ca²⁺ entry and exit across cells or organelles within organisms of early eukaryotic evolution into fungi, plants, and animals. Recent phylogenomics analyses reveal a complex Ca²⁺ signaling machinery in the apusozoan protist *Thecamonas trahens*, a putative unicellular progenitor of Opisthokonta. We compare *T. trahens* Ca²⁺ signaling to that in a marine bikont protist, *Aurantiochytrium limacinum*, and demonstrate the conservation of key Ca²⁺ signaling molecules in the basally diverging alga *Cyanophora paradoxa*. Particularly, our findings reveal the conservation of the CatSper channel complex in *Au. limacinum* and *C. paradoxa*, suggesting that the CatSper complex likely originated from an ancestral Ca²⁺ signaling machinery at the root of early eukaryotic evolution prior to the unikont/bikont split.

Key words: animals, calcium channels, calcium signaling, CatSper, eukaryotes, evolution.

Calcium ions (Ca²⁺) serve as a universal signal to modulate almost every aspect of cellular function in bacteria (Dominguez 2004), plants (Kudla et al. 2010), fungi (Zelter et al. 2004), and animals (Berridge et al. 2003; Clapham 2007). The core principles of Ca²⁺ signaling emerged very early in the life processes of bacteria (Case et al. 2007; Verkhratsky and Parpura 2014). The appearance of eukaryotes with intracellular organelles and the independent evolution of multicellular organisms from diverse ancestral unicellular lineages (Rokas 2008) prompted the refinement of versatile and complex Ca²⁺ signaling systems, which provide precise spatial and temporal control of Ca²⁺ concentration.

Although the basic principles of Ca²⁺ signaling appear to be universal in eukaryotes (Shemarova and Nesterov 2005), the components of the Ca²⁺ signaling machinery exhibit significant differences among distinct groups of organisms such as animals (Clapham 2007), plants (Nagata et al. 2004; Verret et al. 2010), and fungi (Zelter et al. 2004). In large skeletal and cardiac muscle cells, a highly distributed intracellular Ca²⁺ store, released by ryanodine receptor channels, was coupled directly or indirectly to plasma membrane voltage-gated Ca²⁺ (Ca_v) channels (Amador et al. 2013). Another specialized distribution system along the sperm flagella evolved via the sperm-specific CatSper Ca²⁺ channels that are critical for sperm cell hyperactivation (Lishko et al. 2012). It has been argued that in contrast to the complex Ca²⁺ signaling machinery in animals, plants and fungi have adopted more simplified Ca²⁺ signaling cascades to suit their physiological requirements (Nagata et al. 2004; Zelter et al. 2004). The origin and evolution of these distinct Ca²⁺ signaling machineries has

remained a long-standing question—they might have evolved independently of one another, or they might share the same evolutionary origin in the last common unicellular ancestor with subsequent lineage-specific evolution after divergence.

Recent phylogenomics studies of close unicellular relatives of metazoans including two choanoflagellates *Monosiga brevicollis* and *Salpingoeca rosetta* and the filasterean *Capsaspora owczarzaki* have revealed the presence of ancestral signaling molecules previously thought to be restricted to animals (King et al. 2008; Sebe-Pedros et al. 2011; Fairclough et al. 2013; Suga et al. 2013). *Monosiga brevicollis* also has an extensive Ca²⁺ signaling system resembling that in animals (Cai 2008). Animals, fungi, and their unicellular relatives form the eukaryotic supergroup Opisthokonta (Stechmann and Cavalier-Smith 2002; Steenkamp et al. 2006; Ruiz-Trillo et al. 2008). The apusozoan protist, *Thecamonas trahens*, is a putative unicellular progenitor of Opisthokonta (Cavalier-Smith and Chao 2010). We recently identified a complex ancestral Ca²⁺ signaling network in *T. trahens* that thus predates the divergence of animals and fungi (Cai and Clapham 2012).

The eukaryotic root has been hypothesized to be placed between Unikonta, the eukaryotic supergroup composed of Opisthokonta and Amoebozoa, and Bikonta, the eukaryotic supergroup including Archaeplastida (plants and relatives) and Chromalveolata (fig. 1), and other groups (Stechmann and Cavalier-Smith 2002; Burki and Pawlowski 2006; Derelle and Lang 2012). Comparative genomics studies of several plant, algal, and other bikont species showed the conservation




Fig. 1. Distribution of Ca²⁺ signaling molecules in eukaryotes. (A) Comprehensive Ca²⁺ signaling network in the apusozoan protist *Thecamonas trahens* (Cai and Clapham 2012). Shown are representative classes of Ca²⁺ signaling molecules previously known to be animal specific, such as CatSper channels and K⁺-dependent Na⁺/Ca²⁺ exchangers, or fungal/plant specific, such as CAXs, or ubiquitous, such as mitochondrial Letm1 Ca/H exchangers (Tsai et al. 2014), from three recently analyzed genomes—*T. trahens*, the choanoflagellate *Monosiga brevicollis* (Cai 2008), and the basal fungus *Allomyces macrogynus* (Cai and Clapham 2012), as well as genomes of *Homo sapiens*, *Neurospora crassa*, *Arabidopsis thaliana*, and *Cyanidioschyzon merolae* (Wheeler and Brownlee 2008; Verret et al. 2010). (B) Schematic diagram illustrating the evolutionary history of the marine thraustochytrid protist *Aurantiochytrium limacinum* and the basally diverging alga *Cyanophora paradoxa* (Price et al. 2012) in Bikonta. Inferred from the Tree of Life project (<http://www.tolweb.org/>, last accessed July 11, 2014) and recent references on the eukaryotic tree (Burki and Pawlowski 2006; Derelle and Lang 2012). Letm1, leucine zipper-EF-hand containing transmembrane protein 1.

and divergence of many Ca²⁺ signaling molecules such as Ca_v channels, transient receptor potential (TRP) channels, and ligand-gated channels among animals, fungi, plants, and many unicellular organisms (Wheeler and Brownlee 2008; Verret et al. 2010; Chan et al. 2011). Thus, it has been demonstrated that many components of the Ca²⁺ signaling machineries in animals, plants, and fungi likely emerged from early eukaryotic lineages before their divergence.

Several key types of Ca²⁺ signaling molecules conserved in *T. trahens* have not been extensively analyzed. For instance, CatSper channels and Na⁺/Ca²⁺ exchangers are present in *T. trahens*, the basal fungus *Allomyces macrogynus* and most animals but are absent in land plants such as *Arabidopsis thaliana*, *Cyanidioschyzon merolae* (a unicellular red alga), and most fungi (fig. 1A). To elucidate whether the ancestral Ca²⁺ signaling network shown in *T. trahens* is also preserved in bikonts, we examined related genomic databases of organisms in Bikonta to search for homologs of Ca²⁺ signaling molecules.

An Extensive Ca²⁺ Signaling Machinery in *Aurantiochytrium limacinum*

Aurantiochytrium limacinum is a common marine thraustochytrid protist within the class labyrinthulomycetes, which is one of the earliest diverging lineages in the phylum of stramenopiles (Riisberg et al. 2009) (fig. 1B). We found that the genome of *Au. limacinum* encodes Ca²⁺ signaling machinery as comprehensive as that in *T. trahens* (fig. 2 and




Fig. 2. Ca²⁺ signaling machinery in *Aurantiochytrium limacinum*. The evolutionary relationship of *A. limacinum* is inferred from the Tree of Life project (<http://www.tolweb.org/>) and the JGI genome portal (<http://genome.jgi-psf.org/Aurl11/Aurl11.home.html>, last accessed July 11, 2014). *Aurantiochytrium limacinum* life stages include colonies of vegetative cells or zoospores with heterokont flagella (http://syst.bio.konan-u.ac.jp/labybase/Aurantiochytrium_limacinum_life_cycle.html, last accessed July 11, 2014). CatSper, sperm-associated cation channel; CNG, cyclic nucleotide-gated channel; CysLoop receptors, cysteine-loop ligand-gated receptor; IP₃ receptor, inositol 1,4,5-trisphosphate receptor; Letm1, leucine zipper-EF-hand containing transmembrane protein 1; MCU, mitochondrial Ca²⁺ uniporter; MICU, mitochondrial EF hand Ca²⁺ uniporter regulator; NC(K)X, Na⁺/Ca²⁺ (K⁺-dependent) exchanger; P2X receptor, P2X purinergic receptor; PMCA, plasma membrane Ca²⁺ ATPase; SERCA, sarco/endoplasmic reticulum Ca²⁺ ATPase; SPCA, secretory pathway Ca²⁺ ATPase.

supplementary fig. S1, Supplementary Material online) including CatSper channels, Ca_v channels, ligand-gated channels, second messenger-gated channels, and TRP channels. Components of the animal Ca²⁺-release-activated Ca²⁺

channel complex—stromal interaction molecules (STIM) and Orai channels—are absent in *Au. limacinum* and *T. trahens*. Both *Au. limacinum* and *T. trahens* possess intracellular ion channels—inositol 1,4,5-trisphosphate receptors and two-pore channels but not ryanodine receptors. In addition, *Au. limacinum* and *T. trahens* contain a complete set of Ca²⁺ exchange systems—Ca²⁺ ATPases and three classes of the cation/Ca²⁺ exchangers (Cai and Lytton 2004)—the K⁺-independent Na⁺/Ca²⁺ exchangers, the K⁺-dependent Na⁺/Ca²⁺ exchangers, and the Ca²⁺/H⁺ exchangers (CAXs). Here, we focus our analysis on the early evolution of the CatSper channel complex. Description of Ca²⁺ signaling molecules other than the CatSper complex is discussed in [Supplementary Material](#) online.

The CatSper Ca²⁺ Channel Complex

The typical “9 + 2” motile flagellum in human spermatozoa is conserved across species and likely evolved in the last common ancestor of all eukaryotic organisms (Mitchell 2007). Studies of flagellar axonemes have focused on sperm cells from unikont model organisms—marine invertebrates such as sea urchins and tunicates and also in bikont protists such as *Chlamydomonas* and *Paramecium* (Inaba 2011). Basic flagellar motility is controlled by the active sliding of paired outer doublet microtubules coupled with ATP hydrolysis by axonemal dyneins, which does not require the elevation of Ca²⁺ concentration or even an intact plasma membrane. In contrast, Ca²⁺ is necessary for complex flagellar activities such as capacitation, chemotaxis, and hyperactivated motility (Lishko et al. 2012).

As the most biochemically complex ion channel known to date, the CatSper complex is located in the principle piece of the sperm flagellum (fig. 3A). Ca²⁺ influx through CatSper is essential for sperm cell hyperactivated motility and male fertility in mammals (Lishko et al. 2012). CatSper is present in sea urchins and tunicates but have not been identified previously in any bikonts (Cai and Clapham 2008).

Except in the basal fungus *A. macrogynus* (Cai and Clapham 2012), all known CatSper channel complexes are composed of four pore-forming α subunits. Loss of either of the four α subunits by gene knockout results in male infertility and the complete loss of the whole protein complex (Qi et al. 2007). Similarly, numerous cases of lineage-specific simultaneous loss of all four CatSper α subunit and auxiliary subunit genes are present throughout metazoan evolution (Cai and Clapham 2008). The mammalian CatSper complex also contains at least three auxiliary subunits—CatSper- β , CatSper- γ , and CatSper- δ , all of which are sperm-specific transmembrane proteins (Lishko et al. 2012).

We found that *Au. limacinum* contains the same components of the CatSper channel complex in *T. trahens*; these include the pore-forming α subunits 1–4, auxiliary β and γ subunits, and a distantly related homolog of the δ subunit (figs. 2 and 3B and C, [supplementary fig. S1, Supplementary Material](#) online). The loop regions containing the key acidic residue aspartate are highly conserved in CatSper α subunits from *Au. limacinum* to humans (fig. 3D), suggesting the high

Ca²⁺ selectivity observed in mammalian CatSper is likely retained in *Au. limacinum*. The sequence regions outside the transmembrane segments and the pore loop of CatSper are poorly conserved across species, which possibly indicates that modulation of complex flagellar motility differs in a species-specific manner.

The highly conserved composition of the CatSper complex between Unikonta and Bikonta suggests that CatSper might play an important role in regulating flagellar activity in ancestral protists prior to the beginning of eukaryotic radiation, much earlier than previously thought (Cai and Clapham 2008). Alternatively, it remains a possibility that the presence of the CatSper complex in *Au. limacinum* might be caused by a horizontal gene transfer event from basal fungal species. In humans and in *A. macrogynus*, sperm cell motility can be regulated by Ca²⁺ influx in response to progesterone (Lishko et al. 2011; Strunker et al. 2011) or sex pheromones (Pommerville et al. 1990), respectively. Activation of the CatSper complex by environmental signals such as progesterone and pH changes induces Ca²⁺ influx in human sperm.

Thraustochytrid protists reproduce solely by means of biflagellate zoospores (fig. 2) (Raghukumar and Damare 2011). It is conceivable that in thraustochytrid protists such as *Au. limacinum*, modulation of cell motility by the CatSper complex prompts biflagellate zoospores to move toward environmental cues (Fan et al. 2002). The CatSper complex is absent in most bikonts and fungi. Although the basic “9 + 2” flagellar structure is highly conserved across bikonts and unikonts, the protein composition of flagellar axonemes and signaling cascades that regulate flagellar motility may have been modified to fit different physiological environments (Inaba 2011).

The CatSper channel pore-forming α subunits and NC(K)X and CAXs are believed to be closely related to their prokaryotic counterparts, the NaChBac Na⁺ channel (Clapham and Garbers 2005), and the YRBG exchanger (Philipson and Nicoll 2000), respectively. The identification of these genes in *Au. limacinum* and *T. trahens* supports a prokaryotic genesis of ancestral eukaryotic signaling systems (Shpakov and Pertseva 2008). It is speculated that gene duplication gave rise to four unique CatSper channel pore-forming α subunits and three classes of exchangers—K⁺-dependent Na⁺/Ca²⁺ exchangers, K⁺-independent Na⁺/Ca²⁺ exchangers, and CAXs. Indeed, two diatom genomes contain four and three copies of NaChBac-type channels, which are believed to have been acquired through horizontal gene transfer of an ancestral NaChBac-like channel from a prokaryote into the common ancestor of diatoms, followed by subsequent gene duplications (Verret et al. 2010). Notably, these diatom NaChBac-type channels exhibit sequence divergence that might render them more selective for Ca²⁺, a function adapted for the CatSper complex. Recent phylogenetic analyses of ion selectivity between animal and bacterial ion channels also suggest that NaChBac-type bacterial channels are closely related to CatSper channels (Liebeskind et al. 2013).




Fig. 3. Identification of CatSper channels in *Aurantiochytrium limacinum*. (A) Schematic representation of mammalian spermatozoa. The CatSper channel complex is located at the principle piece that is critical for Ca²⁺-modulated sperm motility. (B) Distribution of CatSper in select bikont species. *Aurantiochytrium limacinum* contains CatSper α subunits 1–4, whereas the exact number of CatSper α subunit in *Cyanophora paradoxa* is unknown due to incomplete genome assembly. (C) Phylogenetic relationship of CatSper α subunits. A maximum likelihood phylogenetic tree (LG + I + G + F model) showing the relationship of protist and animal homologs of CatSper α subunits 1–4 from the thraustochytrid protist *A. limacinum* (Ali), the amphioxus *Branchiostoma floridae* (Bfi), *Homo sapiens* (Hsa), *Mus musculus* (Mus), and the apusozoan protist *Thecamonas trahens* (Ttr). NaChBac, a prokaryotic voltage-gated Na⁺ channel isolated from *Bacillus halodurans* (Ren et al. 2001), was used as an outgroup. Bootstrap values above 60 are shown at the nodes. (D) Sequence alignment of the pore loop regions of NaChBac and CatSper α subunits from *A. limacinum* (Ali) and *H. sapiens* (Hsa). The asterisk symbol indicates the location of a key acidic residue important for ion selectivity. CatS, CatSper α subunit.

Conservation of Key Components of Ca²⁺ Signaling System in *Cyanophora paradoxa*

The glaucophyte *Cyano. paradoxa* is a basally diverging, photosynthetic freshwater alga that emerged before the divergence of green plants and red algae (Price et al. 2012). The land plant *Ar. thaliana* and the unicellular red alga *C. merolae* are known to retain very simplified Ca²⁺ signaling networks (Verret et al. 2010) (fig. 1A). We found conservation of several key Ca²⁺ signaling molecules in the *Cyano. paradoxa* genome including homologs of CatSper channels, Ca_v channels, and TRP channels, all of which are absent in *Ar. thaliana* and *C. merolae* (figs. 1A and 3B and supplementary fig. S1, Supplementary Material online). *Cyanophora paradoxa* lacks homologs of P2XRs, inositol 1,4,5-trisphosphate receptors, and NC(K)X exchangers, which are all present in

Au. limacinum and *T. trahens*. The identification of CatSper homologs in two bikont groups—Archaeplastida (*Cyano. paradoxa*) and Chromalveolata (*Au. limacinum*)—further supports the ancestral evolution of the CatSper complex.

In summary, we have demonstrated that the bikont protist *Au. limacinum* possess an extensive Ca²⁺ signaling machinery comparable to that in the unikont protist *T. trahens*. Consistent with previous reports (Wheeler and Brownlee 2008; Verret et al. 2010), our findings also suggest that the extensive network of Ca²⁺ signaling molecules observed in animals originated in ancestral protists before the eukaryotic radiation, prior to the divergence of animals, fungi, and plants. The ancestral Ca²⁺ signaling machinery containing the CatSper channel complex is generally conserved in the lineage leading to animals and can also be occasionally identified in very few protists such as *T. trahens* and *Au. limacinum*.

Although many ancestral Ca²⁺ signaling molecules were subsequently lost during the evolution of modern protists, plants and fungi, traces of relatively conserved Ca²⁺ signaling systems harboring CatSper can still be found in basally divergent species, for instance, the basal chytridiomycete fungus *A. macrogynus* (Cai and Clapham 2012) and the basally diverging glaucophyte *Cyano. paradoxa*. Characterization of the function and regulation of these ancestral homologs will provide novel insight into the understanding of their physiological roles and the evolution of the ancestral Ca²⁺ signaling machinery.

Supplementary Material

Supplementary information and figures S1 and S2 are available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>).

Acknowledgments

The authors thank the Department of Energy Joint Genome Institute (<http://genome.jgi.doe.gov>) and investigators of the genome projects listed on the JGI genome portal and the *C. paradoxa* genome project (Price et al. 2012) for making data publicly available. X.C. also thanks Yanhong Zhang for excellent technical assistance and for critical reading of the manuscript.

References

- Amador FJ, Stathopoulos PB, Enomoto M, Ikura M. 2013. Ryanodine receptor calcium release channels: lessons from structure-function studies. *FEBS J.* 280:5456–5470.
- Berridge MJ, Bootman MD, Roderick HL. 2003. Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol.* 4: 517–529.
- Burki F, Pawlowski J. 2006. Monophyly of Rhizaria and multigene phylogeny of unicellular bikonts. *Mol Biol Evol.* 23:1922–1930.
- Cai X. 2008. Unicellular Ca²⁺ signaling ‘toolkit’ at the origin of metazoa. *Mol Biol Evol.* 25:1357–1361.
- Cai X, Clapham DE. 2008. Evolutionary genomics reveals lineage-specific gene loss and rapid evolution of a sperm-specific ion channel complex: CatSper and CatSper β . *PLoS One* 3:e3569.
- Cai X, Clapham DE. 2012. Ancestral Ca²⁺ signaling machinery in early animal and fungal evolution. *Mol Biol Evol.* 29:91–100.
- Cai X, Lytton J. 2004. The cation/Ca²⁺ exchanger superfamily: phylogenetic analysis and structural implications. *Mol Biol Evol.* 21: 1692–1703.
- Case RM, Eisner D, Gurney A, Jones O, Muallem S, Verkhatsky A. 2007. Evolution of calcium homeostasis: from birth of the first cell to an omnipresent signalling system. *Cell Calcium* 42:345–350.
- Cavalier-Smith T, Chao EE. 2010. Phylogeny and evolution of apusomonadida (protozoa: apusozoa): new genera and species. *Protist* 161: 549–576.
- Chan CX, Reyes-Prieto A, Bhattacharya D. 2011. Red and green algal origin of diatom membrane transporters: insights into environmental adaptation and cell evolution. *PLoS One* 6:e29138.
- Clapham DE. 2007. Calcium signaling. *Cell* 131:1047–1058.
- Clapham DE, Garbers DL. 2005. International union of pharmacology. L nomenclature and structure-function relationships of CatSper and two-pore channels. *Pharmacol Rev.* 57:451–454.
- Derelle R, Lang BF. 2012. Rooting the eukaryotic tree with mitochondrial and bacterial proteins. *Mol Biol Evol.* 29:1277–1289.
- Dominguez DC. 2004. Calcium signalling in bacteria. *Mol Microbiol.* 54: 291–297.
- Fairclough SR, Chen Z, Kramer E, Zeng Q, Young S, Robertson HM, Begovic E, Richter DJ, Russ C, Westbrook MJ, et al. 2013. Premetazoan genome evolution and the regulation of cell differentiation in the choanoflagellate *Salpingoeca rosetta*. *Genome Biol.* 14: R15.
- Fan KW, Vrijmoed LL, Jones EB. 2002. Zoospore chemotaxis of mangrove thraustochytrids from Hong Kong. *Mycologia* 94:569–578.
- Inaba K. 2011. Sperm flagella: comparative and phylogenetic perspectives of protein components. *Mol Hum Reprod.* 17:524–538.
- King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough S, Hellsten U, Isogai Y, Letunic I, et al. 2008. The genome of the choanoflagellate *Monosiga brevicollis* and the origins of metazoan multicellularity. *Nature* 451:783–788.
- Kudla J, Batistic O, Hashimoto K. 2010. Calcium signals: the lead currency of plant information processing. *Plant Cell* 22:541–563.
- Liebeskind BJ, Hillis DM, Zakon HH. 2013. Independent acquisition of sodium selectivity in bacterial and animal sodium channels. *Curr Biol.* 23:R948–R949.
- Lishko PV, Botchkina IL, Kirichok Y. 2011. Progesterone activates the principal Ca²⁺ channel of human sperm. *Nature* 471:387–391.
- Lishko PV, Kirichok Y, Ren D, Navarro B, Chung JJ, Clapham DE. 2012. The control of male fertility by spermatozoan ion channels. *Annu Rev Physiol.* 74:453–475.
- Mitchell DR. 2007. The evolution of eukaryotic cilia and flagella as motile and sensory organelles. *Adv Exp Med Biol.* 607:130–140.
- Nagata T, Iizumi S, Satoh K, Ooka H, Kawai J, Carninci P, Hayashizaki Y, Otomo Y, Murakami K, Matsubara K, et al. 2004. Comparative analysis of plant and animal calcium signal transduction element using plant full-length cDNA data. *Mol Biol Evol.* 21:1855–1870.
- Phillipson KD, Nicoll DA. 2000. Sodium-calcium exchange: a molecular perspective. *Annu Rev Physiol.* 62:111–133.
- Pommerville JC, Strickland JB, Harding KE. 1990. Pheromone interactions and ionic communication in gametes of aquatic fungus *Allomyces macrogynus*. *J Chem Ecol.* 16:121–131.
- Price DC, Chan CX, Yoon HS, Yang EC, Qiu H, Weber AP, Schwacke R, Gross J, Blouin NA, Lane C, et al. 2012. *Cyanophora paradoxa* genome elucidates origin of photosynthesis in algae and plants. *Science* 335:843–847.
- Qi H, Moran MM, Navarro B, Chong JA, Krapivinsky G, Krapivinsky L, Kirichok Y, Ramsey IS, Quill TA, Clapham DE. 2007. All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility. *Proc Natl Acad Sci U S A.* 104:1219–1223.
- Raghukumar S, Damare VS. 2011. Increasing evidence for the important role of Labyrinthulomycetes in marine ecosystems. *Bot Mar.* 54:3–11.
- Ren D, Navarro B, Xu H, Yue L, Shi Q, Clapham DE. 2001. A prokaryotic voltage-gated sodium channel. *Science* 294:2372–2375.
- Risberg I, Orr RJ, Kluge R, Shalchian-Tabrizi K, Bowers HA, Patil V, Edvardsen B, Jakobsen KS. 2009. Seven gene phylogeny of heterokonts. *Protist* 160:191–204.
- Rokas A. 2008. The origins of multicellularity and the early history of the genetic toolkit for animal development. *Annu Rev Genet.* 42: 235–251.
- Ruiz-Trillo I, Roger AJ, Burger G, Gray MW, Lang BF. 2008. A phylogenomic investigation into the origin of metazoa. *Mol Biol Evol.* 25: 664–672.
- Sebe-Pedros A, de Mendoza A, Lang BF, Degnan BM, Ruiz-Trillo I. 2011. Unexpected repertoire of metazoan transcription factors in the unicellular holozoan *Capsaspora owczarzaki*. *Mol Biol Evol.* 28: 1241–1254.
- Shemarova IV, Nesterov VP. 2005. [Evolution of Ca(2+) signaling mechanisms. Role of calcium ions in signal transduction in lower eukaryotes]. *Zh Evol Biokhim Fiziol.* 41:303–313.
- Shpakov AO, Pertseva MN. 2008. Signaling systems of lower eukaryotes and their evolution. *Int Rev Cell Mol Biol.* 269:151–282.
- Stechmann A, Cavalier-Smith T. 2002. Rooting the eukaryote tree by using a derived gene fusion. *Science* 297:89–91.
- Steenkamp ET, Wright J, Baldauf SL. 2006. The protistan origins of animals and fungi. *Mol Biol Evol.* 23:93–106.
- Strunker T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R, Kaupp UB. 2011. The CatSper channel mediates progesterone-induced Ca²⁺ influx in human sperm. *Nature* 471:382–386.

- Suga H, Chen Z, de Mendoza A, Sebé-Pedrós A, Brown MW, Kramer E, Carr M, Kerner P, Vervoort M, Sánchez-Pons N, et al. 2013. The *Capsaspora* genome reveals a complex unicellular prehistory of animals. *Nat Commun.* 4:2325.
- Tsai MF, Jiang D, Zhao L, Clapham D, Miller C. 2014. Functional reconstitution of the mitochondrial $\text{Ca}^{2+}/\text{H}^{+}$ antiporter Letm1. *J Gen Physiol.* 143:67–73.
- Verkhatsky A, Parpura V. 2014. Calcium signalling and calcium channels: evolution and general principles. *Eur J Pharmacol.* 739C:1–3.
- Verret F, Wheeler G, Taylor AR, Farnham G, Brownlee C. 2010. Calcium channels in photosynthetic eukaryotes: implications for evolution of calcium-based signalling. *New Phytol.* 187:23–43.
- Wheeler GL, Brownlee C. 2008. Ca^{2+} signalling in plants and green algae—changing channels. *Trends Plant Sci.* 13:506–514.
- Zelter A, Bencina M, Bowman BJ, Yarden O, Read ND. 2004. A comparative genomic analysis of the calcium signaling machinery in *Neurospora crassa*, *Magnaporthe grisea*, and *Saccharomyces cerevisiae*. *Fungal Genet Biol.* 41:827–841.