

A Study of Clinical Profile of Snake Bite at a Tertiary Care Centre

Gaurav Bhalla, Dhanesh Mhaskar, Anubhav Agarwal

Department of Medicine, Krishna Institution of Medical Sciences, Karad, Satara, Maharashtra, India

ABSTRACT

Background: Snake bite is an important occupational and rural hazard because India has always been a land of Exotic snakes. In Maharashtra, common poisonous snakes are Cobra, Russell's Viper, Saw Scaled Viper, and Krait. It is a fact that inspite of heavy morbidity and mortality, very little attention is paid by the clinicians to this occupational hazard. **Aims:** To study the prevalence of poisonous and non-poisonous snake bites in part of Western Maharashtra with reference to age, sex, occupation, part of body bitten, time of bite and seasonal variation, and the types of poisonous snakes common in this locality and their clinical manifestations along with the systemic envenomation from various types of poisonous snakes and their effective management in reducing the mortality rate. **Materials and Methods:** This was a retrospective study conducted between May 2010 to May 2012 at a tertiary health care center in Maharashtra. **Result:** A total of 150 patients were studied in our hospital. Out of 150, 76 patients were of poisonous snake bite and 74 patients were of non-poisonous snake bite. Out of these 76 poisonous snake bites, 42 were viperine snake bites, 21 were neuroparalytic snake bites and 13 were locally toxic (LT) snake bites. **Conclusion:** Snake bite is a common life-threatening emergency in the study area. Delay in hospitalization is associated with poor prognosis and increased mortality rate due to consumptive coagulopathy, renal failure, and respiratory failure. Unusual complications like pulmonary edema, intracerebral hemorrhage, Disseminated intravascular coagulation (DIC) were observed in present study.

Key words: Non-poisonous bite, neuroparalytic bite, poisonous bite, vasculotoxic bite

INTRODUCTION

In Maharashtra, common poisonous snakes are Cobra, Russell's Viper, Saw Scaled Viper, and Krait.^[1] World mortality from snake bite is estimated as 50,000 to 1,00,000 annually (McNamee 2001) and the greatest number of reported snake bite death occurring in Indian subcontinent is 10,000 to 15,000 annually.^[2] World Health Organization (WHO 1963) reports 40,000 annual deaths in tropical countries. Largest number of deaths reported in India are from Bengal, Uttar

Pradesh (UP), Tamil Nadu, Bihar, and Maharashtra.^[3] It is a fact that inspite of heavy morbidity and mortality, very little attention is paid by the clinicians to this occupational hazard.

MATERIALS AND METHODS

The present descriptive observational study was carried out in medicine wards of Krishna hospital, Karad, Satara, Maharashtra, India during May 2010 to May 2012. A total of 150 cases of snake bite were admitted in medicine wards during the study period. After obtaining consent, data was collected on pre-designed, pretested, and structured questionnaire by interviewing the study subjects who were hospitalized during the study period. A detailed information regarding demographic and epidemiological parameters such as age, sex, residence, occupation, site of bite and place of bite, type of snake if identified, etc., was obtained. Time interval to reach the health facility after

Access this article online

Quick Response Code:



Website:

www.toxicologyinternational.com

DOI:

10.4103/0971-6580.139811

Address for correspondence: Dr. Dhanesh Mhaskar, Department of Medicine, Krishna Institution of Medical Sciences, Karad, Satara, Maharashtra, India. E-mail: drdhaneshmh@yahoo.com

snake bite and first aid received if any was asked to them. Thorough clinical examination was carried out in each case. For identification of type of snake bite (Vasculotoxic, VT, Neuroparalytic, and Non-poisonous) opinion from treating physician was taken. Subsequent information was collected on the day of discharge or death of the patient from the case paper of the patient.

RESULTS

A total of 150 patients were studied in our hospital. Out of 150, 76 patients were of poisonous snake bite and 74 patients were of non-poisonous snake bite. Out of these 76 poisonous snake bites, 42 were viperine snake bites, 21 were neuroparalytic snake bites, and 13 were locally toxic (LT) snake bites [Tables 1-18].

Table 1: Age-wise prevalence of poisonous snake bites

Age-group (years)	No. of poisonous bites (%)	Total no. of snake bites %
14-30	43 (56.57)	89
31-50	28 (36.84)	50
>50	5 (6.57)	11

Table 2: Occupational incidence of snake bites

Occupation	No. of cases	Percentage
Farmer	122	81.33
Housewife	14	9.33
Student	13	8.66
Snake charmer	1	0.66

Table 3: Sex incidence of snake bites

Sex	No. of poisonous cases	No. of non-poisonous cases	% (poison/non-poisonous)
Male	61	38	66
Females	15	36	34

Table 4: Incidence of snake bite with reference to site of bite

Site of bite	No. of non-poisonous cases	No. of poisonous cases
Upper extremity (UE)	14	3
Lower extremity (LE)	60	73
Trunk	-	-

Table 5: Relation of time with snake bite

	No. of non-poisonous cases	No. of poisonous cases
Af ernoon (12.00PM-4.00PM)	12	7
Evening (4.00 PM-8.00 PM)	35	41
Night (8.00 PM-6.00 AM)	18	15
Morning (6.00 AM-12.00 PM)	9	13

SUMMARY AND CONCLUSION

In all, 150 patients of snake bite were studied. Out of 150 patients, 74 patients had non-poisonous snake bites and 76 patients had poisonous snake bites. Maximum incidence of snake bite was found in farmers (81.33%) than in any other occupation. Males (66%) were bitten more than females (34%) among poisonous bites. Maximum incidence of snake bites was found in rural areas (85%) than urban areas (15%). Poisonous snake bites were more prevalent in rural than urban areas. Bites were commonly seen on the lower extremities 88.6% than upper (11.33%). Maximum bites were in the night (72.66%) than day time (27.33%) among which 73.68% were poisonous bites at night and 27% were poisonous bites at day time. Higher incidence

Table 6: Seasonal incidence of snake bite

	No. of non-poisonous cases	No. of poisonous cases
Monsoon (June-Sept)	28	24
Winter (Oct-Feb)	11	10
Summer (March-May)	35	42

Table 7: Incidence of snake seen and brought

	No. of non-poisonous cases	No. of poisonous cases
Seen and brought	5	8
Seen not brought	40	56
Not seen	29	12

Table 8: Incidence of poisonous snake bite with local toxicity

	No. of VT cases	No. of NT cases
With local toxicity	39	15
Without local toxicity	3	6

VT = Vasculotoxic, NT = Neuroparalyt c

Table 9: Loss of time from the bite till the admission

Time	No. of non-poisonous cases	No. of poisonous cases			
		VT	NT	LT	Death
Within 6 hours	48	21	16	1	-
6 to 24 hours	26	15	4	5	2
>24 hours	0	6	1	7	1

VT = Vasculotoxic, NT = Neuroparalyt c, LT = Locally toxic

Table 10: The no. of patients who received first-aid

	No. of non-poisonous cases	No. of poisonous cases		
		VT	NT	LT
Tourniquet	22	23	12	10
ASV	-	7	2	1
False beliefs	-	2	3	-
No. Rx	51	9	4	2

VT = Vasculotoxic, NT = Neuroparalyt c, ASV = Ant venom serum, LT = Locally toxic

Table 11: Incidence of local toxicity in poisonous snake bite

	LT	VT	NT
Pain	9	33	11
Local bleed	11	35	3
Cellulitis	4	24	6
Gangrene	0	8	1

VT = Vasculotoxic, NT = Neuroparalytic, LT = Locally toxic

Table 12: Incidence of symptomatology of neuro paralytic snake bite poisoning

S/s	No. of cases	Percentage
Ptosis	21	100
Dysphagia	9	42.85
Ophthalmoplegia	18	85.71
Unconsciousness	6	28.57
Flaccid limb paralysis	6	28.67
Convulsions	-	0
Cellulitis	8	38.09
Respiratory paralysis	19	90.47

Table 13: Incidence of symptomatology in viperine snake bite

S/s	No. of cases	Percentage
Local bleed	35	83.33
Gum bleed	3	7.14
Hemoptysis (H3)	2	4.76
Epistaxis (E)	1	2.3
Hematemesis (H2)	2	4.76
Malena	2	4.76
Hematuria (H1)	18	42.85
Oliguria	20	47.61
Ecchymosis (ec)	2	4.76
Intracerebral hem.	1	2.3
Subconjunctival hem.	2	4.76
Cellulitis	24	57.14
ARF	20	47.61

ARF = Acute renal failure

Table 14: Incidence of complications of vasculotoxic snake bite

VT	ARF	Sur. inter	DIC	Death
42	20	8	24	4

VT = Vasculotoxic, ARF = Acute renal failure, DIC = Disseminated intravascular coagulation

of snake bites was found in summer (51.33%) i.e. March to May. In eight cases, the snakes were seen/identified. A total of 57.33% patients reported to medical help within 6 hours, among which 24.41% were VT and 18.60% were NT. Sixty-seven patients came with tourniquet application, out of which 45 had poisonous bite, out of which 9 patients had received ASV. Fifty-one cases did not receive any sort of medical help before admission. Five cases had received non-medical treatment. Time of onset of systemic

Table 15: Type of treatment received in vasculotoxic snake bite

ASV	42
Diuretics	13
Hemodialysis	7
Blood transfusion	4

ASV = Antivenom serum

Table 16: Duration of neuroparalytics on ventilator

Duration	No. of cases
Less than 12 hours	10
12-24 hours	4
>24 hours	1

Table 17: Incidence in relation to condition at discharge

	LT	NP	VT
Complete recovery	13	17	10
Complications	0	1	28
Death	0	3	4

VT = Vasculotoxic, NP = Non-poisonous, LT = Locally toxic

Table 18: Comparative mortality in poisonous snake bite

	No. of cases	No. of death	%
Neuroparalytic	21	3	3.94
VT	42	4	5.26
LT	13	0	0

VT = Vasculotoxic, LT = Locally toxic

manifestation was earlier in neuroparalytic snake bites than VT snake bite. VT snake bites (55.26%) were more common than neuroparalytic bites (27.63%). Commonest VT manifestation was local bleed (83.33%) and next on the list was cellulitis (57.14%). Hematuria (42.85%) was the commonest manifestation in those patients who developed ARE. Ptosis (100%) was the commonest and earliest manifestation of neuroparalytic snake bite. Out of VT snake bites, 20 were of ARE, 7 required hemodialysis. Four DIC patients received blood transfusion. Thirteen patients of ARF received diuretics and recovered completely. Thirteen cases were LT; out of them, 4 patients developed cellulitis. Nineteen patients with neuroparalytic bite required artificial ventilation. Seven patients died because they came late to seek medical treatment and by that time, complications were beyond control and were associated with systemic infection. Two patients developed anaphylactic reaction to ASV and were treated conservatively and recovered completely.

DISCUSSION

In India, it is believed that 2 lakh people are bitten by snakes and about 15,000-30,000 cases/year prove fatal.^[2]



Figure 1: Ptosis in neuroparalytic snake bite

In the present study, 150 cases of snake bites including poisonous and non-poisonous bites were studied in the medical wards of Krishna Hospital from 1st May 2010 to 31st May 2012. The number of poisonous cases were 76 and the number of non-poisonous cases were 74. Among the poisonous snake bites, the number of Viperine bites were 42, the number of neuroparalytic bites were 21 and the number of LT cases were 13. In the present study, maximum no. of patients bitten were between the age-group of 14-30 years of age. Next in order of frequency was the age-group 31-50 years. These observations go more in favor of the study of Russel *et al.* (1979)^[4] and Hutchison *et al.* (1929). Among the group of 14-30 years, 43 patients had poisonous snake bite and in the age-group of 31-50 years, 28 patients had poisonous bites. In the age-group above 51, five bites were poisonous in nature.

In the present study, 99 (66%) patients were male victims, whereas the female victims were 51 (34%) in number. All the previous reports mention the same findings. Ahuja and Singh^[5] in 1954 reported the ratio as 4:1 (M: F). Bhat^[6] *et al.* in 1974 reported the incidence as 7:3 (M: F).^[6] Among male patients, 61 were poisonous, whereas in females, 15 were poisonous snake bites. In the present study, 122 were farmers, 13 were students, 14 were housewives, and 1 was a snake charmer. Studies undertaken by Bhat^[6] *et al.* (1974), Saini^[7] *et al.* in (1984), and Sarangi^[8] *et al.* in (1977) showed the incidence in farmers to be 75%, 78%, and 72%, respectively. In the present study, rural prevalence of snake bite was 117 out of 150 cases. Among them, 69 were poisonous snake bites. The prevalence in urban area was 33 out of 150 cases, among which 7 were poisonous cases. Incidence of bites in present studies are as follows. In lower extremities bites are 133 (88.66%) and in upper extremities 17 (11.33%). Among them, upper extremity poisonous cases were 3 and lower extremity poisonous cases were 73. Reid^[9] mentions that most of the bites in tropical countries are on lower extremities because the victims are bitten by treading on or near the snake, while in non-tropical countries most bites are on fingers and hands because the victim deliberately handles

the snake. In the present study, the day was divided into four groups. Maximum number of cases were detected in the interval between 4.00 PM to 8.00 PM. The documented figure was 109 among 150 patients. Next on the list was the time interval between 8.00 PM to 6.00 AM where number of patients bitten was 41. These figures show close resemblance with the observation of Virmani and Dutt.^[10] They have noted 12% cases during day time. In the present study, higher incidence was found in summer (March to May), 77 (51.33%) were total snake bites in summer, among them 42 were poisonous bites and 35 were non-poisonous bites. Next on the list was monsoon where the total number of snake bites was 52, among which 24 were poisonous bites and 28 were non-poisonous. Neuroparalytic snake bites were more at the end of summer and beginning of monsoon. Incidence of VT bites were comparatively more in winter. We defined summer as period of the year from March to May, winter period is between October to February, and Monsoon is June to September. Banerjee^[11] (1978) noted incidence of 70-80% during May to October. In the present study, we have noted one case of a snake catcher from Satara district who was a professional. He had a neuroparalytic cobra bite. Remaining cases were all unprovoked.

In the present study, only eight patients had brought snakes along with them and they were identified in this hospital. Among them three were cobras, two vipers, one was Krait, and two were *Echis carinatus*. In the present study, 86 (57.33%) patients were admitted within first 6 hrs. Among which 21 cases (24.41%) were VT and 16 (18.60%) were neuroparalytic. Seventy-six percent of neuroparalytic bites got admitted within the first 6 hrs. Fifty-two patients were admitted within the first 24 hrs but after 6 hrs, among which 15 were VT and 4 were NT. Two NT bites expired within 1 to 2 hrs after admission. In a series of 135 patient by Lahori *et al.*, 85% patients were admitted in the first 24 hours after bite. Among 150 patients, 67 (44.66%) had applied tourniquet. Out of 67 patients, 45 were poisonous bites and 22 non-poisonous bites. Out of 45 cases of poisonous bites, 23 were VT, 12 were neuroparalytic, and 10 were LT bites. In a study conducted by George Watt^[12] *et al.* (May 1988), tourniquet was applied in 94% patients, 4 were asymptomatic prior to release of their tourniquet, and in 11 patients symptoms worsened precipitously. In the present series, 53 (69.73%) patients had local pain. Among which 9 (16.98%) were LT, 33 (62.26%) were VT and 11 (20.75%) were neuroparalytic bites. In the present study, local edema was present in 64 (84.20%) patients of poisonous snake bite. Out of which 40 patients had VT snake bite, 11 patients had neuroparalytic snake bite, and 13 bites were LT. In VT bite 95.23% patients had local edema while in neuroparalytic snake bite 52.38% patients had local edema. Local cellulitis was seen in 24 (57.14%)

patients of VT snake bite and 6 (28.57%) patients of neuroparalytic snake bites.

A total of 32.66% of the patients in the present study had local bleeding. Out of which 11 patients had LT bite, 35 patients had VT bite and 3 patients had neuroparalytic snake bites. In the present study, 3 (7.14%) patients had gum bleeding but Purohit (1944) described gum bleeding as the commonest manifestation of viperine bite. Corkill^[13] (1956) described hematuria as the commonest manifestation but in the present study, 18 (42.85%) patients had hematuria. Ecchymosis was present in 2 (4.75%) patients in the present study but Bhat (1974) described ecchymosis in 43% of patients. Hematemesis was present in 2 (4.75%) patients in the present study, whereas Bhat^[6] showed that 37 patients among 310 patients had hematemesis. In the present study, 2 (4.75%) patients had haemoptysis and 1 (2.3%) patient had epistaxis. Fundal hemorrhage was seen in one patient (2.3%) who expired within 2 days. Reid^[14] *et al.* in his study of 281 cases of viper bite had not found a single case with fundal haemorrhage. Saini^[7] and Sharma *et al.* (1984) in their study of 200 cases of poisonous bite do not report about fundal hemorrhage. In the present study, 100% patients developed ptosis among 21 cases of Neuroparalytic bite. Ophthalmoplegia was seen in 18 (85.71%) patients of Neuroparalytic bite. Six (28.57%) patients became unconscious, 6 (28.57%) patients with flaccid limb paralysis, 19 (90.47%) patients developed respiratory paralysis, among which 15 patients needed respirator.

ARF was seen in 20/42, 47.61% of patients of VT bite. In all the cases of ARF except one, DIC was the cause for it. Basu *et al.* (1977) observed ARF in 27 cases of viper bite and attributed it to circulatory collapse and shock in 5 cases, direct nephrotoxicity in 4 cases. In the present study, 7 patients died of poisonous snake bites out of which 3 (3.94%) were neuroparalytic bite patients and 4 were VT bite patients. Among neuroparalytic bites, mortality rate (14.28%) was higher than in VT bite (9.52%). All 3 patients died because of delay in receiving respiratory assistance. Among four deaths of VT bites, three patients died of acute renal failure, and one patient died of DIC and intracerebral hemorrhage. Lahori^[14] *et al.* (1981) showed mortality rate to be 2% and 3 deaths were due to central nervous system (CNS) involvement.

Bleeding time, clotting time was raised in all patients of VT snake bites. Similar incidence has been reported by Saini^[7] *et al.* and Lahori *et al.* in their studies. BT, CT, PT gradually returned to normal within 3-4 days after full dose of ASV and blood transfusion in present study. Prothrombin time was abnormal in almost all the VT bites. In the present study, thrombocytopenia was noted in 24/41 patients of VT bite with coagulation defects. Saini^[7] *et al.* (1981) noted thrombocytopenia in only 5% patients and Reid^[9] tioned

that platelet count may be depressed but is often normal in viperine bites.

In the present study, two patients developed anaphylaxis for ASV and they were treated with steroids and adrenaline. In the present study, 76/150 (50.66%) patients received ASV. According to Reid *et al.*, ASV is not very effective against local effects of venom. In the present study, 69/76 reversed completely. Three deaths in neuroparalytics were because of delay in administration of ASV producing respiratory paralysis and delay in ventilatory support who were transferred from other hospital to our hospital. In the present study, four (9.52%) patients of viperine snake bites received blood transfusions. Saini^[7] *et al.* (1984) and Reid *et al.* feel that blood transfusion is not required routinely. In the present series, neostigmine with atropine was given to 13 patients out of 21 patients of neuroparalytic snake bite. Most of the patients responded to it within 4-6 hours. In these patients, first to improve was respiratory paralysis, then ophthalmoplegia and lastly ptosis. Dubay^[15] *et al.* (1981), Banerjee^[16] *et al.*, Dash^[17] *et al.* (1976) are strong supporters of this regime but Reid^[9] had doubts about the benefits of this mode of therapy.

In the present series, ARF was observed in 20 patients among 42 (47.61%) patients of VT snake bite. Eleven patients were treated conservatively with diuretics, renal diet, and fluid restriction. Seven patients needed hemodialysis. Three patients died of renal failure among which 2 patients died after hemodialysis and one died because of acute renal failure. Saini^[7] *et al.* reported 8 cases of ARF and 7 cases recovered with conservative treatment. In the present study, among 21 patients of neuroparalytic bite (71.42%), 15 needed ventilatory support and 4 (19.04%) patients needed only intubation. Among which 3 patients died. Total duration of ventilator was less than 24 hrs for most of the patients except one. Not a single patient needed tracheostomy. In the present study, 9 patients required surgical intervention, out of which 8 patients had viper bite and one patient had Elapid bite. Reid^[17] *et al.* recommends early surgical intervention to prevent extension of infection and development of gangrene.

REFERENCES

1. Warell DA. Injuries, envenoming, poisoning and allergic reactions caused by animals. Oxford Textbook of Med; 1984. p. 6.35-6.40.
2. Swaroop and Grab. Snake bite Mortality in world. Vol. 10. WHO bulletin; 1954. p. 35.
3. Manson Bahr Poisonous snakes, Manson's Tropical Diseases. Ch XL VII; 1968. p. 749-60.
4. Russell FE, Emery CA. Effects of corticosteroids on lethality of Ancistrodon contortrix venom. Am J Med Sci 1961;241:507-11.
5. Ahuja MM. Progress in clinical medicine in India. Second Series; 1978. p. 136-77.

6. Bhat RN. Viperine snake poisoning in Jammu. *J Indian Med Assoc* 1974;63:383-92.
7. Saini RK, Sharma S, Singhs, Pathania NS. Snake bite poisoning: A preliminary report. *J Assoc Physicians India* 1984;32:195-7.
8. Sarangi A, Jana I, Das JP. Clinical profile of snake bite poisoning. *J Assoc Physicians India* 1977;25:55-60.
9. Reid HA, Theakston RD. The management of snake bite. *Bull WHO* 1983;61:885-95.
10. Virmani SK, Dutt OP. A profile of snake bite poisoning in Jammu region. *J Indian Med Assoc* 1987;85:132-4.
11. Banerjee RN. Poisonous snakes of India, progress in, clinical medicine' in India. In: Ahuja MS, editor. 1st ed. 1987. p. 136-1.77.
12. Watt G, Padre L, Tuazon ML, Theakston RD, Laughlin LW. Tourniquet application after cobra bite: Delay in the onset of neurotoxicity and the dangers of sudden release. *Am J Trop Med Hyg* 1988;38:618-22.
13. Corkill NL, Venoms EE, Euckley, Porges N. Washington D.C; 1956. p. 331.
14. Reid HA, Thean PC, Martin WJ. Specific antivenin and prednisolone in viper bite poisoning. *Br Med J* 1963:1378-80.
15. Dubey GK, Joglekar VK, Chaubey BS, Kalamkar RB. Neostigmine in the treatment of snake bite. *J Assoc Physicians India* 1981;29:229-31.
16. Banerjee RN, Siddiqui ZA. Therapeutic advance in the treatment of snake venom poisoning. *Proc. 5th international symposium on Animal plant and Microbial Toxins, Toxicology*; 1976.
17. Dash SC, Ghosh SK, Mathur DC, Jha GN, Prasad U, Grewal KS. Neurotoxic snake bite-dramatic recovery following neostigmine therapy. *J Assoc Physicians India* 1976;24:535-7.

How to cite this article: Bhalla G, Mhaskar D, Agarwal A. A study of clinical profile of snake bite at a tertiary care centre. *Toxicol Int* 2014;21:203-8.

Source of Support: Nil **Conflict of Interest:** None declared.

Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

1) **First Page File:**

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

2) **Article File:**

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1 MB. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

3) **Images:**

Submit good quality color images. Each image should be less than 4096 kb (4 MB) in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

4) **Legends:**

Legends for the figures/images should be included at the end of the article file.