

# Role of YpeB in Cortex Hydrolysis during Germination of *Bacillus anthracis* Spores

Casey B. Bernhards, David L. Popham

Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, USA

The infectious agent of the disease anthrax is the spore of *Bacillus anthracis*. Bacterial spores are extremely resistant to environmental stresses, which greatly hinders spore decontamination efforts. The spore cortex, a thick layer of modified peptidoglycan, contributes to spore dormancy and resistance by maintaining the low water content of the spore core. The cortex is degraded by germination-specific lytic enzymes (GSLEs) during spore germination, rendering the cells vulnerable to common disinfection techniques. This study investigates the relationship between SleB, a GSLE in *B. anthracis*, and YpeB, a protein necessary for SleB stability and function. The results indicate that  $\Delta sleB$  and  $\Delta ypeB$  spores exhibit similar germination phenotypes and that the two proteins have a strict codependency for their incorporation into the dormant spore. In the absence of its partner protein, SleB or YpeB is proteolytically degraded soon after expression during sporulation, rather than escaping the developing spore. The three PepSY domains of YpeB were examined for their roles in the interaction with SleB. YpeB truncation mutants illustrate the necessity of a region beyond the first PepSY domain for SleB stability. Furthermore, site-directed mutagenesis of highly conserved residues within the PepSY domains resulted in germination defects corresponding to reduced levels of both SleB and YpeB in the mutant spores. These results identify residues involved in the stability of both proteins and reiterate their codependent relationship. It is hoped that the study of GSLEs and interacting proteins will lead to the use of GSLEs as targets for efficient activation of spore germination and facilitation of spore cleanup.

Bacterial spores from the *Bacillus* and *Clostridium* genera are metabolically dormant and are known for their extreme resistance to heat, desiccation, UV radiation, chemicals, and other insults (1–3). These resistance properties allow spores to survive in the environment for extended periods and have made eradication from contaminated sites incredibly difficult (4). Spore dormancy and wet heat resistance are largely dependent on spore core dehydration, which is maintained by a thick layer of modified peptidoglycan (PG) known as the cortex (2, 5). While vegetative cell wall PG consists of alternating *N*-acetylglucosamine (NAG) and *N*-acetylmuramic acid (NAM) sugars, approximately 50% of NAM residues in the cortex are converted to muramic- $\delta$ -lactam, while an additional portion of the NAM side chains is generally cleaved to a single L-alanine (6–11).

The spore form of *Bacillus anthracis* is the etiological agent for all types of anthrax infections: inhalational, gastrointestinal, and cutaneous anthrax, as well as the newest form described, injective anthrax (12, 13). When the spore senses the availability of nutrients, such as when it enters a suitable host, germination is triggered, causing a chain of events that ultimately result in a vegetative cell capable of producing deadly toxins (1, 12). After germinant contact with receptors at the spore inner membrane, the spore releases its large pool of  $Ca^{2+}$ -dipicolinate ( $Ca^{2+}$ -DPA) stored in the core, which becomes partially rehydrated through an influx of water (1). This is followed by cortex depolymerization, allowing the spore core to rehydrate to levels necessary for metabolism to resume (14). Spore germination concomitantly results in the loss of resistance properties; thus, triggering this process at a high efficiency is a potentially attractive approach to spore decontamination.

Cortex degradation during spore germination is accomplished by germination-specific lytic enzymes (GSLEs) already present within the dormant spore (15, 16). These enzymes exhibit specificity for PG containing the muramic- $\delta$ -lactam modification, en-

sure that only the cortex PG is broken down (17–21). *Bacillus anthracis* contains the four GSLEs SleB, CwlJ1, CwlJ2, and SleL; SleB and CwlJ1 are partially redundant enzymes responsible for the majority of cortex hydrolysis (22–25). SleB or CwlJ1 alone is sufficient for cortex hydrolysis, and the absence of both proteins results in spores that are unable to degrade the cortex and complete germination (23–25). Identical roles for SleB and CwlJ have been demonstrated in *Bacillus subtilis* (26–28) and *Bacillus megaterium* (29). CwlJ2, a homolog of CwlJ1, appears to play a minor role at best during *B. anthracis* spore germination, while SleL further breaks down PG fragments first generated by SleB and/or CwlJ1 (20, 22–25).

The arrangement of *sleB* upstream of *ypeB* in an operon is highly conserved across *Bacillus* species and in a few *Clostridiales* species possessing *sleB* (16, 23, 30–32). In the *Bacillus* species examined, SleB and YpeB are expressed from the forespore under the control of  $\sigma^G$  during sporulation and are then translocated across the inner membrane by way of their N-terminal signal sequences (28, 30, 33, 34). Unlike the YpeB signal sequence, which is not predicted to be cleaved, the SleB signal sequence is removed during sporulation, and SleB is present in its mature form within the dormant spore (28, 30, 33, 35). The uncleaved signal sequence likely anchors YpeB to the inner membrane, an idea supported by a study with *B. subtilis* that found both SleB and YpeB associated

Received 28 May 2014 Accepted 7 July 2014

Published ahead of print 14 July 2014

Address correspondence to David L. Popham, dpopham@vt.edu.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JB.01899-14>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.01899-14

with the inner membrane of the spore (26). The same study, as well as work with *Bacillus cereus*, placed SleB and YpeB at a second location near the outer region of the cortex (26, 30). Not only do SleB and YpeB appear to localize within the same regions of the spore, but it has been demonstrated in *B. subtilis* that YpeB is required for SleB activity and for stable incorporation of SleB into the spore (26, 34). Additionally, the relationship appears to be mutual, such that SleB is also needed to stabilize YpeB (36).

Far more is known about the structure and function of SleB than about those of YpeB. At the N terminus of the mature protein, SleB has a PG-binding domain (pfam01471), while the C terminus contains a catalytic domain (pfam07486) (21, 28, 33, 37). As anticipated, the N-terminal domain plays the dominant role in PG binding; however, the C-terminal domain appears to be responsible both for the lytic transglycosylase activity of SleB on cortex PG and for its specificity for muramic- $\delta$ -lactam (21, 37, 38). Mutational analysis and crystal structure determination of the SleB C-terminal domain in *B. anthracis* (39) and *B. cereus* (38) have revealed a conserved catalytic glutamate residue and a unique substrate-binding cleft thought to mediate specificity for cortex PG. Apart from its signal sequence, the only recognizable features of YpeB are the three putative PepSY domains positioned in the C-terminal 60% of the protein (40). These domains are best characterized in the M4 family of metalloproteases, where the PepSY domains serve as intramolecular inhibitors of protease activity until the protease is secreted from the cell (40). Like a large number of other proteins containing PepSY domains, YpeB does not show homology with proteases; thus, the function of the PepSY domains in YpeB and the other proteins constituting this group is unknown (40).

The current study investigates the relationship between SleB and YpeB in *B. anthracis* and demonstrates the contribution of the PepSY domains to this interaction. The data solidify the codependent nature of this relationship and reveal that SleB and YpeB are each rapidly degraded following expression in the absence of the other protein. The C terminus of YpeB containing the PepSY domains is critical for the stability and function of SleB, and certain highly conserved residues within the PepSY domains are important for the stability of both proteins. By understanding GSLEs and regulators of their activity, it may be possible to trigger germination at the stage of cortex hydrolysis, resulting in cells that are readily killed.

## MATERIALS AND METHODS

**Bacterial strains and general growth conditions.** The strains and plasmids used in this work are listed in Table 1. All *B. anthracis* strains used in this study are derived from the Sterne strain 34F2 and were grown at 37°C in brain heart infusion (BHI; Difco) with 5  $\mu$ g/ml erythromycin or 10  $\mu$ g/ml tetracycline where appropriate. *B. anthracis* strains maintaining pBKJ236-derived plasmids extrachromosomally were incubated at 25°C. Following plasmid integration into the chromosome (at 42°C in most instances), these strains were grown at 37°C. *Escherichia coli* strains were grown in LB at 37°C with antibiotics (500  $\mu$ g/ml erythromycin or 100  $\mu$ g/ml ampicillin for strains involved in plasmid propagation; 30  $\mu$ g/ml chloramphenicol and 50  $\mu$ g/ml ampicillin for strains used in protein overexpression).

**Mutant construction.** The sequences of all primers used during plasmid construction are listed in Table S1 in the supplemental material. All plasmids created were verified by DNA sequencing. To create a  $\Delta$ ypeB strain, ypeB and approximately 500 bp flanking each side of the gene were PCR amplified from the *B. anthracis* chromosome. The PCR product was

inserted into vector pBKJ236 (41) by digestion with the restriction enzymes SacII and NotI and ligation of the DNA to create pDPV388. Inverse PCR of the plasmid using primers with BglI restriction sites at the 3' ends resulted in a linear PCR product with the majority of ypeB deleted, leaving only three codons from each end of the gene. Subsequent BglI digestion and ligation of the PCR product formed pDPV392. This plasmid, containing the ypeB deletion, was introduced into *B. anthracis* by using the markerless gene replacement strategy as described previously (41). Gene deletion was verified by PCR amplification and sequencing.

The sleB complementation strain published previously (DPBa57) contains an extrachromosomally maintained pBKJ236 derivative with full-length sleB (pDPV346) (23). In order to ensure efficient plasmid partitioning into the forespore during spore formation and thus achieve more-complete complementation of the deletion phenotype, pDPV346 was integrated into the  $\Delta$ sleB chromosome by shifting the temperature to 42°C. Plasmid integration through homologous recombination just upstream of the sleB deletion in the chromosome was verified with PCR, and this new sleB complementation strain was designated DPBa134. To create a ypeB complementation plasmid that includes the native promoter, PCR was performed using the  $\Delta$ sleB chromosome (23) as the template. The resulting PCR product contained the promoter region for the sleB operon, followed by the  $\Delta$ sleB region, ypeB, and approximately 500 bp downstream of ypeB. The DNA fragment was cloned into pBKJ236 by digestion with NotI and ligation to form pDPV416. This ypeB complementation plasmid was introduced into the  $\Delta$ ypeB strain of *B. anthracis* via conjugation as done in the initial stages of the markerless gene replacement procedure (41). Subsequent plasmid integration within the 500-bp homologous region downstream of the ypeB deletion in the chromosome was achieved by shifting the temperature to 42°C and was verified by PCR.

YpeB truncation and internal deletion mutations were designed by using InterPro (42) to predict the locations of PepSY domains and the Protean application of Lasergene, version 10.0 (DNASTar), to choose truncation/deletion sites on the basis of high residue surface probability and residues not predicted to disrupt protein secondary structure. For the truncation mutations, inverse PCR of the ypeB complementation plasmid was performed using sets of two primers. The tails of the forward primers contained codons encoding a hexahistidine tag followed by a stop codon and a BglII restriction enzyme recognition sequence. The tail of the reverse primer used to create all the truncation mutants also had a BglII site added. The linear PCR products from inverse PCRs were digested with BglII and were ligated in order to recircularize the plasmids, generating pDPV421 through pDPV424. The internal deletion mutations were constructed using overlap extension PCR (43), followed by restriction-free cloning (44) to insert the PCR products into the plasmid encoding full-length YpeB with a C-terminal His<sub>6</sub> tag (pDPV424). The YpeB truncation and internal deletion plasmids were moved into the  $\Delta$ ypeB strain of *B. anthracis* and were integrated into the chromosome as described above. The 500-bp homologous region following ypeB in pDPV416 was removed during inverse PCR to create pDPV421 through -424; thus, these plasmids were inserted into the  $\Delta$ ypeB chromosome through homologous recombination upstream of sleB, as confirmed by PCR.

In order to predict residues of the YpeB PepSY domains that might be important for protein-protein interactions, the PepSY domain sequences from *B. anthracis* YpeB were aligned with a PepSY domain consensus sequence (45), other PepSY domain sequences (40, 45, 46), and the YpeB PepSY domain sequences from other *Bacillus* species by using Clustal W sequence alignment software (47). Based on these alignments, highly conserved residues were identified as targets for mutagenesis. Site-directed mutagenesis by overlap extension PCR (43) was used to create point mutations in ypeB. The resulting PCR products were cloned into the ypeB complementation plasmid pDPV416 by using restriction-free cloning (44). Successful plasmid construction was verified by screening for the gain or loss of a restriction site in ypeB that was designed as part of the

TABLE 1 Bacterial strains and plasmids

Strain or plasmid	Relevant genotype/phenotype <sup>a</sup>	Construction <sup>b</sup>	Source or reference
<b>Strains</b>			
<i>E. coli</i>			
DPVE13	BL21 λ(DE3) pLysS (Cm <sup>r</sup> )		Novagen
DPVE440	pDDPV426 (His <sub>6</sub> -MBP-YpeB <sub>21-446</sub> Amp <sup>r</sup> ) Cm <sup>r</sup>	pDDPV426 → DPVE13	This study
<i>B. anthracis</i>			
Sterne 34F2	pXO1 <sup>+</sup> pXO2 <sup>-</sup>		P. Hanna
DPBa38	Δ <i>sleB</i>		23
DPBa57	Δ <i>sleB</i> :pDPV346 ( <i>sleB</i> <sup>+</sup> Er <sup>r</sup> )		23
DPBa89	Δ <i>ypeB</i>	pDPV392 → 34F2	This study
DPBa113	Δ <i>ypeB</i> ::pDPV416 ( <i>ypeB</i> <sup>+</sup> Er <sup>r</sup> )	pDDPV416 → DPBa89	This study
DPBa124	Δ <i>ypeB</i> ::pDPV421 (YpeB <sub>1-208</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV421 → DPBa89	This study
DPBa125	Δ <i>ypeB</i> ::pDPV422 (YpeB <sub>1-283</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV422 → DPBa89	This study
DPBa126	Δ <i>ypeB</i> ::pDPV423 (YpeB <sub>1-368</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV423 → DPBa89	This study
DPBa127	Δ <i>ypeB</i> ::pDPV424 (YpeB <sub>1-446</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV424 → DPBa89	This study
DPBa134	Δ <i>sleB</i> ::pDPV346 ( <i>sleB</i> <sup>+</sup> Er <sup>r</sup> )	DPBa57 integration	This study
DPBa143	Δ <i>ypeB</i> ::pDPV435 (YpeB <sup>Y410A</sup> Er <sup>r</sup> )	pDPV435 → DPBa89	This study
DPBa144	Δ <i>ypeB</i> ::pDPV436 (YpeB <sup>G430A</sup> Er <sup>r</sup> )	pDPV436 → DPBa89	This study
DPBa148	Δ <i>ypeB</i> ::pDPV432 (YpeB <sup>T377A</sup> Er <sup>r</sup> )	pDPV432 → DPBa89	This study
DPBa149	Δ <i>ypeB</i> ::pDPV433 (YpeB <sup>Y254A</sup> Er <sup>r</sup> )	pDPV433 → DPBa89	This study
DPBa150	Δ <i>ypeB</i> ::pDPV434 (YpeB <sup>Y329A</sup> Er <sup>r</sup> )	pDPV434 → DPBa89	This study
DPBa158	Δ <i>ypeB</i> ::pDPV448 (YpeB <sub>Δ25-203</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV448 → DPBa89	This study
DPBa159	Δ <i>ypeB</i> ::pDPV449 (YpeB <sub>Δ67-203</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV449 → DPBa89	This study
DPBa160	Δ <i>ypeB</i> ::pDPV450 (YpeB <sub>Δ119-203</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV450 → DPBa89	This study
DPBa161	Δ <i>ypeB</i> ::pDPV451 (YpeB <sub>Δ156-203</sub> -His <sub>6</sub> Er <sup>r</sup> )	pDPV451 → DPBa89	This study
<b>Plasmids</b>			
pBKJ236	Er <sup>r</sup> <i>ori</i> (Ts)		42
pBKJ223	Tet <sup>r</sup> Amp <sup>r</sup> P <sub><i>amy</i></sub> -I-SceI		42
pDEST-HisMBP-T	His <sub>6</sub> -MBP; Amp <sup>r</sup> Cm <sup>r</sup>		F. Schubot
pDPV346	<i>sleB</i> <sup>+</sup>	pBKJ236:: <i>sleB</i>	23
pDPV388	<i>ypeB</i> <sup>+</sup>	pBKJ236:: <i>ypeB</i>	This study
pDPV392	Δ <i>ypeB</i>	pBKJ236::Δ <i>ypeB</i>	This study
pDPV416	<i>ypeB</i> <sup>+</sup>	pBKJ236::Δ <i>sleB ypeB</i>	This study
pDPV421	YpeB <sub>1-208</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>1-208</sub> -His <sub>6</sub>	This study
pDPV422	YpeB <sub>1-283</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>1-283</sub> -His <sub>6</sub>	This study
pDPV423	YpeB <sub>1-368</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>1-368</sub> -His <sub>6</sub>	This study
pDPV424	YpeB <sub>1-446</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>1-446</sub> -His <sub>6</sub>	This study
pDPV426	His <sub>6</sub> -MBP-YpeB <sub>21-446</sub>	pDEST-HisMBP-T:: <i>ypeB</i> <sub>21-446</sub>	This study
pDPV432	YpeB <sup>T377A</sup>	pBKJ236::Δ <i>sleB ypeB</i> <sup>T377A</sup>	This study
pDPV433	YpeB <sup>Y254A</sup>	pBKJ236::Δ <i>sleB ypeB</i> <sup>Y254A</sup>	This study
pDPV434	YpeB <sup>Y329A</sup>	pBKJ236::Δ <i>sleB ypeB</i> <sup>Y329A</sup>	This study
pDPV435	YpeB <sup>Y410A</sup>	pBKJ236::Δ <i>sleB ypeB</i> <sup>Y410A</sup>	This study
pDPV436	YpeB <sup>G430A</sup>	pBKJ236::Δ <i>sleB ypeB</i> <sup>G430A</sup>	This study
pDPV448	YpeB <sub>Δ25-203</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>Δ25-203</sub> -His <sub>6</sub>	This study
pDPV449	YpeB <sub>Δ67-203</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>Δ67-203</sub> -His <sub>6</sub>	This study
pDPV450	YpeB <sub>Δ119-203</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>Δ119-203</sub> -His <sub>6</sub>	This study
pDPV451	YpeB <sub>Δ156-203</sub> -His <sub>6</sub>	pBKJ236::Δ <i>sleB ypeB</i> <sub>Δ156-203</sub> -His <sub>6</sub>	This study

<sup>a</sup> Cm<sup>r</sup>, chloramphenicol resistance; Amp<sup>r</sup>, ampicillin resistance; Er<sup>r</sup>, erythromycin resistance; *ori*(Ts), temperature-sensitive origin of replication; Tet<sup>r</sup>, tetracycline resistance.

<sup>b</sup> Strains were constructed by conjugation or electroporation. The plasmid name precedes the arrow, while the recipient strain designation follows the arrow. Single strain designations indicate that the existing plasmid was integrated into the chromosome.

mutagenic primers, followed by DNA sequencing. Plasmids were mobilized into the Δ*ypeB* strain of *B. anthracis* and were integrated into the chromosome downstream of the *ypeB* deletion as described above for the *ypeB* complementation plasmid.

**Spore preparation and decoating.** *B. anthracis* strains were sporulated in modified G broth (48) with appropriate antibiotics at 37°C with aeration. After 3 to 4 days, dormant spores were harvested by centrifugation and were repeatedly washed with deionized water. Any remaining vegetative cells were heat killed at 65°C for 25 min. Spores were further purified with a 50% sodium diatrizoate (Sigma) gradient as described previously (49). Decoated spores were prepared by suspending up to 30 optical den-

sity (OD) units of spores in 1 ml decoating solution (50 mM Tris-HCl [pH 8], 8 M urea, 1% SDS, 50 mM dithiothreitol [DTT]). The spores were incubated at 37°C for 1 h and were then centrifuged at 8,000 × *g* for 1 min to remove the decoating solution. The decoating procedure was repeated, followed by five washes with 1 ml deionized water at room temperature. Untreated (native) and decoated spores were stored in deionized water at 4°C until analysis.

**Spore and sporangium sample preparation for Western blotting.** Following spore purification, 7.5 to 10 OD units of dormant, native spores were pelleted, frozen at -80°C, and lyophilized. Dried spores were mechanically disrupted with 100 mg 0.1 mm glass beads and 20 pulses of 30

s each at 4,200 rpm by using a Wig-L-Bug bead beater. Samples were placed on ice between pulses. Proteins were extracted from the broken material by adding 10  $\mu$ l/OD unit of 1 $\times$  sample loading buffer (62.5 mM Tris-HCl [pH 6.8], 2% SDS, 10% glycerol, 5%  $\beta$ -mercaptoethanol, 0.05% bromophenol blue) and heating to 100°C for 5 min. Extracts were centrifuged at 15,800  $\times$  g for 1 min, and volumes of supernatants derived from equal starting OD units of spores were used for Western blot analysis.

Strains grown in modified G broth with appropriate antibiotics were monitored spectrophotometrically until the OD at 600 nm (OD<sub>600</sub>) indicated that the cells had entered stationary phase. This time point, termed  $t_0$ , coincides with the initiation of sporulation, and  $t_2$  through  $t_6$  designate 2 through 6 h, respectively, past  $t_0$ . At  $t_2$  through  $t_6$ , 10-ml sporangium samples were collected and were centrifuged at 10,000  $\times$  g for 10 min at 4°C. Pellets were resuspended in 1 ml of 8 mM NaPO<sub>4</sub> (pH 7.0) and were centrifuged at 15,800  $\times$  g for 1 min. The resulting pellets were flash frozen in liquid N<sub>2</sub> and were lyophilized. Lyophilized sporangia were broken with glass beads, and proteins were extracted as described above, except that 100  $\mu$ l of 1 $\times$  sample loading buffer was used. Supernatants of protein extracts were used for Western blotting. The average OD<sub>600</sub> measured for a particular strain from  $t_2$  through  $t_6$  was used to adjust sample loading between strains.

**Germination assays.** To assess the rate of spore germination and outgrowth, decoated spores were heat activated at 70°C for 30 min and were quenched on ice for 5 min. Heat-activated spores were diluted to an OD<sub>600</sub> of 0.2 in liquid BHI medium at 37°C to initiate germination, and the change in OD<sub>600</sub> was measured over time. For germination efficiency assays, decoated spores at an OD<sub>600</sub> of 0.2 were heat activated at 70°C for 20 min and were quenched on ice. Heat-activated spores were serially diluted in deionized water, plated onto BHI medium without antibiotics, and incubated at 37°C overnight. Colonies were counted to determine CFU/OD unit values. Unpaired, two-tailed Student *t* tests with unequal variance were used for statistical analyses of germination assays.

**Protein expression and purification.** The *ypeB* gene lacking the first 20 codons was amplified by PCR and was inserted into a modified version of pDEST-HisMBP (50) containing a tobacco etch virus (TEV) cleavage site (pDEST-HisMBP-T) by using restriction-free cloning (44). The resulting plasmid encoded an N-terminally His<sub>6</sub>-tagged maltose binding protein (MBP) and YpeB<sub>21–446</sub>, separated by a TEV cleavage site. Successful plasmid construction was verified by sequencing. The His<sub>6</sub>-MBP-YpeB<sub>21–446</sub> fusion protein was overexpressed in *E. coli* BL21(DE3) [ $\lambda$ (DE3) pLysS Cm<sup>r</sup>] (Novagen) grown at 37°C until the OD<sub>600</sub> reached 1.0, at which point isopropyl- $\beta$ -D-thiogalactopyranoside was added to a final concentration of 1 mM and the culture was incubated at 10°C for an additional 16 h. Cells were harvested by centrifugation at 10,000  $\times$  g for 10 min at 4°C, and the pellet was resuspended in 5 ml/g buffer A (50 mM NaCl, 50 mM Tris-HCl [pH 7.5], 5% glycerol, 30 mM imidazole). Resuspended cells were lysed by sonication for 15 min and were then centrifuged at 117,000  $\times$  g for 1 h at 4°C. The fusion protein in the soluble fraction was purified using a Ni-Sepharose HisTrap HP affinity column (GE Healthcare) equilibrated with buffer A. The protein was eluted with a linear gradient of 30 to 500 mM imidazole in buffer A, and the elution fractions were dialyzed in buffer A. The fusion protein was digested with 1 mg His<sub>6</sub>-tagged TEV(S219V) protease (51) per 7 mg fusion protein at 15°C for 16 h, and the mixture was centrifuged at 117,000  $\times$  g for 20 min at 4°C to remove any precipitated protein. The cleavage of the fusion protein was verified by SDS-PAGE. YpeB<sub>21–446</sub> was purified from His<sub>6</sub>-MBP and His<sub>6</sub>-TEV by using a second Ni-Sepharose HisTrap HP affinity column as described above. SleB was overexpressed and purified as described previously (21).


**Antibody preparation and Western blot analysis.** Polyclonal anti-SleB and anti-YpeB antibodies were raised in rabbits (Open Biosystems) using purified SleB<sub>33–253</sub> (21) or YpeB<sub>21–446</sub>. SleB, YpeB, and derivatives were detected in Western blots on Amersham Hybond-P (polyvinylidene difluoride [PVDF]) membranes (GE Healthcare) by using BM Fluor POD substrate, precipitating (Roche), for colorimetric detection. Primary anti-

SleB and anti-YpeB antibodies were generally used at 1:1,000 and 1:3,000 dilutions, respectively. Horseradish peroxidase (HRP)-conjugated secondary goat anti-rabbit antibodies (Bio-Rad) were used at a 1:200,000 dilution. Western blot quantification was performed using Image Lab software (Bio-Rad). While efforts were made to ensure that the total protein load was identical for each strain, there is the inherent possibility of slight variations stemming from unequal protein extraction between samples and unequal sample loading. To account for experimental error, relative values were normalized using a highly reproducible non-YpeB background band detected in all strains, including the  $\Delta ypeB$  strain. Similarly, SleB band intensities were normalized using a background band present on anti-SleB immunoblots.

## RESULTS

**Germination of  $\Delta ypeB$  mutant spores.** As previously reported by Heffron et al., a homolog of the *B. subtilis ypeB* gene (57% identity; 77% similarity) exists as part of a putative tricistronic operon at locus BAS2561 in the *B. anthracis* chromosome (23). The *ypeB* homolog is the second gene in the operon; it is preceded by *sleB* (BAS2562) and followed by the BAS2560 open reading frame (23). BAS2560 shows homology to the *B. subtilis* genes encoding the lipoproteins YlaJ and YhcN; YlaJ is an uncharacterized spore protein, and YhcN plays an unknown role in spore germination or outgrowth (23, 52, 53). In this study, a deletion of the *ypeB* gene (BAS2561) in the *B. anthracis* chromosome was made using the markerless gene replacement strategy (41). The effects of this *ypeB* deletion, and those of an *sleB* deletion, are not due to polar effects on the expression of BAS2560, because (i) a BAS2560 deletion results in no change in spore germination, and specifically in SleB activity (data not shown); (ii) as shown below, all effects of a *ypeB* deletion can be complemented by the *ypeB* gene alone; and (iii) as shown below and by Heffron et al. (23), all effects of an *sleB* deletion can be complemented by *sleB* alone.

In assays in which purified spores were germinated in BHI medium and the change in the OD<sub>600</sub> measured over time, the germination phenotype of  $\Delta ypeB$  spores was essentially identical to that of  $\Delta sleB$  spores (23). This could be seen in assays using native spores, in which  $\Delta ypeB$  and  $\Delta sleB$  spores produced a shallower germination curve, indicative of a slightly less efficient germination response, and also showed a lower rate of outgrowth (data not shown). The germination defect in  $\Delta ypeB$  and  $\Delta sleB$  spores alike was more pronounced in assays using decoated spores, where the other GSLEs localized to the outer periphery of the spore are removed or inactivated, and SleB alone is responsible for cortex hydrolysis during germination (Fig. 1A). Both  $\Delta ypeB$  and  $\Delta sleB$  decoated spores proceeded through stage I of germination similarly to wild-type spores, where the initial OD decreases as the spores take up water and release their large deposits of Ca<sup>2+</sup>-DPA. Decoated wild-type spores continued to lose nearly half of their initial OD as the cortex PG was degraded during stage II of germination, followed by an increase in OD as germination was completed and outgrowth into vegetative cells began. Conversely, germination was arrested for decoated  $\Delta ypeB$  and  $\Delta sleB$  spores at the stage of cortex hydrolysis;  $\Delta ypeB$  and  $\Delta sleB$  spores never lost more than 34% of their initial OD and did not proceed to outgrowth (Fig. 1A). This germination deficiency is also illustrated in spore plating efficiency assays, where decoated  $\Delta ypeB$  and  $\Delta sleB$  spores showed >10<sup>4</sup>-fold reductions in the ability to germinate and form colonies on a rich medium from that of wild-type spores (Table 2). A wild-type germination phenotype in terms of both germination rate and plating efficiency assays was



**FIG 1** Effects of a variety of *ypeB* mutations on the germination and outgrowth of decoated *B. anthracis* spores. Decoated spores were heat activated and were germinated in BHI medium at 37°C. The data shown are averages of results from three independent spore preparations; error bars are omitted for clarity. (A) Germination and outgrowth are blocked in decoated  $\Delta ypeB$  spores. Wild-type (●),  $\Delta sleB$  (□),  $\Delta ypeB$  (▲), and *ypeB* complementation (*ypeB*<sup>+</sup>) (◆) strains were analyzed. The germination rates of  $\Delta sleB$  and  $\Delta ypeB$  spores were statistically indistinguishable ( $P > 0.23$ ), as were the germination rates of wild-type and *ypeB*<sup>+</sup> spores ( $P > 0.08$ ). Both  $\Delta sleB$  and  $\Delta ypeB$  spores were significantly different ( $P < 0.04$ ) from wild-type and *ypeB*<sup>+</sup> spores during stage II of germination, from 45 to 95 min. (B) Truncations removing YpeB PepsY domains block decoated-spore germination.  $\Delta ypeB$  (▲), YpeB<sub>1–208</sub>-His<sub>6</sub> (△), YpeB<sub>1–283</sub>-His<sub>6</sub> (+), YpeB<sub>1–368</sub>-His<sub>6</sub> (□), YpeB<sub>1–446</sub>-His<sub>6</sub> (—), and *ypeB*<sup>+</sup> (◆) strains were analyzed. The germination rates of *ypeB*<sup>+</sup> and

**TABLE 2** Decoated-spore plating efficiencies for *sleB* and *ypeB* mutants


Strain	Genotype/phenotype	CFU/OD <sub>600</sub> value <sup>a</sup>
34F2	Wild type	$8.4 \times 10^7$
DPBa38	$\Delta sleB$	$8.3 \times 10^{3*}$
DPBa89	$\Delta ypeB$	$<5.8 \times 10^{3*}$
DPBa113	<i>ypeB</i> <sup>+</sup>	$8.7 \times 10^7$
DPBa124	YpeB <sub>1–208</sub> -His <sub>6</sub>	$9.7 \times 10^{3*}$
DPBa125	YpeB <sub>1–283</sub> -His <sub>6</sub>	$<3.5 \times 10^{3*}$
DPBa126	YpeB <sub>1–368</sub> -His <sub>6</sub>	$<4.5 \times 10^{3*}$
DPBa127	YpeB <sub>1–446</sub> -His <sub>6</sub>	$1.1 \times 10^8$

<sup>a</sup> Values are averages for three independent spore preparations. Asterisks indicate values significantly different from those of wild-type and *ypeB*<sup>+</sup> spores ( $P < 0.006$  and  $< 0.02$ , respectively).

achieved using spores from a *ypeB* complementation strain (DPBa113 [*ypeB*<sup>+</sup>]), in which *ypeB*, provided on a plasmid downstream of its native promoter, was integrated into the  $\Delta ypeB$  chromosome. High-performance liquid chromatography (HPLC) analyses of mucopeptides collected during spore germination revealed the complete absence of the SleB-specific lytic transglycosylase products identified by Heffron et al. (23) in both  $\Delta sleB$  and  $\Delta ypeB$  spores, while these peaks were restored during the germination of *ypeB*<sup>+</sup> spores (data not shown). In agreement with the results observed from a *ypeB* deletion in *B. subtilis* (34, 36), these results indicate that YpeB is necessary for the lytic activity of SleB on spore cortex PG in *B. anthracis*.

**Stability of SleB and YpeB in spores and sporangia.** Not only is YpeB needed for SleB activity, but Western blotting of  $\Delta ypeB$  spore extracts demonstrated that YpeB is also required for the incorporation of SleB into the spore (Fig. 2). SleB was essentially not detected in  $\Delta ypeB$  spore extracts, and this could be complemented. As in *B. subtilis* (36), this relationship is mutual: SleB was also needed for the stable incorporation of YpeB into the spore (Fig. 2). Immunoblots of  $\Delta sleB$  spore extracts revealed significantly diminished levels of YpeB when SleB was absent, which could also be complemented. Miniscule levels of SleB and YpeB detected in Western blots of  $\Delta ypeB$  and  $\Delta sleB$  spores, respectively, could be attributed either to expression of the proteins during spore formation followed by rapid proteolysis in the absence of the stabilizing partner or to escape of the proteins from the developing spore. The former proposal is supported by Western blot analyses of whole-sporangium samples taken throughout sporu-


YpeB<sub>1–446</sub>-His<sub>6</sub> spores were not statistically different ( $P > 0.08$ ); likewise, the germination rates of  $\Delta ypeB$  and YpeB<sub>1–283</sub>-His<sub>6</sub> spores were statistically indistinguishable ( $P > 0.49$ ). YpeB<sub>1–208</sub>-His<sub>6</sub> and YpeB<sub>1–368</sub>-His<sub>6</sub> spores did not differ ( $P > 0.05$ ) from  $\Delta ypeB$  spores during germination, except at 55 min and from 45 to 55 min, respectively ( $P < 0.05$ ). The germination rates of YpeB<sub>1–208</sub>-His<sub>6</sub>, YpeB<sub>1–283</sub>-His<sub>6</sub>, and YpeB<sub>1–368</sub>-His<sub>6</sub> spores were not statistically different ( $P > 0.11$ ) from 4 min onward. (C) Point mutations in YpeB PepsY domain conserved residues slow decoated-spore germination.  $\Delta ypeB$  (▲), YpeB<sup>Y254A</sup> (■), YpeB<sup>Y329A</sup> (×), YpeB<sup>T377A</sup> (○), YpeB<sup>Y410A</sup> (●), YpeB<sup>G430A</sup> (◇), and *ypeB*<sup>+</sup> (◆) strains were analyzed. Statistical analysis of *ypeB* point mutants is complex due to the greater variability of germination rates of decoated-spore preparations, as well as the fact that spores of some strains are initiating outgrowth while others are still germinating. During stage II of germination, between 45 and 95 min, YpeB<sup>Y329A</sup> and YpeB<sup>T377A</sup> spores were not significantly different from *ypeB*<sup>+</sup> spores ( $P > 0.38$ ). The germination rates of YpeB<sup>Y254A</sup> spores from 80 to 90 min, YpeB<sup>Y410A</sup> spores from 45 to 55 min and 70 to 95 min, and YpeB<sup>G430A</sup> spores at 50 min and from 75 to 95 min were significantly different from that of *ypeB*<sup>+</sup> spores ( $P < 0.05$ ).



**FIG 2** *B. anthracis* SleB and YpeB are codependent for incorporation into dormant spores. Dormant wild-type (WT),  $\Delta sleB$ , *sleB* complementation (*sleB*<sup>+</sup>),  $\Delta ypeB$ , and *ypeB* complementation (*ypeB*<sup>+</sup>) spores were mechanically disrupted, and proteins were extracted with sample loading buffer for Western blot analysis. Blots were probed with anti-YpeB (top) and anti-SleB (bottom) antibodies. The predicted molecular sizes of mature SleB lacking its signal sequence and YpeB are 24.1 and 50.0 kDa, respectively. The positions of molecular size marker proteins (not shown) are indicated on the left.

lation (Fig. 3). In wild-type and *ypeB*<sup>+</sup> strains, significant amounts of SleB and YpeB were detected beginning 3 h after the initiation of sporulation ( $t_3$ ) (Fig. 3A and B). This is consistent with the findings of previous  $\beta$ -galactosidase activity assays performed in *B. anthracis*, which show that *sleB* transcription reaches its peak near  $t_{3.5}$  (23). During spore formation in a  $\Delta sleB$  or  $\Delta ypeB$  strain, some YpeB or SleB, respectively, could be seen at  $t_3$  and  $t_4$  but was not strongly detected thereafter (Fig. 3A and B). The expression of YpeB at  $t_3$  and  $t_4$  coincided with the production of large quantities of two YpeB-specific degradation products in  $\Delta sleB$  sporangia (and to a lesser degree in wild-type and *ypeB*<sup>+</sup> sporangia [data not shown]), indicating that the protein was degraded immediately following its expression (Fig. 3C). Specific SleB degradation products were not observed during the sporulation of any of the strains tested (data not shown). The protease(s) responsible for the degradation of SleB and YpeB during spore formation is unknown. Elimination of SpoIVB, which is known to be active in the forespore intermembrane space in *B. subtilis* (54, 55), had no effect on the stability of SleB in the absence of YpeB (see Fig. S1 in the supplemental material). Elimination of BAS5314 (HtrC), which in *B. subtilis* is expressed in the forespore and possesses a signal sequence/membrane anchor (56), had no effect on the stability of SleB and YpeB in the absence of the partner protein (C. B. Bernhards, Y. Chen, H. Toutkoushian, and D. L. Popham, unpublished data).


**Analyses of YpeB PepSY domain truncation mutants.** InterPro (42) predicted the presence of three PepSY domains in *B. anthracis* YpeB from residues 218 to 282, 291 to 358, and 375 to 436. To assess the individual contributions of the three putative PepSY domains located at the C terminus of YpeB, strains expressing truncated forms of YpeB with a C-terminal His<sub>6</sub> tag were constructed. Plasmids expressing YpeB from its native promoter were integrated into the  $\Delta ypeB$  chromosome; the resulting strains expressed YpeB with no PepSY domains (YpeB<sub>1-208</sub>-His<sub>6</sub>), one PepSY domain (YpeB<sub>1-283</sub>-His<sub>6</sub>), two PepSY domains (YpeB<sub>1-368</sub>-His<sub>6</sub>), or all three PepSY domains (full-length YpeB) (YpeB<sub>1-446</sub>-His<sub>6</sub>). In decoated-spore germination rate assays in liquid BHI medium and in decoated-spore plating efficiency assays, spores containing YpeB with no, one, or two PepSY domains were similar to  $\Delta ypeB$  spores (Fig. 1B and Table 2). Like  $\Delta ypeB$  spores, these spores containing truncated versions of YpeB were arrested during germination and showed a  $>10^3$ -fold reduction in plating efficiency in the case of the YpeB<sub>1-208</sub>-His<sub>6</sub> mutant, but generally a  $>10^4$ -fold decrease. The reason for the similarity was revealed in



**FIG 3** *B. anthracis* SleB and YpeB are each degraded during spore formation in the absence of the partner protein. Wild-type (WT),  $\Delta sleB$ ,  $\Delta ypeB$ , and *ypeB* complementation (*ypeB*<sup>+</sup>) strains were grown in modified G broth at 37°C, and samples collected during sporulation were used for Western blot analysis. The designations  $t_2$  through  $t_6$  represent the numbers of hours since the initiation of sporulation. (A) Sporangia probed with anti-SleB antibodies. (B) Sporangia probed with anti-YpeB antibodies. (C) Sporangia from the  $\Delta sleB$  strain probed with anti-YpeB antibodies. The positions of molecular size marker proteins (not shown), in kilodaltons, are indicated on the left.

Western blots of dormant spore extracts, which show that YpeB is unstable in many of the truncated forms (Fig. 4A). In fact, YpeB and SleB were both stable only in strains possessing all three YpeB PepSY domains. In spores with highly unstable YpeB (YpeB<sub>1-208</sub>-His<sub>6</sub> and YpeB<sub>1-368</sub>-His<sub>6</sub>), SleB was also unstable, as anticipated (Fig. 4A). Interestingly, although YpeB with only the first PepSY domain (YpeB<sub>1-283</sub>-His<sub>6</sub>) appeared somewhat stable in the anti-YpeB Western blot, SleB was unstable (Fig. 4A), and these spores had germination phenotypes like those of  $\Delta ypeB$  spores (Fig. 1B and Table 2). This suggests that the C-terminal region of YpeB, more specifically a region beyond the first PepSY domain, is critical for SleB stability and thus for spore germination.


**Analysis of YpeB internal deletion mutants.** In order to determine what other areas of YpeB may be necessary for YpeB function in stabilizing SleB and SleB activity during germination, internal deletions of different lengths were made between the N-terminal signal sequence and the C-terminal PepSY domains. Deletion endpoints were chosen based on software predictions of transitions between secondary structural elements. The strains created expressed YpeB without residues 25 to 203 (YpeB $\Delta_{25-203}$ -His<sub>6</sub>), 67 to 203 (YpeB $\Delta_{67-203}$ -His<sub>6</sub>), 119 to 203 (YpeB $\Delta_{119-203}$ -His<sub>6</sub>), or 156 to 203 (YpeB $\Delta_{156-203}$ -His<sub>6</sub>). Immunoblots of dormant spore extracts revealed that YpeB $\Delta_{25-203}$ -His<sub>6</sub> and YpeB $\Delta_{67-203}$ -His<sub>6</sub> are at least moderately stable, while YpeB $\Delta_{119-203}$ -His<sub>6</sub> and YpeB $\Delta_{156-203}$ -His<sub>6</sub> are not; however, SleB was not stably incorporated into spores of any of these internal deletion strains (Fig. 4B). Even YpeB $\Delta_{25-203}$ -His<sub>6</sub>, which appears to accumulate in dormant spores to levels similar to those of full-length YpeB



**FIG 4** Effects of *ypeB* truncation and internal deletion mutations on the stability of YpeB and SleB. Dormant spores were mechanically disrupted, and proteins were extracted with sample loading buffer for Western blot analysis. Samples were probed with anti-YpeB (top) and anti-SleB antibodies (bottom) antibodies. Right arrows indicate YpeB derivatives. The positions of molecular mass marker proteins (not shown) are indicated on the left. (A) Truncations removing YpeB PepSY domains destabilize both YpeB and SleB. YpeB<sub>1-208</sub>-His<sub>6</sub>, YpeB<sub>1-283</sub>-His<sub>6</sub>, YpeB<sub>1-368</sub>-His<sub>6</sub>, and YpeB<sub>1-446</sub>-His<sub>6</sub> have predicted molecular sizes of 24.3, 32.5, 42.1, and 50.9 kDa, respectively. (B) Internal deletions in the YpeB N-terminal domain destabilize SleB. Lanes intervening between the 5th and 6th lanes were removed for clarity. YpeB<sub>Δ25-203</sub>-His<sub>6</sub>, YpeB<sub>Δ67-203</sub>-His<sub>6</sub>, YpeB<sub>Δ119-203</sub>-His<sub>6</sub>, YpeB<sub>Δ156-203</sub>-His<sub>6</sub>, and YpeB<sub>1-446</sub>-His<sub>6</sub> have predicted molecular sizes of 30.6, 35.5, 41.2, 45.6, and 50.9 kDa, respectively.

(YpeB<sub>1-446</sub>-His<sub>6</sub>), failed to stabilize SleB. Thus, the N-terminal half of YpeB, likely a region between residues 67 and 203, must also play a role in stabilizing SleB.

**Analyses of YpeB PepSY domain point mutants.** As an alternative approach to studying the PepSY domains while attempting to maintain YpeB stability, site-directed mutagenesis was performed on selected highly conserved residues within the three PepSY domains. Y254 within the first PepSY domain, Y329 in the



**FIG 5** Protein stability and plating efficiency in YpeB point mutant spores. Dried, dormant *ΔypeB*, YpeB<sup>Y254A</sup>, YpeB<sup>Y329A</sup>, YpeB<sup>T377A</sup>, YpeB<sup>Y410A</sup>, YpeB<sup>G430A</sup>, and *ypeB* complementation (*ypeB*<sup>+</sup>) spores were broken, and extracted proteins were analyzed by Western blotting with anti-SleB (α-SleB) or anti-YpeB (α-YpeB) antibodies. Band intensities were quantified and are depicted as the levels of YpeB (open bars) and SleB (shaded bars) present in spores, relative to the native protein levels found in *ypeB*<sup>+</sup> spores, which were set at 100%. Spores of the same strains were heat activated, serially diluted, and plated on BHI medium. Following incubation, colonies were counted, and the percentages of native CFU/OD values (striped bars) were determined by comparison to the CFU/OD value for *ypeB*<sup>+</sup> spores, which was set at 100%. Plating efficiency and Western blot quantification data shown are averages from three independent spore preparations; error bars represent the standard deviations. The Western blots below the graph show representative results from one of the replicates.

second PepSY domain, and T377, Y410, and G430, all in the third and last PepSY domain, were selected to be individually changed to alanine. Spores with YpeB<sup>Y329A</sup> or YpeB<sup>T377A</sup> showed no significant decrease either in the germination rate in liquid BHI medium or in plating efficiency from *ypeB*<sup>+</sup> spores (Fig. 1C and 5). Spores with YpeB<sup>Y254A</sup> or YpeB<sup>G430A</sup> showed a delay in germination, reaching their lowest ODs 40 min and 50 min later than *ypeB*<sup>+</sup> spores, respectively (Fig. 1C). While YpeB<sup>Y254A</sup> and *ypeB*<sup>+</sup> spores each lost 59 to 60% of their initial ODs, YpeB<sup>G430A</sup> spores had a shallower germination curve, losing 51% of their initial OD (Fig. 1C). Both YpeB<sup>Y254A</sup> and YpeB<sup>G430A</sup> spores showed slight but significant ( $P < 0.007$ ) decreases in plating efficiency, but <10-fold reductions in the CFU/OD values were seen (Fig. 5). The most dramatic effect on spore germination was seen for YpeB<sup>Y410A</sup> spores, which showed a significant delay in germination and a shallower germination curve (Fig. 1C). YpeB<sup>Y410A</sup> spores lost only 50% of their initial OD, and this point was reached 105 min after *ypeB*<sup>+</sup> spores lost 59% of their initial OD (Fig. 1C). YpeB<sup>Y410A</sup> spores also showed a significant decrease ( $P < 0.004$ ) in their ability to form colonies on BHI plates, with a nearly 100-fold reduction (Fig. 5).

The trend seen for the YpeB point mutants in germination rate and plating efficiency assays was mirrored by the levels of YpeB and SleB seen in Western blots of dormant spore extracts (Fig. 5). Relative YpeB band intensities for each strain were computed by comparison to the YpeB band from the *ypeB*<sup>+</sup> strain. Similarly, SleB band intensities relative to the SleB band

detected in *ypeB*<sup>+</sup> spores were determined. The strains that appeared most similar to *ypeB*<sup>+</sup> spores in germination rate and plating efficiency assays (the *YpeB*<sup>Y329A</sup> and *YpeB*<sup>T377A</sup> strains) contained 101 and 99% of the *YpeB* protein levels and 102 and 93% of the *SleB* protein levels detected in *ypeB*<sup>+</sup> spores, respectively. Likewise, *YpeB*<sup>Y254A</sup> and *YpeB*<sup>G430A</sup> spores, which showed minor defects during the other assays tested, possessed 69 and 75% of native *YpeB* levels and 70 and 65% of native *SleB* levels, respectively. Immunoblots of dormant *YpeB*<sup>G430A</sup> spore extracts also revealed the presence of two stable *YpeB*-specific degradation products (not shown) similar to those observed during the sporulation of a  $\Delta$ *sleB* strain (Fig. 3C). Among the point mutants, the lowest levels of the two proteins were seen for *YpeB*<sup>Y410A</sup>, with *YpeB* levels equivalent to 60%, and *SleB* levels equivalent to 40%, of those found in *ypeB*<sup>+</sup> spores. This corresponds well with the more severe deficit in germination phenotypes demonstrated by *YpeB*<sup>Y410A</sup> spores. While the correlation between *YpeB* and *SleB* band intensities was not perfect, the trend was quite clear: *YpeB* abundance, *SleB* abundance, and the germination rate decreased in concert in all strains tested.

## DISCUSSION

This study investigated the relationship between *SleB* and *YpeB* in *B. anthracis* for the first time. In agreement with findings for *B. subtilis* (34), the germination of decoated  $\Delta$ *sleB* and  $\Delta$ *ypeB* spores of *B. anthracis* was blocked after the initial steps, these spores exhibited a >10<sup>4</sup>-fold decrease in colony-forming ability, and lytic transglycosylase products were absent in germinating spores, confirming that *YpeB* is needed for the lytic activity of *SleB* on cortex PG *in vivo*. The current work clearly demonstrates that not only is *YpeB* required for *SleB* incorporation into the dormant spore, as in *B. subtilis* (26, 36), but *YpeB* has a reciprocal requirement for *SleB*. This mutual dependency was also discovered recently using a *B. subtilis* strain that expressed *YpeB* ectopically in a *cwlJ sleB ypeB* mutant background (36). In that strain, *YpeB* was detected in Western blots of spore inner membrane fractions only if *SleB* was also expressed (36). It should be noted that since only the inner membrane fraction was analyzed in those immunoblots, it is possible that the absence of *SleB* resulted in the mislocalization of *YpeB*. The Western blotting procedure in the present study utilized entire spore extracts derived from strains in which *sleB* or *ypeB* was deleted or expressed from the native operon. Thus, these results provide more-conclusive evidence that *YpeB* is indeed absent in  $\Delta$ *sleB* spores. Furthermore, it is demonstrated that in the absence of the stabilizing partner, *SleB* and *YpeB* are expressed and degraded during early sporulation, rather than simply failing to localize within the developing spore.

Li et al. suggested that both the N- and C-terminal domains of *YpeB* (*YpeB*<sup>N</sup> and *YpeB*<sup>C</sup>) are necessary for *SleB* lytic activity during spore germination, since a *B. subtilis* strain deficient in *cwlJ*, *sleB*, and *ypeB* and ectopically expressing an individual *ypeB* domain (encoding *YpeB*<sup>N</sup> or *YpeB*<sup>C</sup>) with full-length *sleB* (encoding *SleB*<sup>FL</sup>) could not complement the germination defect in these spores (36). While these results were hampered by an inability to determine if *YpeB*<sup>N</sup> or *YpeB*<sup>C</sup> was stably incorporated into the spore, Li et al. did demonstrate that the C terminus of *YpeB* fused to its signal sequence (memseg-*YpeB*<sup>C</sup>) was stable yet could not complement the defect when expressed with *SleB*<sup>FL</sup>, indicating that the N terminus of *YpeB* is needed (36). The results obtained

in the current study when stretches of residues between the *YpeB* signal sequence and the *PepSY* domains were deleted support the necessity of a region within the N-terminal portion of *YpeB*. This critical region occurs within the area spanning residues 67 to 203, since *YpeB* $\Delta$ <sub>67-203</sub>-His<sub>6</sub> was moderately stable in the dormant spore yet could not stabilize *SleB*. Deletions of regions corresponding to the *PepSY* domains at the C terminus of *YpeB* also add to what is known about the portions of *YpeB* needed for *SleB* stability. While many of the truncated *YpeB* variants were unstable, *YpeB*<sub>1-283</sub>-His<sub>6</sub>, which contained the first predicted *PepSY* domain, was moderately stable, yet *SleB* was not detected in Western blots, and the germination phenotypes resembled those of  $\Delta$ *ypeB* spores. These results clearly show that a region beyond the first *YpeB* *PepSY* domain is also essential for *SleB* stabilization. Taken together with the findings by Li et al. (36), they demonstrate that both the N terminus and the *PepSY* domain-containing C terminus of *YpeB* are required for stable incorporation of *SleB* into the spore.

In studies of the effects of *ypeB* point mutations on *SleB* stability and function, the codependency between the two proteins observed in  $\Delta$ *sleB* and  $\Delta$ *ypeB* spores was reinforced. There was a strong correlation between the *SleB* and *YpeB* protein levels within the spores of individual strains, and protein abundance matched well with observed germination phenotypes. This intrinsic codependency becomes a complicating factor in determining which *YpeB* residues are actually important for the interaction with *SleB* and which are just important for proper *YpeB* folding. In the former scenario, alteration of a residue mediating the interaction between *YpeB* and *SleB* would destabilize the interaction, resulting in decreased levels of both proteins. In the latter scenario, a mutation affecting *YpeB* protein folding would likely decrease *YpeB* stability, subsequently decreasing the stability of *SleB*. Residue Y410 of *YpeB* clearly plays a large role in *YpeB* and *SleB* stability, and this residue is one of the most highly conserved across *PepSY* domains (40, 45). While the Y410A substitution had a large effect on spore germination, the tyrosine residues at similar positions in the other two *PepSY* domains did not prove to be as important. The Y254A substitution in the first *PepSY* domain resulted in a milder spore germination defect, and the Y329A replacement in the second *PepSY* domain produced spores that germinated equally as well as *ypeB*<sup>+</sup> spores. Additionally, of the three mutations made in the region encoding the third *PepSY* domain of *YpeB*, two resulted in observable phenotypes, indicating the relative importance of this domain. The three *ypeB* point mutations that produced phenotypic changes (Y254A, Y410A, G430A) affect residues that are 100% conserved in an alignment of *YpeB* proteins from 13 *Bacillus* species, while those mutations that produced no phenotypic changes affect less-conserved residues. This could suggest that the three *PepSY* domains are unequal, with the third being the most crucial but the first also playing a role. Without an alternative means of stabilizing *YpeB* and *SleB* *in vivo*, e.g., via inhibition of proteolysis within the developing sporangium, further *in vivo* genetic study of the relationship between these proteins is difficult.

Analysis of *YpeB* point mutants also indicated that reduced amounts of *SleB* in spores not only resulted in reduced germination rates but also hindered the ability of spores to complete outgrowth and form colonies. A similar phenomenon was evident in *B. subtilis cwlJ sleB* mutants expressing *sleB* ectopically in addition to a low level of *ypeB* expression through apparent readthrough of



an *sleB* insertion mutation (36). While the amounts of SleB and YpeB present in the spores of these *B. subtilis* strains were not explicitly quantified, immunoblots clearly showed reductions in SleB and YpeB levels. The same spores exhibited a >50% decrease in colony-forming ability (36). One might expect that even limited quantities of SleB within a given spore would allow the spore to germinate, albeit at a lower rate. This finding may suggest heterogeneity within a population of dormant spores from an individual mutant strain with respect to the absolute SleB levels present or the portion of SleB capable of being activated.

The simplest explanation for how SleB and YpeB are able to stabilize each other is that the two proteins interact physically, yet no such direct interaction has been demonstrated. Li et al. did not observe an interaction using *in vitro* affinity pulldown assays involving various forms of *B. cereus* and *B. megaterium* SleB and YpeB purified from *E. coli* and *Lactococcus lactis* (36). Similarly, in conjunction with the work presented here, affinity pulldown assays were performed using His<sub>6</sub>-MBP-YpeB<sub>1-446</sub> or His<sub>6</sub>-MBP-YpeB<sub>21-446</sub> and untagged SleB<sub>33-253</sub>, and reciprocally, His<sub>6</sub>-MBP-SleB<sub>33-253</sub> and untagged YpeB<sub>21-446</sub>. No interaction between SleB and YpeB was observed (data not shown). A logical explanation for the lack of *in vitro* interaction is that if SleB and YpeB do indeed physically interact, the interaction likely occurs during spore formation as the proteins are coexpressed and translocated in their unfolded forms across the inner forespore membrane via the Sec pathway. Thus, the proteins may need to be cotranslocated and/or cofolded, or, as postulated by Li et al., a membrane might be necessary for the interaction to occur (36). Protein coexpression is likely not the only variable missing from the equation, since assays performed using cell extracts from *E. coli* in which SleB and YpeB were coexpressed still failed to show an interaction between the two proteins (36), although it is feasible that the N-terminal affinity tags on both proteins impeded an interaction.

Since YpeB does not have any homologs of known function, one of the major clues for determining the precise nature of the relationship between SleB and YpeB may lie in the PepSY domains at the C terminus of YpeB, which this study has demonstrated are important. While these domains have an unknown function in many proteins, they have been found in the propeptide region of M4 class proteases and function as inhibitors of protease activity (40). Another potential model, where SleB and YpeB do not bind and YpeB instead forms an inhibitory interaction with the protease(s) responsible for SleB degradation in the absence of YpeB, fails to explain how SleB stabilizes YpeB. It is possible that PepSY domains are capable of inhibiting a broader range of enzymatic activities, since they belong to a superfamily of bacterial protein domains sharing a  $\beta$ -lactamase inhibitor protein (BLIP)-like fold (45). Members of this superfamily have been associated with various inhibitory roles, including inhibition of  $\beta$ -lactamases, and have also been proposed to mediate protein-protein interactions (45). The potential for PepSY domains to inhibit enzymatic activities outside their known niche makes YpeB a prime candidate for a way in which SleB is held inactive in the dormant spore. It has been demonstrated in *B. subtilis* that YpeB is cleaved during spore germination (26); such cleavage could release SleB from the inhibitory interaction with YpeB, thereby activating SleB for cortex degradation. Recent *in vitro* experiments using various forms of exogenous SleB and YpeB from *B. cereus* have demonstrated inhibition of SleB activity in the presence of YpeB (36). While the *in vitro* experiments performed by Li et al. (36) support an inhibitory

role for YpeB, more work is needed to further elucidate the relationship between SleB and YpeB. Such studies should aim to clarify the function of the PepSY domains and the mechanism by which YpeB, and specifically YpeB<sup>N</sup>, is able to inhibit SleB activity. YpeB processing during germination should also be investigated to determine if this event is required for SleB activation or is merely part of the process whereby unneeded spore proteins are broken down and recycled for new protein synthesis during outgrowth.

## ACKNOWLEDGMENTS

The research reported in this publication was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award AI060726.

The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## REFERENCES

1. Setlow P. 2003. Spore germination. *Curr. Opin. Microbiol.* 6:550–556. <http://dx.doi.org/10.1016/j.mib.2003.10.001>.
2. Setlow P. 2006. Spores of *Bacillus subtilis*: their resistance to and killing by radiation, heat and chemicals. *J. Appl. Microbiol.* 101:514–525. <http://dx.doi.org/10.1111/j.1365-2672.2005.02736.x>.
3. Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. 2000. Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. *Microbiol. Mol. Biol. Rev.* 64:548–572. <http://dx.doi.org/10.1128/MMBR.64.3.548-572.2000>.
4. Gould GW. 2006. History of science—spores. *J. Appl. Microbiol.* 101:507–513. <http://dx.doi.org/10.1111/j.1365-2672.2006.02888.x>.
5. Gerhardt P, Marquis RE. 1989. Spore thermoresistance mechanisms, p 43–63. In Smith I, Slepceky RA, Setlow P (ed), Regulation of prokaryotic development. American Society for Microbiology, Washington, DC.
6. Warth AD, Strominger JL. 1969. Structure of the peptidoglycan of bacterial spores: occurrence of the lactam of muramic acid. *Proc. Natl. Acad. Sci. U. S. A.* 64:528–535. <http://dx.doi.org/10.1073/pnas.64.2.528>.
7. Atrih A, Zöllner P, Allmaier G, Foster SJ. 1996. Structural analysis of *Bacillus subtilis* 168 endospore peptidoglycan and its role during differentiation. *J. Bacteriol.* 178:6173–6183.
8. Popham DL, Helin J, Costello CE, Setlow P. 1996. Analysis of the peptidoglycan structure of *Bacillus subtilis* endospores. *J. Bacteriol.* 178:6451–6458.
9. Atrih A, Bacher G, Körner R, Allmaier G, Foster SJ. 1999. Structural analysis of *Bacillus megaterium* KM spore peptidoglycan and its dynamics during germination. *Microbiology* 145:1033–1041. <http://dx.doi.org/10.1099/13500872-145-5-1033>.
10. Dowd MM, Orsburn B, Popham DL. 2008. Cortex peptidoglycan lytic activity in germinating *Bacillus anthracis* spores. *J. Bacteriol.* 190:4541–4548. <http://dx.doi.org/10.1128/JB.183.16.4894-4899.2001>.
11. Atrih A, Foster SJ. 2001. Analysis of the role of bacterial endospore cortex structure in resistance properties and demonstration of its conservation amongst species. *J. Appl. Microbiol.* 91:364–372. <http://dx.doi.org/10.1046/j.1365-2672.2001.01394.x>.
12. Mock M, Fouet A. 2001. Anthrax. *Annu. Rev. Microbiol.* 55:647–671. <http://dx.doi.org/10.1146/annurev.micro.55.1.647>.
13. Sweeney DA, Hicks CW, Cui X, Li Y, Eichacker PQ. 2011. Anthrax infection. *Am. J. Respir. Crit. Care Med.* 184:1333–1341. <http://dx.doi.org/10.1164/rccm.201102-0209CI>.
14. Setlow B, Melly E, Setlow P. 2001. Properties of spores of *Bacillus subtilis* blocked at an intermediate stage in spore germination. *J. Bacteriol.* 183:4894–4899. <http://dx.doi.org/10.1128/JB.183.16.4894-4899.2001>.
15. Makino S, Moriyama R. 2002. Hydrolysis of cortex peptidoglycan during bacterial spore germination. *Med. Sci. Monit.* 8:RA119–RA127.
16. Popham DL, Bernhards CB. Spore peptidoglycan. In Driks A, Eichenberger P (ed), The bacterial spore: from molecules to systems, in press. ASM Press, Washington, DC.
17. Popham DL, Helin J, Costello CE, Setlow P. 1996. Muramic lactam in peptidoglycan of *Bacillus subtilis* spores is required for spore outgrowth but not for spore dehydration or heat resistance. *Proc. Natl. Acad. Sci. U. S. A.* 93:15405–15410. <http://dx.doi.org/10.1073/pnas.93.26.15405>.

18. Chen Y, Fukuoka S, Makino S. 2000. A novel spore peptidoglycan hydrolase of *Bacillus cereus*: biochemical characterization and nucleotide sequence of the corresponding gene, *sleL*. *J. Bacteriol.* 182:1499–1506. <http://dx.doi.org/10.1128/JB.182.6.1499-1506.2000>.
19. Chen Y, Miyata S, Makino S, Moriyama R. 1997. Molecular characterization of a germination-specific muramidase from *Clostridium perfringens* S40 spores and nucleotide sequence of the corresponding gene. *J. Bacteriol.* 179:3181–3187.
20. Lambert EA, Sherry N, Popham DL. 2012. *In vitro* and *in vivo* analyses of the *Bacillus anthracis* spore cortex lytic protein SleL. *Microbiology* 158: 1359–1368. <http://dx.doi.org/10.1099/mic.0.056630-0>.
21. Heffron JD, Sherry N, Popham DL. 2011. *In vitro* studies of peptidoglycan binding and hydrolysis by the *Bacillus anthracis* germination-specific lytic enzyme SleB. *J. Bacteriol.* 193:125–131. <http://dx.doi.org/10.1128/JB.00869-10>.
22. Lambert EA, Popham DL. 2008. The *Bacillus anthracis* SleL (YaaH) protein is an *N*-acetylglucosaminidase involved in spore cortex depolymerization. *J. Bacteriol.* 190:7601–7607. <http://dx.doi.org/10.1128/JB.01054-08>.
23. Heffron JD, Orsburn B, Popham DL. 2009. Roles of germination-specific lytic enzymes CwlJ and SleB in *Bacillus anthracis*. *J. Bacteriol.* 191:2237–2247. <http://dx.doi.org/10.1128/JB.01598-08>.
24. Heffron JD, Lambert EA, Sherry N, Popham DL. 2010. Contributions of four cortex lytic enzymes to germination of *Bacillus anthracis* spores. *J. Bacteriol.* 192:763–770. <http://dx.doi.org/10.1128/JB.01380-09>.
25. Giebel JD, Carr KA, Anderson EC, Hanna PC. 2009. The germination-specific lytic enzymes SleB, CwlJ1, and CwlJ2 each contribute to *Bacillus anthracis* spore germination and virulence. *J. Bacteriol.* 191:5569–5576. <http://dx.doi.org/10.1128/JB.00408-09>.
26. Chirakkal H, O'Rourke M, Atrih A, Foster SJ, Moir A. 2002. Analysis of spore cortex lytic enzymes and related proteins in *Bacillus subtilis* endospore germination. *Microbiology* 148:2383–2392.
27. Paidhungat M, Ragkousi K, Setlow P. 2001. Genetic requirements for induction of germination of spores of *Bacillus subtilis* by Ca<sup>2+</sup>-dipicolinate. *J. Bacteriol.* 183:4886–4893. <http://dx.doi.org/10.1128/JB.183.16.4886-4893.2001>.
28. Moriyama R, Hattori A, Miyata S, Kudoh S, Makino S. 1996. A gene (*sleB*) encoding a spore cortex-lytic enzyme from *Bacillus subtilis* and response of the enzyme to L-alanine-mediated germination. *J. Bacteriol.* 178:6059–6063.
29. Setlow B, Peng L, Loshon CA, Li YQ, Christie G, Setlow P. 2009. Characterization of the germination of *Bacillus megaterium* spores lacking enzymes that degrade the spore cortex. *J. Appl. Microbiol.* 107:318–328. <http://dx.doi.org/10.1111/j.1365-2672.2009.04210.x>.
30. Moriyama R, Fukuoka H, Miyata S, Kudoh S, Hattori A, Kozuka S, Yasuda Y, Tochikubo K, Makino S. 1999. Expression of a germination-specific amidase, SleB, of bacilli in the forespore compartment of sporulating cells and its localization on the exterior side of the cortex in dormant spores. *J. Bacteriol.* 181:2373–2378.
31. Christie G, Üstok FI, Lu Q, Packman LC, Lowe CR. 2010. Mutational analysis of *Bacillus megaterium* QM B1551 cortex-lytic enzymes. *J. Bacteriol.* 192:5378–5389. <http://dx.doi.org/10.1128/JB.00830-10>.
32. Popham DL, Heffron JD, Lambert EA. 2012. Degradation of spore peptidoglycan during germination, p 121–142. In Abel-Santos E (ed), *Bacterial spores: current research and applications*. Caister Academic Press, Norwich, United Kingdom.
33. Moriyama R, Kudoh S, Miyata S, Nonobe S, Hattori A, Makino S. 1996. A germination-specific spore cortex-lytic enzyme from *Bacillus cereus* spores: cloning and sequencing of the gene and molecular characterization of the enzyme. *J. Bacteriol.* 178:5330–5332.
34. Boland FM, Atrih A, Chirakkal H, Foster SJ, Moir A. 2000. Complete spore-cortex hydrolysis during germination of *Bacillus subtilis* 168 requires SleB and YpeB. *Microbiology* 146:57–64.
35. Hu K, Yang H, Liu G, Tan H. 2007. Cloning and identification of a gene encoding spore cortex-lytic enzyme in *Bacillus thuringiensis*. *Curr. Microbiol.* 54:292–295. <http://dx.doi.org/10.1007/s00284-006-0430-x>.
36. Li Y, Butzin XY, Davis A, Setlow B, Korza G, Üstok FI, Christie G, Setlow P, Hao B. 2013. Activity and regulation of various forms of CwlJ, SleB, and YpeB proteins in degrading cortex peptidoglycan of spores of *Bacillus* species *in vitro* and during spore germination. *J. Bacteriol.* 195: 2530–2540. <http://dx.doi.org/10.1128/JB.00259-13>.
37. Masayama A, Fukuoka H, Kato S, Yoshimura T, Moriyama M, Moriyama R. 2006. Subcellular localization of a germination-specific cortex-lytic enzyme, SleB, of bacilli during sporulation. *Genes Genet. Syst.* 81:163–169. <http://dx.doi.org/10.1266/ggs.81.163>.
38. Li Y, Jin K, Setlow B, Setlow P, Hao B. 2012. Crystal structure of the catalytic domain of the *Bacillus cereus* SleB protein, important in cortex peptidoglycan degradation during spore germination. *J. Bacteriol.* 194: 4537–4545. <http://dx.doi.org/10.1128/JB.00877-12>.
39. Jing X, Robinson HR, Heffron JD, Popham DL, Schubot FD. 2012. The catalytic domain of the germination-specific lytic transglycosylase SleB from *Bacillus anthracis* displays a unique active site topology. *Proteins* 80:2469–2475. <http://dx.doi.org/10.1002/prot.24140>.
40. Yeats C, Rawlings ND, Bateman A. 2004. The PepSY domain: a regulator of peptidase activity in the microbial environment? *Trends Biochem. Sci.* 29:169–172. <http://dx.doi.org/10.1016/j.tibs.2004.02.004>.
41. Janes BK, Stibitz S. 2006. Routine markerless gene replacement in *Bacillus anthracis*. *Infect. Immun.* 74:1949–1953. <http://dx.doi.org/10.1128/IAI.74.3.1949-1953.2006>.
42. Hunter S, Jones P, Mitchell A, Apweiler R, Attwood TK, Bateman A, Bernard T, Binns D, Bork P, Burge S, de Castro E, Coggill P, Corbett M, Das U, Daugherty L, Duquenne L, Finn RD, Fraser M, Gough J, Haft D, Hulo N, Kahn D, Kelly E, Letunic I, Lonsdale D, Lopez R, Madera M, Maslen J, McAnulla C, McDowall J, McMenamin C, Mi H, Mutowo-Muelleren P, Mulder N, Natale D, Orengo C, Pesset S, Punta M, Quinn AF, Rivoire C, Sangrador-Vegas A, Selengut JD, Sigrist CJ, Scheremetjew M, Tate J, Thimmajananathan M, Thomas PD, Wu CH, Yeats C, Yong SY. 2012. InterPro in 2011: new developments in the family and domain prediction database. *Nucleic Acids Res.* 40:D306–D312. <http://dx.doi.org/10.1093/nar/gkr948>.
43. Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. 1989. Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* 77:51–59. [http://dx.doi.org/10.1016/0378-1119\(89\)90358-2](http://dx.doi.org/10.1016/0378-1119(89)90358-2).
44. van den Ent F, Löwe J. 2006. RF cloning: a restriction-free method for inserting target genes into plasmids. *J. Biochem. Biophys. Methods* 67:67–74. <http://dx.doi.org/10.1016/j.jbbm.2005.12.008>.
45. Das D, Finn RD, Carlton D, Miller MD, Abdubek P, Astakhova T, Axelrod HL, Bakolitsa C, Chen C, Chiu HJ, Chiu M, Clayton T, Deller MC, Duan L, Ellrott K, Ernst D, Farr CL, Feuerhelm J, Grant JC, Grzechnik A, Han GW, Jaroszewski L, Jin KK, Klock HE, Knuth MW, Kozbial P, Krishna SS, Kumar A, Marciano D, McMullan D, Morse AT, Nigoghossian E, Nopakun A, Okach L, Puckett C, Reyes R, Rife CL, Sefcovic N, Tien HJ, Trame CB, van den Bedem H, Weekes D, Wooten T, Xu Q, Hodgson KO, Wooley J, Elslinger MA, Deacon AM, Godzik A, Lesley SA, Wilson IA. 2010. The structure of BVU2987 from *Bacteroides vulgatus* reveals a superfamily of bacterial periplasmic proteins with possible inhibitory function. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.* 66:1265–1273. <http://dx.doi.org/10.1107/S1744309109046788>.
46. Gao X, Wang J, Yu DQ, Bian F, Xie BB, Chen XL, Zhou BC, Lai LH, Wang ZX, Wu JW, Zhang YZ. 2010. Structural basis for the autoprocessing of zinc metalloproteases in the thermolysin family. *Proc. Natl. Acad. Sci. U. S. A.* 107:17569–17574. <http://dx.doi.org/10.1073/pnas.1005681107>.
47. Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignments through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22:4673–4680. <http://dx.doi.org/10.1093/nar/22.22.4673>.
48. Kim HU, Goepfert JM. 1974. A sporulation medium for *Bacillus anthracis*. *J. Appl. Bacteriol.* 37:265–267. <http://dx.doi.org/10.1111/j.1365-2672.1974.tb00438.x>.
49. Nicholson WL, Setlow P. 1990. Sporulation, germination, and outgrowth, p 391–450. In Harwood CR, Cutting SM (ed), *Molecular biological methods for Bacillus*. John Wiley and Sons Ltd., Chichester, England.
50. Austin BP, Nallamsetty S, Waugh DS. 2009. Hexahistidine-tagged maltose-binding protein as a fusion partner for the production of soluble recombinant proteins in *Escherichia coli*. *Methods Mol. Biol.* 498:157–172. [http://dx.doi.org/10.1007/978-1-59745-196-3\\_11](http://dx.doi.org/10.1007/978-1-59745-196-3_11).
51. Kaput RB, Tozser J, Fox JD, Anderson DE, Cherry S, Copeland TD, Waugh DS. 2001. Tobacco etch virus protease: mechanism of autolysis and rational design of stable mutants with wild-type catalytic proficiency. *Protein Eng.* 14:993–1000. <http://dx.doi.org/10.1093/protein/14.12.993>.
52. Kuwana R, Kasahara Y, Fujibayashi M, Takamatsu H, Ogasawara N, Watabe K. 2002. Proteomics characterization of novel spore proteins of *Bacillus subtilis*. *Microbiology* 148:3971–3982.

53. Bagyan I, Noback M, Bron S, Paidhungat M, Setlow P. 1998. Characterization of *yhcN*, a new forespore-specific gene of *Bacillus subtilis*. *Gene* 212:179–188. [http://dx.doi.org/10.1016/S0378-1119\(98\)00172-3](http://dx.doi.org/10.1016/S0378-1119(98)00172-3).
54. Cutting S, Driks A, Schmidt R, Kunkel B, Losick R. 1991. Forespore-specific transcription of a gene in the signal transduction pathway that governs Pro- $\sigma^K$  processing in *Bacillus subtilis*. *Genes Dev.* 5:456–466. <http://dx.doi.org/10.1101/gad.5.3.456>.
55. Wakeley PR, Dorazi R, Hoa NT, Bowyer JR, Cutting SM. 2000. Proteolysis of SpoIVB is a critical determinant in signalling of pro- $\sigma^K$  processing in *Bacillus subtilis*. *Mol. Microbiol.* 36:1336–1348. <http://dx.doi.org/10.1046/j.1365-2958.2000.01946.x>.
56. Fabret C, Hoch JA. 1998. A two-component signal transduction system essential for growth of *Bacillus subtilis*: implications for anti-infective therapy. *J. Bacteriol.* 180:6375–6383.