

Commentary

Open Access

Keeping up with protein folding

François Baneyx*

Address: Department of Chemical Engineering, University of Washington, Box 351750, Seattle, WA 98195, USA

Email: François Baneyx* - baneyx@u.washington.edu

* Corresponding author

Published: 14 May 2004

Received: 22 April 2004

Microbial Cell Factories 2004, **3**:6

Accepted: 14 May 2004

This article is available from: <http://www.microbialcellfactories.com/content/3/1/6>

© 2004 Baneyx; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Protein folding and misfolding in the cellular environment has emerged as a major research theme over the past fifteen years. The implications are broad, ranging from the production of recombinant proteins in a soluble and bio-active form to the understanding of the etiology of neurodegenerative diseases. The basic principle enunciated by Anfinsen – that all the information required for a polypeptide to reach a proper conformation is contained within its amino acid sequence [1] – remains unchallenged. However, it is now well established that many proteins require the assistance of folding modulators to adopt a native structure as they emerge from ribosomes, regain their original conformation following exposure to stress, or traffic to their ultimate destination [2].

One of the most extensively characterized group of folding modulators are molecular chaperones, a ubiquitous class of proteins that help other polypeptides reach a proper conformation or cellular location without affecting folding rates or becoming part of the final structure. Molecular chaperones function by transiently binding hydrophobic stretches of amino acids that should normally be buried within the core of their substrates but have become solvent-exposed due to improper *de novo* folding or imposition of cellular stress (e.g., temperature increase). As a result of the formation of chaperone-substrate complexes, interactive species are shielded from each other and from the solvent and "on-pathway" folding is facilitated.

Three functionally distinct sets of chaperones cooperate in the conformational quality control of the proteome. At the heart of chaperone networks are "folding chaperones" which use conformational changes fueled by ATP hydrolysis to promote the net refolding/unfolding of bound sub-

strates. Archetypal folding chaperones are the DnaK-DnaJ-GrpE (Hsp70-Hsp40) and GroEL-GroES (Hsp60-Hsp10) systems of *E. coli*. "Holding chaperones" (e.g., small heat shock proteins) passively stabilize partially folded species until folding chaperones become available for their refolding. In doing so, they alleviate titration of folding chaperones by unfolding intermediates. Finally, "disaggregating chaperones" such as *E. coli* ClpB and *S. cerevisiae* Hsp104 employ ATP-driven conformational changes to solubilize protein aggregates and transfer partially folded species to folding chaperones for subsequent refolding.

Foldases embody the second group of folding modulators. These enzymes accelerate rate-limiting steps along the folding pathway and include peptidyl-prolyl *cis/trans* isomerases (PPIases), which catalyze the *trans* to *cis* isomerization of peptidyl-prolyl bonds, and thiol-disulfide oxidoreductases, which promote the formation and isomerization of disulfide bridges (e.g., the Dsb proteins of the *E. coli* periplasm). The final participants in the protein quality control machinery are proteases which degrade irreversibly damaged polypeptides that cannot be rescued by the above pathways.

Nevertheless, nature loves to frustrate reductionist efforts and it has recently become apparent that functional lines are often blurred. For instance, PPIases and oxidoreductases can combine chaperone and foldase functions, certain protease components remodel polypeptide substrates and can thus be viewed as chaperones, some proteases have the ability to switch from hydrolase to chaperone function, and there is evidence that certain proteins classified as chaperones may exhibit protease activity (see for instance [3-10]).

To keep readers abreast of the fast moving field of protein folding, *Microbial Cell Factories* has commissioned a series of reviews by experts in the topic. The first of these already appeared in January [11]. In "Unscrambling an egg: protein disaggregation by AAA+ proteins", Weibezahn, Bukau and Mogk discuss the discovery, functional characterization and mechanism of action of Hsp104/ClpB, two orthologous ATPases that have the unique ability to fragment protein aggregates. The remodeling activity of other AAA+ proteins (e.g., ClpA, ClpX and ClpC) which are best known more for their role in proteolysis is also examined. The authors close their thorough review by presenting enticing evidence for the roles of AAA+ proteins in the propagation of neurodegenerative diseases.

In the second paper of the series [12], Miot and Betton review the mechanisms of protein quality control in the periplasmic space of *E. coli*. After a brief discussion of the biogenesis of proteins destined for export from the bacterial cytoplasm, the authors describe the players involved in the periplasmic folding-degradation decision and how cells sense and respond to extracytoplasmic stress. The second half of the review provides a historical perspective on how the maltose binding protein has served as a useful model to understand the mechanisms of conformational quality control in the periplasm. With forthcoming reviews on disulfide bridge formation and in vitro refolding, it is a great time to turn to *Microbial Cell Factories* to keep up with protein folding.

References

- Anfinsen CB: **Principles that govern the folding of protein chains.** *Science* 1973, **181**:223-230.
- Hartl FU, Hayer-Hartl M: **Molecular chaperones in the cytosol: from nascent chain to folded protein.** *Science* 2002, **295**:1852-1858.
- Chen J, Song JL, Zhang S, Wang Y, Cui DF, Wang CC: **Chaperone activity of DsbC.** *J Biol Chem* 1999, **274**:19601-19605.
- Genevaux P, Keppel F, Schwager F, Langendijk-Genevaux PS, Hartl FU, Georgopoulos C: **In vivo analysis of the overlapping functions of DnaK and trigger factor.** *EMBO Rep* 2004, **5**:195-200.
- Hoskins JR, Singh SK, Maurizi MR, Wickner S: **Protein binding and unfolding by the chaperone ClpA and degradation by the protease ClpAP.** *Proc Natl Acad Sci USA* 2000, **97**:8892-8897.
- Kramer G, Patzelt H, Rauch T, Kurz TA, Vorderwulbecke S, Bukau B, Deuerling E: **Trigger factor peptidyl-prolyl cis/trans isomerase activity is not essential for the folding of cytosolic proteins in Escherichia coli.** *J Biol Chem* 2004, **279**:14165-14170.
- Lee SJ, Kim SJ, Kim IK, Ko J, Jeong CS, Kim GH, Park C, Kang SO, Suh PG, Lee HS, et al.: **Crystal structures of human DJ-1 and Escherichia coli Hsp31 that share an evolutionarily conserved domain.** *J Biol Chem* 2003, **278**:44552-44559.
- Portaro FC, Hayashi MA, De Arauz LJ, Palma MS, Assakura MT, Silva CL, de Camargo AC: **The Mycobacterium leprae hsp65 displays proteolytic activity. Mutagenesis studies indicate that the M. leprae hsp65 proteolytic activity is catalytically related to the HslVU protease.** *Biochemistry* 2002, **41**:7400-7406.
- Quigley PM, Korotkov K, Baneyx F, Hol WGJ: **The 1.6-Å crystal structure of the class of chaperones represented by Escherichia coli Hsp31 reveals a putative catalytic triad.** *Proc Natl Acad Sci USA* 2003, **100**:3137-3142.
- Spieß C, Beil A, Ehrmann M: **A temperature-dependent switch from chaperone to protease in a widely conserved heat shock protein.** *Cell* 1999, **97**:339-347.
- Weibezahn J, Bukau B, Mogk A: **Unscrambling an egg: protein disaggregation by AAA+ proteins.** *Microb Cell Fact* 2004, **3**:1.
- Miot M, Betton JM: **Protein quality control in the bacterial periplasm.** *Microb Cell Fact* 2004, **3**:4.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

