

Published in final edited form as:

Neurosci Biobehav Rev. 2014 October ; 46 Pt 3: 345–351. doi:10.1016/j.neubiorev.2014.03.025.

Lifelong opioidergic vulnerability through early life separation: A recent extension of the false suffocation alarm theory of panic disorder

Maurice Preter, MD¹ and Donald F. Klein, MD DSc²

¹Department of Psychiatry, College of Physicians and Surgeons, Columbia University, New York, NY, USA

²Phyllis Green and Randolph Cowen Institute for Pediatric Neuroscience, Department of Child and Adolescent Psychiatry, New York University Langone Medical Center 550 1st Ave, New York, NY 10016, USA. donaldk737@aol.com

We briefly reference previous material on Klein's suffocation false alarm theory (SFA) of panic disorder (Klein, 1993) and its amplification in 2008 (Preter and Klein, 2008). We then discuss a recent finding showing a fundamental difference in opioidergic reactivity to a naloxone challenge in psychiatrically and medically healthy adults, based on the presence or absence of childhood separation/parental loss.

In 1993, Klein published the original SFA theory of panic disorder (Klein, 1993), attempting to integrate the multiplicity of apparently unrelated clinical and laboratory observations. We posited “that a physiologic misinterpretation by a suffocation monitor misfires an evolved suffocation alarm system. This produces sudden respiratory distress followed swiftly by a brief hyperventilation, panic, and the urge to flee. Carbon dioxide hypersensitivity is seen as due to the deranged suffocation alarm monitor. If other indicators of potential suffocation provoke panic, this theoretical extension is supported.” In the original paper, we tested “the theory by examining Ondine's curse as the physiologic and pharmacologic converse of panic disorder, splitting panic in terms of symptomatology and challenge studies, reevaluating the role of hyperventilation, and reinterpreting the contagiousness of sighing and yawning, as well as mass hysteria.” (Klein, 1993) Original SFA focused on relating the observed lactate and carbon dioxide hypersensitivity in panic disorder to a putative dysfunction in a hypothesized suffocation alarm. At the time, the underlying pathophysiology that might connect the apparent disparate “phenomena of panic during relaxation and sleep, late luteal phase dysphoric disorder, pregnancy, childbirth, pulmonary disease, separation anxiety, and treatment”, was unknown.

© 2014 Elsevier Ltd. All rights reserved.

¹ Corresponding Author. Mailing Address: 1160 Fifth Avenue, New York, NY 10029, USA. mp2285@columbia.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Over the intervening decades, much data evolved linking Separation Anxiety, Panic, and respiratory dysfunction. This suggested a potential missing link: “Klein and Fink (1962) posited a developmental pathophysiological link between [clinical levels of] separation anxiety and PD and subsequent agoraphobia, since 50% of hospitalized agoraphobics reported severe early separation anxiety that often prevented school attendance. Further, panic, in this group, was frequently precipitated by bereavement, or separation [...] (Preter and Klein, 2008). This was also noted by others, e.g., Faravelli and Pallanti (1989), Kaunonen et al. (2000), Milrod et al. 2004.

We noted that “[p]atients highly comorbid for multiple anxiety disorders are particularly likely to recall childhood SAD (Lipsitz et al., 1994)”, and that “[c]laims that separation anxiety equivalently antecedes other anxious states (Van der Molen et al., 1989) may be due to diagnostically ambiguous limited symptom attacks and the unreliability of the questionnaire method”. We concluded that “in the only controlled, long-term, direct, blind, clinical interview follow-up of separation- anxious, school-phobic children, the only significant finding was an increased PD rate” (Preter and Klein, 2008) .

Adding further support, Battaglia (1995) showed that “[s]eparation anxiety correlates with increased familial loading and early onset of PD. More recently (2012), as part of their series of brilliant twin studies, Robertson-Nay et al. demonstrated that, “[childhood] separation anxiety disorder and adult onset panic attacks share a common genetic diathesis”.

Panic and comorbid conditions

There has been a regained awareness of the relevance of panic states to other clinical contexts, as Freud suggested in his pioneering statement (Freud, 1895). In the US, nearly half of panic patients are initially seen in the medical emergency room of a hospital. They may undergo extensive diagnostic medical procedures, such as MRI scans of the brain for headaches, or coronary angiograms for chest pain. Cardiovascular symptoms, particularly pseudo-anginal chest pain resembling a heart attack are the most common symptoms in these PD patients’ experience. Accordingly, 25% or more of outpatients seen by a cardiologist have a current diagnosis of PD (Ballenger, 1998). Since primary complaints are of distress, rather than anxiety, they have been termed “non-fearful” panics. This seemingly oxymoronic term nevertheless indicates that fearfulness is not essential to panic disorder (Beitman et al., 1990).

Migraine and other chronic headaches are highly comorbid with panic disorder (Breslau et al., 2001, Hamelsky and Lipton, 2006). Having PD increases the risk of migraine four-fold, and vice versa. This bidirectionality suggests that the migraine- panic association is unlikely to be merely coincidental and that shared environmental and familial factors are involved (Breslau et al., 2001). In a longitudinal study, separation and parental loss early in life increased the risk of both headaches and psychiatric morbidity, mainly anxiety and depression, in adulthood (Fearon and Hotopf, 2001).

Panic disorder is also comorbid with other somatic pain syndromes (Johnson et al., 2006). In a cross-sectional survey of 1,219 female veterans studying the prevalence and frequency of mastalgia, women reporting frequent mastalgia were much more likely to have comorbid

panic disorder (OR 7.1), but also post-traumatic stress disorder, mood disorders, and other somatic pain syndromes, such as fibromyalgia, chronic pelvic pain or irritable bowel syndrome.

Although panic attacks as they occur in panic disorder often prominently feature air hunger, other panic subtypes have prominent vestibular symptoms and unspecific dizziness/lightheadedness. True vertigo was historically recognized as a common presentation of panic disorder (Benedikt 1870; Frommberger et al. 1994). The current rigid distinction between psychiatry and neurology (Staab, 2006) interferes with proper assessment (Preter and Bursztajn, 2009).

In Mandarin Chinese, *tou yun* refers to the disabling sensation of a constant state of movement of oneself or one's surroundings. This dizziness (note that *tou yun* also describes vertigo), is probably the most common expression of panic disorder in Chinese patients (Park and Hinton, 2002), so by sheer numbers, this may well be the most prevalent panic subtype worldwide.

Taken together, the various comorbidities of panic disorder and untreated sequelae massively impact people's quality of life (Sareen et al., 2006).

One prominent characteristic of the panic attack and subthreshold panic-related anxiety is respiratory dysregulation and chaotic breathing. This can be experimentally reproduced in adult panic sufferers, but also in children with separation anxiety disorder (Pine et al. 2000; 2005; Preter and Klein, 2008).

Air hunger and chronic sighing outside of the acute attack are hallmarks of panic that rarely occurs under acute, external-threat initiated fear (Klein, 1993; Preter and Klein, 1998). Increasing hypercapnia is a more salient indicator of potential suffocation than hypoxia, but hypoxia also serves this alarm function. "Beck et al. (1999; 2000) showed that panic patients respond with increased panic symptoms not only to CO₂ inhalation, but also to normocapnic hypoxia, as predicted by SFA." (Preter and Klein, 2008). Unsurprisingly, numerous studies found that panic disorder and lung disease commonly occur together (Goodwin and Eaton 2003; Goodwin et al., 2004; Katon et al., 2004; Nascimento et al., 2002; Klein, 2001; Roy-Byrne et al., 2006; Valenca et al., 2006; Wingate and Hansen-Flaschen, 1997; Yellowlees and Kalucy, 1990; Yellowlees et al., 1988). More specifically, we wrote, "early lung disease, including asthma and COPD may predispose to PD (Craske et al., 2001; Goodwin and Eaton, 2003; Hasler et al., 2005; Karajgi et al., 1990; Perna et al., 1997; Verburg et al., 1995), or present solely with panic symptoms (Edlund et al., 1991; Sietsema et al., 1987). Asthma and PD are both characterized by acute episodes, salient respiratory symptoms and anxiety with avoidance of situations related to acute attacks (Klein, 1993; Yellowlees and Kalucy, 1990). There is a significantly higher (6.5 to 24%) prevalence of PD in asthmatics (Goodwin et al., 2005; Shavitt et al., 1992; Yellowlees et al., 1988) than the 1-3% reported in the general population (Kessler et al., 2006; Weissman, 1988). Perna et al. (1997) found a significantly higher prevalence of PD, sporadic panic attacks, and social phobia in asthmatics than the general population. In 90% of asthmatics with PD, asthma appeared first. Panic symptomatology during the asthmatic attack predicted longer hospitalizations in

asthmatic patients (Baron et al., 1986; Brooks et al., 1989; Jurenc, 1988).” (Preter and Klein, 2008).

The recent amplification of SFA centers on the observation that separation anxiety and suffocation sensitivity are both under endogenous opioidergic control. We amplified the SFA theory by suggesting that PD may be due to an episodic functional endogenous opioid deficit (Preter and Klein, 1998). The following is a necessarily brief explanation.

The endogenous opioid system was discovered in the early 1970's. Electrical stimulation the periaqueductal gray (Mayer et al., 1971) produced analgesia that was reversed by naloxone, suggesting an endogenous opioid system. Opioid molecules are among the oldest evolved signaling substances., functioning in many physiological processes e.g., pain perception, respiration (Stefano et al., 1996). Dyspnea is modulated by central and peripheral opioid levels in both rodents and humans (Santiago and Edelman, 1985).

In mice, exposure to intermittent, severe hypoxia prolonged survival during subsequent lethal suffocation (Mayfield and D'Alecy, 1992). This effect was blocked by naloxone, implying that endogenous opioids increase adaptability to hypoxic environments. Opioid receptors, including ‘non-conventional’ ones, can be found throughout the respiratory tract. Nebulized morphine is an outstanding treatment for chronic dyspnea (Baydur, 2004; Bruera et al., 2005; Zebraski et al., 2000; Mahler, 2011).

In our 2008 paper, we summarized data from developmental psychobiology and neuroanatomy that point to a possible link between separation and the endogenous opioid system, as follows:

“Following birth, mammalian infants cannot survive independently. Survival requires reliable distress signaling mechanisms to elicit parental care and retrieval. Distress vocalizations (DVs) are a primitive form of audio-vocal communication (Panksepp, 1998). A common neuroanatomy subserving DVs may be shared by all mammals, although substantial functional variations depend on the ontogenetic niche. The latter (West and King, 1987) signifies the ecological and social legacies (“the inherited environment”) in which a given set of genes develops. For instance, isolated altricial (developmentally immature) infants do not emit DVs compared to other species, since it is not likely they will stray from the nest (Panksepp et al., 1992).

Immature human infants practically never get lost for their first six months. Despite frequent maternal absence, separation anxiety in humans develops only after their motor system matures. Young rats are not specifically attached to their mother, i.e. any mother will do as heater or feeder. Only once mobile do they socially bond, but their responses do not compare with the vigor seen in other species. Rats also differ from other species, including primates, dogs and chicks in their greater DV suppression by benzodiazepines (Kalin et al., 1987; Panksepp et al., 1992; Scott, 1974). Since benzodiazepines differentially alleviate anticipatory anxiety, social isolation in young rodents, as compared to many other mammals, may activate anxiety mechanisms other than separation distress. Thus, Panksepp emphasizes that when using cross-species analogies, it is important to keep in mind that the type and

degree of social separation distress depends on ecological and developmental parameters (Panksepp et al., 1992).

The developmental phase of separation anxiety serves as a biologic leash for the increasingly mobile, but helpless infant who continually checks for the mother's presence, becomes acutely distressed on discovering her absence, and immediately attempts to elicit retrieval by crying. In humans, separation anxiety usually wanes around age four when the now verbally skilled child can successfully elicit care even from non-relatives.

Using electrical brain stimulation (ESB), DVs have been elicited in many species from homologous areas, including the midbrain, dorsomedial thalamus, ventral septum, preoptic area, and the bed nucleus striae terminalis (BNST). In some higher species, one can obtain separation calls by stimulating the central amygdala and dorsomedial hypothalamus. All these sites have high opioid receptor densities and figure heavily in sexual and maternal behaviors (Panksepp, 1998). Cortically, electrical stimulation of the rostral cingulate gyrus in monkeys consistently elicits distress calls (Jürgens and Ploog, 1970; Ploog, 1981). The cingulate cortex, found exclusively in mammals, is particularly well developed in humans and contains high densities of opioid receptors (Wise and Herkenham, 1982).

Naloxone-blockable opioid agonists reduce isolation-induced distress vocalizations (DVs) across mammalian species (Hofer and Shair, 1978; Kalin et al., 1988; Kehoe and Blass, 1986; Panksepp et al., 1978). In beagles, imipramine, the classic anti-panic agent, and morphine were the only psychotropic drug that yielded specific DV reduction at nonsedating doses (Scott, 1974; Panksepp et al., 1978).

Naloxone given to guinea pigs and young chicks (Panksepp et al., 1978) increased baseline vocalizations (by 600%), but only when the animals were in a group, since isolates already emitted maximum DVs.

Kalin et al. (1988) studied opioid modulation of separation distress in primates, showing morphine (0.1 mg/kg) significantly decreased separation distress vocalizations without changes in autonomic and hormonal activation. Naloxone (0.1 mg/kg) blocked this effect. Sympathetic blockade using the $\alpha(2)$ agonist, clonidine, and the β adrenergic antagonist, propranolol, had no specific effect on separation-induced "coos" in infant rhesus monkeys (Kalin and Shelton, 1988). (Preter and Klein, 2008)

Testing the panic-suffocation-false alarm-endogenous opioid connection

"Panic Disorder is unique among psychiatric disorders in that its salient component, the panic attack, can be reliably incited in laboratory settings by specific chemical challenges as well as having challenges specifically blocked by anti-panic agents, e.g. imipramine. We can experimentally turn panic on and off, producing trenchant causally related data rather than inferences from naturalistic data." (Preter and Klein, 2008). Specifically, sodium lactate infusions and CO₂ inhalation regularly produce panic attacks in patients with panic disorder (Liebowitz et al., 1984b), Gorman et al., 1984, Papp et al. 1993). However, while normal controls or patients with other anxiety disorders rarely show such reactivity (i.e., progress to a full-blown panic attack)(Klein 1993), higher concentrations of inhaled CO₂ are highly

aversive and can produce respiratory panic symptomatology in a dose-dependent fashion (Griez et al., 2007; Esquivel et al., 2010; Leibold et al., 2013).

Both spontaneous and lactate induced panic attacks in panic patients produce air hunger and marked, objective increases in tidal volume (V_t) (Goetz et al., 1993; Martinez et al., 1996). Since sodium lactate infusion causes a metabolic alkalosis, a compensatory decrease in ventilation would be expected. This would homeostatically buffer blood pH, by increasing CO₂ retention. However, the converse actually occurs indicating a specific lactate stimulating effect on respiration.

The usual response of healthy control subjects to a sodium lactate infusion is a minor, but definite increase in V_t (Liebowitz and others 1985). The lesser tidal volume response in lactate challenged normal subjects may be due to buffering by their intact endogenous opioid system.

An open pilot study showed that naloxone infusion (ranging from an initial 0.5mg/kg to a maximum of 2mg/kg) followed by lactate (N+L), caused significant tidal volume increments similar to those observed during clinical and lactate induced panic attacks in 8 of 12 normal subjects, supporting the hypothesis that opioidergic deficiency might be necessary for lactate to produce a marked increase in tidal volume in normal subjects (Sinha et al., 2007).

Based on these initial findings, and cognizant that previous experiments using smaller doses of both intravenous and oral opioid blocking agents had shown little results (Liebowitz et al., 1984a, Esquivel et al., 2009) we decided to conduct a controlled, randomized experimental study to investigate whether high-dose naloxone, an intravenous opioid receptor antagonist, could change the regularly resistant normal controls to become more sensitive to intravenous lactate as a respiratory stimulus to tidal volume increment. Study design and statistical analysis are detailed elsewhere (Preter et al., 2011), but in addition to the usual standard recruitment procedure for healthy research subjects, “eligible volunteers were further interviewed about potentially significant individual and family antecedents and comorbidities of panic, such as near-suffocation, pulmonary disease, and migraine headache. Recent and childhood loss and separation events (parental divorce or death, childhood abuse) were specifically reviewed.” (Preter et al., 2011). Results showed that “normal subjects, usually relatively insensitive to the tidal volume effects of lactate infusion, [...] given opioid antagonist pretreatment, developed tidal volume and respiratory rate increments resembling those occurring in both spontaneous clinical panic attacks and in panic patients who panic during lactate infusions (Liebowitz et al., 1994; Gorman et al., 1984; Papp et al., 1993). The hypothesis that a functioning endogenous opioid system buffers normal subjects from the behavioral and physiological effects of lactate is consonant with these results. [...]

The most interesting aspect of this study is that “for the first time the prolonged physiological effects of actual separations and losses during childhood, i.e. parental death, parental separation or divorce, on the endogenous opioid system of healthy adults have been objectively, experimentally shown. Presence or absence of childhood parental loss (CPL) antecedents determined the response to the naloxone-lactate probe.” (Preter et al., 2011).

A history of CPL decreased the naloxone+lactate effect implying that there was an antecedent decrement in opioidergic activity, so that the naloxone had nothing to block. We however note these normal subjects had been carefully screened for lack of psychiatric or medical disorder. It was the subjects that had not suffered such separation events that showed the expectable naloxone blockade and tidal volume increment.

“The import of these findings is that analyses attempting to relate CPL to other baseline variables may well fail since CPL impact may be specific to challenges to the endogenous opioid system.” (Preter et al., 2011). Also it implies that separation-induced, baseline opioidergic deficiency, while it may not be sufficient to induce overt disease, confers lifetime vulnerability even in healthy adults. Whether it is longitudinally, or cross-sectionally relevant to somatic pain syndromes such as migraine, and to opiate abuse ought to be determined.

Again, we “emphasize that these CPL effects were apparent in a ‘normal’ sample. In a society where divorce rates approach 50%, the results raise the question whether current psychiatric classification and diagnostic scales are sensitive enough to detect the effects of CPL. This applies as well to developing societies like China, where the massive migration of mostly young individuals from the countryside to urban areas has left approximately 30 million small children behind (Liu et al., 2009).

CPL is a risk factor for adult anxious and depressive psychopathology (Kendler et al., 1992; Bandelow et al., 2002). However, its detrimental long-term effect is not limited to psychiatric illness (Shonkoff et al., 2009). Using criteria similar to ours, the Adverse Childhood Experiences (ACE) Study, a CDC supported prospective cohort study of 16,908 adults found a significant relationship between CPL and premature death in adulthood (Brown et al., 2009). Retrospective (e.g., Juang et al., 2004; Kopec and Sayre, 2005) and prospective longitudinal data (Fearon and Hotopf, 2001; Harter et al., 2003; Katerndahl 2008; Jones et al., 2009), link family disruption, physical abuse, separation and maternal loss in early life to chronic physical pain in adulthood. Whether the NL vs. SL probe has a differential effect on pain perception and physiological pain measures, and whether CPL status modifies this interaction should be explored. Unfortunately, our exploratory pain measure was limited to a single item, and in retrospect, was clearly inadequate.

CPL as related to childhood separation anxiety, adult panic disorder (PD) and suffocation hypersensitivity was studied by Battaglia et al. (2009). In a large sample of twins from Norway, CPL accounted in no small part for “the covariation between separation anxiety in childhood, hypersensitivity to CO₂ (as indexed by the anxiety response to a 35% CO₂/65% O₂ mixture), and PD in adulthood”. Note that in Battaglia's study, CPL increased reactivity to the 35% CO₂ probe.” (Preter et al., 2011) [...]

“Testing the specificity of the naloxone-lactate model of clinical panic requires double-blind investigation whether specific anti-panic drugs, but not panic irrelevant drugs, block this effect.” If found, this has practical and heuristic implications.

First, there is currently no specific, screening method for testing putative anti-panic drugs except by the experimental treatment of panic disorder patients. The naloxone+lactate effect

in normal humans may afford such a screening method, and may be extended to preclinical studies. Second, these data offer heuristic support for the theory that an opioidergic dysfunction is the pathophysiological mechanism underlying panic disorder. If so, the appropriateness of opioidergic therapeutic agents comes into question. The use of morphine or other simple agonists would probably be rejected for fear of inducing addiction, although the evidence for addiction during indicated medical treatment is slim. However, recent work with opioidergic mixed agonist-antagonists (Gerra et al., 2006; Wallen et al., 2006), e.g. buprenorphine, may be relevant. The concern about addiction would be mitigated by the fact that higher doses become receptor blockers rather than agonists. Positive results would foster investigations into basic molecular mechanisms. For instance, we note that the dose of naloxone used in our study (2mg/kg) substantially exceeds that needed for μ opioid receptor (MOR) blockade (Sluka et al. 1999), suggesting a role for the δ opioid receptor (DOR). This could spark interest in the development of specific DOR agonists suitable for human use. Currently, such agents have not been developed, although agents suitable for animal use are available.” (Preter et al., 2011)

Since the publication of our paper, exciting new work in panic disorder has emerged, notably from Brazil. Appropriately, the “First World Symposium On Translational Models Of Panic Disorder”, was held in Vitoria, E.S., in November of 2012. Moreira et al. (2013) present a thoughtful review of the use of rodents in panic disorder research. Graeff (2012), studying an animal model of panic disorder found that the inhibitory action of serotonin is connected with activation of endogenous opioids in the periaqueductal gray (PAG). Schenberg and colleagues (Schmitel et al., 2012) suggest “[the PAG] harbors an anoxia-sensitive suffocation alarm system”. Activation precipitates panic attacks and potentiates the subject's responses to hypercapnia. Notably, the resemblance of these effects to panic disorder was supported by their pharmacological parallel to panic disorder treatment. This model was also previously supported by demonstrating a lack of stress hormone release during DPAG stimulation thus paralleling panic disorder (Schenberg et al., 2008). The utility of opioidergic mixed agonist-antagonists in animal models of panic disorder and in treatment refractory patients would seem promising.

The lack of hypothalamic pituitary adrenal (HPA) activation during the panic attack, as it occurs in panic disorder, is a striking peculiarity since it contradicts the belief that the panic is an expression of a hypersensitive fear mechanism that would stimulate the HPA anti-stress response, supposedly reactive to all dangers. This is usually understood as dependent on hyper-responsiveness of the amygdala. For instance, both Stein (2008), and Gorman et al. (2000) neglect or dismiss the incongruity of the lack of HPA response, claiming a supposed amygdala-based hypersensitive fear system as central to panic.

Further damage to the amygdalocentric fear system theory is provided by Feinstein et al. (2013) who studied three patients with amygdala damage produced by Urbach-Wiethe syndrome. It is worth extensive citation:

“A substantial body of evidence has emphasized the importance of the amygdala in fear [...]. In animals, amygdala-restricted manipulations interfere with the acquisition, expression and recall of conditioned fear and other forms of fear and

anxiety-related behaviors. In humans, focal bilateral amygdala lesions are extraordinarily rare, and such cases have been crucial for understanding the role of the human amygdala in fear. [...] The most intensively studied case is patient SM, whose amygdala damage stems from Urbach-Wiethe disease [...] Previous studies have shown that patient SM does not condition to aversive stimuli [...], fails to recognize fearful faces [...] and demonstrates a marked absence of fear during exposure to a variety of fear-provoking stimuli, including life-threatening traumatic events [...]. Patients with similar lesions have largely yielded similar results [...]. One stimulus not previously tested in humans with amygdala damage is CO₂ inhalation. Inhaling CO₂ stimulates breathing and can provoke both air hunger and fear [...] Furthermore, CO₂ can trigger panic attacks, especially in patients with panic disorder [...]. Recent work in mice found that the amygdala directly detects CO₂ and acidosis to produce fear behaviors [...]. Thus, we hypothesized that bilateral amygdala lesions would reduce CO₂-evoked fear in humans.

In contrast with our prediction, patient SM reported fear in response to a 35% CO₂ inhalation challenge. To the best of our knowledge, this was the first time patient SM experienced fear in any setting, laboratory or otherwise, since childhood [...]. To further explore this issue, we tested two additional patients (AM and BG), monozygotic twin sisters with focal bilateral amygdala lesions resulting from Urbach-Wiethe disease [...] As with patient SM, both patients also reported experiencing fear during the CO₂ challenge.” (Feinstein et al., 2013)

These startling observations affirm that the reaction to carbon dioxide must be due to an alternative alarm system, such as has been proposed for possible suffocation. We have suggested that under conditions of threatened asphyxia the activation of the HPA system would produce a counterproductive hyperoxidative state and is therefore inactivated. This is in keeping with the observation that the tachycardia during panic is produced by vagal withdrawal rather than a counterproductive sympathetic oxidative surge (Preter and Klein, 2008).

In conclusion, we objectively, experimentally showed a physiological link between endogenous opioid system deficiency and panic-like suffocation sensitivity in healthy adults. This is consonant with the expanded Suffocation-False Alarm theory of panic suggesting an episodic functional endogenous opioid deficit (Preter and Klein, 1998). The specificity of the naloxone+lactate model of clinical panic should be tested using specific anti-panic components, possibly including opioidergic mixed agonist-antagonists such as buprenorphine. If specific, the naloxone+lactate effect in normal humans affords a screening method for testing putative anti-panic drugs which is currently not available. This could obviate the experimental treatment of panic disorder patients in drug development.

Our data also show for the first time that actual separations and losses during childhood, such as parental death, parental separation or divorce (CPL), effect lifelong alterations in the physiological reactivity of the endogenous opioid system of healthy adults.

This result encourages epigenetic inquiry into the effects of CPL on endogenous opioid systems, and their role in resilience under extreme stress. In addition, a redefinition of what constitutes a (truly) healthy control in clinical research protocols may be called for.

References

- Ballenger JC. Treatment of panic disorder in the general medical setting. *J Psychosom Res.* Jan; 1998 44(1):5–15. [PubMed: 9483459]
- Bandelow B, Spath C, Tichauer GA, Brooks A, Hajak G, Ruther E. Early traumatic life events, parental attitudes, family history, and birth risk factors in patients with panic disorder. *Comprehensive Psychiatry.* 2002; 43:269–278. [PubMed: 12107864]
- Baron C, Lamarre A, Veilleux P, Ducharme G, Spier S, Lapierre JG. Psychomaintenance of childhood asthma: a study of 34 children. *J Asthma.* 1986; 23:69–79. [PubMed: 3745110]
- Battaglia M, et al. Age at onset of panic disorder: influence of familial liability to the disease and of childhood separation anxiety disorder. *Am J Psychiatry.* 1995; 152:1362–4. [PubMed: 7653694]
- Battaglia M, Pesenti-Gritti P, Medland SE, Ogliari A, Tambs K, Spatola CA. A genetically informed study of the association between childhood separation anxiety, sensitivity to CO₂, panic disorder, and the effect of childhood parental loss. *Archives of General Psychiatry.* 2009; 66:64–71. [PubMed: 19124689]
- Baydur A. Nebulized morphine: a convenient and safe alternative to dyspnea relief? *Chest.* 2004; 125:363–5. [PubMed: 14769709]
- Beck JG, Ohtake PJ, Shipherd JC. Exaggerated anxiety is not unique to CO₂ in panic disorder: a comparison of hypercapnic and hypoxic challenges. *J Abnorm Psychol.* 1999; 108:473–82. [PubMed: 10466271]
- Beck JG, Shipherd JC, Ohtake P. Do panic symptom profiles influence response to a hypoxic challenge in patients with panic disorder? A preliminary report. *Psychosom Med.* 2000; 62:678–83. [PubMed: 11020098]
- Beitman BD, Kushner M, Lamberti JW, Mukerji V. Panic disorder without fear in patients with angiographically normal coronary arteries. *J Nerv Ment Dis.* 1990; 178:307–12. [PubMed: 2338538]
- Bellodi, L.; Perna, G., editors. *The Panic Respiration Connection.* MDM Medical Media; 1998.
- Benedikt M. Ueber Platzschwindel. *Allgemeine Wiener Medizinische Zeitung.* 1870; 15:488–490.
- Breslau N, Schultz LR, Stewart WF, Lipton R, Welch KM. Headache types and panic disorder: directionality and specificity. *Neurology.* 2001; 56:350–4. [PubMed: 11171900]
- Briggs AC, Stretch DD, Brandon S. Subtyping of panic disorder by symptom profile. *Br J Psychiatry.* Aug.1993 163:201–9. [PubMed: 8075912]
- Brooks CM, Richards JM Jr, Bailey WC, Martin B, Windsor RA, Soong SJ. Subjective symptomatology of asthma in an outpatient population. *Psychosom Med.* 1989; 51:102–8. [PubMed: 2928458]
- Brown DW, Anda RF, Tiemeier H, Felitti VJ, Edwards VJ, Croft JB, Giles WH. Adverse childhood experiences and the risk of premature mortality. *American Journal of Preventive Medicine.* 2009; 37:389–396. [PubMed: 19840693]
- Craske MG, Poulton R, Tsao JC, Plotkin D. Paths to panic disorder/agoraphobia: an exploratory analysis from age 3 to 21 in an unselected birth cohort. *J Am Acad Child Adolesc Psychiatry.* 2001; 40:556–63. [PubMed: 11349700]
- Edlund MJ, McNamara ME, Millman RP. Sleep apnea and panic attacks. *Compr Psychiatry.* 1991; 32:130–2. [PubMed: 2022111]
- Esquivel G, Fernández-Torre O, Schruers KR, Wijnhoven LL, Griez EJ. The effects of opioid receptor blockade on experimental panic provocation with CO₂. *J Psychopharmacol.* Nov; 2009 23(8):975–8. [PubMed: 18635711]
- Esquivel G, Schruers KR, Maddock RJ, Colasanti A, Griez EJ. Acids in the brain: a factor in panic? *J Psychopharmacol.* May; 2010 24(5):639–47. [PubMed: 19460873]

- Faravelli C, Pallanti S. Recent life events and panic disorder. *Am J Psychiatry*. 1989; 146:622–6. [PubMed: 2712167]
- Fearon P, Hotopf M. Relation between headache in childhood and physical and psychiatric symptoms in adulthood: national birth cohort study. *BMJ*. May 12.2001 322(7295):1145. [PubMed: 11348907]
- Feinstein JS, Buzza C, Hurlmann R, Follmer RL, Dahdaleh NS, Coryell WH, Welsh MJ, Tranel D, Wemmie JA. Fear and panic in humans with bilateral amygdala damage. *Nat Neurosci*. Mar; 2013 16(3):270–2. [PubMed: 23377128]
- Freire RC, Nardi AE. Panic disorder and the respiratory system: clinical subtype and challenge tests. *Rev Bras Psiquiatr*. Jun; 2012 34(Suppl 1):S32–41. [PubMed: 22729448]
- Freud, S. On The Grounds For Detaching A Particular Syndrome From Neurasthenia Under The Description 'Anxiety Neurosis'. 1895.
- Frommberger UH, Tettenborn B, Buller R, Benkert O. Panic disorder in patients with dizziness. *Arch Intern Med*. Mar 14; 1994 154(5):590–1. [PubMed: 8122954]
- Gater R, Tansella M, Korten A, Tiemens BG, Mavreas VG, Olatawura MO. Sex differences in the prevalence and detection of depressive and anxiety disorders in general health care settings: report from the World Health Organization Collaborative Study on Psychological Problems in General Health Care. *Arch Gen Psychiatry*. 1998; 55:405–13. [PubMed: 9596043]
- Goetz RR, Klein DF, Gully D, Kahn J, Liebowitz M, Fyer A, Gorman JM. Panic attacks during placebo procedures in the laboratory : physiology and symptomatology. *Archives of General Psychiatry*. 1993; 50:280–285. [PubMed: 8466389]
- Goodwin RD, Eaton WW. Asthma and the risk of panic attacks among adults in the community. *Psychol Med*. 2003; 33:879–85. [PubMed: 12877402]
- Goodwin RD, Fergusson DM, Horwood LJ. Asthma and depressive and anxiety disorders among young persons in the community. *Psychol Med*. 2004; 34:1465–74. [PubMed: 15724877]
- Goodwin RD, Messineo K, Bregante A, Hoven CW, Kairam R. Prevalence of probable mental disorders among pediatric asthma patients in an inner-city clinic. *J Asthma*. 2005; 42:643–7. [PubMed: 16266954]
- Gorman JM, Kent JM, Sullivan GM, Coplan JD. Neuroanatomical hypothesis of panic disorder, revised. *Am J Psychiatry*. Apr; 2000 157(4):493–505. [PubMed: 10739407]
- Gorman JM, Askanazi J, Liebowitz MR, Fyer AJ, Stein J, Kinney JM, Klein DF. Response to hyperventilation in a group of patients with panic disorder. *American Journal of Psychiatry*. 1984; 141:857–861. [PubMed: 6428243]
- Graeff FG. New perspective on the pathophysiology of panic: merging serotonin and opioids in the periaqueductal gray. *Braz J Med Biol Res*. Apr; 2012 45(4):366–75. [PubMed: 22437485]
- Griez EJ, Colasanti A, van Diest R, Salamon E, Schruers K. Carbon dioxide inhalation induces dose-dependent and age-related negative affectivity. *PLoS One*. Oct 3.2007 2(10):e987. [PubMed: 17912364]
- Hamelsky SW, Lipton RB. Psychiatric comorbidity of migraine. *Headache*. Oct; 2006 46(9):1327–33. [PubMed: 17040330]
- Harter MC, Conway KP, Merikangas KR. Associations between anxiety disorders and physical illness. *Eur Arch Psychiatry Clin Neurosci*. 2003; 253:313–20. [PubMed: 14714121]
- Hasler G, et al. Asthma and panic in young adults: a 20-year prospective community study. *Am J Respir Crit Care Med*. 2005; 171:1224–30. [PubMed: 15764721]
- Hofer MA, Shair H. Ultrasonic vocalization during social interaction and isolation in 2-week-old rats. *Dev Psychobiol*. 1978; 11:495–504. [PubMed: 689298]
- Horwath E, Adams P, Wickramaratne P, Pine D, Weissman MM. Panic disorder with smothering symptoms: evidence for increased risk in first-degree relatives. *Depress Anxiety*. 1997; 6(4):147–53. [PubMed: 9559284]
- Johnson KM, Bradley KA, Bush K, Gardella C, Dobie DJ, Laya MB. Frequency of mastalgia among women veterans. Association with psychiatric conditions and unexplained pain syndromes. *J Gen Intern Med*. Mar; 2006 21(Suppl 3):S70–5. [PubMed: 16637950]

- Jones GT, Power C, Macfarlane GJ. Adverse events in childhood and chronic widespread pain in adult life : results from the 1958 British Birth Cohort Study. *Pain*. 2009; 143:92–96. [PubMed: 19304391]
- Juang KD, Wang SJ, Fuh JL, Lu SR, Chen YS. Association between adolescent chronic daily headache and childhood adversity : a community-based study. *Cephalalgia*. 2004; 24:54–59. [PubMed: 14687014]
- Jurenc GS. Identification of subgroups of childhood asthmatics: a review. *J Asthma*. 1988; 25:15–25. [PubMed: 3053600]
- Jurgens U, Ploog D. Cerebral representation of vocalization in the squirrel monkey. *Exp Brain Res*. 1970; 10:532–54. [PubMed: 4988409]
- Kalin NH, Shelton SE. Effects of clonidine and propranolol on separation-induced distress in infant rhesus monkeys. *Brain Res*. 1988; 470:289–95. [PubMed: 3219586]
- Kalin NH, Shelton SE, Barksdale CM. Separation distress in infant rhesus monkeys: effects of diazepam and Ro 15-1788. *Brain Res*. 1987; 408:192–8. [PubMed: 3036301]
- Kalin NH, Shelton SE, Barksdale CM. Opiate modulation of separation-induced distress in nonhuman primates. *Brain Res*. 1988; 440:285–92. [PubMed: 3359215]
- Karajgi B, Rifkin A, Doddi S, Kolli R. The prevalence of anxiety disorders in patients with chronic obstructive pulmonary disease. *Am J Psychiatry*. 1990; 147:200–1. [PubMed: 2301659]
- Katerndahl DA. Chest pain and its importance in patients with panic disorder : an updated literature review. *Primary Care Companion to the Journal of Clinical Psychiatry*. 2008; 10:376–383.
- Katon WJ, Richardson L, Lozano P, McCauley E. The relationship of asthma and anxiety disorders. *Psychosom Med*. 2004; 66:349–55. [PubMed: 15184694]
- Kaunonen M, Paivi AK, Paunonen M, Erjanti H. Death in the Finnish family: experiences of spousal bereavement. *Int J Nurs Pract*. 2000; 6:127–34. [PubMed: 11249410]
- Kehoe P, Blass EM. Opioid-mediation of separation distress in 10-day-old rats: reversal of stress with maternal stimuli. *Dev Psychobiol*. 1986; 19:385–98. [PubMed: 3732628]
- Kendler KS, Neale MC, Kessler RC, Heath AC, Eaves LJ. Childhood parental loss and adult psychopathology in women. A twin study perspective. *Archives of General Psychiatry*. 1992; 49:109–116. [PubMed: 1550463]
- Kessler RC, Chiu WT, Jin R, Ruscio AM, Shear K, Walters EE. The epidemiology of panic attacks, panic disorder, and agoraphobia in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2006; 63:415–24. [PubMed: 16585471]
- Klein DF. Delineation of two drug-responsive anxiety syndromes. *Psychopharmacologia*. 1964; 5:397–408. [PubMed: 14194683]
- Klein DF. False suffocation alarms, spontaneous panics, and related conditions. An integrative hypothesis. *Arch Gen Psychiatry*. 1993; 50:306–17. [PubMed: 8466392]
- Klein DF. Asthma and psychiatric illness. *JAMA*. 2001; 285:881–2. [PubMed: 11180724]
- Klein DF, Fink M. Psychiatric reaction patterns to imipramine. *Am J Psychiatry*. 1962; 119:432–8. [PubMed: 14033346]
- Klein RG. Is Panic Disorder Associated with Childhood Separation Anxiety Disorder? *Clin Neuropsychopharmacology*. 1995; 18(Suppl 2):7–14.
- Kopec JA, Sayre EC. Stressful experiences in childhood and chronic back pain in the general population. *Clinical Journal of Pain*. 2005; 21:478–483. [PubMed: 16215332]
- Leibold NK, Viechtbauer W, Goossens L, De Cort K, Griez EJ, Myin-Germeyns I, Steinbusch HW, van den Hove DL, Schruers KR. Carbon dioxide inhalation as a human experimental model of panic: the relationship between emotions and cardiovascular physiology. *Biol Psychol*. Oct; 2013 94(2): 331–40. [PubMed: 23816952]
- Liebowitz MR, Gorman JM, Fyer AJ, Dillon DJ, Klein DF. Effects of naloxone on patients with panic attacks. *Am J Psychiatry*. Aug; 1984a 141(8):995–7. [PubMed: 6087689]
- Liebowitz MR, Fyer AJ, Gorman JM, Dillon D, Appleby IL, Levy G, Anderson S, Levitt M, Palij M, Davies SO, Klein DF. Lactate provocation of panic attacks. • I. Clinical and behavioral findings. *Archives of General Psychiatry*. 1984b; 41:764–770. [PubMed: 6742978]

- Lipsitz JD, et al. Childhood separation anxiety disorder in patients with adult anxiety disorders. *Am J Psychiatry*. 1994; 151:927–9. [PubMed: 8185008]
- Liu Z, Li X, Ge X. Left too early : the effects of age at separation from parents on Chinese rural children's symptoms of anxiety and depression. *American Journal of Public Health*. 2009; 99:2049–2054. [PubMed: 19762669]
- Mahler DA. Understanding mechanisms and documenting plausibility of palliative interventions for dyspnea. *Curr Opin Support Palliat Care*. Jun; 2011 5(2):71–6. [PubMed: 21430539]
- Martinez JM, Papp LA, Coplan JD, Anderson DE, Mueller CM, Klein DF, Gorman JM. Ambulatory monitoring of respiration in anxiety. *Anxiety*. 1996; 2:296–302. [PubMed: 9160637]
- Mayer DJ, Wolfle TL, Akil H, Carder B, Liebeskind JC. Analgesia from electrical stimulation in the brainstem of the rat. *Science*. 1971; 174:1351–4. [PubMed: 5167502]
- Mayfield KP, D'Alecy LG. Role of endogenous opioid peptides in the acute adaptation to hypoxia. *Brain Res*. 1992; 582:226–31. [PubMed: 1393544]
- Milrod B, Leon AC, Shear MK. Can interpersonal loss precipitate panic disorder? *Am J Psychiatry*. 2004; 161:758–9. [PubMed: 15056531]
- Moreira FA, Gobira PH, Viana TG, Vicente MA, Zangrossi H, Graeff FG. Modeling panic disorder in rodents. *Cell Tissue Res*. Apr 13.2013
- Nascimento I, et al. Psychiatric disorders in asthmatic outpatients. *Psychiatry Res*. 2002; 110:73–80. [PubMed: 12007595]
- Panksepp, J. *Affective neuroscience: The foundations of human and animal emotions*. Oxford University Press; New York: 1998.
- Panksepp J, Herman B, Conner R, Bishop P, Scott JP. The biology of social attachments: opiates alleviate separation distress. *Biol Psychiatry*. 1978; 13:607–18. [PubMed: 83167]
- Panksepp, J.; Newman, JD.; Insel, TR. Critical conceptual issues in the analysis of separation-distress systems of the brain.. In: Strongman, KT., editor. *International review of studies on emotion*, Vol 2. Wiley; New York: 1992.
- Papp LA, Klein DF, Gorman JM. Carbon dioxide hypersensitivity, hyperventilation, and panic disorder. *American Journal of Psychiatry*. 1993; 150:1149–1157. [PubMed: 8392296]
- Park L, Hinton D. Dizziness and panic in China: associated sensations of zang fu organ disequilibrium. *Cult Med Psychiatry*. Jun; 2002 26(2):225–57. [PubMed: 12211326]
- Perna G, Bertani A, Politi E, Colombo G, Bellodi L. Asthma and panic attacks. *Biol Psychiatry*. 1997; 42:625–30. [PubMed: 9376459]
- Pine DS, et al. Differential carbon dioxide sensitivity in childhood anxiety disorders and nonill comparison group. *Arch Gen Psychiatry*. 2000; 57:960–7. [PubMed: 11015814]
- Pine DS, et al. Response to 5% carbon dioxide in children and adolescents: relationship to panic disorder in parents and anxiety disorders in subjects. *Arch Gen Psychiatry*. 2005; 62:73–80. [PubMed: 15630075]
- Ploog D. Neurobiology of primate audio-vocal behavior. *Brain Res*. 1981; 228:35–61. [PubMed: 7023614]
- Preter M, Bursztajn HJ. Crisis and opportunity-The DSM-V and its neurology quandary. *Asian J Psychiatr*. Dec.2009 2(4):143. [PubMed: 23051093]
- Preter, M.; Klein, DF. Panic disorder and the suffocation false alarm theory: current state of knowledge and further implications for neurobiologic theory testing.. In: Bellodi, L.; Perna, G., editors. *The Panic Respiration Connection*. MDM Medical Media; Milan: 1998.
- Preter M, Klein DF. Panic, suffocation false alarms, separation anxiety and endogenous opioids. *Prog Neuropsychopharmacol Biol Psychiatry*. Apr 1; 2008 32(3):603–12. [PubMed: 17765379]
- Preter M, Lee SH, Petkova E, Vannucci M, Kim S, Klein DF. Controlled cross-over study in normal subjects of naloxone-preceding-lactate infusions;respiratory and subjective responses: relationship to endogenous opioid system,suffocation false alarm theory and childhood parental loss. *Psychol Med*. Feb; 2011 41(2):385–93. [PubMed: 20444308]
- Roberson-Nay R, Eaves LJ, Hetttema JM, Kendler KS, Silberg JL. Childhood separation anxiety disorder and adult onset panic attacks share a common genetic diathesis. *Depress Anxiety*. Apr; 2012 29(4):320–7. [PubMed: 22461084]

- Roberson-Nay R, Kendler KS. Panic disorder and its subtypes: a comprehensive analysis of panic symptom heterogeneity using epidemiological and treatment seeking samples. *Psychol Med.* Nov; 2011 41(11):2411–21. [PubMed: 21557895]
- Roberson-Nay R, Latendresse SJ, Kendler KS. A latent class approach to the external validation of respiratory and non-respiratory panic subtypes. *Psychol Med.* Mar; 2012 42(3):461–74. [PubMed: 21846423]
- Roberson-Nay R, Moruzzi S, Ogliari A, Pezzica E, Tambs K, Kendler KS, Battaglia M. Evidence for distinct genetic effects associated with response to 35% CO₂. *Depression and Anxiety.* 2013; 30(3):259–266. [PubMed: 23349098]
- Roy-Byrne PP, Craske MG, Stein MB. Panic disorder. *Lancet.* 2006; 368:1023–32. [PubMed: 16980119]
- Santiago TV, Edelman NH. Opioids and breathing. *J Appl Physiol.* 1985; 59:1675–85. [PubMed: 2934364]
- Sareen J, Jacobi F, Cox BJ, Belik SL, Clara I, Stein MB. Disability and poor quality of life associated with comorbid anxiety disorders and physical conditions. *Arch Intern Med.* Oct 23; 2006 166(19): 2109–16. [PubMed: 17060541]
- Schenberg LC, Dos Reis AM, Ferreira Póvoa RM, Tufik S, Silva SR. A panic attack-like unusual stress reaction. *Horm Behav.* Nov; 2008 54(5):584–91. [PubMed: 18423636]
- Schimitel FG, De Almeida GM, Pitol DN, Armini RS, Tufik S, Schenberg LC. Evidence Of A Suffocation Alarm System Within The Periaqueductal Gray Matter Of The Rat. *Neuroscience.* 2012; 200:59–73. [PubMed: 22062132]
- Scott, JP. Effects of psychotropic drugs in separation distress in dogs. *Proc IX Congress ECNP Exc Med; Amsterdam:* 1974.
- Shavitt RG, Gentil V, Mandetta R. The association of panic/agoraphobia and asthma. Contributing factors and clinical implications. *Gen Hosp Psychiatry.* 1992; 14:420–3. [PubMed: 1473713]
- Sheikh JI, Leskin GA, Klein DF. Gender differences in panic disorder: findings from the National Comorbidity Survey. *Am J Psychiatry.* 2002; 159:55–8. [PubMed: 11772690]
- Shonkoff JP, Boyce WT, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities : building a new framework for health promotion and disease prevention. *Journal of the American Medical Association.* 2009; 301:2252–2259. [PubMed: 19491187]
- Sietsema KE, Simon JI, Wasserman K. Pulmonary hypertension presenting as a panic disorder. *Chest.* 1987; 91:910–2. [PubMed: 3581939]
- Sinha SS, Goetz RR, Klein DF. Physiological and behavioral effects of naloxone and lactate in normal volunteers with relevance to the pathophysiology of panic disorder. *Psychiatry Res.* 2007; 149:309–14. [PubMed: 17156855]
- Sluka KA, Deacon M, Stibal A, Strissel S, Terpstra A. Spinal blockade of opioid receptors prevents the analgesia produced by TENS in arthritic rats. *Journal of Pharmacology and Experimental Therapeutics.* 1999; 289:840–846. [PubMed: 10215661]
- Staab JP. Chronic dizziness: the interface between psychiatry and neuro-otology. *Curr Opin Neurol.* Feb; 2006 19(1):41–8. [PubMed: 16415676]
- Stefano GB, Scharrer B, Smith EM, Hughes TK Jr, Magazine HI, Bilfinger TV, Hartman AR, Fricchione GL, Liu Y, Makman MH. Opioid and opiate immunoregulatory processes. *Crit Rev Immunol.* 1996; 16:109–44. [PubMed: 8879941]
- Stein G. Panic disorder: The psychobiology of external treat and introceptive distress. *CNS Spectrums.* 2008; 13(Issue 1):26–30. [PubMed: 18204411]
- Valença AM, Falcão R, Freire RC, Nascimento I, Nascentes R, Zin WA, Nardi AE. The relationship between the severity of asthma and comorbidities with anxiety and depressive disorders. *Rev Bras Psiquiatr.* 2006; 28:206–8. [PubMed: 16924350]
- van der Molen GM, van den Hout MA, van Dieren AC, Griez E. Childhood separation anxiety and adult-onset panic disorders. *J Anxiety Disord.* 1989; 3:97–106.
- van Duinen MA, Schruers KR, Maes M, Griez EJ. CO₂ challenge induced HPA axis activation in panic. *Int J Neuropsychopharmacol.* Dec; 2007 10(6):797–804. [PubMed: 17076937]
- Verburg K, Griez E, Meijer J, Pols H. Respiratory disorders as a possible predisposing factor for panic disorder. *J Affect Disord.* 1995; 33:129–34. [PubMed: 7759661]

- Wallen MC, Lorman WJ, Gosciniak JL. Combined buprenorphine and chlonidine for short-term opiate detoxification : patient perspectives. *Journal of Addictive Diseases*. 2006; 25:23–31. [PubMed: 16597570]
- Weissman MM. The epidemiology of anxiety disorders: rates, risks and familial patterns. *J Psychiatr Res*. 1988; 22(Suppl 1):99–114. [PubMed: 3050062]
- Wingate BJ, Hansen-Flaschen J. Anxiety and depression in advanced lung disease. *Clin Chest Med*. 1997; 18:495–505. [PubMed: 9329872]
- Wise SP, Herkenham M. Opiate receptor distribution in the cerebral cortex of the Rhesus monkey. *Science*. 1982; 218:387–9. [PubMed: 6289441]
- Yellowlees PM, Haynes S, Potts N, Ruffin RE. Psychiatric morbidity in patients with life-threatening asthma: initial report of a controlled study. *Med J Aust*. 1988; 149:246–9. [PubMed: 3412214]
- Yellowlees PM, Kalucy RS. Psychobiological aspects of asthma and the consequent research implications. *Chest*. 1990; 97:628–34. [PubMed: 2407451]
- Zebraski SE, Kochenash SM, Raffa RB. Lung opioid receptors: pharmacology and possible target for nebulized morphine in dyspnea. *Life Sci*. 2000; 66:2221–31. [PubMed: 10855942]

Review Highlights

1. Presentations at the First Symposium On Translational Models Of Panic Disorder
2. Physiologic suffocation alarm monitors information about potential suffocation
3. Impaired endogenous opioid system basis for heightened suffocation sensitivity
4. Sudden loss, childhood separation anxiety antecedents of “spontaneous” panic
5. Childhood separations cause lifelong endogenous opioid impairment in **normal** adults