

Research

Open Access

Epigenetic *CRBP* downregulation appears to be an evolutionarily conserved (human and mouse) and oncogene-specific phenomenon in breast cancer

Alice Arapshian^{1,3}, Silvina Bertran¹, Yuvarani S Kuppumbatti^{1,4},
Shigeo Nakajo² and Rafael Mira-y-Lopez*¹

Address: ¹Department of Medicine, Mount Sinai School of Medicine, New York, NY 10029-6574, USA, ²Laboratory of Biological Chemistry, School of Pharmaceutical Sciences, Showa University, Japan, ³Current address: 61-02 79th Street, Middle Village, NY 11379, USA and ⁴Current address: Department of Neurology, University of California, Irvine, CA 92697, USA

Email: Alice Arapshian - aa74ny@yahoo.com; Silvina Bertran - silvina.bertran@mssm.edu; Yuvarani S Kuppumbatti - ykuppumb@uci.edu; Shigeo Nakajo - nakajo@pharm.showa-u.ac.jp; Rafael Mira-y-Lopez* - rafael.mira@mssm.edu

* Corresponding author

Published: 27 April 2004

Received: 05 September 2003

Molecular Cancer 2004, 3:13

Accepted: 27 April 2004

This article is available from: <http://www.molecular-cancer.com/content/3/1/13>

© 2004 Arapshian et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The cellular retinol binding protein I gene (*CRBP*) is downregulated in a subset of human breast cancers and in MMTV-Myc induced mouse mammary tumors. Functional studies suggest that *CRBP* downregulation contributes to breast tumor progression. What is the mechanism underlying *CRBP* downregulation in cancer? Here we investigated the hypothesis that *CRBP* is epigenetically silenced through DNA hypermethylation in human and mouse breast cancer.

Results: Bisulfite sequencing of *CRBP* in a panel of 6 human breast cancer cell lines demonstrated that, as a rule, *CRBP* hypermethylation is closely and inversely related to *CRBP* expression and identified one exception to this rule. Treatment with 5-azacytidine, a DNA methyltransferase inhibitor, led to *CRBP* reexpression, supporting the hypothesis that *CRBP* hypermethylation is a proximal cause of *CRBP* silencing. In some cells *CRBP* reexpression was potentiated by co-treatment with retinoic acid, an inducer of *CRBP*, and trichostatin A, a histone deacetylase inhibitor. Southern blot analysis of a small panel of human breast cancer specimens identified one case characterized by extensive *CRBP* hypermethylation, in association with undetectable *CRBP* mRNA and protein. Bisulfite sequencing of *CRBP* in MMTV-Myc and MMTV-Neu/NT mammary tumor cell lines extended the rule of *CRBP* hypermethylation and silencing (both seen in MMTV-Myc but not MMTV-Neu/NT cells) from human to mouse breast cancer and suggested that *CRBP* hypermethylation is an oncogene-specific event.

Conclusion: *CRBP* hypermethylation appears to be an evolutionarily conserved and principal mechanism of *CRBP* silencing in breast cancer. Based on the analysis of transgenic mouse mammary tumor cells, we hypothesize that *CRBP* silencing in human breast cancer may be associated with a specific oncogenic signature.

Background

Retinol has been implicated in the regulation of adult epi-

thelial cell differentiation in addition to being essential for normal embryonic development, vision and

reproduction [1-3]. Most of the biological effects of retinol are secondary to its metabolism to retinoic acid (RA) and RA activation of members of the RA receptor (RAR) and retinoid X receptor (RXR) family of ligand-dependent transcription factors. As such, the study of RARs and RXRs has gained widespread attention. However, a number of other genes play fundamental roles in regulating RXR-RAR function by regulating retinol and RA metabolism. Among these genes is cellular retinol-binding protein I (CRBP), which has been shown to be indispensable for life when vitamin A levels in the diet are limiting. Thus, CRBP null mice are unable to maintain adequate retinol liver stores and survive only when dietary vitamin A is abundant, a condition that is artificial and does not reflect the dietary habits of mammals in the wild [4].

In addition to being expressed in stellate cells of the liver, CRBP is expressed in many other organs and cell types including epithelia [5-7]. Several observations, including the finding that CRBP expression during mouse embryogenesis overlapped that of RAR β and RAR γ , led to the suggestion that the function of CRBP is to store and release retinol where high levels of RA are developmentally required [8]. *mCRBP* expression is regulated by RA through a RA response element located \sim 1 kb upstream of the transcription start site, suggesting the existence of a positive feedback mechanism for regulating retinoid metabolism and action [9]. Glucocorticoids, cAMP, transforming growth factor β and serum factors have also been shown to impinge on CRBP expression [10-13]. The proximal *hCRBP* promoter region has been cloned, the transcription start site identified, and functional NF1 sites at approximately -250 and +70 demonstrated [14,15] but beyond these earlier studies there have been no functional promoter analyses.

We have recently shown that CRBP expression is downregulated in a subset of human breast cancers [7]. In follow-up studies of the potential significance of this finding, we found that the downregulation of CRBP blocks the differentiation and promotes the growth of SV40-transformed breast epithelial cells ([16] and Farias et al., manuscript submitted). The differentiation-promoting, growth-arresting effect of CRBP, witnessed in both human and mouse mammary epithelial cells, was assigned to CRBP regulation of retinol storage and RAR activation (Farias et al., manuscript submitted). These results suggested that CRBP downregulation is a contributing cause rather than a consequence of breast cancer and led us to investigate the mechanism underlying CRBP downregulation. We report here that CRBP is epigenetically silenced as a result of promoter region hypermethylation in both human and mouse breast cancer cells. Furthermore, we identify one exception to this rule and provide data that suggests that CRBP silencing is an oncogene-specific event at least in

the context of mouse mammary cancer. These results extend those published by Esteller et al. while this work was in progress [17].

Results

hCRBP expression in breast cancer cells varies widely

We first analyzed a panel of human breast cancer cell lines for CRBP expression using a semi-quantitative RT-PCR Southern blot protocol. CRBP expression varied from robust (MDA-MB-468), to weak (MCF-7), to very weak or undetectable (ZR75-1 and MTSV1-7; Fig. 1A). Given that CRBP downregulation occurs with an estimated frequency of 24% in human breast cancer, i.e., it is not a universal event, the identification of cell lines, such as MDA-MB-468, in which CRBP was well expressed, should not be surprising. We wondered whether the intermediate level of CRBP expression in MCF-7 cells was a consequence of clonal heterogeneity and tested this idea by random clonal selection and analysis. In support of this notion, we were able to isolate subclones that mirrored the extremes of CRBP expression, with MCF-7 clone B resembling MDA-MB-468 and MCF-7 clone A resembling ZR75-1 (Fig. 1B, compare with Fig. 1A). After densitometry and normalization to a house keeping gene, we plotted the results for all cell lines on a log scale (Figs. 2A and 3A).

hCRBP methylation, with one exception, is inversely proportional to hCRBP expression

On the basis of preliminary Southern blot analyses that revealed no major *hCRBP* deletions or insertions in breast cancer cells (unpublished data), and taking into account the presence of a CpG island in the *hCRBP* regulatory region [14], we hypothesized that *hCRBP* downregulation might be secondary to gene hypermethylation. This hypothesis was tested by bisulfite sequencing of 26 CpG dinucleotides within the *hCRBP* locus (promoter and 5'-UTR region) in the cell lines whose CRBP expression we had characterized (Fig. 1). We found that all 26 sites were hypomethylated in MDA-MB-468 cells and 14 of the 26 sites were hypermethylated in ZR75-1 cells, providing the first evidence of an inverse association between CRBP methylation and expression (Fig. 2B). However, this notion was challenged by the finding that the same CRBP region was uniformly hypomethylated in MTSV1-7 cells (Fig. 2B). The bisulfite sequencing analysis of CRBP in MCF-7 parental cells and clones A and B proved very useful in that it demonstrated a very tight inverse association between CRBP methylation and expression. Thus, the intermediate level of CRBP expression in parental MCF-7 cells was associated with a mixed pattern of CRBP methylation (of 17 amplicons sequenced, 6 showed uniform hypermethylation and 10 showed uniform hypomethylation), and this mixed pattern resolved into uniform hyper or hypomethylation in clones A and B, respectively (Fig. 3B). These results suggest that, as a rule, CRBP hypermeth-




Figure 1
***hCRBP* expression differs widely among human breast cancer cell lines.** **A.** RT-PCR Southern blot analysis of *CRBP* expression in ZR75-1, MTSV1-7, MCF-7 and MDA-MB-468 cells. **B.** RT-PCR Southern blot analysis of *CRBP* expression in MCF-7 subclones and again MDA-MB-468, to allow comparison with the cell lines studied in A. Cells were either untreated (MDA-MB-468) or treated with RA and TSA as further described in Fig. 4 (MCF-7 clone A and B cells). The data in Fig. 1 were used to generate Figs. 2A and 3A after densitometric analysis and normalization to autoradiographic exposure time (1 to 24 h) and GAPDH expression (loading control).

ylation is silencing, and that MTSV1-7 cells represent an exception to this rule.

Treatment with a demethylating agent rescues *hCRBP* expression

CRBP is a RA-inducible gene [9,18] and trichostatin A (TSA), a histone deacetylase inhibitor, has been shown to synergize with RAR induction of target gene transcription [19]. Cells in culture vary widely in their sensitivity to

retinoids in the serum present in the culture medium; in the case of MDA-MB-468 cells (hypomethylated *CRBP*), there was robust *CRBP* expression in the absence of exogenous RA, whereas in the case of MCF-7 clone B cells (also hypomethylated *CRBP*) there was no *CRBP* expression unless exogenous RA was added and expression increased further in presence of TSA (Fig. 4A). This difference may relate to the fact that MDA-MB-468 cells are competent in RA synthesis from serum retinol whereas MCF-7 cells are not [20]. Based on the data for MCF-7 clone B cells (Fig. 4A), we predicted that treatment of MCF-7 clone A cells (hypermethylated *CRBP*) with 5-azacytidine (azaC), a DNA methyltransferase inhibitor, would result in some *CRBP* reexpression and that this effect would be potentiated by the addition of RA and TSA. We further predicted that in absence of azaC, RA and TSA would be ineffective. The data confirmed these expectations (Fig. 4B), thus supporting our hypothesis that hypermethylation is a proximal cause of *CRBP* silencing. In another cell line with hypermethylated *CRBP* (ZR75-1), reexpression could also be demonstrated to be critically dependent on azaC (Fig. 4A). Although we did not directly probe the effect of azaC on *CRBP* methylation, we showed earlier that the same protocol used here achieved partial demethylation of another hypermethylated gene in MCF-7 cells [21]. The work of Esteller et al. [17] suggest that our data for MCF-7 clone A is rather the exception than the rule in that azaC alone is generally sufficient to achieve *CRBP* reexpression when the gene is hypermethylated, suggesting that endogenous retinoids or other modulators of *CRBP* expression (Introduction) can drive *CRBP* transcription upon azaC-induced demethylation.

Extensive *hCRBP* hypermethylation in association with undetectable *hCRBP* in a human breast cancer specimen

Esteller et al. demonstrated by methylation-sensitive PCR (MSP) that *hCRBP* is hypermethylated and silenced in a subset of human breast cancers [17]. MSP is a highly sensitive technique that will detect methylation of a gene of interest even when only a small percentage of cells within the tissue have methylated alleles [22]. We wished to probe *CRBP* methylation using a less sensitive technique to infer whether *CRBP* methylation was a rare event on a per cell basis or, alternatively, was a tissue-wide event. For this purpose we used a classical Southern blot analysis approach [23] that relied on the use of methylation-sensitive isoschizomers and knowledge of the specific CpG sites that are methylated in human breast cancer cells (Figs. 2B and 3B). To test the validity of this strategy, we first applied it to MDA-MB-468 and ZR57-1 cells. Southern blotting correctly identified *CRBP* hypo and hypermethylation in MDA-MB-468 and ZR57-1 cells, respectively (Fig. 5A). This strategy was then used to investigate *CRBP* methylation in a small panel of human breast cancers (n = 10) and the results obtained for 2 cases are




Figure 2
***hCRBP* hypermethylation explains some but not all cases of *hCRBP* silencing.** **A.** Relative levels of *hCRBP* expression determined by RT-PCR Southern analysis (see Fig. 1). The limit of detection is represented by a relative log value of 1.5 that corresponds to film background after 24 h exposure. Note that *hCRBP* expression in MDA-MB-468 cells is greater than that in ZR75-1 or MTSV1-7 cells by about 3 logs. **B.** *hCRBP* bisulfite sequencing results for the same cell lines as in A. Filled circles designate methylated and open circles designate unmethylated CpGs. Each row displays the methylation status of CpG sites across an individual amplicon; each "column" displays the methylation status of an individual CpG site across amplicons. The approximate position of each CpG site relative to the transcription start site is indicated in the abscissa.




Figure 3
hCRBP methylation explains clonal variations in *hCRBP* expression. **A.** Same as for Fig. 2A. Note that MCF-7 cells express *hCRBP* at a level that is intermediate to the extremes represented by MCF-7 clones A and B, which resemble in this regard ZR75-I and MDA-MB-468 cells, respectively. **B.** *hCRBP* bisulfite sequencing results for the same cell lines as in A. Filled circles designate methylated and open circles designate unmethylated CpGs (see legend to Fig. 2B for further details).




Figure 4
Treatment with a demethylating agent rescues *hCRBP* expression in MCF-7 clone B and ZR75-1 cells. **A.** RT-PCR Southern blot analysis of *hCRBP* expression in MDA-MB-468 and MCF-7 clone B cells treated as indicated, demonstrating different sensitivity to endogenous (serum) retinoids. RA, 1 μ M all-trans RA; TSA, 100 nM trichostatin A (24 h treatment). **B.** RT-PCR Southern analysis of *hCRBP* expression in MCF-7 clone A and ZR75-1 cells treated as indicated, demonstrating that a demethylating agent is required for CRBP reexpression. AzaC, 5 μ M 5-azacytidine (72 h pretreatment, 96 h total); RA and TSA as above (present during last 24 h).

shown in Fig. 5B. One breast cancer specimen (tumor B) exhibited frank *CRBP* hypermethylation in association with low levels of CRBP mRNA, whereas tumor A did not show evidence of *CRBP* methylation by Southern blotting and this correlated with easily detectable CRBP mRNA levels. Based on a densitometric evaluation of the yield of the 205 bp fragment, we estimate that >50% of the cells in tumor B had at least one hypermethylated *CRBP* allele. When viewed in the context of the other data presented here and the work of Esteller et al. [17], this finding is of interest because it suggests that in some breast cancers there is clonal expansion of a cell population in which the *CRBP* gene is hypermethylated. Fig. 5B also demonstrates that there is good agreement between *CRBP* expression at

the mRNA and protein levels. Analysis of the remaining 8 human breast cancer specimens did not identify other examples like tumor B in Fig. 5, either because of the small sample size, the low sensitivity of Southern blotting, or the low cellularity of the tumor specimen (data not shown).

***Myc* but not *Neu* leads to mCRBP hypermethylation and silencing in mouse mammary epithelial cells**

Morrison and Leder have shown that *CRBP* expression in transgenic mouse mammary tumors is a function of the initiating oncogene, with *Myc*-induced tumors lacking and *Neu*-induced tumors displaying *CRBP* expression [24]. We hypothesized that this observation might be




Figure 5

In some human breast cancers, *hCRBP* hypermethylation is widespread. **A.** Sketch outlining the Southern blot strategy used to evaluate *hCRBP* methylation in human breast cancer specimens. The strategy was tested in MDA-MB-468 and ZR75-1 cells whose *hCRBP* methylation status was known beforehand. B, BH and BM stand for, respectively, Bsu36I digestion alone, Bsu36I digestion followed by HpaII digestion, and Bsu36I digestion followed by MspI digestion (MspI is the methylation-insensitive isoschizomer of HpaII). In agreement with the bisulfite sequencing data, Southern blotting revealed *hCRBP* undermethylation in MDA-MB-468 and hypermethylation in MCF-7 clone A cells (note loss of the large Bsu36I fragment after BH digestion of MDA-MB-468 genomic DNA and retention of this fragment in the case of MCF-7 clone A). A *hCRBP* restriction fragment length polymorphism (an extra Bsu36I site within the 205 bp fragment) accounts for the displaced band migration pattern in MDA-MB-468 cells. **B.** Southern, RT-PCR and Western blotting results for 2 human breast cancer specimens. The yield of the 205 bp product was decreased by >50% after BH digestion relative to BM digestion. We infer from this that >50% of the cells in the tumor displayed *hCRBP* hypermethylation. The protein blot was post-stained with Ponceau S and a major band used as loading control.




Figure 6
***mCRBP* hypermethylation and silencing is an oncogene-specific event.** **A.** Northern blot analysis. **B.** *mCRBP* bisulfite sequencing results for the cell lines tested in A. Each row corresponds to one sequenced amplicon. Filled circles designate methylated and open circles designate unmethylated CpGs (see legend to Fig. 2B for further details). **C.** RT-PCR demonstrating effect of azaC treatment (as in Fig. 4) on *mCRBP* expression in 16MB9a cells.

explained by Myc but not Neu promotion of *CRBP* hypermethylation. To test this, we obtained two mammary tumor cell lines established from Myc and two others established from Neu-induced tumors [25] and, as a first step, confirmed that *CRBP* was expressed in the latter but not the former cell lines (Fig. 6A). We then analyzed *CRBP* methylation in the same cell lines. As predicted by our hypothesis, bisulfite sequencing of CpG dinucleotides within the *mCRBP* promoter and 5'-UTR region revealed extensive to moderate *CRBP* hypermethylation in MMTV-Myc as opposed to virtually no methylation in MMTV-Neu mammary tumor cells (Fig. 6B). As shown in Fig. 6C, treatment with azaC rescued *CRBP* expression in MMTV-

Myc (16MB9a) cells. Thus, *CRBP* hypermethylation and silencing appears to represent an evolutionarily conserved event in mammary carcinogenesis. The data in Fig. 6 sets the stage for additional work investigating whether *CRBP* hypermethylation in human breast cancer can be associated with a specific oncogenic signature.

Discussion

Our data extend the work of Esteller et al. [17] in several respects. First, it provides an analysis of the pattern of CpG methylation across the *CRBP* promoter region. One interesting feature revealed by the bisulfite sequencing of *CRBP* in ZR75-1 and MCF-7 clone A cells, both of which express virtually no *CRBP*, is that 2 alternative methylation patterns are equally suppressive. Thus, in ZR75-1 cells, *CRBP* methylation showed intra-allelic variegation, with alternating methylated and non-methylated CpG sites (Fig. 2B). Contrast this with the uniform methylation of all CpG sites in MCF-7 clone A cells (Fig. 3B). As shown in Fig. 4, both of these patterns contributed to *CRBP* down-regulation, suggesting that uniform methylation is not required for silencing, which is consistent with the demonstrated ability of MeCP2 to bind single methylated CpG sites [26]. *CRBP* methylation in MMTV-Myc mammary tumor cells defined yet another methylation pattern characterized by both intra- and inter-allelic variegation and different degrees of methylation penetrance (Fig. 6B). The conditions that lead to the establishment of these variant methylation patterns in the first place are not known.

Second, our data identified the MTSV1-7 cell line as an exception to the rule that *CRBP* silencing is associated with *CRBP* hypermethylation. The mechanism responsible for *CRBP* silencing in MTSV1-7 cells remains to be identified but the ease with which *CRBP* can be ectopically expressed in these cells from a heterologous promoter points to a defect in transcriptional initiation rather than a subsequent event [16]. In view of the association between MMTV-Myc expression and *mCRBP* silencing (Fig. 6), we hypothesize that *c-myc* overexpression may underly *hCRBP* silencing in MTSV1-7 cells. This possibility is supported by our finding that c-myc protein is easily detected in MTSV1-7 cells by direct Western whereas it is normally present in trace levels (Farias and Mira-y-Lopez, unpublished data). c-myc protein levels in MTSV1-7 cells may be elevated either as a result of their long-term passage in vitro [27] and/or as a result of their transduction with the SV40 small t antigen [28], which was recently shown to stabilize c-myc [29]. That possibility is further supported by the finding that immortalized mouse mammary epithelial cells lose endogenous *mCRBP* expression over long term in vitro passage (Farias and Mira-y-Lopez, unpublished data). On the other hand, the fact that MMTV-Myc is associated with *mCRBP* hypermethylation (Fig. 6) but *hCRBP* was not hypermethylated

in MTSV1-7 cells (Fig. 2B) argues against our idea that *c-myc* overexpression underlies *CRBP* silencing in MTSV1-7 cells. However, hypermethylation has been suggested to represent a mechanism for the maintenance not the initial induction of transcriptional silencing [30]. Therefore, it is conceivable that the *hCRBP* locus in MTSV1-7 cells is repressed in a *c-myc*-dependent manner and is susceptible to but has not yet undergone hypermethylation.

Third, based on a small sample of 10 human breast cancer specimens, we succeeded in identifying one instance of widespread *CRBP* hypermethylation and silencing. Thus, hypermethylation was easily detected in total tumor DNA using a low sensitivity technique (Southern blotting) and this correlated with the lack of any detectable *CRBP* mRNA and protein in total tumor RNA and tumor lysates, respectively. Although only a single such tumor was identified, it seems reasonable to extrapolate, based on the strength of the combined evidence ([17] and this report), that a small subset of human breast cancers are characterized by pervasive *CRBP* methylation.

Fourth, we have found that *mCRBP* methylation is associated with *Myc* but not *Neu*-induced mammary tumors, suggesting that *mCRBP* silencing may be a requirement for MMTV-*Myc* tumorigenesis. We plan to test this hypothesis by monitoring *CRBP* expression and methylation by *in situ* hybridization and MSP, respectively, over the period of time during which premalignant changes accumulate in MMTV-*Myc* mammary tissue. We expect to find that *CRBP* downregulation precedes or coincides with the appearance of preneoplasia. We further expect that ectopic *CRBP* will prevent or delay the onset of mammary tumorigenesis. One theoretical argument against our hypothesis is that MMTV-*Myc* may transform a subpopulation of cells that do not express *CRBP* to begin with. We find this unlikely because *CRBP* mRNA is uniformly expressed in both layers of human breast ducts, *i.e.*, the epithelium itself and the underlying myoepithelium [7]. Interestingly, MMTV-*Myc* tumors were recently identified as having both epithelial and myoepithelial components [31], suggesting that MMTV-*Myc* transforms a pluripotent cell population. For the reason just stated above, we hypothesize that this population expresses *CRBP*.

Finally, our study of *CRBP* methylation in mouse mammary tumor cell lines derived from *Myc* and *Neu*-induced tumors provides a hypothetical explanation for why *CRBP* hypermethylation is characteristic of only a subset of human breast cancers. It should be noted that *CRBP* is not unique in this regard, *i.e.*, several other genes shown to be epigenetically silenced in breast cancer are silenced in a subset of, not all breast cancers [32]. The data in Fig. 6 suggests that subset-specific *CRBP* silencing might be

explained by the nature of the initiating oncogenic event. Specifically, we propose that most human breast cancers in which *c-myc* is amplified and/or overexpressed might be characterized by *CRBP* hypermethylation and, conversely, most breast cancers with *c-neu* amplification and/or overexpression might be characterized by *CRBP* hypomethylation. Interestingly, the incidence of *c-myc* amplification and *CRBP* hypermethylation in breast cancer have both been estimated at 16% [17,33]. If further work were to validate this notion, then it would be of interest to understand how *c-myc* overexpression leads to *CRBP* silencing and how *c-neu* bypasses the anti-tumor function of *CRBP*. The finding of other defects in retinol and RA metabolism in breast cancer as well as alterations in RAR isoform expression suggest that several bypassing mechanisms may exist [34-37].

Conclusions

We conclude that: (i) *CRBP* hypermethylation is, as a rule, a proximal cause of *CRBP* silencing in a subset of human and mouse breast cancers; (ii) exceptions to this rule may exist in some human breast cancers, as indicated by our results for the MTSV1-7 cell line, in which *CRBP* is silenced despite being hypomethylated; (iii) the intra-allelic pattern of *CRBP* hypermethylation may vary and tumors may consist of mixtures of cells with hyper and hypomethylated *CRBP*, but in at least some cases most tumor cells (>50%) display *CRBP* hypermethylation; (iv) at least in the context of mouse mammary cancer, *CRBP* hypermethylation appears to be an oncogene-specific event. The significance of these findings should be viewed against the background of our functional analyses of *CRBP*, which suggest that *CRBP* silencing is a contributing cause rather than a consequence of breast carcinogenesis [16].

Methods

Cells and reagents

Human breast cancer cell lines, with the exception of MTSV1-7 cells [16,28], were obtained from the American Type Culture Collection (Rockville, MD) and grown as recommended. 16MB9a and M1011 (MMTV-*Myc*) and SMF and NAF (MMTV-*Neu/NT*) mammary tumor cells were cultured as described [24]. Gel purified primers were obtained from Genelink (Hawthorne, NY). RA, TSA and azaC were obtained from Sigma (St.Louis, MO). Sodium metabisulfite was obtained from BDH, England.

RNA analyses

For RT-PCR, 2 μ g total RNA (Purescript kit, Gentra, Minneapolis, MN) were reverse transcribed using the Super-script Preamplification System (Gibco BRL Life Technologies, Grand Island, NY). This RNA input was chosen after preliminary experiments demonstrated a dose-dependent increase in *CRBP* RT-PCR product within

the range of 0.5 to 4 µg total RNA. The PCR mixture consisted of 10X PCR Buffer, 1 mM MgCl₂, 200 µM dNTPs, 0.4 µM each primer, 10% cDNA, and 5U of Taq (Promega, Madison, WI) in a total volume of 50 µl. hCRBP cDNA (320 bp) was amplified with sense primer TTGTGGCCAAACTGGCTCCA and antisense primer AACTGGAGCTTGTCTCCGT and the following cycling parameters: 94 C for 5 min; 26 cycles of 94 C for 1 min, 48 C for 1 min, 72 C for 1 min; 72 C for 5 min. 18S rRNA (489 bp) was amplified using the same protocol as above except that 5% cDNA, a 3:7 mix of primers:competimers (0.4 µM total), and an annealing temperature of 57°C were used (18S rRNA primers and competimers were from Ambion, Austin, TX). GAPDH RT-PCR was performed as described [38]. The PCR products were separated on 1% agarose, stained with ethidium bromide (18S rRNA) or blotted overnight and probed with full length CRBP cDNA or a commercial GAPDH cDNA fragment (Ambion). Northern blot analyses were performed using standard methods and a full length CRBP cDNA probe as described earlier [7].

Bisulfite sequencing

Genomic DNA was isolated using the Puregene kit (Gentra Minneapolis, MN) and digested with XbaI. The bisulfite method established by Frommer et al. [39,40] was carried out on 5 µg genomic DNA digest with the following modifications. Digested DNA was denatured with 0.5 M NaOH at 75 C for 15 min, followed by deamination with freshly prepared 4.0 M, pH 5.0 sodium metabisulfite and incubated at 55 C for 8 hours in a total volume of 1.2 ml under mineral oil. The bisulfite-treated DNA was then desalted using the Wizard DNA clean-up system (Promega), desulfonated with 0.3 M NaOH at 37 C for 15 min, neutralized with 0.1 volume of 3 M NH₄AC and ethanol precipitated with 2 volumes of cold ethanol. The pellet was resuspended in 50 µl of sterile H₂O. A 780 bp *hCRBP* fragment was amplified using 2 rounds of fully-nested PCR, which in our experience was required to yield reproducible results. First round PCR was carried out in a 50 µl reaction mixture of 10X PCR buffer, 1.2 mM MgCl₂, 160 µM dNTPs, 0.4 µM each primer, 10 µl of bisulfite-treated DNA, and 5 U of Taq (Promega). The nested PCR reactions (rounds 2 and 3) were carried out in the same way using 5% of the product obtained in the preceding round as template. The primers used were as follows: 1st round PCR, GTTGAATTTTTAGTTTTTTGATTTTAGT (sense) and ACACCAAAAAATTAACAAAAAACT (antisense); 2nd round, AGTTTTAGATAAAGTTTTGTAAGTGG (sense) and TCCCAAAATAACAAAACCCAAAAA (antisense); 3rd round, GTTTTTAATTTTTGAGTGGTTGT (sense) and ATCTACAACCTAAAACTACCCTAAAA (antisense). Cycling parameters were: 94 C for 5 min; 40 (round 1) or 20 cycles (rounds 2 and 3) of 94 C for 1 min, 45 C (rounds 1 and

3) or 50 C (round 2) for 1 min, 72 C for 1 min; 72 C for 5 min. The amplified PCR product was gel purified, subcloned into the TA cloning vector (Invitrogen, Faraday, CA), transformed into TOPO kit cells (Invitrogen) using a blue-white colony X-gal selection, and individual clones submitted for sequencing. mCRBP bisulfite sequencing was performed essentially as described above for *hCRBP* but using semi-nested PCR and the following primers and conditions: 1st round, TGAAGGATTTTAAGGGAAGTAAGGAG (sense) and AAATTCTCATTACTCAACATCTTCCAATAC (antisense), 40 cycles, annealing at 52 C; 2nd round, ATCCACAAACATTTTTAAAACAACTA (sense) and same antisense primer as above, 42 cycles, annealing at 46 C.

Southern blot analysis

Human breast carcinoma specimens that had been flash frozen within 1 h of surgery were obtained through the Cooperative Human Tissue Network <http://www.chtn.ims.nci.nih.gov> under Institutional Review Board approval. All specimens were diagnosed as invasive breast carcinoma, with or without a carcinoma in situ component, except for one specimen which was an adenoid cystic breast carcinoma (not shown in Fig. 5). Approximately 200 mg of tissue were pulverized under liquid nitrogen and RNA and DNA isolated using Qiagen's RNA/DNA kit according to manufacturer's instructions (Qiagen, Valencia, CA). Tumor DNA (45 µg) was digested with Bsu36I, ethanol precipitated, and divided into 3 parts that were either not digested further, digested with HpaII, or digested with MsPI (restriction enzymes were from New England Biolabs, Beverly, MA). After ethanol precipitation, Southern blots were prepared and hybridized to a 205 bp *hCRBP* probe amplified from a cloned *hCRBP* genomic fragment using CCGGTCTCCTCTTCTTTGTAGGGG (sense) and GGGACAGGGGCTCTGCGGGG (antisense) primers.

Immunoblot analysis

A small amount of pulverized tissue was homogenized in 10 mM Tris.Cl pH 7.4, 1 mM EDTA, 7 µM beta-mercaptoethanol, the homogenate was centrifuged at 10,000 × g for 10 min, and the supernatant centrifuged further at 100,000 × g for 30 min to yield cytosol. Protein concentration was analyzed using the method of Bradford [41] with Bio-Rad reagents (Bio-Rad, Hercules, CA) and 50 µg protein were electrophoresed per lane in a 15% sodium dodecyl sulfate-polyacrylamide gel. After blotting, CRBP protein was visualized by reaction with anti-CRBP antibody followed by secondary antibody and enhanced chemiluminescence detection. The anti-CRBP antibody was raised in rabbits against a keyhole limpet hemocyanin-conjugated CRBP-specific peptide [42] and was characterized elsewhere [16].

List of abbreviations

CRBP cellular retinol binding protein type I

MMTV mouse mammary tumor virus

RA retinoic acid

RAR RA receptor

RXR retinoid x receptor

RT-PCR reverse transcription-coupled polymerase chain reaction

ER estrogen receptor

TSA trichostatin A

azaC 5-azacytidine

MSP methylation-sensitive PCR

Authors' contributions

AA established the bisulfite sequencing protocol for *hCRBP*, applied it to most of the human breast cancer cell lines tested, generated data toward Figs. 1 and 4, established the Southern blotting strategy used in Fig 5 and generated the data in Fig. 5A. SB performed the analysis of *hCRBP* methylation in MTSV1-7 cells, extended the bisulfite sequencing method to *mCRBP* (Fig. 6), and generated data toward Figs. 1,4 and 6C. YSK established the semi-quantitative CRBP RT-PCR method and contributed toward Fig. 1. SN generated the CRBP peptide antibody used in the experiment of Fig. 5. RML initiated and supervised all aspects of the project, helped perform some of the experiments, and wrote the manuscript.

Acknowledgements

We thank the following investigators for reagent gifts: Joyce Taylor-Papadimitriou (MTSV1-7 cells), Phil Leder (16MB9a and SMF cells), Pierre Chambon (*mCRBP* cDNA) and Winnie Eskild (*hCRBP* cDNA). Tissue samples were provided by the Cooperative Human Tissue Network which is funded by the National Cancer Institute. Other investigators may have received specimens from the same subjects. This work was supported by NCI grant CA54273 to RML, the Samuel Waxman Cancer Research Foundation, the Norman and Rosita Winston Foundation and the Chemotherapy Foundation.

References

- Chambon P: **A decade of molecular biology of retinoic acid receptors.** *Faseb J* 1996, **10**:940-54.
- Wolbach SB, Howe PR: **Tissue changes following deprivation of fat soluble A vitamin.** *J Exp Med* 1925, **42**:753-777.
- Thompson JN, Howell JM, Pitt GAJ: **Vitamin A and reproduction in rats.** *Proc Royal Soc* 1964, **159**:510-535.
- Ghyselinck NB, Bavik C, Sapin V, Mark M, Bonnier D, Hindelang C, Dierich A, Nilsson CB, Hakansson H, Sauvart P, et al: **Cellular retinol-binding protein I is essential for vitamin A homeostasis.** *Embo J* 1999, **18**:4903-14.
- Bok D: **The retinal pigment epithelium: a versatile partner in vision.** *J Cell Sci Suppl* 1993, **17**:189-95.
- Ylikoski J, Pirvola U, Eriksson U: **Cellular retinol-binding protein type I is prominently and differentially expressed in the sensory epithelium of the rat cochlea and vestibular organs.** *J Comp Neurol* 1994, **349**:596-602.
- Kuppumbatti YS, Bleiweiss IJ, Mandeli JP, Waxman S, Mira-y-Lopez R: **Cellular retinol-binding protein expression and breast cancer.** *J Natl Cancer Inst* 2000, **92**:475-80.
- Ruberte E, Dolle P, Chambon P, Morriss-Kay G: **Retinoic acid receptors and cellular retinoid binding proteins. II. Their differential pattern of transcription during early morphogenesis in mouse embryos.** *Development* 1991, **111**:45-60.
- Smith WC, Nakshatri H, Leroy P, Rees J, Chambon P: **A retinoic acid response element is present in the mouse cellular retinol binding protein I (*mCRBP*) promoter.** *Embo J* 1991, **10**:2223-30.
- Eskild W, Oyen O, Beebe S, Jahnsen T, Hansson V: **Regulation of mRNA levels for cellular retinol binding protein in rat Sertoli cells by cyclic AMP and retinol.** *Biochem Biophys Res Commun* 1988, **152**:1504-10.
- Kroepelien CF, Knutsen HK, Haugen TB, Hansson V, Eskild W: **Serum factors induce messenger ribonucleic acid levels for cellular retinol-binding protein in rat Sertoli cells.** *Endocrinology* 1993, **132**:968-74.
- Rush MG, Ul-Haq R, Chytil F: **Opposing effects of retinoic acid and dexamethasone on cellular retinol-binding protein ribonucleic acid levels in the rat.** *Endocrinology* 1991, **129**:705-9.
- Xu G, Bochaton-Piallat ML, Andreutti D, Low RB, Gabbiani G, Neuville P: **Regulation of alpha-smooth muscle actin and CRBP-I expression by retinoic acid and TGF-beta in cultured fibroblasts.** *J Cell Physiol* 2001, **187**:315-25.
- Nilsson MH, Spurr NK, Lundvall J, Rask L, Peterson PA: **Human cellular retinol-binding protein gene organization and chromosomal location.** *Eur J Biochem* 1988, **173**:35-44.
- Eskild W, Simard J, Hansson V, Guerin SL: **Binding of a member of the NF1 family of transcription factors to two distinct cis-acting elements in the promoter and 5'-flanking region of the human cellular retinol binding protein I gene.** *Mol Endocrinol* 1994, **8**:732-45.
- Kuppumbatti YS, Rexer B, Nakajo S, Nakaya K, Mira-y-Lopez R: **CRBP suppresses breast cancer cell survival and anchorage-independent growth.** *Oncogene* 2001, **20**:7413-9.
- Esteller M, Guo M, Moreno V, Peinado MA, Capella G, Galm O, Baylin SB, Herman JG: **Hypermethylation-associated inactivation of the Cellular Retinol-Binding-Protein I Gene in Human Cancer.** *Cancer Res* 2002, **62**:5902-5.
- Fisher GJ, Reddy AP, Datta SC, Kang S, Yi JY, Chambon P, Voorhees JJ: **All-trans retinoic acid induces cellular retinol-binding protein in human skin in vivo.** *J Invest Dermatol* 1995, **105**:80-6.
- Minucci S, Horn V, Bhattacharyya N, Russanova V, Ogryzko VV, Gabriele L, Howard BH, Ozato K: **A histone deacetylase inhibitor potentiates retinoid receptor action in embryonal carcinoma cells.** *Proc Natl Acad Sci USA* 1997, **94**:11295-300.
- Mira-y-Lopez R, Zheng WL, Kuppumbatti YS, Rexer B, Jing Y, Ong DE: **Retinol conversion to retinoic acid is impaired in breast cancer cell lines relative to normal cells.** *J Cell Physiol* 2000, **185**:302-9.
- Farias EF, Arapshian A, Bleiweiss IJ, Waxman S, Zelent A, Mira-y-Lopez R: **Retinoic acid receptor alpha2 is a growth suppressor epigenetically silenced in MCF-7 human breast cancer cells.** *Cell Growth Differ* 2002, **13**:335-41.
- Herman JG, Graff JR, Myohanen S, Nelkin BD, Baylin SB: **Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands.** *Proc Natl Acad Sci USA* 1996, **93**:9821-6.
- Singer J, Roberts-Ems J, Riggs AD: **Methylation of mouse liver DNA studied by means of the restriction enzymes msp I and hpa II.** *Science* 1979, **203**:1019-21.
- Morrison BVV, Leder P: **neu and ras initiate murine mammary tumors that share genetic markers generally absent in c-myc and int-2-initiated tumors.** *Oncogene* 1994, **9**:3417-26.
- Morrison AJ, Herrera RE, Heinsohn EC, Schiff R, Osborne CK: **Dominant Negative N-CoR Relieves Transcriptional Inhibition of Retinoic Acid Receptor but Does Not Alter the Agonist/Antagonist Activities of the Tamoxifen-bound Estrogen Receptor.** *Mol Endocrinol* 2003.

26. Nan X, Campoy FJ, Bird A: **MeCP2 is a transcriptional repressor with abundant binding sites in genomic chromatin.** *Cell* 1997, **88**:471-81.
27. Wang J, Hannon GJ, Beach DH: **Risky immortalization by telomerase.** *Nature* 2000, **405**:755-6.
28. Bartek J, Bartkova J, Kyprianou N, Lalani EN, Staskova Z, Shearer M, Chang S, Taylor-Papadimitriou J: **Efficient immortalization of luminal epithelial cells from human mammary gland by introduction of simian virus 40 large tumor antigen with a recombinant retrovirus.** *Proc Natl Acad Sci U S A* 1991, **88**:3520-4.
29. Yeh E, Cunningham M, Arnold H, Chasse D, Monteith T, Ivaldi G, Hahn WC, Stukenberg PT, Shenolikar S, Uchida T, et al.: **A signalling pathway controlling c-Myc degradation that impacts oncogenic transformation of human cells.** *Nat Cell Biol* 2004, **6**:308-18.
30. Baylin S, Bestor TH: **Altered methylation patterns in cancer cell genomes: cause or consequence?** *Cancer Cell* 2002, **1**:299-305.
31. Li Y, Welm B, Podsypanina K, Huang S, Chamorro M, Zhang X, Rowlands T, Egeblad M, Cowin P, Werb Z, et al.: **Evidence that transgenes encoding components of the Wnt signaling pathway preferentially induce mammary cancers from progenitor cells.** *Proc Natl Acad Sci U S A* 2003, **100**:15853-8.
32. Yang X, Yan L, Davidson NE: **DNA methylation in breast cancer.** *Endocr Relat Cancer* 2001, **8**:115-27.
33. Deming SL, Nass SJ, Dickson RB, Trock BJ: **C-myc amplification in breast cancer: a meta-analysis of its occurrence and prognostic relevance.** *Br J Cancer* 2000, **83**:1688-95.
34. Guo X, Ruiz A, Rando RR, Bok D, Gudas LJ: **Esterification of all-trans-retinol in normal human epithelial cell strains and carcinoma lines from oral cavity, skin and breast: reduced expression of lecithin :retinol acyltransferase in carcinoma lines.** *Carcinogenesis* 2000, **21**:1925-33.
35. Rexer BN, Zheng WL, Ong DE: **Retinoic acid biosynthesis by normal human breast epithelium is via aldehyde dehydrogenase 6, absent in MCF-7 cells.** *Cancer Res* 2001, **61**:7065-70.
36. Triano EA, Slusher LB, Atkins TA, Beneski JT, Gestl SA, Zolfaghari R, Polavarapu R, Fraumeni E, Weisz J: **Class I Alcohol Dehydrogenase Is Highly Expressed in Normal Human Mammary Epithelium but not in Invasive Breast Cancer: Implications for Breast Carcinogenesis.** *Cancer Res* 2003, **63**:3092-100.
37. Chen LI, Sommer KM, Swisshelm K: **Downstream codons in the retinoic acid receptor beta -2 and beta -4 mRNAs initiate translation of a protein isoform that disrupts retinoid-activated transcription.** *J Biol Chem* 2002, **277**:35411-21.
38. Arapshian A, Kuppumbatti YS, Mira-y-Lopez R: **Methylation of conserved CpG sites neighboring the beta retinoic acid response element may mediate retinoic acid receptor beta gene silencing in MCF-7 breast cancer cells.** *Oncogene* 2000, **19**:4066-70.
39. Frommer M, McDonald LE, Millar DS, Collis CM, Watt F, Grigg GW, Molloy PL, Paul CL: **A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual DNA strands.** *Proc Natl Acad Sci USA* 1992, **89**:1827-31.
40. Clark SJ, Harrison J, Paul CL, Frommer M: **High sensitivity mapping of methylated cytosines.** *Nucleic Acids Res* 1994, **22**:2990-7.
41. Bradford MM: **A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.** *Anal Biochem* 1976, **72**:248-54.
42. Busch C, Sakena P, Funa K, Nordlinder H, Eriksson U: **Tissue distribution of cellular retinol-binding protein and cellular retinoic acid-binding protein: use of monospecific antibodies for immunohistochemistry and cRNA for in situ localization of mRNA.** *Methods Enzymol* 1990, **189**:315-24.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

