

Preferences for treatment of lobectomy in Chinese lung cancer patients: video-assisted thoracoscopic surgery or open thoracotomy?

Bo Yang¹
Fang Zhao¹
Zhenfeng Zong¹
Jun Yuan¹
Xiang Song¹
Mingming Ren¹
Qingjun Meng¹
Guoguang Dai¹
Fanyi Kong¹
Shumin Xie²
Siyang Cheng²
Tianwen Gao²

¹Hebei Cangzhou Central Hospital, Department of Thoracic Surgery, Hebei, ²Xiangya Medical School of Central-South University, Changsha, People's Republic of China

Background: This study was designed to investigate the preferences for treatment of lobectomy in Chinese lung cancer patients and differences in the psychological and social factors that influence treatment decision-making.

Methods: One hundred and forty patients with stage I lung cancer were recruited from Hebei Cangzhou Central Hospital. Before surgery, the patients completed a questionnaire that surveyed their preferences for treatment and the relevant influencing factors. Differences in psychological and social characteristics were compared between lung cancer patients who chose video-assisted thoracoscopic surgery (VATS) and those who opted for open thoracotomy.

Results: Among the 135 valid questionnaires, 79 patients preferred VATS and 56 patients chose open thoracotomy. Potential side effects, doctors' recommendation, the prognosticated chance for cure, cosmesis, and financial burden influenced the patients' decisions.

Conclusion: The minimally invasive advantages of VATS, including lesser trauma to the chest wall, earlier remission of postoperative pain, faster recovery, less bleeding, and improved cardiopulmonary function made VATS more attractive to patients needing lobectomy for lung cancer. However, the choice of VATS over open thoracotomy is still influenced by the degree of prognosticated cure and the feasibility of surgery.

Keywords: lobectomy, treatment decision-making, Chinese, oncologists

Introduction

Lung cancer has been the leading cause of cancer death since the mid 20th century, with increasing morbidity and mortality around the world.¹ Since the 1930s, surgery has been the preferred treatment for lung cancer, in the form of conventional open thoracotomy. With the advent of minimally invasive surgery, video-assisted thoracoscopic surgery (VATS) has been widely adopted as an alternative surgical approach for thoracic disease.² VATS is a newly developed minimally invasive thoracic procedure that allows complicated operations to be done through a minimal incision and assisted by a video camera. In the past 2 decades, evidence has shown that, compared with open thoracotomy, VATS has several advantages, including earlier remission of postoperative pain, less release of inflammatory factors, less negative effect on pulmonary function, fewer complications, and faster recovery, without compromising the long-term survival and disease-free survival.³⁻⁶ However, differences still exist with regard to preference for and acceptance of these two surgical approaches among patients. Plenty of studies have focused on the efficacy of the two surgical treatments but few studies have investigated the preferences for treatment decision-making and the factors that influence the patient's treatment decision. This study focuses on the preferences

Correspondence: Bo Yang
Hebei Cangzhou Central Hospital,
Department of Thoracic Surgery,
Hebei 061000, People's Republic of China
Tel +86 031 7207 5624
Fax +86 031 7207 5624
Email yangbtmu@163.com

for treatment of lobectomy in Chinese lung cancer patients according to differences in performance status, psychological factors, and social factors.

Materials and methods

Participants

One hundred and forty lung cancer patients awaiting lobectomy at Hebei Cangzhou Central Hospital from February 2012 to April 2013 were enrolled. The patients were assessed to be suitable for VATS by preoperative tests. Patients had to be diagnosed with stage I lung cancer and be able to consent to surgical treatment. Patients who were in hospital for less than 5 days, who received neoadjuvant chemotherapy, and who had pleural effusion were excluded. Before the survey, all patients were informed about VATS and open thoracotomy procedures, and were asked to sign an informed consent form for surgery. A total of 140 questionnaires were distributed, with five excluded due to more than three missing values. The statistical analysis included the 135 patients who completed the questionnaires fully.

All patients were explicitly informed of their diagnosis, the possibility of stage I lung cancer seen on computed tomography, emission computed tomography, or magnetic resonance imaging, the need for surgery, and the advantages and disadvantages of VATS and open thoracotomy. Sufficient time was allowed for patients to consider these two different surgical approaches, and the preferred surgical approach was put down on the informed consent form for surgery. The patients were divided into a VATS group and an open thoracotomy group based on their preferred surgical approach. Patient demographics, psychological factors, and social factors contributing to the patients' decisions were analyzed.

Demographic characteristics included age, sex, marital status, education level, performance status score, family income, and employment status. Based on an earlier study that examined treatment decisions in cancer patients⁷ and our own clinical experience, seven factors relevant to treatment decision-making were assessed: the best chance for cure, doctor's recommendation of the best treatment, side effects, invasiveness, knowing people who were satisfied after having undergone open thoracotomy or VATS, pain, and affordability. All patients self-administered the questionnaire, and the top three factors were chosen and analyzed (Figure S1).

Statistical analysis

All data were analyzed using Stata version 10.0 software. Chi-square and Wilcoxon rank sum tests were conducted

to analyze the difference in treatment decision-making according to demographic characteristics. $P < 0.05$ was considered to be statistically significant with a two-sided test.

Results

Sixty-six (48.9%) of the 135 patients were male and 69 (51.1%) were female, with an average age of 56 ± 12.9 years. One hundred and twenty-seven (94.1%) were married, 98 (72.6%) were employed, and a high number (58.5%) had completed college or higher education. Family income was $< 3,000$ RMB per month, 3,000–6,000 RMB per month, and $> 6,000$ RMB per month for 49 (36.3%), 41 (30.4%), and 45 (33.3%) of patients, respectively.

Comparison of demographics and characteristics between two groups

The 135 valid questionnaires indicated that 79 patients preferred VATS and 56 patients chose open thoracotomy as their treatment. There was a statistically significant effect of sex and age on treatment decision-making ($P < 0.05$). Fewer males ($n=30$, 38%) than females ($n=49$, 62%) chose VAT, and both younger patients (aged < 35 years) and older patients (> 65 years) preferred VATS, while middle-aged patients (aged 35–65 years) preferred open thoracotomy. Family income played a large role in the decision regarding treatment, given that patients in the People's Republic of China have to pay for their surgery, and are then reimbursed a portion of the medical expenses by an insurance company. Most patients ($n=33$, 59%) with a low family income ($< 3,000$ RMB per month) chose open thoracotomy, while those with a high family income were more likely to choose VATS (Table 1).

Factors affecting treatment decision-making

Of the factors deemed to influence patient decisions, having the fewest side effects ($n=80$, 59%) was rated as the most important reason for choosing treatment, followed by doctor's recommendation ($n=77$, 57%) and best chance for the treatment being curative ($n=67$, 50%). Each treatment preference group had different reasons/factors influencing their decision. Best chance of cure ($n=49$, 88%) and affordability ($n=41$, 73%) were the two most important factors influencing the patient's decision in the open thoracotomy group, while fewest side effects ($n=60$, 76%), doctor's recommendation ($n=54$, 68%), and less invasiveness were regarded as important in the VATS group ($n=47$, 59%; Table 2). Interestingly, predicted patient satisfaction and pain were the two factors that were rated similarly between the two groups.

Table 1 Comparison of demographic and clinical characteristics between two groups

	Treatment decision-making		P-value
	Open thoracotomy	VATS	
Sex	56	79	0.0026 ^a
Male	36 (64%)	30 (38%)	
Female	20 (36%)	4 (62%)	
Age, years	56	79	0.0164 ^b
<35	5 (9%)	44 (56%)	^c
35–65	45 (80%)	10 (13%)	
>65	6 (11%)	25 (32%)	^c
Marital status	56	79	0.3821 ^a
Married	51 (91%)	76 (96%)	
Not married	5 (9%)	3 (4%)	
Education level	56	79	0.3260 ^a
High school graduate or less	26 (46%)	30 (38%)	
College graduate or higher	30 (54%)	49 (62%)	
PS score (0–4)	56	79	0.5528 ^b
0	11 (20%)	13 (16%)	
1	21 (38%)	27 (34%)	
2	10 (18%)	19 (24%)	
3	9 (16%)	12 (15%)	
4	5 (9%)	8 (10%)	
Family income (RMB per month)	56	79	<0.001 ^{b,d}
<3,000	33 (59%)	16 (20%)	
3,000–6,000	17 (30%)	24 (30%)	
>6,000	6 (11%)	39 (49%)	
Employment status	56	79	0.0092 ^a
Unemployment	22 (39%)	15 (19%)	
Employment	34 (61%)	64 (81%)	

Notes: ^aChi-square test; ^bWilcoxon rank sum test; ^cstatistical difference existed in age between the two groups; ^dstatistical difference existed in family income between the two groups.

Abbreviations: PS, performance status; VATS, video-assisted thoracoscopic surgery.

Discussion

A number of studies have analyzed differences in the application and clinical efficacy of VATS and open thoracotomy for lobectomy.^{8–10} However, few studies have looked at factors that influence the patient's decision to choose one or the other procedure.

With the development of minimally invasive surgery, more attention has been paid to the clinical application of VATS. VATS emerged as a new technique for thoracic surgery in the 1990s, and since then has been shown to be useful in lobectomy for lung cancer.^{5,6,8–10} Previous studies have shown significant advantages of VATS over open thoracotomy, including less trauma to the chest wall, faster remission of postoperative pain and recovery, less bleeding, less compromise of cardiopulmonary function, and fewer postoperative complications. As far as we know, patients are less familiar with VATS, and depending on age, sex, and cultural background, they have different attitudes toward VATS versus open thoracotomy.

VATS has been shown to have an important role in the treatment of lung cancer,^{11,12} and has the potential to become the main type of surgery performed for lung cancer in the future. Patients' preference for each type of surgery is impacted by various factors. In this study, we discovered a significant difference in age, sex, and income between patients who choose VATS versus open thoracotomy. Middle-aged patients (n=45) were more inclined toward open thoracotomy ($P<0.05$), while younger patients (n=44) and older patients (n=25) were more likely to opt for VATS. Ninety percent of young patients, 18% of middle-aged patients, and 81% of older patients preferred VATS. These findings might be explained by the fact that middle-aged patients tend to hold traditional views. Further, young patients may attach much importance to decreased invasiveness resulting in improved cosmesis while older patients may value decreased invasiveness resulting in improved recovery, given that older patients are poorer surgical candidates to start with. Further, it was found that female patients (n=49, 62%) chose VATS while male patients (n=30, 38%) preferred open thoracotomy ($P<0.05$). It is possible that female patients choose VATS because of improved cosmesis. Finally, the financial burden of VATS influences the patient's decision; patients who chose VATS had a higher family income than

Table 2 Factors influencing treatment decision-making

Reasons for treatment decision-making	Open thoracotomy		VATS		P-value	Total	
	n	%	n	%		n	%
Best chance for cure	49	88	18	23	<0.001	67	50
Doctors' recommendation	23	41	54	68	<0.001	77	57
Fewest side effects	20	36	60	76	<0.001	80	59
Less invasive	5	9	47	59	<0.001	52	39
Knowing people satisfied with it	14	25	22	28	0.712	36	27
Least painful	16	29	26	46	0.175	55	31
Most affordable	41	73	10	13	<0.001	61	38

Abbreviation: VATS, video-assisted thoracoscopic surgery.

those who chose open thoracotomy, with a family income of less than 3,000 RMB per month in a majority of the open thoracotomy group (n=33, 59%).

In this study, several factors were demonstrated to influence a patient's treatment choice, including side effects (59%), doctor's recommendation (57%), prognosticated chance for cure (50%), less invasiveness (39%), and financial burden (38%). The choice of open thoracotomy was significantly influenced by the potential for cure and a decreased risk of recurrence of cancer, given that the open approach allows for complete lymph node dissection and tumor resection. Another significant reason why lower income patients choose open thoracotomy was its improved affordability over VATS. In the VATS group, fewer side effects, doctor's recommendation, and less invasiveness influenced patient choice.

Previous studies have shown that VATS is at least as safe as open thoracotomy.^{13,14} However, in our study, we found that some patients were concerned about the safety of VATS. In order to shift this misconception, further research is needed to confirm that VATS is a safe surgical approach, and this information needs to be disseminated to educate patients. Finally, because of its superior cosmesis, VATS is preferred by young patients and females. Given that VATS meets patients' requirements regarding cosmesis and safety, it has the potential to become the main surgical approach in the future.¹⁵

The limitations of this study are as follows. In the early phase of the study, during which the patients were learning about VATS versus thoracotomy, different patients received information from different doctors who may have had a bias toward one or the other procedure, which in turn would influence patients' decision-making. Another limitation is the fact that the subjects were all recruited from the Hebei Cangzhou Central Hospital, and the regional culture could have affected personal preferences, psychology, and intellectual capacity.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Liu J, Song ZQ. A review on lung cancer. *Journal of Guangdong Medical College*. 2011;5:536–539.
2. Hai Q. Minimally invasive treatment of lung cancer. *Chin Oncol*. 2005;3:213–216.
3. Lian Y. Effect comparison of thoracoscopic lung resection surgery with conventional thoracotomy. *Chinese Journal of Medical Guide*. 2014;16:56–57.
4. Hu XC. Comparative study on complete video-assisted thoracic surgery and the racotomy lobectomy in the treatment of non-small cell lung cancer. *China Modern Medicine*. 2013;20:67–68.
5. McKenna RJ Jr, Houck W, Fuller CB. Video-assisted thoracic surgery lobectomy: experience with 1,100 cases. *Ann Thorac Surg*. 2006;81:421–425.
6. Zhu R, Weng GX. Comparison of surgical trauma between video-assisted thoracoscopic surgery and conventional thoracotomy. *Chin J Clin Thorac Cardiovasc Surg*. 2009;5:357–361.
7. Palmer NR, Tooze JA, Turner AR, Xu J, Avis NE. African American prostate cancer survivors' treatment decision-making and quality of life. *Patient Educ Couns*. 2013;90:61–68.
8. Chang JH, You QJ, Weng Y, Cai M, Chang Q. Mini-incision video-assisted thoracoscopic versus conventional surgery for lung cancer. *Chin J Min Inv Surg*. 2007;7:412–414.
9. Jiang G, Yang F, Li X, et al. Video-assisted thoracoscopic surgery is more favorable than thoracotomy for administration of adjuvant chemotherapy after lobectomy for non-small cell lung cancer. *World J Surg Oncol*. 2011;9:170.
10. Cao C, Manganas C, Ang SC, Yan TD. A meta-analysis of unmatched and matched patients comparing video-assisted thoracoscopic lobectomy and conventional open lobectomy. *Ann Cardiothorac Surg*. 2012;1:16–23.
11. Xu L, Zhang Z. Video-assisted thoracoscopic surgery for lung cancer. *Chin J Surg Oncol*. 2013;5:137–140.
12. Liu LX. Open chest operation will gradually “upgrade” to a supplement to the minimally invasive thoracoscopic operation. *Chin J Clin Thorac Cardiovasc Surg*. 2012;19:109–112.
13. He Y, Coonar A, Gelvez-Zapata S, Sastry P, Page A. Evaluation of a robot-assisted video-assisted thoracoscopic surgery programme. *Exp Ther Med*. 2014;7:873–876.
14. Linden D, Linden K, Oparka J. In patients with resectable non-small-cell lung cancer, is video-assisted thoracoscopic segmentectomy a suitable alternative to thoracotomy and segmentectomy in terms of morbidity and equivalence of resection. *Interact Cardiovasc Thorac Surg*. 2014;19:107–110.
15. Zhao XG, Peng CJ, Cong B. The application of thoracoscopic lobectomy in treating lung cancer. *Medicine and Philosophy (Clinical Decision Making Forum Edition)*. 2011;9:14–15.

Supplementary material

Hospital number:

Treatment decision-making: Video-assisted thoracoscopic surgery Open thoracotomy

Sociodemographic characteristics of the sample:

Name:

Sex: Male Female

Age:

Marital status: Married Single

Highest level of education: Less than compulsory University level

PS: 0 1 2 3 4

Salary: <3,000 RMB 3,000–6,000 RMB >6,000 RMB

Employment status: Unemployed Employed

Reasons for treatment decision-making: (choose the three answers with the most consideration)

- Best chance for cure
- Doctor's recommendation
- Fewest side effects
- Less invasive
- Knowing people who were satisfied with it
- Least painful
- Most affordable

Figure S1 Questionnaire used in this study.

Abbreviation: PS, performance status.

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.