



Published in final edited form as:

*Eur Med J Oncol.* 2013 November 1; 1: 106–115.

## Yoga as Treatment for Insomnia Among Cancer Patients and Survivors: A Systematic Review

Karen M. Mustian, PhD

### Abstract

Many cancer patients and survivors, between 15 to 90%, report some form of insomnia or sleep quality impairment during and post-treatment, such as excessive daytime napping, difficulty falling asleep, difficulty staying asleep, and waking up too early. Insomnia and sleep quality impairment are among the most prevalent and distressing problems reported by cancer patients and survivors, and can be severe enough to increase cancer mortality. Despite the ubiquity of insomnia and sleep quality impairment, they are under-diagnosed and under-treated in cancer patients and survivors. When sleep problems are present, providers and patients are often hesitant to prescribe or take pharmaceuticals for sleep problems due to poly pharmacy concerns, and cognitive behavioral therapy for insomnia can be very difficult and impractical for patients to adhere to throughout the cancer experience. Research suggests yoga is a well-tolerated exercise intervention with promising evidence for its efficacy in improving insomnia and sleep quality impairment among survivors. This article provides a systematic review of existing clinical research on the effectiveness of yoga for treating insomnia and sleep quality impairment among cancer patients and survivors.

### Keywords

Yoga; sleep; insomnia; cancer; survivorship; exercise

## INTRODUCTION

Between 15 and 90% of cancer patients and survivors report some form of sleep quality impairment both during and post-treatment, such as excessive daytime napping, difficulty falling asleep, difficulty staying asleep, and waking up too early.<sup>1–10</sup> These sleep quality impairments are also symptoms of insomnia, which is defined by one or more of these symptoms (e.g., difficulty falling asleep or difficulty staying asleep) in severe and persistent forms (e.g., 3 or more days per week for one month or longer).<sup>11</sup> Insomnia and sleep quality impairment are among the most prevalent and distressing problems reported by cancer patients and survivors and can increase the risk of cancer-related fatigue and depression, impair cancer-treatment adherence, physical function and quality of life, and, when severe, increase cancer mortality.<sup>1–10,12,13</sup> Despite the ubiquity of insomnia and sleep quality impairment, they are under-diagnosed and under-treated in cancer patients and survivors.<sup>1–10,12,14</sup>

---

**Conflict of Interest Disclosures:** There are no disclosures to report by the author.

Treatment options for insomnia and sleep quality impairment include: 1) pharmaceuticals, which do not cure insomnia and can lead to toxicities, negative interactions with cancer therapeutics, dependency, and rebound impairment after discontinuation; 2) traditional exercise, which is recommended in treatment guidelines, but not widely implemented in survivorship care plans beyond the use of generalised statements, in which survivors are encouraged to be physically active and exercise; and 3) psychobehavioral interventions.<sup>1–10,13</sup> Yoga is a well-tolerated exercise intervention with promising evidence for its efficacy in improving insomnia and sleep quality impairment among survivors.

## **YOGA: A HOLISTIC MIND-BODY MODE OF EXERCISE**

Yoga is an increasingly popular mind-body practice and is also characterised as a mindfulness mode of exercise.<sup>15–18</sup> There are many different styles and types of yoga. They are based on Eastern traditions from India (e.g. Classical, Advaita Vedanta, Tantra), Tibet (e.g. Tibetan), and China (e.g. Chi Kung, Tai Chi).<sup>15,19,20</sup> The word yoga is derived from its Sanskrit root “yuj,” which literally means “to yoke” or join together. In this case, yoga refers to joining the mind and the body.<sup>15,19,20</sup> The earliest forms of yoga were firmly rooted in physical and mindful (breathing and meditative) practices and led, to what is known today as classical yoga which forms the basis for most of the yoga currently taught today.<sup>19</sup> Hatha yoga, the foundation of all yoga styles and the most popular form, includes both Gentle Hatha and Restorative yoga, and is growing in acceptance for therapeutic use in traditional Western medicine.<sup>13,15–18,21–24</sup> Gentle Hatha yoga focuses on physical aspects and is part of many styles of yoga, including Iyengar, Anusara, and others.<sup>15–18,21</sup> Restorative yoga focuses on full relaxation and is part of the Iyengar style.<sup>25,26</sup> The combination of Gentle Hatha and Restorative yoga may provide an effective approach for improving sleep, because it utilises a holistic sequence of meditative, breathing, and physical alignment exercises requiring both the active and passive engagement of skeletal muscles.<sup>15,16,21,22,25,26</sup> Existing scientific evidence suggests that yoga is effective for improving insomnia and sleep quality impairment in cancer patients and survivors.<sup>13,20,27–36</sup>

## **EXISTING SCIENTIFIC EVIDENCE ON YOGA FOR THE TREATMENT OF INSOMNIA AND SLEEP QUALITY IMPAIRMENT (SEE TABLE 1)**

Research suggests that yoga is helpful in treating depression, anxiety, fatigue and other conditions associated with sleep disorders among healthy individuals and those with cancer.<sup>37–39</sup> Herein, we review the extant literature on yoga and its use in the treatment of sleep problems among cancer patients.

### **Yoga Programme Evaluations**

Four evaluations of community yoga programmes for cancer patients and survivors suggest that yoga may improve insomnia and sleep quality impairment.<sup>40–43</sup> For example, Joseph et al. conducted an early study comparing yoga, support therapy and meditation interventions among cancer survivors undergoing radiation therapy where participants in the yoga group reported improvements in sleep, treatment tolerance, mood, appetite, and quality of life.<sup>40</sup> These yoga programmes were based in cancer centres or community-based yoga studios,

and offered yoga classes specifically for cancer patients receiving treatment and survivors who had completed at least primary treatments. The yoga classes included a wide variety of postures and mindfulness exercises from different styles and types, and they were offered one-two times a week for 60–90 minutes. Participants in two of these programmes also attributed improvements in strength, physical function, and physical fitness to their yoga practice.<sup>42,43</sup> However, these reports utilised convenience samples and did not use rigorous research methods designed to answer specific scientific questions about the effects of yoga on sleep quality impairment. These programmes also did not use standardised yoga interventions that can be accurately and consistently replicated for dissemination and explicitly prescribed for the treatment of insomnia or sleep quality impairment. Finally, all of these studies used only patient-reports of insomnia or sleep quality impairment, some of which have not undergone rigorous validation, and no objective assessments of sleep such as polysomnography or actigraphy.

### Phase I and II Pilot Clinical Trials

Although limitations exist (see Limitations of Existing Scientific Data on Yoga and Sleep section to follow), one phase I and seven phase II studies provide preliminary support for the safety, feasibility and efficacy of yoga for improving insomnia and sleep quality impairment among cancer patients and survivors.<sup>20,25,27–30,34–36</sup> Cohen et al. published the first study investigating yoga and sleep using a validated measure of sleep quality impairment with defined clinical cut offs (i.e., Pittsburg Sleep Quality Index) among survivors post-adjuvant treatment.<sup>20</sup> These studies assessed a range of yoga doses from one-five sessions/week with classes lasting 50–120 minutes using a variety of different styles and types of yoga over 4–26 weeks. The interventions included a variety of postures and mindfulness exercises. The interventions were deemed safe and feasible for cancer patients receiving treatment and survivors. Participants enjoyed the yoga interventions and in five studies reported improvements in insomnia and sleep quality impairment; three studies showed no changes in insomnia or sleep quality impairment.<sup>27,34,35</sup> Bower et al. published the first study testing the efficacy of yoga for treating sleep problems that both blinded participants to the study hypotheses and used a rigorous time and attention control condition.<sup>27</sup> Six of the phase II randomised controlled trials (RCTs) compared yoga to a waitlist control, one to a support therapy control condition and one to a health education control condition.<sup>27,30</sup> The latter two studies suggest that yoga may be more effective for improving insomnia and sleep quality impairment than counselling, health education, time and attention.

### Phase III Randomised Controlled Clinical Trials

Recently, Mustian et al. published the first and only multicentre, phase III, randomised controlled trial examining the effects of yoga on insomnia and sleep quality impairment assessed both via validated patient-report measures and objective actigraphy measures.<sup>13</sup> This clinical trial is the most definitive trial to date, and demonstrates that yoga is effective for improving insomnia and sleep quality impairment when compared to a usual care waitlist control condition.<sup>13</sup> The trial compared a standardised yoga intervention (YOCAS<sup>®</sup>: 4 weeks, two times a week, 75 minutes/session; Gentle Hatha and Restorative Yoga) to a usual care waitlist control condition among 410 cancer survivors from 12 community oncology practices throughout the United States. Participants in the yoga condition

demonstrated significant moderate to large improvements in patient-reported outcomes of insomnia and sleep quality impairment as well as significant improvements on objective actigraphy assessments of sleep outcomes including wake after sleep onset and sleep efficiency. Yoga participants also significantly decreased their sleep medication use by 21%, while control participants increased their sleep medication use by 5%. Adherence to YOCAS<sup>®</sup> was good at 80%, and there were no study-related adverse events. All (100%) participants found the YOCAS<sup>®</sup> programme useful and would recommend it to other cancer survivors experiencing sleep problems.<sup>13</sup> Although positive, results are not generalizable to all types of yoga (e.g., yoga in a heating room, vigorously aerobic yoga), the majority of participants were women, white and well-educated and there were no long term follow ups to determine if the benefits of yoga on sleep lasted beyond the immediate post-intervention period.

## LIMITATIONS OF EXISTING SCIENTIFIC DATA ON YOGA AND SLEEP

While very promising, this body of scientific literature needs to be interpreted with caution due to design limitations. None of the phase I-II studies was a definitive phase III RCT that was planned and powered a priori to test the effects of yoga on insomnia or sleep quality impairment as a primary outcome. Many studies did not use validated patient reports of insomnia or sleep quality impairment or objective assessments of sleep problems. The sample sizes were small, ranging from 20–88. They did not screen for or require a specific level of insomnia or sleep quality impairment as part of participant eligibility. The studies did not blind participants with the exception of the Bower study.<sup>27</sup> Yoga interventions were not standardized and were highly variable in content, type, intensity and duration of yoga, making it impossible to determine the actual dose of yoga needed to improve insomnia or sleep quality impairment. The yoga interventions were not described in great detail, making repeatability and standardized dissemination impossible. While general comments suggested the interventions were safe and that participants enjoyed them, no specific details were provided on the rate of adverse events. Information on participant attendance, compliance and attrition, details of the prescribed yoga dose versus the actual dose achieved (e.g., mode, frequency, intensity, duration), and information on sustainability of improvements in sleep quality impairment stemming from yoga were limited.

The phase III clinical trial addressed many of the limitations of the phase I-II clinical trials. For example, the phase III trial was appropriately a priori designed and powered to test sleep as the primary outcome with a sample of 410 survivors, screened for a pre-defined baseline level of sleep quality impairment, used validated patient report and objective measures of sleep, rigorously standardised the yoga intervention and checked for intervention quality, fidelity and drift. The yoga prescription in the intervention was fully detailed in the publication along with accurate reporting of adverse events, attendance, compliance and attrition as well as the achieved dose of yoga versus the prescribed dose of yoga. To date, we could find no studies that have compared yoga to a gold-standard treatment for insomnia, or sleep quality impairment such as pharmaceuticals or cognitive behavioral therapy for insomnia — a required next step in clinical research if yoga is to be considered as such a treatment. In addition, no studies examined the individual components of yoga (i.e. physical postures, breathing and mindfulness activities) to determine which single component, if any,

is primarily responsible for the positive effects stemming from yoga or any possible biological mechanisms (e.g. circadian, muscular, cardiovascular, pulmonary, neurological, immunological or neuroendocrine). Yoga may improve insomnia or sleep quality impairment, but an important body of knowledge needs to be developed in order to better tailor yoga prescriptions to improve sleep problems, and meet the unique needs of individual cancer patients and survivors. Finally, these studies include primarily Caucasian, well-educated, middle to upper middle class women; they have very little racial, economic, social, cultural, gender or age diversity in the sample populations limiting external validity. Importantly, this limits the ability to determine which patient profile may be best suited for and have the best response to yoga therapy. For example, what about the impact of being male, having a cancer diagnosis other than breast cancer, or being non-white, socially isolated or unemployed.

## CLINICAL IMPLICATIONS

While yoga is increasingly popular throughout the world, and there are many books and DVDs as well as cancer centre and community programmes marketed toward cancer survivors (e.g. “Gentle Yoga for Cancer Patients,” “Yoga for Breast Cancer Patients and Survivors, and “Healing Yoga”), there is little, if any, scientific evidence as to the efficacy of these programmes for improving insomnia or sleep quality impairment among cancer survivors. These yoga programmes are not professionally regulated with respect to instructor qualifications and licensure, or adherence to best practice, standard of care or evidence-based therapeutic guidelines, resulting in significant variability as to what is offered to cancer patients and survivors. For example, some yoga programmes focus on very gentle, low-intensity, meditative practices (e.g. Restorative, Integral, Svaroopa), while others focus on vigorous practices (e.g. Power, Ashtanga), and yet others focus on both (e.g. Hatha, Iyengar, Kundalini).<sup>44</sup> Some programmes modify the yoga environment by using heaters and humidifiers (e.g. Bikram) or props such as straps, blocks, ropes and chairs (e.g. Iyengar).<sup>44</sup> Class structure varies considerably with some classes focusing only on physical postures and no mindfulness exercises, while others only include mindfulness exercises and no physical postures. The small number of studies examining the safety and effectiveness of only limited styles and types of yoga for improving insomnia and sleep quality impairment among survivors coupled with the lack of regulation and wide variability of yoga offerings substantially increases the chance that patients and survivors may spend a sizeable amount of time, energy and money participating in yoga programmes that may not be safe or effective. For example, yoga in a room heated to over 100 degrees may be contraindicated for some survivors, and vigorous yoga may result in excessive muscle soreness and joint pain, increasing insomnia or sleep quality impairment. With this in mind, oncology practitioners can play an important role in helping cancer patients and survivors safely and effectively participate in yoga.

Despite their limitations, these phase I-III studies collectively suggest that: 1) cancer patients and survivors can safely participate in yoga during and after cancer treatments; 2) yoga interventions are feasible in a variety of cancer centers and community-based yoga studios; 3) cancer patients and survivors participating in these yoga programmes enjoy them and find them beneficial; 4) participation in low to moderate intensity yoga that incorporates Gentle

Hatha and Restorative postures, breathing and meditation exercises ranging from one-five sessions/week for 50–120 minutes per session over a period of 4–26 weeks may lead to improvements in insomnia and sleep quality impairment; and 5) participation in standardised yoga programmes designed explicitly for cancer patients and survivors experiencing sleep problems, such as YOCAS<sup>®</sup>, will reduce insomnia and sleep quality impairment they experience.

Clinicians can provide important information to help cancer patients and survivors understand how they can safely begin or continue an exercise programme — in this specific case yoga – during and after treatments.<sup>45</sup> Patients and survivors can benefit from knowing potential contraindications (e.g. orthopedic, cardiopulmonary and oncologic) that might affect their exercise safety and tolerance.<sup>46</sup> Contraindications do not necessarily mean that a cancer patient or survivor cannot participate in yoga at all; in fact, this is rarely the case. In most instances, contraindications simply require specific modifications to the yoga regimen so that the individual can safely and effectively participate and achieve physical and mental health benefits. The American College of Sports Medicine Exercise Guidelines for Cancer Patients and Survivors — the only guidelines currently based on scientific evidence — provide an excellent resource regarding recommendations for screening and evaluation of cancer patients and survivors prior to participation in yoga (See Tables 2 and 3).<sup>46</sup>

In addition, referral resources that can help patients and survivors connect with the most qualified and competent yoga instructors in their community, particularly those who have special training and experience working with cancer patients and survivors, or individuals with other medical conditions. Patients and survivors with interest in yoga may also benefit from understanding that the styles and types of yoga that have been tested and shown safe and effective for improving sleep among cancer patients and survivors include primarily Gentle Hatha or Restorative postures combined with breathing and meditation exercises, and they are of low to moderate intensity. When screening patients and survivors for sleep problems and making clinical recommendations about the use of yoga for managing sleep problems, research suggests yoga is effective for individuals who reported mild to moderate sleep quality impairment as well as clinical insomnia, continue to report sleep problems after trying pharmaceutical treatments, demonstrate greater than 1 hour of wakefulness in the middle of the night, and have very poor sleep efficiency (60% or lower), or some combination of these characteristics. Patients and survivors with these characteristics were shown to derive the greatest benefits from participation in yoga — specifically, improved sleep with reduced medication use.

## SUMMARY AND FUTURE RESEARCH DIRECTIONS

Although a definitive phase III randomized clinical trial has been published and positive results were noted from this study and the other smaller phase I-II studies preceding it, the variability across studies and methodological limitations in the published literature continue to limit the extent to which yoga can be considered effective for treating insomnia or sleep quality impairment among cancer patients and survivors. Further research is needed to determine whether yoga is equivalent or superior to existing gold standard pharmaceutical and cognitive behavioral treatments for insomnia and sleep quality impairment. Studies need

to employ validated gold-standard patient reported outcomes of insomnia and sleep quality impairment along with objective measures of insomnia and sleep quality impairment. Studies also need to conduct long term follow up assessments (e.g., 3, 6, 9, 12 months post intervention) to determine the duration and magnitude of any sleep benefits derived from yoga. In addition, a wider variety of yoga types and intensities need to be examined for their safety and efficacy in treating sleep problems among cancer patients and survivors. Dismantling trials are needed to determine which component of yoga (e.g. postures, breathing or meditation) is primarily responsible for its effectiveness along with mechanistic studies to determine the biopsychosocial pathways through which yoga exerts a positive influence on sleep and other toxicities related to cancer and its treatments such as fatigue, functional decline, cognitive impairment, and deregulated immune function, among others. Trials are needed to elucidate effective ways to increase yoga participation among racially, economically, socially and culturally diverse patients and survivors, as well as older and male cancer patients and survivors. Trials are needed to determine which patients and survivors are most likely to benefit yoga as a treatment for sleep problems. Trials are also needed to compare the efficacy of yoga to other modes of exercise, such as walking and resistance training, for improving sleep. Finally, trials are needed that extend yoga to the cancer patients' and survivors' care partners (e.g. sister, brother, mother, father, child, spouse, or friend) who provide unpaid care and support to the patient at significant expense to their own health throughout the cancer experience, and without whom effective treatment, support and recovery would not be possible for these patients — these trials may show yoga to be a low cost-effective therapeutic intervention for both patient and care partner in dyads — with even greater benefits to the patient or survivor than when interventions are directed solely at the individual.

## Acknowledgments

**Funding/Support:** Funding was provided by NCI U10CA037420, K07CA120025

## References

1. Ancoli-Israel S. Recognition and treatment of sleep disturbances in cancer. *J Clin Oncol.* 2009; 27:5864–5866. [PubMed: 19884528]
2. Savard J, Ivers H, Villa J, Caplette-Gingras A, Morin CM. Natural course of insomnia comorbid with cancer: an 18-month longitudinal study. *J Clin Oncol.* 2011; 29:3580–3586. [PubMed: 21825267]
3. Palesh OG, Roscoe JA, Mustian KM, et al. Prevalence, demographics, and psychological associations of sleep disruption in patients with cancer: University of Rochester Cancer Center-Community Clinical Oncology Program. *J Clin Oncol.* 2010; 28:292–298. [PubMed: 19933917]
4. Berger AM, Farr LA, Kuhn BR, Fischer P, Agrawal S. Values of sleep/wake, activity/rest, circadian rhythms, and fatigue prior to adjuvant breast cancer chemotherapy. *Journal of Pain & Symptom Management.* 2007; 33:398–409. [PubMed: 17397701]
5. Ancoli-Israel S, Liu L, Marler MR, et al. Fatigue, sleep, and circadian rhythms prior to chemotherapy for breast cancer. *Supportive Care in Cancer.* 2006; 14:201–209. [PubMed: 16010529]
6. Savard J, Simard S, Blanchet J, Ivers H, Morin CM. Prevalence, clinical characteristics, and risk factors for insomnia in the context of breast cancer. *Sleep.* 2001; 24:583–590. [PubMed: 11480655]

7. Schultz PN, Klein MJ, Beck ML, Stava C, Sellin RV. Breast cancer: relationship between menopausal symptoms, physiologic health effects of cancer treatment and physical constraints on quality of life in long-term survivors. *J Clin Nurs*. 2005; 14:204–211. [PubMed: 15669929]
8. Berger AM, Grem JL, Visovsky C, Marunda HA, Yurkovich JM. Fatigue and other variables during adjuvant chemotherapy for colon and rectal cancer. *Oncol Nurs Forum*. 2010; 37:E359–E369. [PubMed: 21059569]
9. Savard J, Morin CM. Insomnia in the context of cancer: a review of a neglected problem. *J Clin Oncol*. 2001; 19:895–908. [PubMed: 11157043]
10. Pinto AC, de Azambuja E. Improving quality of life after breast cancer: dealing with symptoms. *Maturitas*. 2011; 70:343–348. [PubMed: 22014722]
11. Palesh O, Peppone L, Innominato PF, et al. Prevalence, putative mechanisms, and current management of sleep problems during chemotherapy for cancer. *Nat Sci Sleep*. 2012; 4:151–162. [PubMed: 23486503]
12. Partinen M, HC. *Epidemiology of sleep disorders*. Philadelphia: Elsevier; 2005.
13. Mustian KM, Sprod LK, Janelsins M, et al. Multicenter, Randomized Controlled Trial of Yoga for Sleep Quality Among Cancer Survivors. *J Clin Oncol*. 2013; 26:3233–3241. [PubMed: 23940231]
14. Palesh O, Peppone L, Innominato P, et al. Prevalence, putative mechanisms, and current management of sleep problems during chemotherapy for cancer. *Nature and Science of Sleep*. 2012; 4:151–162.
15. Bower JE, Woolery A, Sternlieb B, Garett D. Yoga for Cancer Patients and Survivors. *Cancer Control*. 2005; 12:165–171. [PubMed: 16062164]
16. Elkins G, Fisher W, Johnson A. Mind-body therapies in integrative oncology. *Curr Treat Options Oncol*. 2010; 11:128–140. [PubMed: 21116746]
17. Mustian KM, Morrow GR, Carroll JK, Figueroa-Moseley CD, Jean-Pierre P, Williams GC. Integrative nonpharmacologic behavioral interventions for the management of cancer-related fatigue. *The Oncologist*. 2007; 12:52–67. [PubMed: 17573456]
18. Mustian KM, Sprod LK, Palesh OG, et al. Exercise for the management of side effects and quality of life among cancer survivors. *Curr Sports Med Rep*. 2009; 8:325–330. [PubMed: 19904073]
19. Kirk, M.; Boon, B. *Hatha Yoga for greater strength, flexibility, and focus*. Champaign, IL: Human Kinetics; 2003.
20. Cohen L, Warneke C, Fouladi RT, Rodriguez MA, Chaoul-Reich A. Psychological adjustment and sleep quality in a randomized trial of the effects of a Tibetan yoga intervention in patients with lymphoma. *Cancer*. 2004; 100:2253–2260. [PubMed: 15139072]
21. Saper RB, Eisenberg DM, Davis RB, Culpepper L, Phillips RS. Prevalence and patterns of adult yoga use in the United States: results of a national survey.[comment]. *Alternative Therapies in Health & Medicine*. 2004; 10:44–49. [PubMed: 15055093]
22. Smith KB, Pukall CF. An evidence-based review of yoga as a complementary intervention for patients with cancer. *Psycho-oncology*. 2009; 18:465–475. [PubMed: 18821529]
23. Muktibodhananda, S. *Hatha Yoga Pradipika*. Poughkeepsie, NY: Nesma Books India; 2000.
24. Lasater, J. *Relax and Renew Restful Yoga for Stressful Times*. Berkeley, CA: Publishers Group West; 1995.
25. Danhauer SC, Mihalko SL, Russell GB, et al. Restorative yoga for women with breast cancer: findings from a randomized pilot study. *Psycho-oncology*. 2009; 18:360–368. [PubMed: 19242916]
26. Danhauer SC, Tooze JA, Farmer DF, et al. Restorative yoga for women with ovarian or breast cancer: findings from a pilot study. *Journal of the Society for Integrative Oncology*. 2008; 6:47–58. [PubMed: 18544284]
27. Bower JE, Garett D, Sternlieb B, et al. Yoga for persistent fatigue in breast cancer survivors: A randomized controlled trial. *Cancer*. 2011
28. Carson JW, Carson KM, Porter LS, Keefe FJ, Seewaldt VL. Yoga of Awareness program for menopausal symptoms in breast cancer survivors: results from a randomized trial. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer*. 2009; 17:1301–1309. [PubMed: 19214594]



29. Ulger O, Yagl NV. Effects of yoga on the quality of life in cancer patients. *Complement Ther Clin Pract.* 2010; 16:60–63. [PubMed: 20347834]
30. Vadiraja SH, Rao MR, Nagendra RH, et al. Effects of yoga on symptom management in breast cancer patients: A randomized controlled trial. *International journal of yoga.* 2009; 2:73–79. [PubMed: 20842268]
31. Mustian KM, Janelins M, Sprod L, et al. YOCAS<sup>®</sup> Yoga Significantly Improves Circadian Rhythm, Anxiety, Mood and Sleep: A Randomized, Controlled Clinical Trial Among 410 Cancer Survivors. *Supportive Care in Cancer.* 2011; 19:317–318.
32. Mustian KM, Palesh OG, Sprod LK, et al. YOCAS<sup>®</sup> Yoga Significantly Improves Sleep, Fatigue and Quality of Life: A URCC CCOP Randomized, Controlled Clinical Trial Among 410 Cancer Survivors. *Journal of Clinical Oncology.* 2010; 28:639.
33. Mustian KM, Palesh OG, Sprod LK, et al. YOCAS<sup>®</sup> Yoga Improves Insomnia Among 410 Cancer Survivors. *International Journal of Behavioral Medicine.* 2010; 17:209.
34. Chandwani KD, Thornton B, Perkins GH, et al. Yoga improves quality of life and benefit finding in women undergoing radiotherapy for breast cancer. *Journal of the Society for Integrative Oncology.* 2010; 8:43–55. [PubMed: 20388445]
35. Cadmus-Bertram L, Littman AJ, Ulrich CM, et al. Predictors of Adherence to a 26-Week Viniyoga Intervention Among Post-Treatment Breast Cancer Survivors. *J Altern Complement Med.* 2013
36. Dhruva A, Miaskowski C, Abrams D, et al. Yoga breathing for cancer chemotherapy-associated symptoms and quality of life: results of a pilot randomized controlled trial. *J Altern Complement Med.* 2012; 18:473–479. [PubMed: 22525009]
37. Balasubramaniam M, Telles S, Doraiswamy PM. Yoga on our minds: a systematic review of yoga for neuropsychiatric disorders. *Frontiers in psychiatry.* 2012; 3:117. [PubMed: 23355825]
38. Buffart LM, van Uffelen JG, Riphagen II, et al. Physical and psychosocial benefits of yoga in cancer patients and survivors, a systematic review and meta-analysis of randomized controlled trials. *BMC cancer.* 2012; 12:559. [PubMed: 23181734]
39. Cramer H, Lauche R, Langhorst J, Dobos G. Yoga for Depression: A Systematic Review and Meta-Analysis. *Depress Anxiety.* 2013
40. Joseph CD. Psychological supportive therapy for cancer patients. *Indian journal of cancer.* 1983; 20:268–270. [PubMed: 6392068]
41. Rosenbaum E, Gautier H, Fobair P, et al. Cancer supportive care, improving the quality of life for cancer patients. A program evaluation report. *Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer.* 2004; 12:293–301. [PubMed: 14991388]
42. Duncan MD, Leis A, Taylor-Brown JW. Impact and outcomes of an Iyengar yoga program in a cancer centre. *Curr Oncol.* 2008; 15(Suppl 2):s109, es72–es78. [PubMed: 18769575]
43. Speed-Andrews AE, Stevinson C, Belanger LJ, Mirus JJ, Courneya KS. Pilot evaluation of Iyengar yoga program for breast cancer survivors. *Cancer Nursing.* 2010; 33:369–381. [PubMed: 20467310]
44. Ross A, Thomas S. The health benefits of yoga and exercise: a review of comparison studies. *J Altern Complement Med.* 2010; 16:3–12. [PubMed: 20105062]
45. Mustian KM, Sprod LK, Janelins M, Peppone LJ, Mohile S. Exercise Recommendations for Cancer-Related Fatigue, Cognitive Impairment, Sleep problems, Depression, Pain, Anxiety, and Physical Dysfunction: A Review. *Oncology & hematology review.* 2012; 8:81–88. [PubMed: 23667857]
46. Schmitz KH, Courneya KS, Matthews C, et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc.* 2010; 42:1409–1426. [PubMed: 20559064]

**Table 1**  
Published phase I-III clinical research trials investigating the efficacy of yoga for treating insomnia and sleep quality impairment

Phase I	Trial Design	Sample	Treatment Status	Type of Yoga	Dose Information	Outcomes
Ulger et al., 2010	1-Arm	Breast Cancer (N=20)	Post-adjuvant treatment	Classical	Frequency: 2x s/week Duration: 60 min/session 4 weeks Intensity: Low to Moderate	Sleep * ↑ Measure(s): Turkish Version of Nottingham Health Profile
<b>Phase II</b>						<b>Outcomes</b>
Cohen et al., 2004	2-Arm RCT w/ SC Waitlist	Lymphoma (N=39)	Mixed receiving active treatment and within 12 months post-adjuvant treatment	Tibetan	Frequency: 1x/week Duration: NA min/session 7 weeks Intensity: Low	Sleep * ↑ Sleep Meds * ↑ Measure(s): Pittsburg Sleep Quality Index
Danhauer et al., 2009	2-Arm RCT w/ SC Waitlist	Breast Cancer (N = 44)	Mixed receiving active treatment and within 2 to 24 months post-adjuvant treatment	Integral (Restorative Yoga Postures)	Frequency: 1x/week Duration: 75 min/session 10 weeks Intensity: Low	Sleep * ↑ Measure(s): Pittsburg Sleep Quality Index
Carson et al., 2009	2-Arm RCT w/ SC Waitlist	Breast Cancer (N=37)	Post-adjuvant treatment	Yoga of Awareness	Frequency: 1x/week Duration: 120 min/session 8 weeks Intensity: Low to Moderate	Sleep * ↑ Measure(s): Daily Diary Sleep Rating
Vidraja et al., 2009	2-Arm RCT w/ SC + Support Therapy	Breast Cancer (N=88)	During adjuvant radiotherapy	Integrated	Frequency: 3x s/week Duration: 50 min/session 6 weeks Intensity: Low to Moderate	Sleep * ↑ Measure(s): European Organization for Research in the Treatment of Cancer QOL C30

Phase I	Trial Design	Sample	Treatment Status	Type of Yoga	Dose Information	Outcomes
Chandwani et al., 2010	2-Arm RCT w/ SC Waitlist	Breast Cancer (N=61)	During adjuvant radiotherapy	Patanjali from VYASA	Frequency: 2x s/week Duration: 60 min/session 6 weeks Intensity: Low to Moderate	Sleep NC Measure(s): Pittsburgh Sleep Quality Index
Bower et al., 2011	2-Arm RCT w/ SC + Health Education	Breast Cancer (N=31)	Survivors post-adjuvant treatment	Iyengar	Frequency: 2x s/week Duration: 90 min/session 12 weeks Intensity: Low to Moderate	Sleep NC Measure(s): Pittsburgh Sleep Quality Index
Dhruva et al., 2012	2-Arm RCT w/ SC Waitlist	Mixed types of Cancer (N=23)	Receiving intravenous chemotherapy	Pranayama	Frequency: 1x/week Duration: 60 min/session 12 months Intensity: Low	Sleep * ↑ Measure(s): General Sleep Disturbance Scale
Cadmus-Bertram et al., 2013	2-Arm BRCT w/ SC Waitlist	Breast Cancer (N=32)	Post-adjuvant treatment	Viniyoga	Frequency: 5x s/week Duration: 75 min/session 6 months Intensity: Low to Moderate	Sleep NC Measure(s): Pittsburgh Sleep Quality Index
<b>Phase III</b>	<b>Trial Design</b>	<b>Sample</b>	<b>Treatment Status</b>	<b>Type of Yoga</b>	<b>Dose Information</b>	<b>Outcomes</b>
Mustian et al., 2010	2-Arm RCT w/ SC Waitlist	Mixed Cancer Survivors (N=410)	Survivors post-adjuvant treatment	YOCAS®	Frequency: 2x s/week Duration: 75 min/session 4-weeks Intensity: Low to Moderate	Sleep * ↑ Sleep Meds * ↑ Measure(s): Pittsburgh Sleep Quality Index & Actigraphy

Notes: An arrow with an asterisk pointing upward indicates a statistically significant improvement in study outcome (i.e., less insomnia or sleep quality impairment and less sleep medication use).

RCT = Randomized, Controlled Trial. BRCT = Block-Randomized, Controlled Trial. NC = No Change, SC = Standard Care, HE = Health Education, NA = Not Available in published article

**Table 2**Exercise Guidelines for Cancer Patients and Survivors Adapted from ACSM<sup>45,46</sup>

<b>MODE OF EXERCISE</b>	<b>RECOMMENDATION</b>
<u>Aerobic Exercise</u>	Achieve a weekly volume of 150 minutes of moderate intensity exercise or 75 minutes of vigorous intensity exercise, or some combination of the two.
<u>Resistance Exercise</u>	Perform strength training exercises 2-3 times per week. Include exercises that target all of the major muscle groups.
<u>Flexibility Exercises</u>	Include stretching exercises for all of the major muscle groups on all the days that other exercises are performed.
<u>Additional Information</u>	Return to normal activity as soon as possible during and following cancer treatment. Some exercise is better than none. Start slowly and progressively increase. Strive to achieve the recommended levels of exercise. See a medical professional if any questions or concerns arise. See an exercise oncology professional for assistance with exercise testing, prescription and monitoring.

**Table 3**Examples of Exercise Contraindications Among Cancer Patients and Survivors Adapted from ACSM<sup>45,46</sup>

Examples of Cancer-Specific Concerns	Examples of Recommendations
Extreme fatigue, anemia, and ataxia	Refer to medical specialist and exercise oncology professional to determine if exercise is safe. If determined to be safe, exercise at a low intensity, as tolerated, preferably under the supervision of an exercise oncology professional.
Surgery	Allow sufficient time to heal after surgery before commencing exercise.
Pain at surgery site	Refer to surgeon and/or physical therapist for clearance prior to exercise. Use exercises that do not involve that area of the body until pain is appropriately managed.
Limited mobility at surgery site	Refer to surgeon and/or physical therapist for clearance prior to exercise. Consider physical activity that does not involve that area of the body.
Risk of hernia due to ostomy	Avoid contact sports and exercises that increase intra-abdominal pressures.
Swelling and lymphedema	Refer to oncologist and physical therapist for clearance prior to exercise. Monitor limb circumference and stop exercise and seek medical evaluation if circumference changes in patients/survivors. Patients at increased risk can wear a compression garment when exercising.
Peripheral neuropathy	Refer to neurologist, physical therapist and exercise oncology professional. Monitor closely for balance impairments. Include exercises that improve balance.
Cardiovascular toxicities	Refer to cardiologist for and exercise oncology professional to determine if exercise is safe.
Compromised immune function	Refer to exercise oncology professional. Prescribe exercise at a low to moderate intensity. Ensure facility is clean to reduce infection risk.
Increased fracture risk	Avoid exercises that put excessive stress on bones, including high impact activities.